

PROF. ANTONIO DI BIAGIO (Orcid ID : 0000-0003-1436-5089)

DR. DANIELE ROBERTO GIACOBBE (Orcid ID : 0000-0003-2385-1759)

DR. MATTEO BASSETTI (Orcid ID : 0000-0002-0145-9740)

Article type : Letter

Corresponding author mail id: antonio.dibiagio@hsanmartino.it

High doses of hydroxychloroquine do not affect viral clearance in patients with SARS-CoV-2 infection.

Lucia Taramasso¹, Antonio Di Biagio^{1,2}, Malgorzata Mikulska^{1,2}, Daniele Roberto Giacobbe¹, Antonio Vena¹, Chiara Dentone¹, Andrea De Maria^{1,2}, Emanuele Delfino¹, Marco Berruti², Chiara Russo², Andrea Orsi^{3,4}, Bianca Bruzzone⁴ and Matteo Bassetti^{1,2}

1 Infectious Diseases Clinic, Policlinico San Martino Hospital-IRCCS, Genoa, Italy

2 Infectious Diseases Clinic, Department of Health Sciences, University of Genoa, Genoa, Italy.

3 Department of Health Sciences, University of Genoa, Genoa, Italy

4 Hygiene Unit, Policlinico San Martino Hospital-IRCCS, Genoa, Italy

Corresponding author

Antonio Di Biagio, MD

Infectious Diseases Clinic,

Department of Health Sciences,

University of Genoa,

Via Antonio Pastore 1,

Genova 16132,

This article has been accepted for publication and undergone full peer review but has not been through the copyediting, typesetting, pagination and proofreading process, which may lead to differences between this version and the [Version of Record](#). Please cite this article as [doi: 10.1111/ECL.13358](#)

This article is protected by copyright. All rights reserved

Introduction

The use of hydroxychloroquine (HCQ) has been considered a therapeutic option by international guidelines and expert opinions during the first phase of COVID-19 pandemics, ¹⁻¹⁰ although scientific evidence remained too scarce to make a definitive recommendation ¹¹. While HCQ use has now been questioned by recent data ⁷, some previous works draw attention to contrasting results on enhanced viral clearance after HCQ treatment ^{1-6,10}. In our hospital, we started using high dose of HCQ at the beginning of the epidemics, associated with tocilizumab and/or methylprednisolone in case of severe pneumonia. In this context, we analysed the data on the rate of nasal swab clearance of SARS-CoV-2, 7 days after HCQ initiation.

Materials and methods

All consecutive patients diagnosed with confirmed SARS-CoV-2 and respiratory symptoms admitted to IRCCS Policlinico San Martino Hospital from 9 March 2020 to 29 March 2020 were enrolled in the present prospective observational study. All enrolled patients gave their consent for off-label HCQ use and for inclusion in the study. Diagnosis of SARS-CoV-2 was made on the basis of a positive reverse-transcriptase–polymerase-chain-reaction (RT-PCR) assay performed on nasal swabs at the time of hospitalization. HCQ was administered at the dose of 400 mg q12 h for 5 to 20 days, based on clinicians' judgment of clinical improvement. Laboratory and clinical data, including time of symptoms onset before HCQ administration were collected. The primary endpoint was viral clearance at day 7(± 2 days), defined as the achievement of the first negative RT-PCR for SARS-CoV-2 at nasal swab. The secondary endpoints were viral clearance at day 10 and median time from symptom onset to viral clearance. We described categorical data as percentages and continuous data as median and range. Nonparametric comparative test for continuous data and χ^2 test for categorical data were used to compare variables between groups.

$P < 0.05$ was considered statistically significant. The study was conducted in accordance with the International Conference on Harmonization guidelines, applicable regulations, and the principles of the Declaration of Helsinki. The study protocol has been approved by the Ligurian Ethical Committee on March 2020.

Results

Out of 140 patients included in the study at the time of writing (19 April 2020), 97 had an available nasal swab seven days after initiation of HCQ treatment. Twenty-four (25%) were female and median age was 64.6 years (range 32.0-89.8). At first clinical presentation, they had median $\text{PaO}_2/\text{FiO}_2$ of 167 (range 57-423) at the time of hospitalization. All were treated with HCQ 400 mg twice daily, 95 received tocilizumab at 8 mg/kg (85 single dose, 12 two doses) and 71 also received methylprednisolone (1 mg/kg). Only 6 patients received azithromycin. After 7 (± 2) days of HCQ treatment, only 22 patients (23%) had a negative nasal swab, while the remaining 75 (77%) still carried SARS-CoV-2. The proportion of 7-day viral clearance was not statistically different in patients who received or not methylprednisolone (20% and 31% respectively, $p = 0.25$). Among 56 patients who had an available nasal swab ten days after initiation of HCQ, 12 were negative (21%) and 44 still positive (79%). Of the six patients treated with azithromycin, 2 had negative nasal swab at day 7. Moreover, at 7-day evaluation, we found that the median time from symptom onset was longer in people who cleared the virus (15 days, range 10-31) than in people who did not (13 days, range 6-21), $p = 0.024$.

Discussion

These findings suggest that the time from symptoms onset, and not that from HCQ initiation, is an important driver of viral clearance. In our study, the exposure to high doses of HCQ did not bring benefits on viral clearance. Stopping the viral shedding could be considered as an important outcome in terms of global health, although not yet clearly related with a clinical outcome in patients with COVID-19. Indeed, it is reasonable to think that patients who clear SARS-CoV-2 stop to be contagious for other people, and all international consensus are based on this assumption to allow the hospital discharge or discontinuation of quarantine¹¹. Then, finding

a drug capable of guaranteeing 7-day viral clearance would be of pivotal importance for stopping the cycle of viral transmission in the general population. However, according to our data, HCQ did not allow to achieve this important outcome in a high proportion of patients, even if given at a higher dosage than previously reported ^{3,5}. The study is limited by the fact that we do not have a control group and thus we cannot exclude that the viral shedding could be even longer without HCQ treatment. Additionally, because of the discomfort associated with collection of consecutive nasopharyngeal swabs, we did not perform a daily collection of specimens; as a consequence, we were not able to estimate the overall median duration of viral shedding. Also, we did not analyze viral load. Moreover, the study was conducted in patients treated with steroids or tocilizumab in association to HCQ and we cannot exclude that the immunosuppression given by these drugs might have slowed down the time to viral clearance. However, most patients hospitalized with severe forms of COVID-19 require immune modulatory treatment and thus data collected in this population likely reflect those of hospitalized patients in real life context. With these limitations, in our study, a relevant proportion of patients with COVID-19 did not achieve rapid viral clearance after 7-day course of high dose of HCQ treatment.

Author Contributions

Lucia Taramasso, Antonio Di Biagio had the idea for and designed the study and had full access to all data. Malgorzata Mikulska, Antonio Vena, Marco Berruti and Chiara Russo take responsibility for the integrity of the data and the accuracy of the data analysis. Lucia Taramasso, Antonio Di Biagio, Malgorzata Mikulska, Daniele Roberto Giacobbe, Antonio Vena, Chiara Dentone, Andrea De Maria, Marco Berruti and Chiara Russo contributed to writing of the report. All authors contributed to data acquisition, data analysis, or data interpretation, and reviewed and approved the final version.

Acknowledgments

We want to acknowledge all the members of **GECOVID-19 Study group**

Conflict of interest

Authors have no conflict of interest to declare.

References

1. Bhimraj A et al. Infectious Diseases Society of America Guidelines on the Treatment and Management of Patients with COVID-19 Infection.
<https://www.idsociety.org/COVID19guidelines>
2. Multicenter collaboration group of Department of Science and Technology of Guangdong Province and Health Commission of Guangdong Province for chloroquine in the treatment of novel coronavirus pneumonia. [Expert consensus on chloroquine phosphate for the treatment of novel coronavirus pneumonia]. *Zhonghua Jie He He Hu Xi Za Zhi*. 2020;43(3):185-188. doi:10.3760/cma.j.issn.1001-0939.2020.03.009
3. Gautret P, Lagier J-C, Parola P, et al. Hydroxychloroquine and azithromycin as a treatment of COVID-19: results of an open-label non-randomized clinical trial. *International Journal of Antimicrobial Agents*. 2020 Mar 20:105949. doi: 10.1016/j.ijantimicag.2020.105949. Online ahead of print.
4. Singh AK, Singh A, Shaikh A, Singh R, Misra A. Chloroquine and hydroxychloroquine in the treatment of COVID-19 with or without diabetes: A systematic search and a narrative review with a special reference to India and other developing countries. *Diabetes & Metabolic Syndrome: Clinical Research & Reviews*. 2020;14(3):241-246. doi:10.1016/j.dsx.2020.03.011
5. Molina JM, Delaugerre C, Goff JL, et al. No Evidence of Rapid Antiviral Clearance or Clinical Benefit with the Combination of Hydroxychloroquine and Azithromycin in Patients with Severe COVID-19 Infection. *Médecine et Maladies Infectieuses*. Published online March 2020:S0399077X20300858. doi:10.1016/j.medmal.2020.03.006
6. Gautret P, Lagier J-C, Parola P, et al. Clinical and microbiological effect of a combination of hydroxychloroquine and azithromycin in 80 COVID-19 patients with at least a six-day follow up: A pilot observational study. *Travel Med Infect Dis*. Published online April 11, 2020:101663. doi:10.1016/j.tmaid.2020.101663
7. Mehra MR, Desai SS, Ruschitzka F, Patel AN. Hydroxychloroquine or chloroquine with or without a macrolide for treatment of COVID-19: a multinational registry analysis. *The Lancet*. Published online May 2020:S0140673620311806. doi:10.1016/S0140-6736(20)31180-6

- Accepted Article
8. The Lancet Editors. Expression of concern: Hydroxychloroquine or chloroquine with or without a macrolide for treatment of COVID-19: a multinational registry analysis. *The Lancet*. Published online June 2020:S0140673620312903. doi:10.1016/S0140-6736(20)31290-3
 9. Geleris J, Sun Y, Platt J, et al. Observational Study of Hydroxychloroquine in Hospitalized Patients with Covid-19. *New England Journal of Medicine* June 18, 2020 382(25):2411-2418. doi:10.1056/NEJMoa2012410
 10. Weehuizen JM, Hoepelman AIM. An Open-Label Cluster-Randomized Controlled Trial of Chloroquine, Hydroxychloroquine or Only Supportive Care in Patients Admitted With Moderate to Severe COVID-19 (ARCHAIC)-Protocol Publication. *European Journal of Clinical Investigation*. Published 31 May 2020. doi.org/10.1111/eci.13297
 11. Alhazzani W, Møller MH, Arabi YM, et al. Surviving Sepsis Campaign: Guidelines on the Management of Critically Ill Adults with Coronavirus Disease 2019 (COVID-19). *Intensive Care Med* 46, 854–887 (2020)
 12. ECDC, European Centre for Disease Prevention and Control. Discharge criteria for confirmed COVID-19 cases. :5. <https://www.ecdc.europa.eu/sites/default/files/documents/COVID-19-Discharge-criteria.pdf>