

# Partial Vandermonde Problems and PASS Encrypt

Katharina Boudgoust<sup>1</sup>   Amin Sakzad<sup>2</sup>   Ron Steinfeld<sup>2</sup>

<sup>1</sup>Aarhus University, Denmark

<sup>2</sup>Faculty of Information Technology, Monash University, Australia


Journées C2, 11th April 2022

# Lattice-Based Cryptography

Provably secure public-key cryptography needs well-defined assumptions in the form of mathematical problems.

(Main) Lattice Problems for Crypto:

- Short Integer Solution [Ajt96]
- NTRU [HPS98]
- Learning With Errors [Reg05]




# Lattice-Based Cryptography

Provably secure public-key cryptography needs well-defined assumptions in the form of mathematical problems.

(Main) Lattice Problems for Crypto:

- Short Integer Solution [Ajt96]
- NTRU [HPS98]
- Learning With Errors [Reg05]
- Partial Vandermonde Problems [HPS<sup>+</sup>14]

↗ today



# Partial Vandermonde Problems

## Partial Vandermonde Transform [HPS<sup>+</sup>14, LZA18]

Let  $R = \mathbb{Z}[x]/\langle x^n + 1 \rangle$ , with  $n$  a power-of-two ( $2n$ -th cyclotomic ring)

## Partial Vandermonde Transform [HPS<sup>+</sup>14, LZA18]

Let  $R = \mathbb{Z}[x]/\langle x^n + 1 \rangle$ , with  $n$  a power-of-two ( $2n$ -th cyclotomic ring)

Choose  $q$  prime such that  $q \equiv 1 \pmod{2n}$ , then:

- it exists  $2n$ -th root of unity  $\omega \in \mathbb{Z}_q$
- $x^n + 1 = \prod_{j \in \mathbb{Z}_{2n}^\times} (x - \omega^j)$

## Partial Vandermonde Transform [HPS<sup>+</sup>14, LZA18]

Let  $R = \mathbb{Z}[x]/\langle x^n + 1 \rangle$ , with  $n$  a power-of-two ( $2n$ -th cyclotomic ring)

Choose  $q$  prime such that  $q \equiv 1 \pmod{2n}$ , then:

- it exists  $2n$ -th root of unity  $\omega \in \mathbb{Z}_q$
- $x^n + 1 = \prod_{j \in \mathbb{Z}_{2n}^\times} (x - \omega^j)$

Write  $\{\omega_j\}_{j=1,\dots,n}$  for  $\{\omega^k : k \in \mathbb{Z}_{2n}^\times\}$ . This defines the **Vandermonde transform**  $\mathbf{V}: R \rightarrow \mathbb{Z}_q^n$

$$\mathbf{V} \cdot a = \begin{bmatrix} 1 & \omega_1 & \cdots & \omega_1^{n-1} \\ 1 & \omega_2 & \cdots & \omega_2^{n-1} \\ 1 & \omega_3 & \cdots & \omega_3^{n-1} \\ \vdots & & & \vdots \\ 1 & \omega_n & \cdots & \omega_n^{n-1} \end{bmatrix} \cdot \begin{bmatrix} a_1 \\ a_2 \\ a_3 \\ \vdots \\ a_n \end{bmatrix} = \begin{bmatrix} b_1 \\ b_2 \\ b_3 \\ \vdots \\ b_n \end{bmatrix} = b \pmod{q}.$$

## Partial Vandermonde Transform [HPS<sup>+</sup>14, LZA18]

Let  $R = \mathbb{Z}[x]/\langle x^n + 1 \rangle$ , with  $n$  a power-of-two ( $2n$ -th cyclotomic ring)

Choose  $q$  prime such that  $q \equiv 1 \pmod{2n}$ , then:

- it exists  $2n$ -th root of unity  $\omega \in \mathbb{Z}_q$
- $x^n + 1 = \prod_{j \in \mathbb{Z}_{2n}^\times} (x - \omega^j)$

Write  $\{\omega_j\}_{j=1,\dots,n}$  for  $\{\omega^k : k \in \mathbb{Z}_{2n}^\times\}$ . This defines the **Vandermonde transform**  $\mathbf{V}: R \rightarrow \mathbb{Z}_q^n$

$$\mathbf{V} \cdot a = \begin{bmatrix} 1 & \omega_1 & \cdots & \omega_1^{n-1} \\ 1 & \omega_2 & \cdots & \omega_2^{n-1} \\ 1 & \omega_3 & \cdots & \omega_3^{n-1} \\ \vdots & & & \vdots \\ 1 & \omega_n & \cdots & \omega_n^{n-1} \end{bmatrix} \cdot \begin{bmatrix} a_1 \\ a_2 \\ a_3 \\ \vdots \\ a_n \end{bmatrix} = \begin{bmatrix} b_1 \\ b_2 \\ b_3 \\ \vdots \\ b_n \end{bmatrix} = b \pmod{q}.$$

**Observation:**  $b$  uniquely defines  $a \pmod{q}$  and vice versa. ( $\mathbf{V}^{-1}$  exists)



# Partial Vandermonde Transform [HPS<sup>+</sup>14, LZA18]

Let  $R = \mathbb{Z}[x]/\langle x^n + 1 \rangle$ , with  $n$  a power-of-two ( $2n$ -th cyclotomic ring)

Choose  $q$  prime such that  $q \equiv 1 \pmod{2n}$ , then:

- it exists  $2n$ -th root of unity  $\omega \in \mathbb{Z}_q$
- $x^n + 1 = \prod_{j \in \mathbb{Z}_{2n}^\times} (x - \omega^j)$

Write  $\{\omega_j\}_{j=1,\dots,n}$  for  $\{\omega^k : k \in \mathbb{Z}_{2n}^\times\}$ . This defines the **Vandermonde transform**  $\mathbf{V}: R \rightarrow \mathbb{Z}_q^n$

$$\mathbf{V} \cdot a = \begin{bmatrix} 1 & \omega_1 & \cdots & \omega_1^{n-1} \\ 1 & \omega_2 & \cdots & \omega_2^{n-1} \\ 1 & \omega_3 & \cdots & \omega_3^{n-1} \\ \vdots & \vdots & \ddots & \vdots \\ 1 & \omega_n & \cdots & \omega_n^{n-1} \end{bmatrix} \cdot \begin{bmatrix} a_1 \\ a_2 \\ a_3 \\ \vdots \\ a_n \end{bmatrix} = \begin{bmatrix} b_1 \\ b_2 \\ b_3 \\ \vdots \\ b_n \end{bmatrix} = b \pmod{q}.$$

**Observation:**  $b$  uniquely defines  $a \pmod{q}$  and vice versa. ( $\mathbf{V}^{-1}$  exists)

**Question:** What happens if we only provide  $t$  out of  $n$  coefficients? (say half)

# Partial Vandermonde Transform [HPS<sup>+</sup>14, LZA18]

Let  $R = \mathbb{Z}[x]/\langle x^n + 1 \rangle$ , with  $n$  a power-of-two ( $2n$ -th cyclotomic ring)

Choose  $q$  prime such that  $q \equiv 1 \pmod{2n}$ , then:

- it exists  $2n$ -th root of unity  $\omega \in \mathbb{Z}_q$
- $x^n + 1 = \prod_{j \in \mathbb{Z}_{2n}^\times} (x - \omega^j)$

Write  $\{\omega_j\}_{j=1,\dots,n}$  for  $\{\omega^k : k \in \mathbb{Z}_{2n}^\times\}$ . This defines the **Vandermonde transform**  $\mathbf{V}: R \rightarrow \mathbb{Z}_q^n$

$$\mathbf{V} \cdot a = \begin{bmatrix} 1 & \omega_1 & \dots & \omega_1^{n-1} \\ 1 & \omega_2 & \dots & \omega_2^{n-1} \\ \vdots & \vdots & \ddots & \vdots \\ 1 & \omega_n & \dots & \omega_n^{n-1} \end{bmatrix} \cdot \begin{bmatrix} a_1 \\ a_2 \\ a_3 \\ \vdots \\ a_n \end{bmatrix} = \begin{bmatrix} b_1 \\ b_2 \\ b_3 \\ \vdots \\ b_n \end{bmatrix} = b \pmod{q}.$$

**Observation:**  $b$  uniquely defines  $a \pmod{q}$  and vice versa. ( $\mathbf{V}^{-1}$  exists)


**Question:** What happens if we only provide  $t$  out of  $n$  coefficients? (say half)

**Note:** For  $\Omega \subseteq \{1, \dots, n\}$  write  $\mathbf{V}_\Omega \cdot a = b$ . (**partial Vandermonde transform**)

# Partial Vandermonde Problems

Choose a random subset  $\Omega \subseteq \{\omega_j\}_{j=1,\dots,n}$  of size  $|\Omega| = t$ .

**Partial Vandermonde knapsack problem (PV-Knap):** Sample  $\mathbf{e} \sim \text{DistrE}$  over  $\mathbb{Z}^n$  defining




The diagram illustrates the equation  $\mathbf{V}_\Omega \mathbf{e} = \mathbf{b} \pmod{q}$ . On the left, the label  $\mathbf{V}_\Omega$  has a dashed arrow pointing to a 10x10 grid of colored squares. The colors in the grid transition from purple at the top to red at the bottom, and from blue on the left to yellow on the right. To the right of the grid is a vertical purple bar labeled  $\mathbf{e}$ . This is followed by an equals sign, a gray bar labeled  $\mathbf{b}$ , and the text  $\pmod{q}$ .

Search: find  $\mathbf{e}$

## Partial Vandermonde Problems

Choose a random subset  $\Omega \subseteq \{\omega_j\}_{j=1,\dots,n}$  of size  $|\Omega| = t$ .

**Partial Vandermonde knapsack problem (PV-Knap):** Sample  $e \sim \text{DistrE}$  over  $\mathbb{Z}^n$  defining




The diagram illustrates the PV-Knap problem. On the left, the label  $V_\Omega$  has a dashed arrow pointing to a grid representing a Vandermonde matrix. The grid has 10 columns and 10 rows, with columns colored in a rainbow gradient from red to purple. A vertical purple bar labeled  $e$  is positioned to the right of the grid. To the right of the bar is an equals sign, followed by a gray vertical bar labeled  $b$ , and then the text  $\text{mod } q$ .

$$V_\Omega e = b \pmod{q}.$$

Search: find  $e$

**Partial Vandermonde Learning With Errors (PV-LWE):** Sample  $s \sim \text{DistrS}$  over  $\mathbb{Z}^t$  and

$e \sim \text{DistrE}$  over  $\mathbb{Z}^n$  defining



The diagram illustrates the PV-LWE problem. On the left, the label  $V_\Omega^T$  has a dashed arrow pointing to a grid representing a Vandermonde matrix. The grid has 10 columns and 10 rows, with columns colored in a rainbow gradient from red to purple. To the right of the grid is a yellow vertical bar labeled  $s$ , followed by a plus sign, a vertical purple bar labeled  $e$ , an equals sign, a gray vertical bar labeled  $b$ , and then the text  $\text{mod } q$ .


$$V_\Omega^T s + e = b \pmod{q}.$$

Search: find  $e$  (and secret  $s$ )

## Partial Vandermonde Problems

Choose a random subset  $\Omega \subseteq \{\omega_j\}_{j=1,\dots,n}$  of size  $|\Omega| = t$ .

**Partial Vandermonde knapsack problem (PV-Knap):** Sample  $\mathbf{e} \sim \text{DistrE}$  over  $\mathbb{Z}^n$  defining




The diagram illustrates the PV-Knap problem. On the left, the label  $\mathbf{V}_\Omega$  has a dashed arrow pointing to a grid representing a submatrix of a Vandermonde matrix. The grid has 10 columns and 10 rows, with columns colored in a rainbow gradient from red to purple. To the right of the grid is a vertical purple bar labeled  $\mathbf{e}$ . Further right is an equals sign, followed by a vertical grey bar labeled  $\mathbf{b}$ , and then the text  $\text{mod } q$ .

$$\mathbf{V}_\Omega \mathbf{e} = \mathbf{b} \pmod{q}.$$

Search: find  $\mathbf{e}$

**Partial Vandermonde Learning With Errors (PV-LWE):** Sample  $\mathbf{s} \sim \text{DistrS}$  over  $\mathbb{Z}^t$  and

$\mathbf{e} \sim \text{DistrE}$  over  $\mathbb{Z}^n$  defining



The diagram illustrates the PV-LWE problem. On the left, the label  $\mathbf{V}_\Omega^T$  has a dashed arrow pointing to a grid representing a submatrix of a Vandermonde matrix. The grid has 10 columns and 10 rows, with columns colored in a rainbow gradient from red to purple. To the right of the grid is a vertical yellow bar labeled  $\mathbf{s}$ . This is followed by a plus sign, then a vertical purple bar labeled  $\mathbf{e}$ , then an equals sign, then a vertical grey bar labeled  $\mathbf{b}$ , and finally the text  $\text{mod } q$ .

$$\mathbf{V}_\Omega^T \mathbf{s} + \mathbf{e} = \mathbf{b} \pmod{q}.$$

Search: find  $\mathbf{e}$  (and secret  $\mathbf{s}$ )

**Conjecture:** Hard to solve if  $\text{DistrE}$  provides elements of small norm.

# Equivalence of PV-Knap and PV-LWE

Let  $t = n/2$  and set  $\mathcal{P}_t = \{\Omega \subseteq \{\omega_j\}_{j=1,\dots,n} : |\Omega| = t\}$ .

**Property 1:**  $\mathbf{V}_\Omega$  defines a ring homomorphism from  $R$  to  $\mathbb{Z}_q^t$ :

$$\mathbf{V}_\Omega(a \cdot b) = (\mathbf{V}_\Omega a) \circ (\mathbf{V}_\Omega b)$$

(component-wise multiplication  $\circ$ )

# Equivalence of PV-Knap and PV-LWE

Let  $t = n/2$  and set  $\mathcal{P}_t = \{\Omega \subseteq \{\omega_j\}_{j=1,\dots,n} : |\Omega| = t\}$ .

**Property 1:**  $\mathbf{V}_\Omega$  defines a ring homomorphism from  $R$  to  $\mathbb{Z}_q^t$ :

$$\mathbf{V}_\Omega(a \cdot b) = (\mathbf{V}_\Omega a) \circ (\mathbf{V}_\Omega b)$$

(component-wise multiplication  $\circ$ )

**Property 2:**  $\Omega^c = \{\omega_j\}_j \setminus \Omega$  defines the **complement** partial Vandermonde transform  $\mathbf{V}_{\Omega^c}$ .

Given  $\mathbf{V}_\Omega a$  and  $\mathbf{V}_{\Omega^c} a$ , we can recover  $a \bmod q$ .

# Equivalence of PV-Knap and PV-LWE

Let  $t = n/2$  and set  $\mathcal{P}_t = \{\Omega \subseteq \{\omega_j\}_{j=1,\dots,n} : |\Omega| = t\}$ .

**Property 1:**  $\mathbf{V}_\Omega$  defines a ring homomorphism from  $R$  to  $\mathbb{Z}_q^t$ :

$$\mathbf{V}_\Omega(a \cdot b) = (\mathbf{V}_\Omega a) \circ (\mathbf{V}_\Omega b)$$


(component-wise multiplication  $\circ$ )

**Property 2:**  $\Omega^c = \{\omega_j\}_j \setminus \Omega$  defines the **complement** partial Vandermonde transform  $\mathbf{V}_{\Omega^c}$ .

Given  $\mathbf{V}_\Omega a$  and  $\mathbf{V}_{\Omega^c} a$ , we can recover  $a \bmod q$ .

**Property 3:** For every  $\Omega \in \mathcal{P}_t$ , there exists a  $\Omega' \in \mathcal{P}_t$  such that

$$\mathbf{V}_{\Omega'} \cdot \mathbf{V}_\Omega^T = 0 \in \mathbb{Z}_q^{t \times t}.$$

(parity check matrix,  only for power-of-two cyclotomics)



# Equivalence of PV-Knap and PV-LWE

Let  $t = n/2$  and set  $\mathcal{P}_t = \{\Omega \subseteq \{\omega_j\}_{j=1,\dots,n} : |\Omega| = t\}$ .

**Property 1:**  $\mathbf{V}_\Omega$  defines a ring homomorphism from  $R$  to  $\mathbb{Z}_q^t$ :

$$\mathbf{V}_\Omega(a \cdot b) = (\mathbf{V}_\Omega a) \circ (\mathbf{V}_\Omega b)$$

(component-wise multiplication  $\circ$ )

**Property 2:**  $\Omega^c = \{\omega_j\}_j \setminus \Omega$  defines the **complement** partial Vandermonde transform  $\mathbf{V}_{\Omega^c}$ .

Given  $\mathbf{V}_\Omega a$  and  $\mathbf{V}_{\Omega^c} a$ , we can recover  $a \bmod q$ .

**Property 3:** For every  $\Omega \in \mathcal{P}_t$ , there exists a  $\Omega' \in \mathcal{P}_t$  such that

$$\mathbf{V}_{\Omega'} \cdot \mathbf{V}_\Omega^T = 0 \in \mathbb{Z}_q^{t \times t}.$$

(parity check matrix,  only for power-of-two cyclotomics)

## Lemma (Adapted [MM11, Sec. 4.2])


Let  $\psi$  denote a distribution over  $\mathbb{Z}^n \cong R$ . There is an efficient reduction from  $\text{PV-LWE}_\psi$  to  $\text{PV-Knap}_\psi$ , and vice versa.

**Idea:** Given  $(\mathbf{V}_\Omega, b)$ , with  $b = \mathbf{V}_\Omega^T s + e$ . Compute  $\Omega'$  such that  $\mathbf{V}_{\Omega'} \cdot \mathbf{V}_\Omega^T = 0$ . Then,  $b' := \mathbf{V}_{\Omega'} b = \mathbf{V}_{\Omega'} e$  is an instance of PV-Knap.

# Hidden Ideal Lattice 1/2

Choose a random subset  $\Omega \subseteq \{\omega_j\}_{j=1,\dots,n}$  of size  $|\Omega| = t$ .

**Partial Vandermonde knapsack problem (PV-Knap):** Sample  $\mathbf{e} \sim \text{DistrE}$  over  $\mathbb{Z}^n$  defining



$$\begin{bmatrix} \text{grid of colored squares} \end{bmatrix} \mathbf{e} = \mathbf{b} \pmod{q}.$$

Search: find  $\mathbf{e}$

# Hidden Ideal Lattice 1/2

Choose a random subset  $\Omega \subseteq \{\omega_j\}_{j=1,\dots,n}$  of size  $|\Omega| = t$ .

**Partial Vandermonde knapsack problem (PV-Knap):** Sample  $\mathbf{e} \sim \text{DistrE}$  over  $\mathbb{Z}^n$  defining


$$\mathbf{V}_\Omega \mathbf{e} = \mathbf{b} \pmod{q}.$$

Search: find  $\mathbf{e}$


The matrix  $\mathbf{V}_\Omega$  defines an **ideal lattice**:

$$\Lambda_q^\perp(\mathbf{V}_\Omega) = \{a \in R : \mathbf{V}_\Omega a = 0 \pmod{q}\}$$

## Hidden Ideal Lattice 1/2

Choose a random subset  $\Omega \subseteq \{\omega_j\}_{j=1,\dots,n}$  of size  $|\Omega| = t$ .

**Partial Vandermonde knapsack problem (PV-Knap):** Sample  $\mathbf{e} \sim \text{DistrE}$  over  $\mathbb{Z}^n$  defining


$$\mathbf{V}_\Omega \mathbf{e} = b \pmod{q}.$$

Search: find  $\mathbf{e}$

The matrix  $\mathbf{V}_\Omega$  defines an **ideal lattice**:

$$\Lambda_q^\perp(\mathbf{V}_\Omega) = \{a \in R : \mathbf{V}_\Omega a = 0 \pmod{q}\}$$


Idea:

- 1) Solve  $\mathbf{V}_\Omega y = b \pmod{q}$  for the unknown  $y$  in  $R$  (in general not in the support of  $\text{DistrE}$ )
- 2) Find a **closest vector**  $v$  of  $y$  in  $\Lambda_q^\perp(\mathbf{V}_\Omega)$ , i.e.,  $\|y - v\|$  smallest
- 3) The element  $e := y - v$  is a solution to PV-Knap

⚠ Promise variant of the closest vector problem, called **Bounded Distance Decoding (BDD)**

## Hidden Ideal Lattice 2/2


**Partial Vandermonde Learning With Errors (PV-LWE):** Sample  $\mathbf{s} \sim \text{DistrS}$  over  $\mathbb{Z}^t$  and  $\mathbf{e} \sim \text{DistrE}$  over  $\mathbb{Z}^n$  defining


$$\text{Matrix} + \mathbf{s} = \mathbf{b} \pmod{q}$$

Search: find  $\mathbf{e}$  (and secret  $\mathbf{s}$ )

## Hidden Ideal Lattice 2/2

**Partial Vandermonde Learning With Errors (PV-LWE):** Sample  $\mathbf{s} \sim \text{DistrS}$  over  $\mathbb{Z}^t$  and  $\mathbf{e} \sim \text{DistrE}$  over  $\mathbb{Z}^n$  defining


$$\mathbf{V}_\Omega \mathbf{s} + \mathbf{e} = \mathbf{b} \pmod{q}.$$

Search: find  $\mathbf{e}$  (and secret  $\mathbf{s}$ )

This is an instance of **BDD** in the **ideal lattice**

$$\Lambda_q(\mathbf{V}_\Omega) = \{a \in R : a = \mathbf{V}_\Omega^T s \pmod{q} \text{ for some } s \in \mathbb{Z}_q^t\}$$

# PASS Encrypt

# PASS Encrypt [HS15]

[HS15]	Our work
deterministic	randomized
without proof of security	with proof of security
fixed $\mathbf{V}_\Omega$	random $\mathbf{V}_\Omega$



# PASS Encrypt [HS15]

[HS15]	Our work
deterministic	randomized
without proof of security	with proof of security
fixed $\mathbf{V}_\Omega$	random $\mathbf{V}_\Omega$

Let  $p \ll q$  be two primes,  $m \in \{0, 1\}^n$ ,  $\psi$  a distribution over  $\mathbb{Z}^n$  and  $t = n/2$ .

**KeyGen**( $1^\lambda$ ): sample  $f \leftarrow \psi$  and  $\Omega \leftarrow \text{Unif}(\mathcal{P}_t)$ ; return  $\text{sk} = f$  and  $\text{pk} = (\Omega, \mathbf{V}_\Omega f)$

**Enc**( $\text{pk}, m$ ): sample  $r, s \leftarrow \psi$ ; set  $r' = pr$  and  $s' = m + ps$

$$e_1 = (\text{pk} \circ \mathbf{V}_\Omega r') + \mathbf{V}_\Omega s'$$

$$e_2 = \mathbf{V}_\Omega^c r'$$

$$e_3 = \mathbf{V}_\Omega^c s'$$

return  $c = (e_1, e_2, e_3)$

**Dec**( $\text{sk}, c$ ): compute  $c' = (\mathbf{V}_\Omega^c \text{sk} \circ e_2) + e_3$  and combine with  $e_1$  to  $c'' \in \mathbb{Z}_q^n$ ;

return  $\mathbf{V}^{-1} c'' \bmod p$ .

# PASS Encrypt [HS15]

[HS15]	Our work
deterministic	randomized
without proof of security	with proof of security
fixed $\mathbf{V}_\Omega$	random $\mathbf{V}_\Omega$

Let  $p \ll q$  be two primes,  $m \in \{0, 1\}^n$ ,  $\psi$  a distribution over  $\mathbb{Z}^n$  and  $t = n/2$ .

**KeyGen**( $1^\lambda$ ): sample  $f \leftarrow \psi$  and  $\Omega \leftarrow \text{Unif}(\mathcal{P}_t)$ ; return  $\text{sk} = f$  and  $\text{pk} = (\Omega, \mathbf{V}_\Omega f)$

**Enc**( $\text{pk}, m$ ): sample  $r, s \leftarrow \psi$ ; set  $r' = pr$  and  $s' = m + ps$

$$e_1 = (\text{pk} \circ \mathbf{V}_\Omega r') + \mathbf{V}_\Omega s'$$

$$e_2 = \mathbf{V}_{\Omega^c} r'$$

$$e_3 = \mathbf{V}_{\Omega^c} s'$$

return  $c = (e_1, e_2, e_3)$

**Dec**( $\text{sk}, c$ ): compute  $c' = (\mathbf{V}_{\Omega^c} \text{sk} \circ e_2) + e_3$  and combine with  $e_1$  to  $c'' \in \mathbb{Z}_q^n$ ;

return  $\mathbf{V}^{-1} c'' \bmod p$ .

# PASS Encrypt [HS15]

[HS15]	Our work
deterministic	randomized
without proof of security	with proof of security
fixed $\mathbf{V}_\Omega$	random $\mathbf{V}_\Omega$

Let  $p \ll q$  be two primes,  $m \in \{0, 1\}^n$ ,  $\psi$  a distribution over  $\mathbb{Z}^n$  and  $t = n/2$ .

**KeyGen**( $1^\lambda$ ): sample  $f \leftarrow \psi$  and  $\Omega \leftarrow \text{Unif}(\mathcal{P}_t)$ ; return  $\text{sk} = f$  and  $\text{pk} = (\Omega, \mathbf{V}_\Omega f)$

**Enc**( $\text{pk}, m$ ): sample  $r, s \leftarrow \psi$ ; set  $r' = pr$  and  $s' = m + ps$

$$e_1 = (\text{pk} \circ \mathbf{V}_\Omega r') + \mathbf{V}_\Omega s'$$

$$e_2 = \mathbf{V}_{\Omega^c} r'$$

$$e_3 = \mathbf{V}_{\Omega^c} s'$$

return  $c = (e_1, e_2, e_3)$

**Dec**( $\text{sk}, c$ ): compute  $c' = (\mathbf{V}_{\Omega^c} \text{sk} \circ e_2) + e_3$  and combine with  $e_1$  to  $c'' \in \mathbb{Z}_q^n$ ;

return  $\mathbf{V}^{-1} c'' \bmod p$ .

# PASS Encrypt [HS15]

[HS15]	Our work
deterministic	randomized
without proof of security	with proof of security
fixed $\mathbf{V}_\Omega$	random $\mathbf{V}_\Omega$

Let  $p \ll q$  be two primes,  $m \in \{0, 1\}^n$ ,  $\psi$  a distribution over  $\mathbb{Z}^n$  and  $t = n/2$ .

**KeyGen**( $1^\lambda$ ): sample  $f \leftarrow \psi$  and  $\Omega \leftarrow \text{Unif}(\mathcal{P}_t)$ ; return  $\text{sk} = f$  and  $\text{pk} = (\Omega, \mathbf{V}_\Omega f)$

**Enc**( $\text{pk}, m$ ): sample  $r, s \leftarrow \psi$ ; set  $r' = pr$  and  $s' = m + ps$

$$e_1 = (\text{pk} \circ \mathbf{V}_\Omega r') + \mathbf{V}_\Omega s'$$

$$e_2 = \mathbf{V}_{\Omega^c} r'$$

$$e_3 = \mathbf{V}_{\Omega^c} s'$$

return  $c = (e_1, e_2, e_3)$

**Dec**( $\text{sk}, c$ ): compute  $c' = (\mathbf{V}_{\Omega^c} \text{sk} \circ e_2) + e_3$  and combine with  $e_1$  to  $c'' \in \mathbb{Z}_q^n$ ;

return  $\mathbf{V}^{-1} c'' \bmod p$ .

Recall:  $\mathbf{V}_\Omega$  and  $\mathbf{V}_{\Omega^c}$  define  $\mathbf{V}$  and  $\mathbf{V}^{-1}$ .

# PASS Encrypt [HS15]

[HS15]	Our work
deterministic	randomized
without proof of security	with proof of security
fixed $\mathbf{V}_\Omega$	random $\mathbf{V}_\Omega$

Let  $p \ll q$  be two primes,  $m \in \{0, 1\}^n$ ,  $\psi$  a distribution over  $\mathbb{Z}^n$  and  $t = n/2$ .

**KeyGen**( $1^\lambda$ ): sample  $f \leftarrow \psi$  and  $\Omega \leftarrow \text{Unif}(\mathcal{P}_t)$ ; return  $\text{sk} = f$  and  $\text{pk} = (\Omega, \mathbf{V}_\Omega f)$

**Enc**( $\text{pk}, m$ ): sample  $r, s \leftarrow \psi$ ; set  $r' = pr$  and  $s' = m + ps$

$$e_1 = (\text{pk} \circ \mathbf{V}_\Omega r') + \mathbf{V}_\Omega s'$$

$$e_2 = \mathbf{V}_{\Omega^c} r'$$

$$e_3 = \mathbf{V}_{\Omega^c} s'$$

return  $c = (e_1, e_2, e_3)$

**Dec**( $\text{sk}, c$ ): compute  $c' = (\mathbf{V}_{\Omega^c} \text{sk} \circ e_2) + e_3$  and combine with  $e_1$  to  $c'' \in \mathbb{Z}_q^n$ ;

return  $\mathbf{V}^{-1} c'' \bmod p$ .

Recall:  $\mathbf{V}_\Omega$  and  $\mathbf{V}_{\Omega^c}$  define  $\mathbf{V}$  and  $\mathbf{V}^{-1}$ .

**Correctness:**

$$\left. \begin{aligned} e_1 &= (\mathbf{V}_\Omega f \circ \mathbf{V}_\Omega r') + \mathbf{V}_\Omega s' = \mathbf{V}_\Omega (f \cdot r' + s') \\ c' &= (\mathbf{V}_{\Omega^c} \text{sk} \circ (\mathbf{V}_{\Omega^c} r')) + \mathbf{V}_{\Omega^c} s' = \mathbf{V}_{\Omega^c} (f \cdot r' + s') \end{aligned} \right\} \begin{array}{l} \text{ring} \\ \text{homomorphism} \end{array}$$

# PASS Encrypt [HS15]

[HS15]	Our work
deterministic	randomized
without proof of security	with proof of security
fixed $\mathbf{V}_\Omega$	random $\mathbf{V}_\Omega$

Let  $p \ll q$  be two primes,  $m \in \{0, 1\}^n$ ,  $\psi$  a distribution over  $\mathbb{Z}^n$  and  $t = n/2$ .

**KeyGen**( $1^\lambda$ ): sample  $f \leftarrow \psi$  and  $\Omega \leftarrow \text{Unif}(\mathcal{P}_t)$ ; return  $\text{sk} = f$  and  $\text{pk} = (\Omega, \mathbf{V}_\Omega f)$

**Enc**( $\text{pk}, m$ ): sample  $r, s \leftarrow \psi$ ; set  $r' = pr$  and  $s' = m + ps$

$$e_1 = (\text{pk} \circ \mathbf{V}_\Omega r') + \mathbf{V}_\Omega s'$$

$$e_2 = \mathbf{V}_{\Omega^c} r'$$

$$e_3 = \mathbf{V}_{\Omega^c} s'$$

return  $c = (e_1, e_2, e_3)$

**Dec**( $\text{sk}, c$ ): compute  $c' = (\mathbf{V}_{\Omega^c} \text{sk} \circ e_2) + e_3$  and combine with  $e_1$  to  $c'' \in \mathbb{Z}_q^n$ ;

return  $\mathbf{V}^{-1} c'' \bmod p$ .

Recall:  $\mathbf{V}_\Omega$  and  $\mathbf{V}_{\Omega^c}$  define  $\mathbf{V}$  and  $\mathbf{V}^{-1}$ .

**Correctness:**

$$\left. \begin{aligned} e_1 &= (\mathbf{V}_\Omega f \circ \mathbf{V}_\Omega r') + \mathbf{V}_\Omega s' = \mathbf{V}_\Omega (f \cdot r' + s') \\ c' &= (\mathbf{V}_{\Omega^c} \text{sk} \circ (\mathbf{V}_{\Omega^c} r')) + \mathbf{V}_{\Omega^c} s' = \mathbf{V}_{\Omega^c} (f \cdot r' + s') \end{aligned} \right\} \begin{array}{l} \text{ring} \\ \text{homomorphism} \end{array}$$

$$\mathbf{V}^{-1}(e_1 || c') = \mathbf{V}^{-1}(\mathbf{V}(f \cdot r' + s')) = f \cdot pr + ps + m = m \bmod p$$

if  $f, r$  and  $s$  are small enough

# PASS Encrypt [HS15]

[HS15]	Our work
deterministic	randomized
without proof of security	with proof of security
fixed $\mathbf{V}_\Omega$	random $\mathbf{V}_\Omega$

Let  $p \ll q$  be two primes,  $m \in \{0, 1\}^n$ ,  $\psi$  a distribution over  $\mathbb{Z}^n$  and  $t = n/2$ .

**KeyGen**( $1^\lambda$ ): sample  $f \leftarrow \psi$  and  $\Omega \leftarrow \text{Unif}(\mathcal{P}_t)$ ; return  $\text{sk} = f$  and  $\text{pk} = (\Omega, \mathbf{V}_\Omega f)$

**Enc**( $\text{pk}, m$ ): sample  $r, s \leftarrow \psi$ ; set  $r' = pr$  and  $s' = m + ps$

$$e_1 = (\text{pk} \circ \mathbf{V}_\Omega r') + \mathbf{V}_\Omega s'$$

$$e_2 = \mathbf{V}_{\Omega^c} r'$$

$$e_3 = \mathbf{V}_{\Omega^c} s'$$

return  $c = (e_1, e_2, e_3)$

**Dec**( $\text{sk}, c$ ): compute  $c' = (\mathbf{V}_{\Omega^c} \text{sk} \circ e_2) + e_3$  and combine with  $e_1$  to  $c'' \in \mathbb{Z}_q^n$ ;

return  $\mathbf{V}^{-1} c'' \bmod p$ .

# PASS Encrypt [HS15]

[HS15]	Our work
deterministic	randomized
without proof of security	with proof of security
fixed $\mathbf{V}_\Omega$	random $\mathbf{V}_\Omega$

Let  $p \ll q$  be two primes,  $m \in \{0, 1\}^n$ ,  $\psi$  a distribution over  $\mathbb{Z}^n$  and  $t = n/2$ .

**KeyGen**( $1^\lambda$ ): sample  $f \leftarrow \psi$  and  $\Omega \leftarrow \text{Unif}(\mathcal{P}_t)$ ; return  $\text{sk} = f$  and  $\text{pk} = (\Omega, \mathbf{V}_\Omega f)$

**Enc**( $\text{pk}, m$ ): sample  $r, s \leftarrow \psi$ ; set  $r' = pr$  and  $s' = m + ps$

$$e_1 = (\text{pk} \circ \mathbf{V}_\Omega r') + \mathbf{V}_\Omega s' = \mathbf{V}_\Omega(f \cdot r' + s')$$

$$e_2 = \mathbf{V}_{\Omega^c} r'$$

$$e_3 = \mathbf{V}_{\Omega^c} s'$$

return  $c = (e_1, e_2, e_3)$

**Dec**( $\text{sk}, c$ ): compute  $c' = (\mathbf{V}_{\Omega^c} \text{sk} \circ e_2) + e_3$  and combine with  $e_1$  to  $c'' \in \mathbb{Z}_q^n$ ;

return  $\mathbf{V}^{-1} c'' \bmod p$ .

## Security:

$e_1 = \mathbf{V}_\Omega(f \cdot r' + s')$  defines an instance of PV-Knap



# PASS Encrypt [HS15]

[HS15]	Our work
deterministic	randomized
without proof of security	with proof of security
fixed $\mathbf{V}_\Omega$	random $\mathbf{V}_\Omega$

Let  $p \ll q$  be two primes,  $m \in \{0, 1\}^n$ ,  $\psi$  a distribution over  $\mathbb{Z}^n$  and  $t = n/2$ .

**KeyGen**( $1^\lambda$ ): sample  $f \leftarrow \psi$  and  $\Omega \leftarrow \text{Unif}(\mathcal{P}_t)$ ; return  $\text{sk} = f$  and  $\text{pk} = (\Omega, \mathbf{V}_\Omega f)$

**Enc**( $\text{pk}, m$ ): sample  $r, s \leftarrow \psi$ ; set  $r' = pr$  and  $s' = m + ps$

$$e_1 = (\text{pk} \circ \mathbf{V}_\Omega r') + \mathbf{V}_\Omega s' = \mathbf{V}_\Omega(f \cdot r' + s')$$

$$e_2 = \mathbf{V}_{\Omega^c} r'$$

$$e_3 = \mathbf{V}_{\Omega^c} s'$$

return  $c = (e_1, e_2, e_3)$

**Dec**( $\text{sk}, c$ ): compute  $c' = (\mathbf{V}_{\Omega^c} \text{sk} \circ e_2) + e_3$  and combine with  $e_1$  to  $c'' \in \mathbb{Z}_q^n$ ;

return  $\mathbf{V}^{-1} c'' \bmod p$ .

## Security:

$e_1 = \mathbf{V}_\Omega(f \cdot r' + s')$  defines an instance of PV-Knap with  $\text{pk}$ ,  $e_2$  and  $e_3$  as additional information.

$\Rightarrow$  leaky variant of **PV-Knap**, that we call the **PASS problem**.

# PASS Encrypt [HS15]

[HS15]	Our work
deterministic	randomized
without proof of security	with proof of security
fixed $\mathbf{V}_\Omega$	random $\mathbf{V}_\Omega$

Let  $p \ll q$  be two primes,  $m \in \{0, 1\}^n$ ,  $\psi$  a distribution over  $\mathbb{Z}^n$  and  $t = n/2$ .

**KeyGen**( $1^\lambda$ ): sample  $f \leftarrow \psi$  and  $\Omega \leftarrow \text{Unif}(\mathcal{P}_t)$ ; return  $\text{sk} = f$  and  $\text{pk} = (\Omega, \mathbf{V}_\Omega f)$

**Enc**( $\text{pk}, m$ ): sample  $r, s \leftarrow \psi$ ; set  $r' = pr$  and  $s' = m + ps$

$$e_1 = (\text{pk} \circ \mathbf{V}_\Omega r') + \mathbf{V}_\Omega s' = \mathbf{V}_\Omega(f \cdot r' + s')$$

$$e_2 = \mathbf{V}_\Omega^c r'$$

$$e_3 = \mathbf{V}_\Omega^c s'$$

return  $c = (e_1, e_2, e_3)$

**Dec**( $\text{sk}, c$ ): compute  $c' = (\mathbf{V}_\Omega^c \text{sk} \circ e_2) + e_3$  and combine with  $e_1$  to  $c'' \in \mathbb{Z}_q^n$ ;

return  $\mathbf{V}^{-1} c'' \bmod p$ .

## Security:

$e_1 = \mathbf{V}_\Omega(f \cdot r' + s')$  defines an instance of PV-Knap with  $\text{pk}$ ,  $e_2$  and  $e_3$  as additional information.

$\Rightarrow$  leaky variant of **PV-Knap**, that we call the **PASS problem**.



?

PASS problem is tailored to PASS Encrypt!  
Reduce it from some more general problem?

# Properties of PASS Encrypt

## Homomorphic properties:

**Addition:**  $\text{Enc}(\text{pk}, m_1) + \text{Enc}(\text{pk}, m_2) = \text{Enc}(\text{pk}, m_1 + m_2)$

**Multiplication:**  $\text{Enc}(\text{pk}, m_1) \circ \text{Enc}(\text{pk}, m_2) = \text{Enc}(\text{pk}, m_1 \cdot m_2)$

# Properties of PASS Encrypt

## Homomorphic properties:

**Addition:**  $\text{Enc}(\text{pk}, m_1) + \text{Enc}(\text{pk}, m_2) = \text{Enc}(\text{pk}, m_1 + m_2)$

**Multiplication:**  $\text{Enc}(\text{pk}, m_1) \circ \text{Enc}(\text{pk}, m_2) = \text{Enc}(\text{pk}, m_1 \cdot m_2)$

 For  $\circ$ , need of 1 additional cross-term and the decryption algorithm has to be changed.

# Properties of PASS Encrypt

## Homomorphic properties:

**Addition:**  $\text{Enc}(\text{pk}, m_1) + \text{Enc}(\text{pk}, m_2) = \text{Enc}(\text{pk}, m_1 + m_2)$

**Multiplication:**  $\text{Enc}(\text{pk}, m_1) \circ \text{Enc}(\text{pk}, m_2) = \text{Enc}(\text{pk}, m_1 \cdot m_2)$

 For  $\circ$ , need of 1 additional cross-term and the decryption algorithm has to be changed.

## Efficiency:

Scheme	NTRU [ <a href="#">HPS98</a> ]	P-LWE Regev [ <a href="#">LP11</a> ]	PASS Encrypt
$\frac{ c + \text{pk} }{ m }$	$2 \log_2 q$	$3 \log_2 q$	$2.5 \log_2 q$

# Properties of PASS Encrypt

## Homomorphic properties:

**Addition:**  $\text{Enc}(\text{pk}, m_1) + \text{Enc}(\text{pk}, m_2) = \text{Enc}(\text{pk}, m_1 + m_2)$

**Multiplication:**  $\text{Enc}(\text{pk}, m_1) \circ \text{Enc}(\text{pk}, m_2) = \text{Enc}(\text{pk}, m_1 \cdot m_2)$

 For  $\circ$ , need of 1 additional cross-term and the decryption algorithm has to be changed.

## Efficiency:

Scheme	NTRU [ <a href="#">HPS98</a> ]	P-LWE Regev [ <a href="#">LP11</a> ]	PASS Encrypt
$\frac{ c + \text{pk} }{ m }$	$2 \log_2 q$	$3 \log_2 q$	$2.5 \log_2 q$

# Properties of PASS Encrypt

## Homomorphic properties:

**Addition:**  $\text{Enc}(\text{pk}, m_1) + \text{Enc}(\text{pk}, m_2) = \text{Enc}(\text{pk}, m_1 + m_2)$

**Multiplication:**  $\text{Enc}(\text{pk}, m_1) \circ \text{Enc}(\text{pk}, m_2) = \text{Enc}(\text{pk}, m_1 \cdot m_2)$

⚠ For  $\circ$ , need of 1 additional cross-term and the decryption algorithm has to be changed.

## Efficiency:

Scheme	NTRU [HPS98]	P-LWE Regev [LP11]	PASS Encrypt
$\frac{ c + \text{pk} }{ m }$	$2 \log_2 q$	$3 \log_2 q$	$2.5 \log_2 q$

## Concrete Security:

**Known:** key recovery and randomness recovery attacks [HS15, DHSS20]

**New:** plaintext recovery using hints attacks

💡 make use of leaky LWE estimator of Dachman-Soled et al. [DDGR20]

# Conclusion and Perspectives



# Open Questions and Perspectives

## Follow-ups

- Construct encryption scheme based only on PV-LWE / PV-Knap

## Questions ?

- Hardness of partial Vandermonde problems
  - ▶ Cryptanalysis?
  - ▶ Worst-case to average-case reductions as for LWE?
- More cryptographic applications

Thank you.



Miklós Ajtai.

Generating hard instances of lattice problems (extended abstract).

In *STOC*, pages 99–108. ACM, 1996.



Dana Dachman-Soled, Léo Ducas, Huijing Gong, and Mélissa Rossi.

LWE with side information: Attacks and concrete security estimation.

In *CRYPTO (2)*, volume 12171 of *Lecture Notes in Computer Science*, pages 329–358. Springer, 2020.



Yarkin Doröz, Jeffrey Hoffstein, Joseph H. Silverman, and Berk Sunar.

MMSAT: A scheme for multimessage multiuser signature aggregation.

*IACR Cryptol. ePrint Arch.*, page 520, 2020.



Jeffrey Hoffstein, Jill Pipher, and Joseph H. Silverman.

NTRU: A ring-based public key cryptosystem.

In *ANTS*, volume 1423 of *Lecture Notes in Computer Science*, pages 267–288. Springer, 1998.



Jeffrey Hoffstein, Jill Pipher, John M. Schanck, Joseph H. Silverman, and William Whyte.

Practical signatures from the partial fourier recovery problem.

In *ACNS*, volume 8479 of *Lecture Notes in Computer Science*, pages 476–493. Springer, 2014.



Jeffrey Hoffstein and Joseph H. Silverman.

Pass-encrypt: a public key cryptosystem based on partial evaluation of polynomials.

*Des. Codes Cryptogr.*, 77(2-3):541–552, 2015.



Richard Lindner and Chris Peikert.

Better key sizes (and attacks) for LWE-based encryption.

In *CT-RSA*, volume 6558 of *Lecture Notes in Computer Science*, pages 319–339. Springer, 2011.



Xingye Lu, Zhenfei Zhang, and Man Ho Au.

Practical signatures from the partial fourier recovery problem revisited: A provably-secure and gaussian-distributed construction.

In *ACISP*, volume 10946 of *Lecture Notes in Computer Science*, pages 813–820. Springer, 2018.



Daniele Micciancio and Petros Mol.

Pseudorandom knapsacks and the sample complexity of LWE search-to-decision reductions.

In *CRYPTO*, volume 6841 of *Lecture Notes in Computer Science*, pages 465–484. Springer, 2011.



Oded Regev.

On lattices, learning with errors, random linear codes, and cryptography.

In *STOC*, pages 84–93. ACM, 2005.



Peter W. Shor.

Polynomial-time algorithms for prime factorization and discrete logarithms on a quantum computer.

