

有限矩发散条件下的统计分析

A Statistical Analysis of Diverging Moments

郭方健


richardkwo@gmail.com

指导教师: 周涛


英才实验学院 2009 级 1 班

2013 年 6 月 7 日

Overview



Overview




Question

- How do diverging moments grow with sample size?



Overview



Question


- How do diverging moments grow with sample size?

Method

- EPM (Equiprobable Partition Method) \rightarrow *Asymptotics*
- The precision of EPM moment estimators



Overview



Question

- How do diverging moments grow with sample size?

Method

- EPM (Equiprobable Partition Method) \rightarrow *Asymptotics*
- The precision of EPM moment estimators

Application

- Theoretical approximation for statistics with several moments
- Memory constraints for power-law series



Moments

Moments are statistics that characterize

- the **shape** of **distributions** (e.g. *mean, variance, skewness*)
- the **interdependence** among variables (e.g. *Pearson's correlation coefficient*)



Moments

Moments are statistics that characterize

- the **shape** of **distributions** (e.g. *mean, variance, skewness*)
- the **interdependence** among variables (e.g. *Pearson's correlation coefficient*)

A moment corresponds to a **sample moment** and a **population moment**.

k -th moment

Sample moment (algebraic)

$$m_k(c) = \frac{1}{n} \sum_{i=1}^n (x_i - c)^k$$

Population moment (integral)

$$\mu_k(c) = E[(X - c)^k] = \int_{-\infty}^{+\infty} (x - c)^k dF(x)$$



Moments

Moments are statistics that characterize

- the **shape** of **distributions** (e.g. *mean, variance, skewness*)
- the **interdependence** among variables (e.g. *Pearson's correlation coefficient*)

A moment corresponds to a **sample moment** and a **population moment**.

k -th moment

Sample moment (*algebraic*)

$$m_k(c) = \frac{1}{n} \sum_{i=1}^n (x_i - c)^k$$

Population moment (*integral*)

$$\mu_k(c) = E[(X - c)^k] = \int_{-\infty}^{+\infty} (x - c)^k dF(x)$$

- $c = 0$: **raw moment**
- $c = \text{mean}$: **central moment**



Relation between two moments: convergent case

For i.i.d. $\{x_1, x_2, \dots, x_n\}$, when the population moment (*Riemann-Stieltjes integral*) **converges**, we know that the corresponding *sample moment* is an **unbiased estimators** for the *population moment*, i.e.

$$E[m_k(c)] = \mu_k(c).$$



Relation between two moments: convergent case

For i.i.d. $\{x_1, x_2, \dots, x_n\}$, when the population moment (*Riemann-Stieltjes integral*) **converges**, we know that the corresponding *sample moment* is an **unbiased estimators** for the *population moment*, i.e.

$$E[m_k(c)] = \mu_k(c).$$

Then, by the *Weak Law of Large Numbers*, we have

Theorem

A sample moment $m_k(c)$ **converges in probability towards** the corresponding population moment $\mu_k(c)$ when the sample size $n \rightarrow \infty$, namely

$$\lim_{n \rightarrow \infty} P(|m_k(c) - \mu_k(c)| > \epsilon) = 0, \quad \forall \epsilon > 0.$$

given that μ'_k exists (is convergent).

In other words, the estimator is **more accurate with more data**.



Yet, what if population moments diverge?

But a population moment may ***diverge*** for a **heavy-tailed distribution** (e.g. *power-law*, *log-normal*, *Weibull*).


$$\int_{-\infty}^{+\infty} x^k f(x) dx = \infty, \text{ for } k > k_0.$$




Yet, what if population moments diverge?

But a population moment may **diverge** for a **heavy-tailed distribution** (e.g. *power-law, log-normal, Weibull*).

$$\int_{-\infty}^{+\infty} x^k f(x) dx = \infty, \text{ for } k > k_0.$$




Yet, what if population moments diverge?



Yet, what if population moments diverge?

Objective

- Find the asymptotics for diverging moments in the form of $n^\gamma g(n)$.




Yet, what if population moments diverge?

Objective

- Find the asymptotics for diverging moments in the form of $n^\gamma g(n)$.

Application

- A theoretical reference for diverging moments.




Yet, what if population moments diverge?

Objective

- Find the asymptotics for diverging moments in the form of $n^\gamma g(n)$.

Application

- A theoretical reference for diverging moments.
- Approximation for statistics involving several moments, e.g.



$$A = \frac{m_{k_1}}{m_{k_2}} \approx \frac{g_1(n)n^{\gamma_1}}{g_2(n)n^{\gamma_2}} \quad (n \rightarrow \infty).$$



Equiprobable Partition Method (EPM)



Equiprobable Partition Method (EPM)

Equiprobable Partition Method is presented as a tool for constructing the **asymptotics of diverging sample moments**, i.e., answering — “How do diverging sample moments **grow** with n ?”



Equiprobable Partition Method (EPM)

Equiprobable Partition Method is presented as a tool for constructing the **asymptotics of diverging sample moments**, i.e., answering — “How do diverging sample moments **grow** with n ?”

EPM consists of 3 steps:

- ① Constructing equiprobable partitions
- ② Forming moment estimators
(substituting samples with representative points)
- ③ Rewriting into the asymptotic form




Step 1: Equiprobable partitions

Definition

Let X be a continuous random variable defined on the interval $[a, b] \in \mathbb{R}$ with the PDF $f(x)$ ($a \leq x \leq b$), then its **n -separated equiprobable partition** $\mathcal{P}_n(a, b)$ is defined as $\mathcal{P}_n(a, b) = (\hat{t}_0, \hat{t}_1, \dots, \hat{t}_n)$, where $\hat{t}_0 = a$, $\hat{t}_n = b$ and it satisfies that

$$\int_{\hat{t}_i}^{\hat{t}_{i+1}} f(x) dx = \frac{1}{n} \quad (i = 1, 2, \dots, n-1).$$




Step 1: Equiprobable partitions

Definition

Let X be a continuous random variable defined on the interval $[a, b] \in \mathbb{R}$ with the PDF $f(x)$ ($a \leq x \leq b$), then its **n -separated equiprobable partition** $\mathcal{P}_n(a, b)$ is defined as $\mathcal{P}_n(a, b) = (\hat{t}_0, \hat{t}_1, \dots, \hat{t}_n)$, where $\hat{t}_0 = a$, $\hat{t}_n = b$ and it satisfies that

$$\int_{\hat{t}_i}^{\hat{t}_{i+1}} f(x) dx = \frac{1}{n} \quad (i = 1, 2, \dots, n-1).$$



$$\blacksquare F(\hat{t}_i) = \frac{i}{n} \quad (i = 0, 1, \dots, n)$$




Step 1: Equiprobable partitions

Definition

Let X be a continuous random variable defined on the interval $[a, b] \in \mathbb{R}$ with the PDF $f(x)$ ($a \leq x \leq b$), then its **n -separated equiprobable partition** $\mathcal{P}_n(a, b)$ is defined as $\mathcal{P}_n(a, b) = (\hat{t}_0, \hat{t}_1, \dots, \hat{t}_n)$, where $\hat{t}_0 = a$, $\hat{t}_n = b$ and it satisfies that


$$\int_{\hat{t}_i}^{\hat{t}_{i+1}} f(x) dx = \frac{1}{n} \quad (i = 1, 2, \dots, n-1).$$




- $F(\hat{t}_i) = \frac{i}{n} \quad (i = 0, 1, \dots, n)$
- $b = +\infty$ for a right-heavy-tailed distribution



Step 2: Moment estimators via sample substitution



Step 2: Moment estimators via sample substitution




Sample moment

$$m_k(c) = \frac{1}{n} \sum_{i=1}^n (x_i - c)^k$$



Step 2: Moment estimators via sample substitution



Sample moment

$$m_k(c) = \frac{1}{n} \sum_{i=1}^n (x_i - c)^k$$

Substituting $x_{(i)}$ with \hat{t}_{i-1} ,
EPM moment estimator

$$\hat{m}_k(n; c) = \frac{1}{n} \sum_{i=1}^n (\hat{t}_{i-1} - c)^k$$



Step 3: Rewriting into the asymptotic form

Then, we rewrite $\hat{m}_k(n)$ (c is left out for simplicity) into the form of

$$\hat{m}_k(n) = n^\gamma g(n),$$



Step 3: Rewriting into the asymptotic form

Then, we rewrite $\hat{m}_k(n)$ (c is left out for simplicity) into the form of

$$\hat{m}_k(n) = n^\gamma g(n),$$

where

- n^γ ($\gamma > 0$) is the **leading order** that characterizes **the speed of convergence**.



Step 3: Rewriting into the asymptotic form

Then, we rewrite $\hat{m}_k(n)$ (c is left out for simplicity) into the form of

$$\hat{m}_k(n) = n^\gamma g(n),$$

where

- n^γ ($\gamma > 0$) is the **leading order** that characterizes **the speed of convergence**.
- $g(n)$ is a function that satisfies

$$0 < \lim_{n \rightarrow \infty} |g(n)| < \infty,$$

which is the “**convergent remainder**” left by taking off the effect of divergence from the sample moment.



Why equiprobable?

Rationales



Why equiprobable?

Rationales

- ① EPM moments estimators **coincide** with population moments in the **convergent** case.

Theorem

Let X be a random variable defined on the interval $[a, b]$ with its PDF $f(x)$ ($a \leq x \leq b$). Supposing there exists some M that it holds that $\forall x \in [a, b], f(x) < M$, we have

$$\lim_{n \rightarrow \infty} \hat{m}'_k(n) = \mu'_k.$$



Why equiprobable?

Rationales

- 1 EPM moments estimators **coincide** with population moments in the **convergent** case.

Theorem

Let X be a random variable defined on the interval $[a, b]$ with its PDF $f(x)$ ($a \leq x \leq b$). Supposing there exists some M that it holds that $\forall x \in [a, b], f(x) < M$, we have

$$\lim_{n \rightarrow \infty} \hat{m}'_k(n) = \mu'_k.$$

- 2 If viewing *partition* as **discretizing** a continuous distribution, then EPM maximizes the entropy $H = - \sum_{i=1}^n p_i \log(p_i)$.



Why equiprobable?

Rationales

- 1 EPM moments estimators **coincide** with population moments in the **convergent** case.

Theorem

Let X be a random variable defined on the interval $[a, b]$ with its PDF $f(x)$ ($a \leq x \leq b$). Supposing there exists some M that it holds that $\forall x \in [a, b], f(x) < M$, we have

$$\lim_{n \rightarrow \infty} \hat{m}'_k(n) = \mu'_k.$$

- 2 If viewing *partition* as **discretizing** a continuous distribution, then EPM maximizes the entropy $H = -\sum_{i=1}^n p_i \log(p_i)$.
- 3 Getting \hat{t}_i is easy with $F^{-1}(x)$.



Example: Asymptotics for moments of power-law

EPM asymptotics for diverging moments of the power-law distribution.



Example: Asymptotics for moments of power-law

EPM asymptotics for diverging moments of the power-law distribution.
Power-law with $x_{\min} = 1$ and $\alpha > 1$,

$$\text{PDF } f(x) = (\alpha - 1)x^{-\alpha}, \quad \text{CDF } F(x) = 1 - x^{1-\alpha}.$$



Example: Asymptotics for moments of power-law

EPM asymptotics for diverging moments of the power-law distribution.
Power-law with $x_{\min} = 1$ and $\alpha > 1$,

$$\text{PDF } f(x) = (\alpha - 1)x^{-\alpha}, \quad \text{CDF } F(x) = 1 - x^{1-\alpha}.$$

1. n -separated equiprobable partition $\mathcal{P}_n(1, +\infty) = \{\hat{t}_1, \hat{t}_2, \dots, \hat{t}_n\}$.

From

$$F(\hat{t}_i) = 1 - (\hat{t}_i)^{1-\alpha} = \frac{i}{n},$$

we have

$$\hat{t}_i = \left(1 - \frac{i}{n}\right)^{-c} \quad (i = 0, 1, \dots, n-1),$$

where $c = \frac{1}{\alpha-1} > \frac{1}{2}$ is a shorthand.



Example: Asymptotics for moments of power-law

2. The EPM estimator for m'_k is

$$\hat{m}'_k(n) = \frac{1}{n} \sum_{i=0}^{n-1} (\hat{t}_i)^k = \frac{1}{n} \sum_{i=0}^{n-1} \left(1 - \frac{i}{n}\right)^{-ck} \quad \left(c = \frac{1}{\alpha - 1}\right).$$



Example: Asymptotics for moments of power-law

2. The EPM estimator for m'_k is

$$\hat{m}'_k(n) = \frac{1}{n} \sum_{i=0}^{n-1} (\hat{t}_i)^k = \frac{1}{n} \sum_{i=0}^{n-1} \left(1 - \frac{i}{n}\right)^{-ck} \quad \left(c = \frac{1}{\alpha - 1}\right).$$

3. Rewriting it into

$$\hat{m}'_k(n) = \frac{1}{n} \sum_{i=0}^{n-1} n^{ck} (n - i)^{-ck} = n^{ck-1} \sum_{i=1}^n \frac{1}{i^{ck}},$$



Example: Asymptotics for moments of power-law

2. The EPM estimator for m'_k is

$$\hat{m}'_k(n) = \frac{1}{n} \sum_{i=0}^{n-1} (\hat{t}_i)^k = \frac{1}{n} \sum_{i=0}^{n-1} \left(1 - \frac{i}{n}\right)^{-ck} \quad \left(c = \frac{1}{\alpha - 1}\right).$$

3. Rewriting it into

$$\hat{m}'_k(n) = \frac{1}{n} \sum_{i=0}^{n-1} n^{ck} (n - i)^{-ck} = n^{ck-1} \sum_{i=1}^n \frac{1}{i^{ck}},$$

where

- n^{ck-1} is the leading order,



Example: Asymptotics for moments of power-law

2. The EPM estimator for m'_k is

$$\hat{m}'_k(n) = \frac{1}{n} \sum_{i=0}^{n-1} (\hat{t}_i)^k = \frac{1}{n} \sum_{i=0}^{n-1} \left(1 - \frac{i}{n}\right)^{-ck} \quad \left(c = \frac{1}{\alpha - 1}\right).$$

3. Rewriting it into

$$\hat{m}'_k(n) = \frac{1}{n} \sum_{i=0}^{n-1} n^{ck} (n - i)^{-ck} = n^{ck-1} \sum_{i=1}^n \frac{1}{i^{ck}},$$

where

- n^{ck-1} is the leading order,
- $\sum_{i=1}^n \frac{1}{i^{ck}} \rightarrow \text{Riemann } \zeta(ck) \ (n \rightarrow \infty)$ is the remaining “convergent term”.



Example: Asymptotics for moments of power-law




Figure : The comparison between numerical sample moments m_2 , m_3 , m_4 (circle) and their EPM asymptotics (line) for power-law with $\alpha = 2.5$.



Time series and their distributions



Time series and their distributions

Time series — an ordered sequence of random variables

$$(t_1, t_2, \dots, t_n).$$



Time series and their distributions

Time series — an ordered sequence of random variables

$$(t_1, t_2, \dots, t_n).$$

Distribution characterization

- ① **Marginal** distribution, often assumed to be **stable**

$$f_{t_1}(t) = f_{t_2}(t) = \dots = f_{t_n}(t).$$



Time series and their distributions

Time series — an ordered sequence of random variables

$$(t_1, t_2, \dots, t_n).$$

Distribution characterization

- 1 **Marginal** distribution, often assumed to be **stable**

$$f_{t_1}(t) = f_{t_2}(t) = \dots = f_{t_n}(t).$$

- 2 **Interdependence** among elements.

Memory (*1st-order autocorrelation*) is defined as the Pearson's correlation coefficient between $\{t_2, t_3, \dots, t_n\}$ and its lag-1 counterpart $\{t_1, t_2, \dots, t_{n-1}\}$, i.e.

$$M = \frac{1}{n-1} \sum_{i=1}^{n-1} \frac{(t_i - m_1)(t_{i+1} - m_2)}{\sigma_1 \sigma_2} \quad (-1 \leq M \leq 1).$$



Correlated series with a specified marginal



Correlated series with a specified marginal

Modeling correlated series with marginal distribution $f(t)$.



Correlated series with a specified marginal

Modeling correlated series with marginal distribution $f(t)$.

- 1 Sampling i.i.d. (t_1, t_2, \dots, t_n) from the same marginal $f(t)$.



Correlated series with a specified marginal

Modeling correlated series with marginal distribution $f(t)$.

- ① Sampling i.i.d. (t_1, t_2, \dots, t_n) from the same marginal $f(t)$.
 - ② Sorting samples to $t_{(1)} \leq t_{(2)} \leq \dots \leq t_{(n)}$.
-



Correlated series with a specified marginal

Modeling correlated series with marginal distribution $f(t)$.

- 1 Sampling i.i.d. (t_1, t_2, \dots, t_n) from the same marginal $f(t)$.
- 2 Sorting samples to $t_{(1)} \leq t_{(2)} \leq \dots \leq t_{(n)}$.
- 3 Modifying interdependence with a **permutation**
 $\theta : \{1, \dots, n\} \leftrightarrow \{1, \dots, n\}$, so we have $(t_{(\theta_1)}, t_{(\theta_2)}, \dots, t_{(\theta_n)})$.

¹M. Hallin, G. Melard, X. Milhaud. Permutational extreme values of autocorrelation coefficients and a Pitman test against serial dependence. *Annals of Statistics*, 1992.



Correlated series with a specified marginal

Modeling correlated series with marginal distribution $f(t)$.

- 1 Sampling i.i.d. (t_1, t_2, \dots, t_n) from the same marginal $f(t)$.
- 2 Sorting samples to $t_{(1)} \leq t_{(2)} \leq \dots \leq t_{(n)}$.
- 3 Modifying interdependence with a **permutation**
 $\theta : \{1, \dots, n\} \leftrightarrow \{1, \dots, n\}$, so we have $(t_{(\theta_1)}, t_{(\theta_2)}, \dots, t_{(\theta_n)})$.

Interestingly, there are **fixed** θ_{\max} and θ_{\min} that maximizes and minimizes M for the permuted series¹.

¹M. Hallin, G. Melard, X. Milhaud. Permutational extreme values of autocorrelation coefficients and a Pitman test against serial dependence. *Annals of Statistics*, 1992.



Correlated series with a specified marginal

Modeling correlated series with marginal distribution $f(t)$.

- ① Sampling i.i.d. (t_1, t_2, \dots, t_n) from the same marginal $f(t)$.
- ② Sorting samples to $t_{(1)} \leq t_{(2)} \leq \dots \leq t_{(n)}$.
- ③ Modifying interdependence with a **permutation**
 $\theta : \{1, \dots, n\} \leftrightarrow \{1, \dots, n\}$, so we have $(t_{(\theta_1)}, t_{(\theta_2)}, \dots, t_{(\theta_n)})$.

Interestingly, there are **fixed** θ_{\max} and θ_{\min} that maximizes and minimizes M for the permuted series ¹.


$$\theta_{\max} : t_{(1)}, t_{(3)}, \dots, t_{(2l-1)}, t_{(2l)}, t_{(2l-2)}, \dots, t_{(4)}, t_{(2)} \quad (n = 2l),$$

$$\theta_{\min} : t_{(2l)}, t_{(1)}, t_{(2l-2)}, \dots, t_{(2l-3)}, t_{(2)}, t_{(2l-1)} \quad (n = 2l).$$


¹M. Hallin, G. Melard, X. Milhaud. Permutational extreme values of autocorrelation coefficients and a Pitman test against serial dependence. *Annals of Statistics*, 1992.



Correlated series with a specified marginal



Correlated series with a specified marginal




Therefore, we define the **(tight) bounds** for memory as

$$M_{\max} = \lim_{n \rightarrow \infty} \mathbb{E}[M(t_{\theta_{\max}})] \text{ and } M_{\min} = \lim_{n \rightarrow \infty} \mathbb{E}[M(t_{\theta_{\min}})].$$



Correlated series with a specified marginal



Therefore, we define the **(tight) bounds** for memory as

$$M_{\max} = \lim_{n \rightarrow \infty} \mathbb{E}[M(t_{\theta_{\max}})] \text{ and } M_{\min} = \lim_{n \rightarrow \infty} \mathbb{E}[M(t_{\theta_{\min}})].$$

For **Gaussian** and **uniform** distributions, we have

$$M_{\max} = 1 \text{ and } M_{\min} = -1 \text{ (trivial, as } |M| \leq 1 \text{)}.$$



Probabilistic method for the $\alpha > 3$ case

$\alpha > 3$ is necessary for the population variance $\sigma(\alpha)^2$ to converge.

$$\mathbb{E}[M(t_{(\theta)})] = \frac{1}{\sigma(\alpha)^2} \left(\frac{1}{n-1} \sum_{i=1}^{n-1} \mathbb{E}[t_{(\theta_i)} t_{(\theta_{i+1})}] - m(\alpha)^2 \right),$$



Probabilistic method for the $\alpha > 3$ case

$\alpha > 3$ is necessary for the population variance $\sigma(\alpha)^2$ to converge.

$$\mathbb{E}[M(t_{(\theta)})] = \frac{1}{\sigma(\alpha)^2} \left(\frac{1}{n-1} \sum_{i=1}^{n-1} \mathbb{E}[t_{(\theta_i)} t_{(\theta_{i+1})}] - m(\alpha)^2 \right),$$

where

$$m(\alpha) = \int_1^{+\infty} x f(x) dx = \frac{\alpha - 1}{\alpha - 2},$$

$$\sigma(\alpha)^2 = \int_1^{+\infty} x^2 f(x) dx - m(\alpha)^2 = \frac{\alpha - 1}{\alpha - 3} - m(\alpha)^2.$$



Probabilistic method for the $\alpha > 3$ case

$\alpha > 3$ is necessary for the population variance $\sigma(\alpha)^2$ to converge.

$$\mathbb{E}[M(t_{(\theta)})] = \frac{1}{\sigma(\alpha)^2} \left(\frac{1}{n-1} \sum_{i=1}^{n-1} \mathbb{E}[t_{(\theta_i)} t_{(\theta_{i+1})}] - m(\alpha)^2 \right),$$

where

$$m(\alpha) = \int_1^{+\infty} x f(x) dx = \frac{\alpha - 1}{\alpha - 2},$$

$$\sigma(\alpha)^2 = \int_1^{+\infty} x^2 f(x) dx - m(\alpha)^2 = \frac{\alpha - 1}{\alpha - 3} - m(\alpha)^2.$$

We can use the **distribution for order statistics** to derive

$$M_{\max} = 1,$$

$$M_{\min} = \frac{1}{\sigma(\alpha)^2} \left[2B\left(\frac{1}{2}; \frac{1}{m(\alpha)}, \frac{1}{m(\alpha)}\right) - m(\alpha)^2 \right] \quad (\alpha > 3).$$



EPM for the $\alpha \leq 3$ case

When $\alpha \leq 3$, m and σ^2 may diverge, where we can use EPM estimators to tackle the diverging moments in the limit of large n .



EPM for the $\alpha \leq 3$ case

When $\alpha \leq 3$, m and σ^2 may diverge, where we can use EPM estimators to tackle the diverging moments in the limit of large n .

Rewriting M to

$$M = \frac{s - m^2}{t^2 - m^2},$$



EPM for the $\alpha \leq 3$ case

When $\alpha \leq 3$, m and σ^2 may diverge, where we can use EPM estimators to tackle the diverging moments in the limit of large n .

Rewriting M to

$$M = \frac{s - m^2}{\bar{t}^2 - m^2},$$

where

- $s = \frac{1}{n-1} S = \frac{1}{n-1} \sum_{i=1}^{n-1} t_i t_{i+1}$,
- $\bar{t}^2 = \frac{1}{n} \sum_{i=1}^n t_i^2$,
- and $m^2 = (\frac{1}{n} \sum_{i=1}^n t_i)^2$

are the moments in concern.



EPM for the $\alpha \leq 3$ case

$$M = \frac{s - m^2}{t^2 - m^2}.$$



EPM for the $\alpha \leq 3$ case

$$M = \frac{s - m^2}{t^2 - m^2}.$$

For θ_{\max} , we have EPM estimators

$$\hat{s}_{\theta_{\max}} = n^{2c-1} \left[\sum_{k=2}^{n-1} (k^2 - 1)^{-c} + 2^{-c} \right] \quad \left(c = \frac{1}{\alpha - 1} \right),$$

$$\overline{\hat{t}^2}_{\theta_{\max}} = n^{2c-1} \sum_{k=0}^{n-1} (k+1)^{-2c},$$



EPM for the $\alpha \leq 3$ case

$$M = \frac{s - m^2}{\bar{t}^2 - m^2}.$$

For θ_{\max} , we have EPM estimators

$$\hat{s}_{\theta_{\max}} = n^{2c-1} \left[\sum_{k=2}^{n-1} (k^2 - 1)^{-c} + 2^{-c} \right] \quad \left(c = \frac{1}{\alpha - 1} \right),$$

$$\overline{\hat{t}^2}_{\theta_{\max}} = n^{2c-1} \sum_{k=0}^{n-1} (k+1)^{-2c},$$

while

$$\hat{m}^2 \sim n^{2c-2}.$$

\hat{m}^2 can be thrown away.



EPM for the $\alpha \leq 3$ case

Therefore, we have

$$M_{\max} \approx \frac{\lim_{n \rightarrow \infty} \sum_{k=2}^{n-1} (k^2 - 1)^{-c} + 2^{-c}}{\zeta(2c)} \quad \left(c = \frac{1}{\alpha - 1}, 1 < \alpha \leq 3\right).$$



EPM for the $\alpha \leq 3$ case

Therefore, we have

$$M_{\max} \approx \frac{\lim_{n \rightarrow \infty} \sum_{k=2}^{n-1} (k^2 - 1)^{-c} + 2^{-c}}{\zeta(2c)} \quad \left(c = \frac{1}{\alpha - 1}, 1 < \alpha \leq 3\right).$$

For θ_{\min} , in $M = \frac{s-m^2}{t^2-m^2}$, $\overline{t^2}_{\theta_{\min}} \sim n^{2c-1}$ is the term diverging with the **biggest** order of n .



EPM for the $\alpha \leq 3$ case

Therefore, we have

$$M_{\max} \approx \frac{\lim_{n \rightarrow \infty} \sum_{k=2}^{n-1} (k^2 - 1)^{-c} + 2^{-c}}{\zeta(2c)} \quad (c = \frac{1}{\alpha - 1}, 1 < \alpha \leq 3).$$

For θ_{\min} , in $M = \frac{s-m^2}{t^2-m^2}$, $\overline{t^2}_{\theta_{\min}} \sim n^{2c-1}$ is the term diverging with the **biggest** order of n .

Because the term with the biggest order only appears in the **denominator**, we have

$$M_{\min} \approx \lim_{n \rightarrow \infty} \hat{M}_{\theta_{\min}} = 0 \quad (1 < \alpha \leq 3).$$



Numerical simulation vs. theory




Figure : Theoretical bounds for M_{\min} and M_{\max} and numerical simulations with different series lengths. Each point in simulation is produced by averaging 1000 independent runs.



Empirical results




Figure : Memory for power-law distributed **inter-event time series** from empirical inter-event time series, where each series is represented by a point and theoretical bounds are drawn as solid curves. Left: *MovieLens* dataset for online movie rating. Right: *Twitter* dataset for sending tweets. Power-law is examined with KS-test and α is fitted with MLE ³.

³A. Clauset, C. Shalizi, M. Newman. Power-Law Distributions in Empirical Data. *SIAM Review*, 2009.

Summary



Summary

Contributions

- EPM asymptotics for diverging moments.



Summary

Contributions

- EPM asymptotics for diverging moments.
- Non-trivial bounds for the memory of power-law series (paper in preparation).



Summary

Contributions

- EPM asymptotics for diverging moments.
- Non-trivial bounds for the memory of power-law series (paper in preparation).

Future work

- Proving the *convergence in probability* of the EPM asymptotics, i.e.

$$\lim_{n \rightarrow \infty} P \left(\left| \frac{1}{n^{\gamma+1}} \sum_{i=1}^n x_i^k - g(n) \right| \geq \epsilon \right) = 0, \quad \forall \epsilon > 0.$$



Summary

Contributions

- EPM asymptotics for diverging moments.
- Non-trivial bounds for the memory of power-law series (paper in preparation).

Future work

- Proving the *convergence in probability* of the EPM asymptotics, i.e.

$$\lim_{n \rightarrow \infty} P \left(\left| \frac{1}{n^{\gamma+1}} \sum_{i=1}^n x_i^k - g(n) \right| \geq \epsilon \right) = 0, \quad \forall \epsilon > 0.$$

Thanks!

