
Causal Inference

a summary

Contents

1	General	3	2.2 G-Methods	4
2	Models	4	2.2.1 Time-varying A	5
	2.1 Traditional Methods	4	3 Longitudinal Data	6

1 General

Causal Roadmap (Petersen and van der Laan, 2014)

systematic approach linking causality to statistical procedures

1. Specifying Knowledge. structural causal model (unifying counterfactual language, structural equations, & causal graphs): a set of possible data-generating processes, expresses background knowledge and its limits

2. Linking Data. specifying measured variables and sampling specifics (latter can be incorporated into the model)

3. Specifying Target. define hypothetical experiment: decide

1. variables to intervene on: one (point treatment), multiple (longitudinal, censoring/missing, (in)direct effects)
2. intervention scheme: static, dynamic, stochastic
3. counterfactual summary of interest: absolute or relative, marginal structural models, interaction, effect modification
4. population of interest: whole, subset, different population

4. Assessing Identifiability. are knowledge and data sufficient to derive estimand and if not, what else is needed?

5. Select Estimand. current best answer: knowledge-based assumptions + which minimal convenience-based assumptions (transparency) gets as close as possible

6. Estimate. choose estimator by statistical properties, nothing causal here

7. Interpret. hierarchy: statistical, counterfactual, feasible intervention, randomized trial

Average Causal Effect $E[Y^{a=1}] \neq E[Y^{a=0}]$

$$E[Y^a] = \sum_y yp_{Y^a}(y) \quad (\text{discrete})$$

$$= \int y f_{Y^a}(y) dy \quad (\text{continuous})$$

individual causal effect $Y_i^{a=1} \neq Y_i^{a=0}$ generally unidentifiable

null hypothesis: no average causal effect

sharp null hypothesis: no causal effect for any individual

notation A, Y : random variables (differ for individuals); a, y : particular values; counterfactual $Y^{a=1}$: Y under treatment $a = 1$

stable unit treatment value assumption (SUTVA) Y_i^a is well-defined: no interference between individuals, no multiple versions of treatment (weaker: treatment variation irrelevance)

causal effect measures typically based on means

risk difference: $\Pr[Y^{a=1} = 1] - \Pr[Y^{a=0} = 1]$

risk ratio: $\frac{\Pr[Y^{a=1}=1]}{\Pr[Y^{a=0}=1]}$

odds ratio: $\frac{\Pr[Y^{a=1}=1]/\Pr[Y^{a=1}=0]}{\Pr[Y^{a=0}=1]/\Pr[Y^{a=0}=0]}$

number needed to treat (NNT) to save 1 life: $-1/\text{risk difference}$

sources of random error: sampling variability (use consistent estimators), nondeterministic counterfactuals

association compares $E[Y|A = 1]$ and $E[Y|A = 0]$, **causation** compares $E[Y^{a=1}]$ and $E[Y^{a=0}]$ (whole population)

Target Trial emulating an ideal randomized experiment

explicitly formulate target trial & show how it is emulated → less vague causal question, helps spot issues

missing data problem unknown counterfactuals

randomized experiments: missing completely at random →

exchangeability (= exogeneity as treatment is exogenous)

ideal randomized experiment: no censoring, double-blind,

well-defined treatment, & adherence → association is causation

pragmatic trial: no placebo/blindness, realistic monitoring

PICO (population, intervention, comparator, outcome): some components of target trial

three types of causal effects:

intention-to-treat effect (effect of treatment assignment)

per-protocol effect (usually dynamic when toxicity arises)

other intervention effect (strategy changed during follow-up)

controlled direct effects: effect of A on Y not through B

natural direct effect A on Y if $B^{a=0}$ (cross-world quantity)

principal stratum effect A on Y for subset with $B^{a=0} = B^{a=1}$

crossover experiment: sequential treatment & outcome $t=0, 1$

individual causal effect $Y_{it}^{a_t=1} - Y_{it}^{a_t=0}$ only identifiable if: no

carryover effect, effect $\perp\!\!\!\perp$ time, outcome $\perp\!\!\!\perp$ time

time zero if eligibility at multiple t (observational data):

earliest, random t , all t (adjust variance with bootstrapping)

grace periods: usually treatment starts x months after first

eligible, if death before: randomly assign strategy/copy into both

Identifiability Conditions hold in ideal experiments

consistency counterfactuals correspond to data $Y = Y^A$: if

$A = a$, then $Y^a = Y$ for each individual

positivity $\Pr[A = a|L = l] > 0 \forall l$ with $\Pr[L = l] > 0$

exchangeability $Y^a \perp\!\!\!\perp A$

(conditionally) randomized = conditionally exchangeable

$Y^a \perp\!\!\!\perp A|L$

no guarantee: without randomization unmeasured

confounders U may exist

most of 3

positivity: p. 155, p. 162

additional conditions: chapter 13.5

exchangeability: p 172f, p16-19

positivity: $f_L(l) \neq 0 \Rightarrow f_{A|L}(a|l) > 0 \forall a, l$

consistency: technical point 3.2

effect modification chapter 4

interaction chapter 5

causal diagrams chapter 6, include swigs from 7.5 and that one technical point

more on SWIGS p 242ff

confounding chapter 7

selection bias chapter 8

measurement bias chapter 9

random variability chapter 10

2 Models

Modeling data are a sample from the target population

estimand: quantity of interest, e. g. $E[Y|A = a]$
estimator: function to use, e. g. $\hat{E}[Y|A = a]$
estimate: apply function to data, e. g. 4.1

model: a priori restriction of joint distribution/dose-response curve; *assumption*: no model misspecification (usually wrong)

non-parametric estimator: no restriction (saturated model) = *Fisher consistent estimator* (entire population data \rightarrow true value)

parsimonious model: few parameters estimate many quantities

bias-variance trade-off:

wiggleness $\uparrow \rightarrow$ misspecification bias \downarrow , CI width \uparrow

2.1 Traditional Methods

Outcome regression chapter 15

instrumental variable estimation chapter 16

causal survival analysis chapter 17 (and technical point 22.3)

Variable Selection can induce bias if L includes:

(descendant of) collider: *selection bias under the null*
 noncollider effect of A : *selection bias under the alternative*
 mediator: *overadjustment for mediators*

temporal ordering is not enough to conclude anything

bias amplification: e.g. by adjusting for an instrument Z (can also reduce bias)

Machine Learning L is high-dimensional

use lasso or ML for IP weighting/standardization

but: ML does not guarantee elimination of confounding and has largely unknown statistical properties

\rightarrow **doubly robust estimator**: consistent if bias $< \frac{1}{\sqrt{n}}$

sample splitting: train estimators on training sample, use resulting estimators for doubly robust method on estimation sample (CIs on estimation sample are valid, but n halved)

cross-fitting: do again the other way round, average the two estimates, get CI via bootstrapping

problems: unclear choice of algorithm, is bias small enough?

2.2 G-Methods

G-Methods generalized treatment contrasts: adjust for (surrogate) confounders L

- **standardization** two types of g-formula
- **IP weighting** also g-formula
- **g-estimation**: not needed unless longitudinal

Standardization plug-in (or parametric if so) g-formula

$$E[Y^a] = \overbrace{E[E[Y|A=a, L=l]]}^{\text{conditional expectation}} = \overbrace{\int E[Y|L=l, A=a] f_L[l] dl}^{\text{joint density estimator}}$$

weighted average of stratum-specific risks; unknowns can be estimated non-parametrically or modeled

no need to estimate $f_L[l]$ /integrate as empirical distribution can be used: estimate outcome model \rightarrow predict counterfactuals on whole dataset \rightarrow average the results (\rightarrow CI by bootstrapping)

for discrete L $E[Y|A=a] = \sum_l E[Y|L=l, A=a] \Pr[L=l]$

time-varying standardize over all possible \bar{l} -histories
 simulates joint distribution of counterfactuals $(Y^{\bar{a}}, \bar{L}^{\bar{a}})$ for \bar{a}
joint density estimator (jde)

$$\text{discrete: } E[Y^{\bar{a}}] = \sum_{\bar{l}} E[Y|\bar{A}=\bar{a}, \bar{L}=\bar{l}] \prod_{k=0}^K f(l_k|\bar{a}_{k-1}, \bar{l}_{k-1})$$

$$\text{continuous: } \int f(y|\bar{a}, \bar{l}) \prod_{k=0}^K f(l_k|\bar{a}_{k-1}, \bar{l}_{k-1}) dl$$

for *stochastic strategies* multiply with $\prod_{k=0}^K f^{int}(a_k|\bar{a}_{k-1}, \bar{l}_k)$

estimation (Young et al., 2011; Schomaker et al., 2019)

1. model $f(l_k|\bar{a}_{k-1}, \bar{l}_{k-1})$ and $E[Y|\bar{A}=\bar{a}, \bar{L}=\bar{l}]$
2. simulate data forward in time:
 at $k=0$: use empirical distribution of L_0 (observed data)
 at $k>0$: set $\bar{A}=\bar{a}$, draw from models estimated in 1.
3. calculate mean of $\hat{Y}_{K,i}^{\bar{a}}$ (bootstrap for CI)

iterated conditional expectation (ice)

$$E[Y_T^{\bar{a}}] = E[E[E[...E[Y_T|\bar{A}_{T-1}=\bar{a}_{T-1}, \bar{L}_T]...|\bar{A}_0=a_0, L_1]|L_0]]$$

estimation (Schomaker et al., 2019)

1. model inside out: $Q_T = E[Y_T|\bar{A}_{T-1}, \bar{L}_T]$ to $Q_0 = E[Q_1|\bar{L}_0]$, predict Q_t with $\bar{A}=\bar{a}$ in each step
2. calculate mean of $\hat{Q}_{0,i}^{\bar{a}}$ (bootstrap for CI)

g-null paradox even if the sharp null holds, model misspecification can lead to it being falsely rejected

Proof: for $L_0 \rightarrow A_0 \rightarrow Y_0 \rightarrow L_1 \rightarrow A_1 \rightarrow Y_1$, $\bar{a} = (a_0, a_1)$
 $E[Y_1^{\bar{a}}] \stackrel{\text{CE}}{=} E[E[Y_1^{\bar{a}} | A_0=a_0, L_0]]$

$$\begin{aligned} (\text{ice}) & \stackrel{\text{CE}^*}{=} E[E[E[Y_1 | \bar{L}, \bar{A}=\bar{a}, Y_0] | A_0=a_0, L_0]] \\ & \stackrel{\text{LTP}}{=} E\left[\sum_{l_1} E[Y_1 | A_0=a_0, \bar{L}, Y_0] \Pr[l_1 | a_0, l_0, y_0]\right] \\ & \stackrel{\text{LTP}}{=} \sum_{l_0} \left[\sum_{l_1} E[Y_1 | A_0=a_0, \bar{L}, Y_0] \Pr[l_1 | a_0, l_0, y_0]\right] \Pr[l_0] \\ (\text{jde}) & \stackrel{\text{sum}}{=} \sum_i E[Y_i | A_0=a_0, \bar{L}, Y_0] \Pr[l_1 | a_0, l_0] \Pr[l_0] \end{aligned}$$

CE: conditional expectation; *: exchangeability;
LTP: law of total probability

Marginal Structural Models association is causation
in the IP weighted pseudo-population

associational model $E[Y|A]$ = causal model $E[Y^a]$

step 1: estimate/model $f[A|L]$ (and $f[A]$) \rightarrow get $(SW)^A$

step 2: estimate regression parameters for pseudo-population
effect modification variables V can be included (e.g.

$\beta_0 + \beta_1 a + \beta_2 V a + \beta_3 V$; technically not marginal anymore),
 $SW^A(V) = \frac{f[A|V]}{f[A|L]}$ more efficient than SW^A

Censoring measuring joint effect of A and C

$$E[Y^{a,c=0}] \text{ is of interest}$$

standardization $E[Y|A=a] = \int E[Y|L=l, A=a, C=0] dF_L[l]$

IP weights $W^{A,C} = W^A \times W^C$ (uses n) or
 $SW^{A,C} = SW^A \times SW^C$ (uses $n^{c=0}$)

g-estimation can only adjust for confounding, not selection bias
 \rightarrow use IP weights

G-Estimation (additive) structural nested models

$$\text{logit Pr}[A=1|H(\psi^\dagger), L] = \alpha_0 + \alpha_1 H(\psi^\dagger) + \alpha_2 L$$

$$H(\psi^\dagger) = Y - \psi_1 A$$

find ψ^\dagger which renders $\alpha_1 = 0$; 95%-CI: all ψ^\dagger for which $p > 0.05$
closed-form solution for linear models

derivation: $H(\psi^\dagger) = Y^{a=0}$

$$\text{logit Pr}[A=1|Y^{a=0}, L] = \alpha_0 + \alpha_1 Y^{a=0} + \alpha_2 L$$

$Y^{a=0}$ unknown, but because of exchangeability α_1 should be zero

$$Y^{a=0} = Y^a - \psi_1 a$$

equivalent to $Y^{a=0} = Y^{a=1} - \psi_1$, but using no counterfactuals

structural nested mean model

$$\text{additive: } E[Y^a - Y^{a=0} | A=a, L] = \beta_1 a + \beta_2 a L$$

$$\text{multiplicative: } \log\left(\frac{E[Y^a | A=a, L]}{E[Y^{a=0} | A=a, L]}\right) = \beta_1 a + \beta_2 a L$$

multiplicative is preferred if Y always positive, but does not
extend to longitudinal case

semi-parametric: agnostic about β_0 and effect of $L \rightarrow$ robust \uparrow

no time-varying: no nesting; model equals marginal structural
models with missing β_0, β_3 (unspecified “no treatment”)

sensitivity analysis: unmeasured confounding ($\alpha_1 \neq 0$) can be
examined: do procedure for different values of $\alpha_1 \rightarrow$ plot α_1 vs.
 $\psi^\dagger \rightarrow$ how sensitive is estimate to unmeasured confounding?

effect modification: add V in both g-estimation equations

doubly robust estimators exist

IP Weighting inverse probability of treatment (g-formula)

$$E[Y^a] = E\left[\frac{I(A=a)Y}{f[A|L]}\right]; W^A = \frac{1}{f[A|L]}; SW^A = \frac{f(A)}{f[A|L]}$$

unknowns can be estimated non-parametrically or modeled

pseudo-population: everyone is treated & untreated ($L \not\rightarrow A$)

FRCISTG (fully randomized causally interpreted structured
graph): probability tree for $L \rightarrow A \rightarrow Y$, can be used to
calculate/visualize simulation of values for A
for discrete A, L $f[a|l] = \Pr[A=a, L=l]$
estimators: Horvitz-Thompson; Hajek (modified version)
stabilized weights SW^A should have an average of 1 (check!)
 \rightarrow pseudo-population same size \rightarrow CI width \downarrow

Standardization and IP Weighting are equivalent,
but if modeled, different “no misspecification” assumptions:
standardization: outcome model
IP weighting: treatment model

doubly robust estimators: reduce model misspecification bias,
consistent if either model is correct; **e.g.:**
1. fit outcome regression with variable $R = \begin{cases} +W^A & \text{if } A=1 \\ -W^A & \text{if } A=0 \end{cases}$
2. standardize by averaging

2.2.1 Time-varying A

IP Weighting

$$W^{\bar{A}} = \prod_{k=0}^K \frac{1}{f(A_k | \bar{A}_{k-1}, \bar{L}_k)}$$

$$SW^{\bar{A}} = \prod_{k=0}^K \frac{f(A_k | \bar{A}_{k-1}, \bar{L}_k)}{f(A_k | \bar{A}_{k-1}, \bar{L}_k)}$$

Doubly Robust Estimator sequential estimation

1. estimate $\hat{f}(A_m | \bar{A}_{m-1}, \bar{L}_m)$ (e.g. logistic model), use it to
calculate at each time m : $\widehat{W}^{\bar{A}_m} = \prod_{k=0}^m \frac{1}{\hat{f}(A_k | \bar{A}_{k-1}, \bar{L}_k)}$ and
modified IP weights at m : $\widehat{W}^{\bar{A}_{m-1}, a_m} = \frac{\widehat{W}^{\bar{A}_{m-1}}}{\hat{f}(a_m | \bar{A}_{m-1}, \bar{L}_m)}$
 2. with $\widehat{T}_{K+1} := Y$, recursively for $m = K, K-1, \dots, 0$:
(a) fit outcome regression on \widehat{T}_{m+1} with variable $\widehat{W}^{\bar{A}_m}$
(b) calculate \widehat{T}_m using the outcome model with $\widehat{W}^{\bar{A}_{m-1}, a_m}$
 3. calculate standardized mean outcome $\widehat{E}[Y^{\bar{a}}] = E[\widehat{T}_0]$
- valid, if** treatment or outcome model correct, or treatment
correct until k and outcome otherwise ($k+1$ robustness)

G-Estimation nested equations: for each time k

structural nested mean models separate effect of each a_k

$$E[Y^{\bar{a}_{k-1}, a_k, 0_{k+1}} - Y^{\bar{a}_{k-1}, 0_{k+1}} | \bar{L}^{\bar{a}_{k-1}} = \bar{L}_k, \bar{A}_{k-1} = \bar{a}_{k-1}] =$$

$$a_k \gamma_k(\bar{a}_{k-1}, \bar{L}_k, \beta)$$

calculations

$$H_k(\psi^\dagger) = Y - \sum_{j=k}^K A_j \gamma_j(\bar{A}_{j-1}, \bar{L}_j, \psi^\dagger)$$

function γ_j can be, e.g. constant (ψ_1), time-varying only
($\psi_1 + \psi_2 k$), or dependent on treatment/covariate history

$$\text{logit Pr}[A_k=1|H_k(\psi^\dagger), \bar{L}_k, \bar{A}_{k-1}] =$$

$$\alpha_0 + \alpha_1 H_k(\psi^\dagger) + \alpha_2 w_k(\bar{L}_k, \bar{A}_{k-1})$$

find α_1 that is closest to zero

closed form estimator exists for the linear case

Censoring \bar{C} : monotonic type of missing data

standardization:

$$\int f(y|\bar{a}, \bar{c}=0, \bar{l}) \prod_{k=0}^K dF(l_k | \bar{a}_{k-1}, c_{k-1}=0, \bar{l}_{k-1})$$

IP weighting:

$$SW^{\bar{C}} = \prod_{k=1}^{K+1} \frac{1 \cdot \Pr(C_k=0 | \bar{A}_{k-1}, C_{k-1}=0)}{\Pr(C_k=0 | \bar{A}_{k-1}, C_{k-1}=0, \bar{L}_k)}$$

3 Longitudinal Data

Time-Varying Treatments compare 2 treatments

treatment history up to k : $\bar{A}_k = (A_0, A_1, \dots, A_k)$

shorthand: always treated $\bar{A} = \bar{1}$, never treated $\bar{A} = (\bar{0})$

static strategy: $g = [g_0(\bar{a}_{-1}), \dots, g_K(\bar{a}_{K-1})]$

dynamic strategy: $g = [g_0(\bar{l}_0), \dots, g_K(\bar{l}_K)]$

stochastic strategy: non-deterministic g

optimal strategy is where $E[Y^g]$ is maximized (if high is good)

Sequential Identifiability sequential versions of

exchangability: $Y^g \perp\!\!\!\perp A_k | \bar{A}_{k-1} \quad \forall g, k = 0, 1, \dots, K$

conditional exchangeability:

$$(Y^g, L_{k+1}^g) \perp\!\!\!\perp A_k | \bar{A}_{k-1} = g(\bar{L}_k), \bar{L}^k \quad \forall g, k = 0, 1, \dots, K$$

positivity: $f_{\bar{A}_{k-1}, \bar{L}_k}(\bar{a}_{k-1}, \bar{l}_k) \neq 0 \Rightarrow$

$$f_{A_k | \bar{A}_{k-1}, \bar{L}_k}(a_k | \bar{a}_{k-1}, \bar{l}_k) > 0 \quad \forall (\bar{a}_{k-1}, \bar{l}_k)$$

consistency:

$$Y^{\bar{a}} = Y^{\bar{a}^*} \quad \text{if } \bar{a} = \bar{a}^*; \quad Y^{\bar{a}} = Y \quad \text{if } \bar{A} = \bar{a};$$

$$\bar{L}_k^{\bar{a}} = \bar{L}_k^{\bar{a}^*} \quad \text{if } \bar{a}_{k-1} = \bar{a}_{k-1}^*; \quad \bar{L}_k^{\bar{a}} = \bar{L}_k \quad \text{if } \bar{A}_{k-1} = \bar{a}_{k-1}$$

generalized backdoor criterion (static strategy): all backdoors into A_k (except through future treatments) are blocked $\forall k$

static sequential exchangeability for $Y^{\bar{a}}$

$$Y^{\bar{a}} \perp\!\!\!\perp A_k | \bar{A}_{k-1}, \bar{L}_k \quad \text{for } k = 0, 1, \dots, K$$

use SWIGs to visually check d-separation

time-varying confounding $E[Y^{\bar{a}} | L_0] \neq E[Y | A = \bar{a}, L_0]$

Treatment-Confounder Feedback $A_0 \rightarrow L_1 \rightarrow A_1$:

an unmeasured U influencing L_1 and Y turns L_1 into a collider;

traditional adjustment (e.g. stratification) biased: use g-methods

g-null test sequential exchangeability & sharp null true \Rightarrow

$Y^g = Y \quad \forall g \Rightarrow Y \perp\!\!\!\perp A_0 | L_0 \text{ \& } Y \perp\!\!\!\perp A_1 | A_0, L_0, L_1$; therefore:

if last two independences don't hold, one assumption is violated

g-null theorem: $E[Y^g] = E[Y]$, if the two independences hold

(\Rightarrow sharp null: only if strong faithfulness (no effect cancelling))

References

If no citation is given, the information is taken from the book (Hernán and Robins, 2020)

Hernán, M. A. and Robins, J. M. (2020). *Causal inference: what if*. Boca Raton: Chapman & Hall/CRC.

Petersen, M. L. and van der Laan, M. J. (2014). Causal models and learning from data: integrating causal modeling and statistical estimation. *Epidemiology (Cambridge, Mass.)*, 25(3):418–426.

Schomaker, M., Luque-Fernandez, M. A., Leroy, V., and Davies, M.-A. (2019). Using longitudinal targeted maximum likelihood estimation in complex settings with dynamic interventions. *Statistics in medicine*, 38(24):4888–4911. ISBN: 0277-6715 Publisher: Wiley Online Library.

Young, J. G., Cain, L. E., Robins, J. M., O'Reilly, E. J., and Hernán, M. A. (2011). Comparative effectiveness of dynamic treatment regimes: an application of the parametric g-formula. *Statistics in biosciences*, 3:119–143.

