

# CS & IT ENGINEERING



## Database Management System

Relational model and Normal forms

DPP\_01 Discussion Notes



By- Anjnee Bhatnagar

#Q. Consider the student relation shown below with schema stud (Sname, Sage, Smail, Smarks),

**Stud**

<u>Sname</u>	<u>Sage</u>	<u>Smail</u>	<u>Smarks</u>
<u>Rohit</u>	<u>28</u>	<u>R@pw.live</u>	<u>68</u>
<u>Kanika</u>	<u>25</u>	<u>K@pw.live</u>	<u>75</u>
<u>Pankaj</u>	<u>25</u>	<u>K@pw.live</u>	<u>75</u>
<u>Rohit</u>	<u>28</u>	<u>R@pw.live</u>	<u>88</u>
<u>Anjali</u>	<u>26</u>	<u>A@pw.live</u>	<u>75</u>



1

For the above given instance how many set of attributes of size two can determine each row uniquely?

Sname Sage X

Sname Smarks ✓



Sname Smail X

Sage Smail X

Smail Smarks X

Smarks Sage X

#Q. Consider a relation schema  $R(\underline{A}, \underline{B}, \underline{C}, \underline{D}, \underline{E}, \underline{F}, \underline{H})$  with the given Functional dependency set:

$\{A \rightarrow BC, C \rightarrow AD, DE \rightarrow F, C \rightarrow F\}$

The attribute closure that contains all the attributes of the relation R is?

☐ A  $\underline{AE}^+ = \{A, E, B, C, D, F\}$  ✗

☐ B  $\underline{CE}^+ = \{C, E, F, A, D\}$  ✗

☒ C  $\underline{AEH}^+ = \{A, E, B, C, D, F, H\}$

☐ D All of the above

# [MCQ]



#Q. Consider the following set of FD's:  
 $\{V \rightarrow W, W \rightarrow XZ, X \rightarrow YZ\}$  for relation  
 $R(\underline{V}, \underline{W}, \underline{X}, \underline{Y}, \underline{Z})$   $\angle$   
Then the attribute closure of  $YZ^+$  contains how many elements?

$$YZ^+ = \{Y, Z\}$$

A

0

B

1

C

2

D

3

# [MCQ]



#Q. For the given FD set:  $\{P \rightarrow QT, Q \rightarrow SU, V \rightarrow U\}$  of a relation  $R(P, Q, T, S, U, V)$ . Find the set of attributes that is Super key but not a Candidate key?

A PTQ

B PV

~~C PQV~~

~~D QV~~

$PV^+ = \{P, V, Q, T, S, U\}$  ✓  
Candidate  
key



A Venn diagram illustrating the relationship between candidate keys and super keys. It features two concentric ellipses. The inner ellipse is labeled 'Candidate key' and is completely contained within the outer ellipse, which is labeled 'super key'. This visualizes that every candidate key is a super key, but not every super key is a candidate key.

Candidate key

super key

#Q. Choose the correct statement from the following.

A The cardinality is defined as the number of ~~attributes~~ in a relation. X

B Degree of the relation is the number of tuples in the relation. X

☒ C Relation instance is the set of tuples of a relation at a particular instance of time.

D All of the above X



#Q. Choose the correct statement from the following:

A There can be many primary keys for a relation. ~~X~~

B There can be many alternate keys for a relation.


C All the candidate keys are also super keys.

D All the super keys are also the candidate keys. ~~X~~

minimal set  
of attributes  
that uniquely  
identify a tuple  
in a relation.

#Q. Consider the following statements:

- $S_1$ : A key in DBMS is an attribute (or a set of attributes) that helps in uniquely identifying each tuple (or row) in a relation (or table). *True*
- $S_2$ : There should be only one candidate key in relation, which is chosen as the primary key.



A

Only  $S_1$  is true.

B

Only  $S_2$  is true.

C

Both  $S_1$  and  $S_2$  are true.

D

Neither  $S_1$  nor  $S_2$  is true.

#Q. Consider the following statements:

$S_1$ : Primary key has no duplicate values it has only unique values. ✓

$S_2$ : Primary key is not necessarily formed using a single column of the table, more than one column of the table can also be used to form a primary key of the table. ✓

A

Only  $S_1$  is true.

B

Only  $S_2$  is true.

C

Both  $S_1$  &  $S_2$  are true. ✓

D

Neither  $S_1$  nor  $S_2$  are true.

#Q. Assume a relation R (P, Q, R, S, T). If PR and RT are the only candidate keys of the relation R, then how many total super keys exist in relation R.

$$\begin{array}{ccc}
 PR & & RT \\
 \downarrow & & \downarrow \\
 5-2 & & 5-2 \quad 5-3 \\
 2 & + & 2 \quad - \quad 2 \\
 \hline
 2^3 & + & 2^3 \quad - \quad 2^2 \Rightarrow 8 + 8 - 4 \\
 \Rightarrow & & \Rightarrow \underline{12}
 \end{array}$$

#Q. Assume a relation R (P, Q, R, S, T, U, V).  
If  $\overline{PQ}$ ,  $\overline{RS}$ , and  $\overline{TU}$  are the only three candidate keys of relation R, then how many total super keys exist in relation R?

$\overline{PQ}$

↓

$$\Rightarrow 2^{7-2} + 2^{7-2} + 2^{7-2} - (2^{7-4} + 2^{7-4} + 2^{7-4})$$

$$\Rightarrow 2^5 + 2^5 + 2^5 - (2^3 + 2^3 + 2^3) + 2$$

$$32 + 32 + 32 - (24) + 2$$



$$64 + 8 + 2 \Rightarrow 64 + 10 = \underline{\underline{74}}.$$

**THANK - YOU**

