

To: National Cancer Institute-Nagpur
Khasra No 25
Outer Hingna ring Road
Nagpur- 441108
Maharashtra
Report Of: Mrs. JAYASHREE SADANANDRAO
POHEKAR



Sample ID : 2300109621
Patient ID : 1002350016
Collected on : 04-07-2023 12:54:37
Received on : 06-07-2023 14:34:10
Reported on : 11-07-2023 22:20:15
Ref By : National Cancer Institute-
Nagpur

MOLECULAR GENETICS REPORT

Patient Name	: Mrs. JAYASHREE SADANANDRAO POHEKAR	Age	: 40 Years
Physician Name	: DR.ANAND PATHAK	Gender	: Female
Provisional	: CML-CP	Specimen	: Ok
Diagnosis		Status	
Specimen Type	: Peripheral Blood (PB)	Disease Status:	Followup
Test Requested	: BCR-ABL1 RT-qPCR for p210 (IS)		

Test : BCR-ABL1 RT-qPCR for p210 (IS) Analysis.

Result: Positive

BCR-ABL1 RT-qPCR for p210 (IS) Analysis:

BCR-ABL1 Type p210	ABL1 gene absolute copy number (aCN)	Normalized BCR-ABL1/ ABL1 ratio%	BCR-ABL1 International Scale (IS) %	Molecular Response
Detected	2,48,597	14.1558 %	9.9515 %	No MMR

Conclusion: The specimen tested positive for the BCR-ABL1 p210 transcript with the value of International Scale (IS %) **9.9515 %**. This indicates that the patient has not achieved major molecular response (No MMR). Kindly correlate with the clinical findings.

Mrs. JAYASHREE SADANANDRAO POHEKAR

Sample ID: 2300109621

Methodology:


Total RNA was extracted from the specimen, reverse transcribed and the common p210 fusion transcripts (b2a2 & b3a2) are measured with breakpoint probes using real-time polymerase chain reaction. ABL1 serves as an endogenous reference to monitor RNA quality and to report the percent relative ratio of BCR-ABL1 to ABL1. Levels of BCR-ABL1 are reported on a standardized International Scale (IS). On the IS, a major molecular response (MMR) represents a 3-log reduction in BCR-ABL and is defined as $\leq 0.1\%$ IS. Deep molecular response-4 (DMR-4) represents a 4-log reduction from the IRIS baseline ($\leq 0.01\%$ IS); DMR-4.5 represents a 4.5-log reduction from IRIS baseline ($\leq 0.0032\%$ IS); DMR-5 represents a 5-log reduction from IRIS baseline ($\leq 0.001\%$ IS). Alternate/rare breakpoints are not detected/ quantitated in this assay. The limit of detection of this assay is 3 copies of the BCR-ABL1 transcript. The reproducibility of this assay is such that results within 0.5 log should be considered equivalent.

References:

1. Yoshida et al: Validation of a rapid one-step high sensitivity real-time quantitative PCR system for detecting major BCR-ABL1 mRNA on an International Scale. Springer Plus 2016; 5: 569-575.
2. Cross N.C.P. et.al: Laboratory recommendations for scoring deep molecular responses following treatment for chronic myeloid leukemia. Leukemia 2015; 29: 999-1003.
3. Baccarani et al: European Leukemia Net recommendations for the management of chronic myeloid leukemia: Blood. 2013;122(6):872-884.
4. Branford et al: Desirable performance characteristics for BCR-ABL measurement on an international reporting scale to allow consistent interpretation of individual patient response and comparison of response rate between clinical trials. Blood 2008; 112: 3330-3338.
5. Jerald PR et al: Chronic Myeloid Leukemia, Version 1.2019, NCCN Clinical Practice Guidelines in Oncology J Natl Compr Canc Netw. 2018;16(9):1108-1135.
6. NCCN Guidelines of monitoring Chronic Myeloid leukemia, 2020

Mrs. JAYASHREE SADANANDRAO POHEKAR

Sample ID: 2300109621



BCR-ABL1 qPCR p210 (IS%)

Prepared By : **Dr. Aniket Prabhudesai**
Verified By : **Mr. Rohit Waghole**

Mrs. JAYASHREE SADANANDRAO POHEKAR

Sample ID: 2300109621



Dr. P. S. Kadam Amare
Oncogeneticist
Chief & Lab Director "Cancer &
Clinical Genetics"
Lilac Insights Pvt. Ltd.



Dr. Aniket Prabhudesai
Sr. Scientific Officer
Molecular Oncology Dept
Lilac Insights Pvt. Ltd.

- End of Report -

Conditions of Reporting/Disclaimer:

- This report is based on the sample received in the Lilac Insights laboratory; the analysis is based on the assumption that samples received are representative of the patient mentioned on the test requisition form and the sample. When samples are received from various referral centers, it is presumed that patient demographics are verified at the point of sample collection.
- All samples for molecular studies must be collected in EDTA tubes (lavender cap). A sample where the test requisition requires RNA and subsequent cDNA conversion, must be maintained and transported at 4°C until it reaches Lilac Insights Pvt. Ltd. within 24 hours to prevent degradation of the RNA in the sample. FFPE tissue blocks often yield fragmented DNA of low concentrations which can impact result quality.
- Assays are performed in accordance with standard procedure on receipt. The reported results are dependent on individual assay methods, equipment used, method specificity, sensitivity and quality of specimen(s) received.
- Despite all the necessary precautions and stringency adopted whilst performing DNA tests, the currently available data indicates that the technical error rate associated with all types of DNA analysis, is approximately 2%.
- Although molecular testing is highly accurate, rarely false-positive & false-negative diagnostic errors may occur due to improper quality control during sample collection, cellular integrity of sample, selection of inappropriate specimen and/or presence of PCR inhibitors. PCR primer binding site polymorphisms or mutations might lead to allele dropout & cause false negative results.
- It is important that all clinicians or persons requesting DNA diagnostic tests are aware of these data before acting upon these results. As with all diagnostic tests, the laboratory report must be interpreted in conjunction with the presenting clinical profile of the patient and evaluation of all reports.
- In sequencing based tests sometimes variants of unknown significance (VUCS) are detected that have either not been reported before, and/or whose effect cannot be determined based on the current knowledge standards and reporting guidelines. In such cases, we recommend periodic review of these variants to determine any change in classification based on new published research.
- Lilac Insights Pvt. Ltd. has policy to return the FFPE blocks within one month after final reporting with proper documentation of the dispatch of the block to customer from accession dept, Lilac. After dispatch, if there is no intimation from customer within two weeks, Lilac will not be responsible for the dispatched FFPE block.
- Soft copies of reports are sent to customer by office mail ID. Also, Hardcopies are sent to customer only on the address provided by client.