

* Mathematics - 3 (M3) - Assignment Number - 2

Name:- Kaustubh Shrikant Kabra.

Class:- Second Year Engineering.

Div:- A

Batch:-

Department:- Computer Department.

College:- AISSMS, IOIT.

Roll Number:- ERP No.:- 34, Teas Sr. No.:- 20

Q-1. Solve the integral equation (May 2019)

$$\int_0^{\infty} f(x) \cos(\lambda x) dx = 1 - \lambda, \quad 0 \leq \lambda < 1$$

$$= 0, \quad \lambda \geq 1$$

$$\rightarrow F(\lambda) = \int_0^{\infty} f(u) \cos(\lambda u) du = \begin{cases} 1 - \lambda, & 0 \leq \lambda < 1 \\ 0, & \lambda \geq 1 \end{cases}$$

$$f(x) = \frac{2}{\pi} \int_0^{\infty} F(\lambda) \cos(\lambda x) d\lambda$$

$$= \frac{2}{\pi} \left[\int_0^1 (1 - \lambda) \cos(\lambda x) d\lambda + \int_1^{\infty} (0) \cos(\lambda x) d\lambda \right]$$

$$= \frac{2}{\pi} \left[(1 - \lambda) \left(\frac{\sin(\lambda x)}{x} \right) - (-1) \left(-\frac{\cos(\lambda x)}{x^2} \right) \right]_0^1$$

$$= \frac{2}{\pi} \left[-\frac{\cos x}{x^2} + \frac{1}{x^2} \right]$$

$$= \frac{2}{\pi} \left[\frac{1 - \cos x}{x^2} \right]$$

$$\therefore \boxed{f(x) = \frac{2}{\pi} \left[\frac{1 - \cos x}{x^2} \right]}$$

Q-2. Find F.T of $f(x) = \begin{cases} 1-x^2, & |x| \leq 1 \\ 0, & \text{otherwise} \end{cases}$ (May 2018)

→ Here, $f(x) = \begin{cases} 1-x^2, & -1 \leq x \leq 1 \\ 0, & \text{otherwise} \end{cases}$

F.T for $f(x)$ is given by

$$F(\lambda) = \int_{-\infty}^{\infty} f(u) \cdot e^{-i\lambda u} \cdot du$$

$$= \int_{-\infty}^{-1} f(u) \cdot e^{-i\lambda u} \cdot du + \int_{-1}^1 f(u) \cdot e^{-i\lambda u} \cdot du + \int_1^{\infty} f(u) \cdot e^{-i\lambda u} \cdot du$$

$$= \int_{-\infty}^{-1} 0 \cdot du + \int_{-1}^1 (1-u^2) \cdot e^{-i\lambda u} \cdot du + \int_1^{\infty} 0 \cdot du$$

$$= \int_{-1}^1 (1-u^2) (\cos \lambda u - i \sin \lambda u) \cdot du$$

$$= \int_{-1}^1 \underbrace{(1-u^2)}_{\text{even}} \underbrace{(\cos \lambda u)}_{\text{even}} \cdot du - i \int_{-1}^1 \underbrace{(1-u^2)}_{\text{even}} \underbrace{(\sin \lambda u)}_{\text{odd}} \cdot du$$

$$= 2 \int_0^1 (1-u^2) \cos(\lambda u) \cdot du - i(0)$$

$$= 2 \left[\frac{(1-u^2) \sin \lambda u}{\lambda} - \frac{2u \cos \lambda u}{\lambda^2} + \frac{2 \sin \lambda u}{\lambda^3} \right]_0^1$$

$$= 2 \left[\frac{-2 \cos \lambda}{\lambda^2} + \frac{2 \sin \lambda}{\lambda^3} \right]$$

$$\therefore \text{F.T of } f(x) = \frac{4 \sin \lambda}{\lambda^3} - \frac{4 \cos \lambda}{\lambda^2}$$

Q-3. Find FIR of $f(x) = \begin{cases} 1, & |x| < 1 \\ 0, & |x| > 1 \end{cases}$ [May 2017]

Hence evaluate $\int_0^\infty \frac{\sin \lambda}{\lambda} \cos \lambda x \cdot dx \cdot d\lambda$ and $\int_0^\infty \frac{\sin \lambda}{\lambda} \cdot d\lambda$

$$\rightarrow f(x) = \begin{cases} 1, & -1 < x < 1 \\ 0, & \text{otherwise} \end{cases}$$

$$F(\lambda) = \int_{-\infty}^{\infty} f(u) \cdot e^{-i\lambda u} \cdot du$$

$$= \int_{-\infty}^{-1} f(u) \cdot e^{-i\lambda u} \cdot du + \int_{-1}^1 f(u) \cdot e^{-i\lambda u} \cdot du + \int_1^{\infty} f(u) \cdot e^{-i\lambda u} \cdot du$$

$$= 0 + \int_{-1}^1 1 \cdot e^{-i\lambda u} \cdot du + 0$$

$$= \int_{-1}^1 (\cos \lambda u - i \sin \lambda u) \cdot du$$

$$= \int_{-1}^1 \underbrace{\cos \lambda u}_{\text{even}} \cdot du - i \int_{-1}^1 \underbrace{\sin \lambda u}_{\text{odd}} \cdot du$$

$$= 2 \left[\frac{\sin \lambda u}{\lambda} \right]_0^1$$

$$F(\lambda) = \frac{2 \sin \lambda}{\lambda}$$

For FIR, take inverse F.T of $F(\lambda)$

$$f(x) = \frac{1}{2\pi} \int_{-\infty}^{\infty} F(\lambda) \cdot e^{-i\lambda x} \cdot d\lambda$$

$$f(x) = \frac{1}{2\pi} \int_{-\infty}^{\infty} \frac{2 \sin \lambda}{\lambda} \cdot e^{-i\lambda x} \cdot d\lambda$$

$$= \frac{1}{\pi} \left[\int_{-\infty}^{\infty} \underbrace{\frac{\sin \lambda}{\lambda}}_{\text{even}} \cdot \underbrace{\cos \lambda x}_{\text{even}} \cdot d\lambda + i \int_{-\infty}^{\infty} \underbrace{\frac{\sin \lambda}{\lambda}}_{\text{even}} \cdot \underbrace{\sin \lambda x}_{\text{odd}} \cdot d\lambda \right]$$

$$f(x) = \frac{2}{\pi} \int_0^{\infty} \frac{\sin \lambda}{\lambda} \cdot \cos \lambda x \cdot d\lambda$$

↳ FIR of $f(x)$.

Hence, $f(x) = \frac{2}{\pi} \int_0^{\infty} \frac{\sin \lambda}{\lambda} \cdot \cos \lambda x \cdot d\lambda$

$$\begin{aligned} \int_0^{\infty} \frac{\sin \lambda}{\lambda} \cdot \cos \lambda x \cdot d\lambda &= \frac{\pi}{2} \cdot f(x) \\ &= \frac{\pi}{2} \cdot \begin{cases} 1, & |x| < 1 \\ 0, & |x| > 1 \end{cases} \end{aligned}$$

$$= \begin{cases} \pi/2, & |x| < 1 \\ 0, & |x| > 1 \end{cases}$$

put $x = 0$, $\therefore \int_0^{\infty} \frac{\sin \lambda}{\lambda} \cdot d\lambda = \boxed{\frac{\pi}{2}}$

Q-4. Find $F(z) = \frac{1}{(z-3)(z-4)}$, $|z| > 4$ [Nov. 2017]

$$\rightarrow F(z) = \frac{1}{(z-3)(z-4)}$$

$$= \frac{-1}{(z-3)} + \frac{1}{(z-4)}$$

$$z^{-1}(F(z)) = z^{-1} \left(\frac{-1}{(z-3)} \right) + z^{-1} \left(\frac{1}{(z-4)} \right)$$

$$= \frac{1}{3} z^{-1} \left[\frac{-3}{z-3} \right] + \frac{1}{4} z^{-1} \left[\frac{4}{z-4} \right]$$

$|z| > 3 \qquad \qquad \qquad |z| > 4$

$$= \frac{-1}{3} \cdot 3^k \cdot \quad + \quad \frac{1}{4} \cdot 4^k$$

$k > 0 \qquad \qquad \qquad k > 0$

$$\therefore z^{-1} \left\{ \frac{1}{(z-3)(z-4)} \right\} = \begin{cases} 0 & , k \leq 0 \\ 4^{k-1} - 3^{k-1} & , k > 0 \end{cases}$$

Q-5. Solve the following difference equation:-

$$12f(k+2) - 7f(k+1) + f(k) = 0, \quad k \geq 0, \quad f(0) = 0, \quad f(1) = 3$$

$$\rightarrow \text{Given DE} \rightarrow 12f(k+2) - 7f(k+1) + f(k) = 0.$$

Applying z-transform, we get

$$12z \{f(k+2)\} - 7z \{f(k+1)\} + z \{f(k)\} = z \{0\}$$

$$z \{f(k+2)\} = z^2 F(z) - z^2 f(0) - z f(1) = z^2 F(z) - 3z$$

$$z \{f(k+1)\} = z F(z) - z f(0) = z F(z)$$

$$\therefore 12 [z^2 F(z) - 3z] - 7 [z F(z) + F(z)] = 0$$

$$F(z) = \frac{36z}{12z^2 - 7z + 1}$$

$$= \frac{36z}{(4z-1)(3z-1)}$$

$$= \frac{36z}{4(z-1/4)3(z-1/3)}$$

$$= \frac{3z}{(z-1/4)(z-1/3)} \Rightarrow \text{Poles are } 1/4 \text{ and } 1/3$$

$$F(z)z^{k-1} = \frac{3z}{(z-1/4)(z-1/3)} \cdot z^{k-1} = \frac{3z^k}{(z-1/4)(z-1/3)}$$

Residue of $F(z)z^{k-1}$ at pole $z = 1/4$ is

$$= \frac{3 \cdot (1/4)^k}{\left[\frac{1}{4} - \frac{1}{3} \right]} = \frac{-36}{4^k} = \frac{-9}{4^{k-1}}$$

Residue of $F(z) \cdot z^{k-1}$ at pole $z = 1/3$ is

$$\left[(z - 1/3) \cdot F(z) \cdot z^{k-1} \right]_{z=1/3} = \frac{3 \left(\frac{1}{3} \right)^k}{\left[\frac{1}{3} - \frac{1}{4} \right]} = \frac{36}{3^k} = \frac{4}{3^{k-2}}$$

$\therefore z^{-1} \{F(z)\} = \text{Sum of residue}$

$$= \frac{36}{3^k} - \frac{36}{4^k}$$

$$= 36 \left[\left(\frac{1}{3} \right)^k - \left(\frac{1}{4} \right)^k \right]$$

$$\boxed{f(k) = 36 \left[\left(\frac{1}{3} \right)^k - \left(\frac{1}{4} \right)^k \right]}$$