

Plan

- Dual of a Primal
- Fundamental theorem of Duality
- The Complementary Slackness Theorem

Dual of a Primal

- Given an LPP (P) (*)

Max $\mathbf{c}^T \mathbf{x}$

subject to $A_{m \times n} \mathbf{x} \leq \mathbf{b}_{m \times 1}$, $\mathbf{x} \geq \mathbf{0}$

Dual of a Primal

- Given an LPP (P) (*)
Max $\mathbf{c}^T \mathbf{x}$
subject to $A_{m \times n} \mathbf{x} \leq \mathbf{b}_{m \times 1}$, $\mathbf{x} \geq \mathbf{0}$
- $Fea(P) = \{\mathbf{x} \in \mathbb{R}^n : A\mathbf{x} \leq \mathbf{b}, \mathbf{x} \geq \mathbf{0}\}$

Dual of a Primal

- Given an LPP (P) (*)
Max $\mathbf{c}^T \mathbf{x}$
subject to $A_{m \times n} \mathbf{x} \leq \mathbf{b}_{m \times 1}$, $\mathbf{x} \geq \mathbf{0}$
- $Fea(P) = \{\mathbf{x} \in \mathbb{R}^n : A\mathbf{x} \leq \mathbf{b}, \mathbf{x} \geq \mathbf{0}\}$
- The Dual of this problem is given by (D) (**)
Min $\mathbf{b}^T \mathbf{y}$

Dual of a Primal

- Given an LPP (P) (*)
Max $\mathbf{c}^T \mathbf{x}$
subject to $A_{m \times n} \mathbf{x} \leq \mathbf{b}_{m \times 1}, \mathbf{x} \geq \mathbf{0}$
- $Fea(P) = \{\mathbf{x} \in \mathbb{R}^n : A\mathbf{x} \leq \mathbf{b}, \mathbf{x} \geq \mathbf{0}\}$
- The Dual of this problem is given by (D) (**)
Min $\mathbf{b}^T \mathbf{y}$
subject to $A_{n \times m}^T \mathbf{y} \geq \mathbf{c}_{n \times 1}, \mathbf{y} \geq \mathbf{0}$

Dual of a Primal

- Given an LPP (P) (*)
Max $\mathbf{c}^T \mathbf{x}$
subject to $A_{m \times n} \mathbf{x} \leq \mathbf{b}_{m \times 1}, \mathbf{x} \geq \mathbf{0}$
- $Fea(P) = \{\mathbf{x} \in \mathbb{R}^n : A\mathbf{x} \leq \mathbf{b}, \mathbf{x} \geq \mathbf{0}\}$
- The Dual of this problem is given by (D) (**)
Min $\mathbf{b}^T \mathbf{y}$
subject to $A_{n \times m}^T \mathbf{y} \geq \mathbf{c}_{n \times 1}, \mathbf{y} \geq \mathbf{0}$
- $Fea(D) = \{\mathbf{y} \in \mathbb{R}^m : A^T \mathbf{y} \geq \mathbf{c}, \mathbf{y} \geq \mathbf{0}\}$

Dual of a Primal

- Given an LPP (P) (*)
Max $\mathbf{c}^T \mathbf{x}$
subject to $A_{m \times n} \mathbf{x} \leq \mathbf{b}_{m \times 1}, \mathbf{x} \geq \mathbf{0}$
- $Fea(P) = \{\mathbf{x} \in \mathbb{R}^n : A\mathbf{x} \leq \mathbf{b}, \mathbf{x} \geq \mathbf{0}\}$
- The Dual of this problem is given by (D) (**)
Min $\mathbf{b}^T \mathbf{y}$
subject to $A_{n \times m}^T \mathbf{y} \geq \mathbf{c}_{n \times 1}, \mathbf{y} \geq \mathbf{0}$
- $Fea(D) = \{\mathbf{y} \in \mathbb{R}^m : A^T \mathbf{y} \geq \mathbf{c}, \mathbf{y} \geq \mathbf{0}\}$
- The LPP(*) is called a Primal problem

- **Theorem 1:** If $\mathbf{x} \in \text{Fea}(P)$ and $\mathbf{y} \in \text{Fea}(D)$, then
 $\mathbf{c}^T \mathbf{x} \leq \mathbf{b}^T \mathbf{y}$

- **Theorem 1:** If $\mathbf{x} \in \text{Fea}(P)$ and $\mathbf{y} \in \text{Fea}(D)$, then $\mathbf{c}^T \mathbf{x} \leq \mathbf{b}^T \mathbf{y}$
- **Corollary 1:** If both the **Primal** and the **Dual** have **feasible solutions** then both have **optimal solutions**

- **Theorem 1:** If $\mathbf{x} \in \text{Fea}(P)$ and $\mathbf{y} \in \text{Fea}(D)$, then $\mathbf{c}^T \mathbf{x} \leq \mathbf{b}^T \mathbf{y}$
- **Corollary 1:** If both the **Primal** and the **Dual** have **feasible solutions** then both have **optimal solutions**
- **Corollary 2:** Let $\mathbf{x}_0 \in \text{Fea}(P)$ and $\mathbf{y}_0 \in \text{Fea}(D)$, be such that $\mathbf{c}^T \mathbf{x}_0 = \mathbf{b}^T \mathbf{y}_0$, then $\mathbf{x}_0$ and $\mathbf{y}_0$ are **optimal** for the **Primal** and the **Dual**, respectively

- **Theorem 1:** If $\mathbf{x} \in \text{Fea}(P)$ and $\mathbf{y} \in \text{Fea}(D)$, then $\mathbf{c}^T \mathbf{x} \leq \mathbf{b}^T \mathbf{y}$
- **Corollary 1:** If both the **Primal** and the **Dual** have **feasible solutions** then both have **optimal solutions**
- **Corollary 2:** Let $\mathbf{x}_0 \in \text{Fea}(P)$ and $\mathbf{y}_0 \in \text{Fea}(D)$, be such that $\mathbf{c}^T \mathbf{x}_0 = \mathbf{b}^T \mathbf{y}_0$, then $\mathbf{x}_0$ and $\mathbf{y}_0$ are **optimal** for the **Primal** and the **Dual**, respectively
- **Theorem 3:** The **Dual** of the **Dual** (D) (of the Primal (P)) is the **Primal** (P)

- **Example 1:** Consider the problem **(P)**
Max $5x_1 + 2x_2$ subject to

- **Example 1:** Consider the problem **(P)**

Max $5x_1 + 2x_2$ subject to

$$3x_1 + 2x_2 \leq 6$$

- **Example 1:** Consider the problem **(P)**

Max $5x_1 + 2x_2$ subject to

$$3x_1 + 2x_2 \leq 6$$

$$x_1 + 2x_2 \leq 4$$

- **Example 1:** Consider the problem (P)

Max $5x_1 + 2x_2$ subject to

$$3x_1 + 2x_2 \leq 6$$

$$x_1 + 2x_2 \leq 4$$

$$x_1 \geq 0, x_2 \geq 0$$

- **Example 1:** Consider the problem **(P)**

Max $5x_1 + 2x_2$ subject to

$$3x_1 + 2x_2 \leq 6$$

$$x_1 + 2x_2 \leq 4$$

$$x_1 \geq 0, x_2 \geq 0$$

- The dual of **(P)** is given by **(D)**

- **Example 1:** Consider the problem **(P)**

Max $5x_1 + 2x_2$ subject to

$$3x_1 + 2x_2 \leq 6$$

$$x_1 + 2x_2 \leq 4$$

$$x_1 \geq 0, x_2 \geq 0$$

- The dual of **(P)** is given by **(D)**

Min $6y_1 + 4y_2$

- **Example 1:** Consider the problem **(P)**

Max $5x_1 + 2x_2$ subject to

$$3x_1 + 2x_2 \leq 6$$

$$x_1 + 2x_2 \leq 4$$

$$x_1 \geq 0, x_2 \geq 0$$

- The dual of **(P)** is given by **(D)**

Min $6y_1 + 4y_2$

subject to

- **Example 1:** Consider the problem **(P)**

Max $5x_1 + 2x_2$ subject to

$$3x_1 + 2x_2 \leq 6$$

$$x_1 + 2x_2 \leq 4$$

$$x_1 \geq 0, x_2 \geq 0$$

- The dual of **(P)** is given by **(D)**

Min $6y_1 + 4y_2$

subject to

$$3y_1 + y_2 \geq 5$$

- **Example 1:** Consider the problem **(P)**

Max $5x_1 + 2x_2$ subject to

$$3x_1 + 2x_2 \leq 6$$

$$x_1 + 2x_2 \leq 4$$

$$x_1 \geq 0, x_2 \geq 0$$

- The dual of **(P)** is given by **(D)**

Min $6y_1 + 4y_2$

subject to

$$3y_1 + y_2 \geq 5$$

$$2y_1 + 2y_2 \geq 2$$

- **Example 1:** Consider the problem **(P)**

Max $5x_1 + 2x_2$ subject to

$$3x_1 + 2x_2 \leq 6$$

$$x_1 + 2x_2 \leq 4$$

$$x_1 \geq 0, x_2 \geq 0$$

- The dual of **(P)** is given by **(D)**

Min $6y_1 + 4y_2$

subject to

$$3y_1 + y_2 \geq 5$$

$$2y_1 + 2y_2 \geq 2$$

$$y_1 \geq 0, y_2 \geq 0$$

- **Example 1:** Consider the problem **(P)**

Max $5x_1 + 2x_2$ subject to

$$3x_1 + 2x_2 \leq 6$$

$$x_1 + 2x_2 \leq 4$$

$$x_1 \geq 0, x_2 \geq 0$$

- The dual of **(P)** is given by **(D)**

Min $6y_1 + 4y_2$

subject to

$$3y_1 + y_2 \geq 5$$

$$2y_1 + 2y_2 \geq 2$$

$$y_1 \geq 0, y_2 \geq 0$$

- $\mathbf{x} = [2, 0]^T$ is a feasible solution of **(P)** and $\mathbf{y} = [\frac{5}{3}, 0]^T$ is a feasible solution of **(D)**

- **Example 1:** Consider the problem **(P)**

Max $5x_1 + 2x_2$ subject to

$$3x_1 + 2x_2 \leq 6$$

$$x_1 + 2x_2 \leq 4$$

$$x_1 \geq 0, x_2 \geq 0$$

- The dual of **(P)** is given by **(D)**

Min $6y_1 + 4y_2$

subject to

$$3y_1 + y_2 \geq 5$$

$$2y_1 + 2y_2 \geq 2$$

$$y_1 \geq 0, y_2 \geq 0$$

- $\mathbf{x} = [2, 0]^T$ is a feasible solution of **(P)** and $\mathbf{y} = [\frac{5}{3}, 0]^T$ is a feasible solution of **(D)**
- $\mathbf{b}^T \mathbf{y} = \mathbf{c}^T \mathbf{x} = 10$, where $\mathbf{c} = [5, 2]^T$ and $\mathbf{b} = [6, 4]^T$

- **Example 1:** Consider the problem **(P)**

Max $5x_1 + 2x_2$ subject to

$$3x_1 + 2x_2 \leq 6$$

$$x_1 + 2x_2 \leq 4$$

$$x_1 \geq 0, x_2 \geq 0$$

- The dual of **(P)** is given by **(D)**

Min $6y_1 + 4y_2$

subject to

$$3y_1 + y_2 \geq 5$$

$$2y_1 + 2y_2 \geq 2$$

$$y_1 \geq 0, y_2 \geq 0$$

- $\mathbf{x} = [2, 0]^T$ is a feasible solution of **(P)** and $\mathbf{y} = [\frac{5}{3}, 0]^T$ is a feasible solution of **(D)**
- $\mathbf{b}^T \mathbf{y} = \mathbf{c}^T \mathbf{x} = 10$, where $\mathbf{c} = [5, 2]^T$ and $\mathbf{b} = [6, 4]^T$
- $\mathbf{x} = [2, 0]^T$ is **optimal** for **Primal (P)** and $\mathbf{y} = [\frac{5}{3}, 0]^T$ is **optimal** for **Dual (D)**

- **Theorem 2: (Fundamental Theorem of Duality (FTD))**

- **Theorem 2: (Fundamental Theorem of Duality (FTD))**
 1. If **both** **Primal (P)** and **Dual (D)** of Primal (P) have **feasible solutions** then **both** have **optimal solutions** and the **optimal value** of (P) and (D) are equal ($\text{Min } \mathbf{b}^T \mathbf{y} = \text{Max } \mathbf{c}^T \mathbf{x}$)

- **Theorem 2: (Fundamental Theorem of Duality (FTD))**
 1. If **both** **Primal (P)** and **Dual (D)** of Primal (P) have **feasible solutions** then **both** have **optimal solutions** and the **optimal value** of (P) and (D) are equal ($\text{Min } \mathbf{b}^T \mathbf{y} = \text{Max } \mathbf{c}^T \mathbf{x}$)
 2. If any one of them (the Primal or the Dual) does not have a **feasible solution** then the other does not have an **optimal solution**

- **Theorem 2: (Fundamental Theorem of Duality (FTD))**
 1. If **both** **Primal (P)** and **Dual (D)** of Primal (P) have **feasible solutions** then **both** have **optimal solutions** and the **optimal value** of (P) and (D) are equal ($\text{Min } \mathbf{b}^T \mathbf{y} = \text{Max } \mathbf{c}^T \mathbf{x}$)
 2. If any one of them (the Primal or the Dual) does not have a **feasible solution** then the other does not have an **optimal solution**
- The **fundamental theorem of duality** implies that either **both** P and D has **optimal solution/s** or **none** has **optimal solution/s**
- To complete the proof of **FTD** it is enough to show:

- **Theorem 2: (Fundamental Theorem of Duality (FTD))**
 1. If **both** **Primal (P)** and **Dual (D)** of Primal (P) have **feasible solutions** then **both** have **optimal solutions** and the **optimal value** of (P) and (D) are equal ($\text{Min } \mathbf{b}^T \mathbf{y} = \text{Max } \mathbf{c}^T \mathbf{x}$)
 2. If any one of them (the Primal or the Dual) does not have a **feasible solution** then the other does not have an **optimal solution**
- The **fundamental theorem of duality** implies that either **both** P and D has **optimal solution/s** or **none** has **optimal solution/s**
- To complete the proof of **FTD** it is enough to show:
 1. If both (P) and (D) have **feasible solutions** then their **optimal values** are equal

- **Theorem 2: (Fundamental Theorem of Duality (FTD))**
 1. If **both** **Primal (P)** and **Dual (D)** of Primal (P) have **feasible solutions** then **both** have **optimal solutions** and the **optimal value** of (P) and (D) are equal ($\text{Min } \mathbf{b}^T \mathbf{y} = \text{Max } \mathbf{c}^T \mathbf{x}$)
 2. If any one of them (the Primal or the Dual) does not have a **feasible solution** then the other does not have an **optimal solution**
- The **fundamental theorem of duality** implies that either **both** P and D has **optimal solution/s** or **none** has **optimal solution/s**
- To complete the proof of **FTD** it is enough to show:
 1. If both (P) and (D) have **feasible solutions** then their **optimal values** are equal
 2. If any one of (P) or (D) **does not** have a **feasible solution** then the other **does not** have an **optimal solution**

- **Example 2:** Consider (P)

- **Example 2:** Consider (P)

$$\text{Min } -x + 2y$$

- **Example 2:** Consider (P)

Min $-x + 2y$

subject to $x + 2y \geq 1$

- **Example 2:** Consider (P)

$$\text{Min } -x + 2y$$

$$\text{subject to } x + 2y \geq 1$$

$$-x + y \leq 1$$

$$x \geq 0, y \geq 0$$

- **Example 2:** Consider (P)

$$\text{Min } -x + 2y$$

$$\text{subject to } x + 2y \geq 1$$

$$-x + y \leq 1$$

$$x \geq 0, y \geq 0$$

- (P) has a feasible solution but **does not** an optimal solution

- **Example 2:** Consider (P)

$$\text{Min } -x + 2y$$

$$\text{subject to } x + 2y \geq 1$$

$$-x + y \leq 1$$

$$x \geq 0, y \geq 0$$

- (P) has a **feasible solution** but **does not** an **optimal solution**
($\mathbf{d} = [1, 0]^T$ is a direction of Fea(P), (P) is a minimization problem and $\mathbf{c}^T \mathbf{d} = -1$)

- **Example 2:** Consider (P)

$$\text{Min } -x + 2y$$

$$\text{subject to } x + 2y \geq 1$$

$$-x + y \leq 1$$

$$x \geq 0, y \geq 0$$

- (P) has a **feasible solution** but **does not** an **optimal solution**
($\mathbf{d} = [1, 0]^T$ is a direction of Fea(P), (P) is a minimization problem and $\mathbf{c}^T \mathbf{d} = -1$)

On converting the second inequality to $x - y \geq -1$

- **Example 2:** Consider (P)

$$\text{Min } -x + 2y$$

$$\text{subject to } x + 2y \geq 1$$

$$-x + y \leq 1$$

$$x \geq 0, y \geq 0$$

- (P) has a **feasible solution** but **does not** an **optimal solution**
($\mathbf{d} = [1, 0]^T$ is a direction of Fea(P), (P) is a minimization problem and $\mathbf{c}^T \mathbf{d} = -1$)

On converting the second inequality to $x - y \geq -1$

- The dual (D) of (P) is given by

$$\text{Max } u - v$$

- **Example 2:** Consider (P)

$$\text{Min } -x + 2y$$

$$\text{subject to } x + 2y \geq 1$$

$$-x + y \leq 1$$

$$x \geq 0, y \geq 0$$

- (P) has a **feasible solution** but **does not** an **optimal solution**
($\mathbf{d} = [1, 0]^T$ is a direction of Fea(P), (P) is a minimization problem and $\mathbf{c}^T \mathbf{d} = -1$)

On converting the second inequality to $x - y \geq -1$

- The dual (D) of (P) is given by

$$\text{Max } u - v$$

$$\text{subject to } u + v \leq -1$$

- **Example 2:** Consider (P)

$$\text{Min } -x + 2y$$

$$\text{subject to } x + 2y \geq 1$$

$$-x + y \leq 1$$

$$x \geq 0, y \geq 0$$

- (P) has a **feasible solution** but **does not** an **optimal solution**
($\mathbf{d} = [1, 0]^T$ is a direction of Fea(P), (P) is a minimization problem and $\mathbf{c}^T \mathbf{d} = -1$)

On converting the second inequality to $x - y \geq -1$

- The dual (D) of (P) is given by

$$\text{Max } u - v$$

$$\text{subject to } u + v \leq -1$$

$$2u - v \leq 2$$

- **Example 2:** Consider (P)

$$\text{Min } -x + 2y$$

$$\text{subject to } x + 2y \geq 1$$

$$-x + y \leq 1$$

$$x \geq 0, y \geq 0$$

- (P) has a **feasible solution** but **does not** an **optimal solution**
($\mathbf{d} = [1, 0]^T$ is a direction of Fea(P), (P) is a minimization problem and $\mathbf{c}^T \mathbf{d} = -1$)

On converting the second inequality to $x - y \geq -1$

- The dual (D) of (P) is given by

$$\text{Max } u - v$$

$$\text{subject to } u + v \leq -1$$

$$2u - v \leq 2$$

$$u \geq 0, v \geq 0$$

- **Example 2:** Consider (P)

$$\text{Min } -x + 2y$$

$$\text{subject to } x + 2y \geq 1$$

$$-x + y \leq 1$$

$$x \geq 0, y \geq 0$$

- (P) has a **feasible solution** but **does not** an **optimal solution**
($\mathbf{d} = [1, 0]^T$ is a direction of Fea(P), (P) is a minimization problem and $\mathbf{c}^T \mathbf{d} = -1$)

On converting the second inequality to $x - y \geq -1$

- The dual (D) of (P) is given by

$$\text{Max } u - v$$

$$\text{subject to } u + v \leq -1$$

$$2u - v \leq 2$$

$$u \geq 0, v \geq 0$$

- Clearly (D) **does not** have any **feasible solution**, since the first constraint cannot be satisfied by any non negative u, v

- **Example 3:** The LPP (P)

Max $-x + 2y$

- **Example 3:** The LPP (P)

Max $-x + 2y$

subject to $x + 2y \leq 1$

● **Example 3:** The LPP (P)

$$\text{Max } -x + 2y$$

$$\text{subject to } x + 2y \leq 1$$

$$-x + y \geq 1$$

$$x, y \geq 0$$

- **Example 3:** The LPP (P)

$$\text{Max } -x + 2y$$

$$\text{subject to } x + 2y \leq 1$$

$$-x + y \geq 1$$

$$x, y \geq 0$$

- (P) **does not** have a feasible solution

- **Example 3:** The LPP (P)

$$\text{Max } -x + 2y$$

$$\text{subject to } x + 2y \leq 1$$

$$-x + y \geq 1$$

$$x, y \geq 0$$

- (P) **does not** have a feasible solution

On converting the second inequality to $x - y \leq -1$

- **Example 3:** The LPP (P)

$$\text{Max } -x + 2y$$

$$\text{subject to } x + 2y \leq 1$$

$$-x + y \geq 1$$

$$x, y \geq 0$$

- (P) **does not** have a feasible solution

On converting the second inequality to $x - y \leq -1$

- The dual of (P) is given by (D)

$$\text{Min } u - v$$

- **Example 3:** The LPP (P)

$$\text{Max } -x + 2y$$

$$\text{subject to } x + 2y \leq 1$$

$$-x + y \geq 1$$

$$x, y \geq 0$$

- (P) **does not** have a feasible solution

On converting the second inequality to $x - y \leq -1$

- The dual of (P) is given by (D)

$$\text{Min } u - v$$

$$\text{subject to } u + v \geq -1$$

- **Example 3:** The LPP (P)

$$\text{Max } -x + 2y$$

$$\text{subject to } x + 2y \leq 1$$

$$-x + y \geq 1$$

$$x, y \geq 0$$

- (P) **does not** have a feasible solution

On converting the second inequality to $x - y \leq -1$

- The dual of (P) is given by (D)

$$\text{Min } u - v$$

$$\text{subject to } u + v \geq -1$$

$$2u - v \geq 2$$

- **Example 3:** The LPP (P)

$$\text{Max } -x + 2y$$

$$\text{subject to } x + 2y \leq 1$$

$$-x + y \geq 1$$

$$x, y \geq 0$$

- (P) **does not** have a feasible solution

On converting the second inequality to $x - y \leq -1$

- The dual of (P) is given by (D)

$$\text{Min } u - v$$

$$\text{subject to } u + v \geq -1$$

$$2u - v \geq 2$$

$$u \geq 0, v \geq 0$$

- **Example 3:** The LPP (P)

$$\text{Max } -x + 2y$$

$$\text{subject to } x + 2y \leq 1$$

$$-x + y \geq 1$$

$$x, y \geq 0$$

- (P) **does not** have a feasible solution

On converting the second inequality to $x - y \leq -1$

- The dual of (P) is given by (D)

$$\text{Min } u - v$$

$$\text{subject to } u + v \geq -1$$

$$2u - v \geq 2$$

$$u \geq 0, v \geq 0$$

(D) has a feasible solution but **no** optimal solution

- **Example 3:** The LPP (P)

$$\text{Max } -x + 2y$$

$$\text{subject to } x + 2y \leq 1$$

$$-x + y \geq 1$$

$$x, y \geq 0$$

- (P) **does not** have a feasible solution

On converting the second inequality to $x - y \leq -1$

- The dual of (P) is given by (D)

$$\text{Min } u - v$$

$$\text{subject to } u + v \geq -1$$

$$2u - v \geq 2$$

$$u \geq 0, v \geq 0$$

(D) has a feasible solution but **no optimal solution**

($\mathbf{d} = [1, 2]^T$ is a direction of Fea(D) and $\mathbf{b}^T \mathbf{d} = -1 < 0$)

- **Definition:** A nonempty set $T \subseteq \mathbb{R}^n$ is said to be a **cone** if $\mathbf{x} \in T$ implies $\lambda \mathbf{x} \in T$ for all $\lambda \geq 0$

- **Definition:** A nonempty set $T \subseteq \mathbb{R}^n$ is said to be a **cone** if $\mathbf{x} \in T$ implies $\lambda \mathbf{x} \in T$ for all $\lambda \geq 0$
- So a cone always contains the **origin**

- **Definition:** A nonempty set $T \subseteq \mathbb{R}^n$ is said to be a **cone** if $\mathbf{x} \in T$ implies $\lambda \mathbf{x} \in T$ for all $\lambda \geq 0$
- So a cone always contains the **origin**
- Any straight line passing through the origin is a cone

- **Definition:** A nonempty set $T \subseteq \mathbb{R}^n$ is said to be a **cone** if $\mathbf{x} \in T$ implies $\lambda \mathbf{x} \in T$ for all $\lambda \geq 0$
- So a cone always contains the **origin**
- Any straight line passing through the origin is a cone
Union of a finite number of straight lines passing through the origin is also a cone

- **Definition:** A nonempty set $T \subseteq \mathbb{R}^n$ is said to be a **cone** if $\mathbf{x} \in T$ implies $\lambda \mathbf{x} \in T$ for all $\lambda \geq 0$
- So a cone always contains the **origin**
- Any straight line passing through the origin is a cone
Union of a finite number of straight lines passing through the origin is also a cone
- A **cone** T is said to be a **convex cone** if it is also a **convex subset** of $\mathbb{R}^n$.

- **Definition:** A nonempty set $T \subseteq \mathbb{R}^n$ is said to be a **cone** if $\mathbf{x} \in T$ implies $\lambda \mathbf{x} \in T$ for all $\lambda \geq 0$
- So a cone always contains the **origin**
- Any straight line passing through the origin is a cone
Union of a finite number of straight lines passing through the origin is also a cone
- A **cone** T is said to be a **convex cone** if it is also a **convex subset** of $\mathbb{R}^n$.
- **Exercise:** If $\mathbf{a}_1, \mathbf{a}_2, \dots, \mathbf{a}_k$ are all vectors in $\mathbb{R}^n$ then check that

- **Definition:** A nonempty set $T \subseteq \mathbb{R}^n$ is said to be a **cone** if $\mathbf{x} \in T$ implies $\lambda \mathbf{x} \in T$ for all $\lambda \geq 0$
- So a cone always contains the **origin**
- Any straight line passing through the origin is a cone
Union of a finite number of straight lines passing through the origin is also a cone
- A **cone** T is said to be a **convex cone** if it is also a **convex subset** of $\mathbb{R}^n$.
- **Exercise:** If $\mathbf{a}_1, \mathbf{a}_2, \dots, \mathbf{a}_k$ are all vectors in $\mathbb{R}^n$ then check that

$$T = \{\alpha_1 \mathbf{a}_1 + \alpha_2 \mathbf{a}_2 + \dots + \alpha_k \mathbf{a}_k : \alpha_i \geq 0 \text{ for all } i = 1, \dots, k\}$$
is a **convex cone**

- **Definition:** A nonempty set $T \subseteq \mathbb{R}^n$ is said to be a **cone** if $\mathbf{x} \in T$ implies $\lambda \mathbf{x} \in T$ for all $\lambda \geq 0$
- So a cone always contains the **origin**
- Any straight line passing through the origin is a cone
Union of a finite number of straight lines passing through the origin is also a cone
- A **cone** T is said to be a **convex cone** if it is also a **convex subset** of $\mathbb{R}^n$.
- **Exercise:** If $\mathbf{a}_1, \mathbf{a}_2, \dots, \mathbf{a}_k$ are all vectors in $\mathbb{R}^n$ then check that

$$T = \{\alpha_1 \mathbf{a}_1 + \alpha_2 \mathbf{a}_2 + \dots + \alpha_k \mathbf{a}_k : \alpha_i \geq 0 \text{ for all } i = 1, \dots, k\}$$
is a **convex cone**
 $\mathbf{a}_1, \mathbf{a}_2, \dots, \mathbf{a}_k$ are called **generators** of the **convex cone** T

Fundamental Theorem of Duality (revisited):

- **Theorem 4:** (**Farka's Lemma**) Exactly **one** of the following two systems has a solution

$$A_{m \times n} \mathbf{x} = \mathbf{b}, \mathbf{x} \geq \mathbf{0} \quad (1)$$

Fundamental Theorem of Duality (revisited):

- **Theorem 4: (Farka's Lemma)** Exactly **one** of the following two systems has a solution

$$A_{m \times n} \mathbf{x} = \mathbf{b}, \mathbf{x} \geq \mathbf{0} \quad (1)$$

$$\mathbf{y}^T A \geq \mathbf{0}_{1 \times n}, \mathbf{y}^T \mathbf{b} < 0 \quad (2)$$

Fundamental Theorem of Duality (revisited):

- **Theorem 4:** (**Farka's Lemma**) Exactly **one** of the following two systems has a solution

$$A_{m \times n} \mathbf{x} = \mathbf{b}, \mathbf{x} \geq \mathbf{0} \quad (1)$$

$$\mathbf{y}^T A \geq \mathbf{0}_{1 \times n}, \mathbf{y}^T \mathbf{b} < 0 \quad (2)$$

- **Corollary 4:** Exactly **one** of the following two systems has a solution

Fundamental Theorem of Duality (revisited):

- **Theorem 4:** (**Farka's Lemma**) Exactly **one** of the following two systems has a solution

$$A_{m \times n} \mathbf{x} = \mathbf{b}, \mathbf{x} \geq \mathbf{0} \quad (1)$$

$$\mathbf{y}^T A \geq \mathbf{0}_{1 \times n}, \mathbf{y}^T \mathbf{b} < 0 \quad (2)$$

- **Corollary 4:** Exactly **one** of the following two systems has a solution

$$A_{m \times n} \mathbf{x} \leq \mathbf{b}, \mathbf{x} \geq \mathbf{0} \quad (1)$$

Fundamental Theorem of Duality (revisited):

- **Theorem 4:** (**Farka's Lemma**) Exactly **one** of the following two systems has a solution

$$A_{m \times n} \mathbf{x} = \mathbf{b}, \mathbf{x} \geq \mathbf{0} \quad (1)$$

$$\mathbf{y}^T A \geq \mathbf{0}_{1 \times n}, \quad \mathbf{y}^T \mathbf{b} < 0 \quad (2)$$

- **Corollary 4:** Exactly **one** of the following two systems has a solution

$$A_{m \times n} \mathbf{x} \leq \mathbf{b}, \mathbf{x} \geq \mathbf{0} \quad (1)$$

$$\mathbf{y}^T A \geq \mathbf{0}, \quad \mathbf{y}^T \mathbf{b} < 0, \quad \mathbf{y} \geq \mathbf{0} \quad (2)$$

- Since $Fea(D) = \{\mathbf{y} \in \mathbb{R}^m : A^T \mathbf{y} \geq \mathbf{c}, \mathbf{y} \geq \mathbf{0}\}$

- Since $Fea(D) = \{\mathbf{y} \in \mathbb{R}^m : A^T \mathbf{y} \geq \mathbf{c}, \mathbf{y} \geq \mathbf{0}\}$
the set of all directions of $Fea(D)$ is given by

- Since $Fea(D) = \{\mathbf{y} \in \mathbb{R}^m : A^T \mathbf{y} \geq \mathbf{c}, \mathbf{y} \geq \mathbf{0}\}$
the set of all directions of $Fea(D)$ is given by
- $D_D = \{\mathbf{d} \in \mathbb{R}^m : A^T \mathbf{d} \geq \mathbf{0}, \mathbf{d} \geq \mathbf{0}\}$

- Since $Fea(D) = \{\mathbf{y} \in \mathbb{R}^m : A^T \mathbf{y} \geq \mathbf{c}, \mathbf{y} \geq \mathbf{0}\}$
the set of all directions of $Fea(D)$ is given by
- $D_D = \{\mathbf{d} \in \mathbb{R}^m : A^T \mathbf{d} \geq \mathbf{0}, \mathbf{d} \geq \mathbf{0}\}$
- The **Dual** (D) has an optimal solution if and only if either

- Since $Fea(D) = \{\mathbf{y} \in \mathbb{R}^m : A^T \mathbf{y} \geq \mathbf{c}, \mathbf{y} \geq \mathbf{0}\}$
the set of all directions of $Fea(D)$ is given by
- $D_D = \{\mathbf{d} \in \mathbb{R}^m : A^T \mathbf{d} \geq \mathbf{0}, \mathbf{d} \geq \mathbf{0}\}$
- The **Dual** (D) has an optimal solution if and only if either
(i) $Fea(D)$ is **bounded**, or

- Since $Fea(D) = \{\mathbf{y} \in \mathbb{R}^m : A^T \mathbf{y} \geq \mathbf{c}, \mathbf{y} \geq \mathbf{0}\}$
the set of all directions of $Fea(D)$ is given by
- $D_D = \{\mathbf{d} \in \mathbb{R}^m : A^T \mathbf{d} \geq \mathbf{0}, \mathbf{d} \geq \mathbf{0}\}$
- The **Dual** (D) has an optimal solution if and only if either
 - $Fea(D)$ is **bounded**, or
 - $Fea(D)$ is **unbounded** and $\mathbf{b}^T \mathbf{d} \geq 0$ for all $\mathbf{d} \in D_D$.

- Since $Fea(D) = \{\mathbf{y} \in \mathbb{R}^m : A^T \mathbf{y} \geq \mathbf{c}, \mathbf{y} \geq \mathbf{0}\}$
the set of all directions of $Fea(D)$ is given by
- $D_D = \{\mathbf{d} \in \mathbb{R}^m : A^T \mathbf{d} \geq \mathbf{0}, \mathbf{d} \geq \mathbf{0}\}$
- The **Dual** (D) has an optimal solution if and only if either
 - (i) $Fea(D)$ is **bounded**, or
 - (ii) $Fea(D)$ is **unbounded** and $\mathbf{b}^T \mathbf{d} \geq 0$ for all $\mathbf{d} \in D_D$.
- **Corollary 4** implies that if the **Primal** does **not** have a feasible solution then the **Dual** does not have an optimal solution

- To complete the proof of Fundamental Theorem of Duality we need to show that if $\text{Fea}(P) \neq \phi$ and $\text{Fea}(D) \neq \phi$, then the following system **(1)** has a solution

- To complete the proof of Fundamental Theorem of Duality we need to show that if $\text{Fea}(P) \neq \phi$ and $\text{Fea}(D) \neq \phi$, then the following system **(1)** has a solution

$$A_{m \times n} \mathbf{x} \leq \mathbf{b}, \mathbf{x} \geq \mathbf{0} \quad (1)$$

- To complete the proof of Fundamental Theorem of Duality we need to show that if $\text{Fea}(P) \neq \phi$ and $\text{Fea}(D) \neq \phi$, then the following system **(1)** has a solution

$$A_{m \times n} \mathbf{x} \leq \mathbf{b}, \mathbf{x} \geq \mathbf{0} \quad (1)$$

$$A^T \mathbf{y} \geq \mathbf{c}, \mathbf{y} \geq \mathbf{0}$$

- To complete the proof of Fundamental Theorem of Duality we need to show that if $\text{Fea}(P) \neq \phi$ and $\text{Fea}(D) \neq \phi$, then the following system **(1)** has a solution

$$A_{m \times n} \mathbf{x} \leq \mathbf{b}, \mathbf{x} \geq \mathbf{0} \quad (1)$$

$$A^T \mathbf{y} \geq \mathbf{c}, \mathbf{y} \geq \mathbf{0}$$

$$\mathbf{c}^T \mathbf{x} = \mathbf{b}^T \mathbf{y}$$

- To complete the proof of Fundamental Theorem of Duality we need to show that if $Fea(P) \neq \phi$ and $Fea(D) \neq \phi$, then the following system (1) has a solution

$$A_{m \times n} \mathbf{x} \leq \mathbf{b}, \mathbf{x} \geq \mathbf{0} \quad (1)$$

$$A^T \mathbf{y} \geq \mathbf{c}, \mathbf{y} \geq \mathbf{0}$$

$$\mathbf{c}^T \mathbf{x} = \mathbf{b}^T \mathbf{y}$$

- If $\mathbf{x} \in Fea(P)$ and $\mathbf{y} \in Fea(D)$ then $\mathbf{c}^T \mathbf{x} \leq \mathbf{b}^T \mathbf{y}$, hence system (1) has a solution $\Leftrightarrow$ system (1)'' has a solution

$$A_{m \times n} \mathbf{x} \leq \mathbf{b}, \mathbf{x} \geq \mathbf{0} \quad (1)''$$

$$A^T \mathbf{y} \geq \mathbf{c}, \mathbf{y} \geq \mathbf{0}$$

$$\mathbf{c}^T \mathbf{x} \geq \mathbf{b}^T \mathbf{y}$$

- System (1)'' can be written as

$$\begin{bmatrix} A_{m \times n} & \mathbf{0}_{m \times m} \\ \mathbf{0}_{n \times n} & -A_{n \times m}^T \\ -\mathbf{c}_{1 \times n}^T & \mathbf{b}_{1 \times m}^T \end{bmatrix} \begin{bmatrix} \mathbf{x} \\ \mathbf{y} \end{bmatrix} \leq \begin{bmatrix} \mathbf{b} \\ -\mathbf{c} \\ 0 \end{bmatrix}, \quad \begin{bmatrix} \mathbf{x} \\ \mathbf{y} \end{bmatrix} \geq \begin{bmatrix} \mathbf{0}_{n \times 1} \\ \mathbf{0}_{m \times 1} \end{bmatrix}$$

- System (1)'' can be written as

$$\begin{bmatrix} A_{m \times n} & \mathbf{0}_{m \times m} \\ \mathbf{0}_{n \times n} & -A_{n \times m}^T \\ -\mathbf{c}_{1 \times n}^T & \mathbf{b}_{1 \times m}^T \end{bmatrix} \begin{bmatrix} \mathbf{x} \\ \mathbf{y} \end{bmatrix} \leq \begin{bmatrix} \mathbf{b} \\ -\mathbf{c} \\ 0 \end{bmatrix}, \quad \begin{bmatrix} \mathbf{x} \\ \mathbf{y} \end{bmatrix} \geq \begin{bmatrix} \mathbf{0}_{n \times 1} \\ \mathbf{0}_{m \times 1} \end{bmatrix}$$

- By **corollary 4**, exactly one of the two systems, (1'') (given above) and (2'') (given below) has a solution.

$$\begin{bmatrix} \mathbf{z}_{1 \times m}^T & \mathbf{w}_{1 \times n}^T & a_{1 \times 1} \end{bmatrix} \begin{bmatrix} A_{m \times n} & \mathbf{0}_{m \times m} \\ \mathbf{0}_{n \times n} & -A_{n \times m}^T \\ -\mathbf{c}_{1 \times n}^T & \mathbf{b}_{1 \times m}^T \end{bmatrix} \geq \begin{bmatrix} \mathbf{0}_{1 \times n} & \mathbf{0}_{1 \times m} \end{bmatrix},$$

$$\begin{bmatrix} \mathbf{z}_{1 \times m}^T & \mathbf{w}_{1 \times n}^T & a_{1 \times 1} \end{bmatrix} \begin{bmatrix} \mathbf{b} \\ -\mathbf{c} \\ 0 \end{bmatrix} < 0 \quad \begin{bmatrix} \mathbf{z}_{1 \times m}^T & \mathbf{w}_{1 \times n}^T & a_{1 \times 1} \end{bmatrix} \geq \mathbf{0}_{1 \times (n+m)}$$

- We show that given $Fea(P) \neq \phi$ and $Fea(D) \neq \phi$, system (2)'' does not have a solution, which implies system (1)'' has solution

- We show that given $Fea(P) \neq \phi$ and $Fea(D) \neq \phi$, system (2)'' does not have a solution, which implies system (1)'' has solution
- **Theorem 5 (Complementary Slackness Theorem):** Let $\mathbf{x} \in Fea(P)$ and $\mathbf{y} \in Fea(D)$, then $\mathbf{x}$ and $\mathbf{y}$ are optimal for the primal and the dual respectively if and only if

- We show that given $Fea(P) \neq \phi$ and $Fea(D) \neq \phi$, system (2)'' does not have a solution, which implies system (1)'' has solution
- **Theorem 5 (Complementary Slackness Theorem):** Let $\mathbf{x} \in Fea(P)$ and $\mathbf{y} \in Fea(D)$, then $\mathbf{x}$ and $\mathbf{y}$ are optimal for the primal and the dual respectively if and only if $x_j = 0$ whenever $(A^T \mathbf{y})_j > c_j, j = 1, 2, \dots, n$ (1) and

- We show that given $Fea(P) \neq \phi$ and $Fea(D) \neq \phi$, system (2)'' does not have a solution, which implies system (1)'' has solution
- **Theorem 5 (Complementary Slackness Theorem):** Let $\mathbf{x} \in Fea(P)$ and $\mathbf{y} \in Fea(D)$, then $\mathbf{x}$ and $\mathbf{y}$ are optimal for the primal and the dual respectively if and only if
 $x_j = 0$ whenever $(A^T \mathbf{y})_j > c_j, j = 1, 2, \dots, n$ (1)
 and
 $y_i = 0$ whenever $(A\mathbf{x})_i < b_i, i = 1, 2, \dots, m.$ (1*)

- **Example 1:** Consider the following primal problem (P):

- **Example 1:** Consider the following primal problem (P):

$$\text{Max } 4x_1 + 4x_2 + 2x_3$$

subject to

$$2x_1 + 3x_2 + 4x_3 \leq 10 \quad (i)$$

- **Example 1:** Consider the following primal problem (P):

$$\text{Max } 4x_1 + 4x_2 + 2x_3$$

subject to

$$2x_1 + 3x_2 + 4x_3 \leq 10 \quad (i)$$

$$2x_1 + x_2 + 3x_3 \leq 4 \quad (ii)$$

- **Example 1:** Consider the following primal problem (P):

$$\text{Max } 4x_1 + 4x_2 + 2x_3$$

subject to

$$2x_1 + 3x_2 + 4x_3 \leq 10 \quad (i)$$

$$2x_1 + x_2 + 3x_3 \leq 4 \quad (ii)$$

$$x_1, x_2, x_3 \geq 0$$

- **Example 1:** Consider the following primal problem (P):

$$\text{Max } 4x_1 + 4x_2 + 2x_3$$

subject to

$$2x_1 + 3x_2 + 4x_3 \leq 10 \quad (i)$$

$$2x_1 + x_2 + 3x_3 \leq 4 \quad (ii)$$

$$x_1, x_2, x_3 \geq 0$$

The dual (D) of the above problem is given by:

- **Example 1:** Consider the following primal problem (P):

$$\text{Max } 4x_1 + 4x_2 + 2x_3$$

subject to

$$2x_1 + 3x_2 + 4x_3 \leq 10 \quad (i)$$

$$2x_1 + x_2 + 3x_3 \leq 4 \quad (ii)$$

$$x_1, x_2, x_3 \geq 0$$

The dual (D) of the above problem is given by:

$$\text{Min } 10y_1 + 4y_2$$

- **Example 1:** Consider the following primal problem (P):

$$\text{Max } 4x_1 + 4x_2 + 2x_3$$

subject to

$$2x_1 + 3x_2 + 4x_3 \leq 10 \quad (i)$$

$$2x_1 + x_2 + 3x_3 \leq 4 \quad (ii)$$

$$x_1, x_2, x_3 \geq 0$$

The dual (D) of the above problem is given by:

$$\text{Min } 10y_1 + 4y_2$$

subject to

$$2y_1 + 2y_2 \geq 4 \quad (i)'$$

- **Example 1:** Consider the following primal problem (P):

$$\text{Max } 4x_1 + 4x_2 + 2x_3$$

subject to

$$2x_1 + 3x_2 + 4x_3 \leq 10 \quad (i)$$

$$2x_1 + x_2 + 3x_3 \leq 4 \quad (ii)$$

$$x_1, x_2, x_3 \geq 0$$

The dual (D) of the above problem is given by:

$$\text{Min } 10y_1 + 4y_2$$

subject to

$$2y_1 + 2y_2 \geq 4 \quad (i)'$$

$$3y_1 + y_2 \geq 4 \quad (ii)'$$

- **Example 1:** Consider the following primal problem (P):

$$\text{Max } 4x_1 + 4x_2 + 2x_3$$

subject to

$$2x_1 + 3x_2 + 4x_3 \leq 10 \quad (i)$$

$$2x_1 + x_2 + 3x_3 \leq 4 \quad (ii)$$

$$x_1, x_2, x_3 \geq 0$$

The dual (D) of the above problem is given by:

$$\text{Min } 10y_1 + 4y_2$$

subject to

$$2y_1 + 2y_2 \geq 4 \quad (i)'$$

$$3y_1 + y_2 \geq 4 \quad (ii)'$$

$$4y_1 + 3y_2 \geq 2 \quad (iii)'$$

- **Example 1:** Consider the following primal problem (P):

$$\text{Max } 4x_1 + 4x_2 + 2x_3$$

subject to

$$2x_1 + 3x_2 + 4x_3 \leq 10 \quad (i)$$

$$2x_1 + x_2 + 3x_3 \leq 4 \quad (ii)$$

$$x_1, x_2, x_3 \geq 0$$

The dual (D) of the above problem is given by:

$$\text{Min } 10y_1 + 4y_2$$

subject to

$$2y_1 + 2y_2 \geq 4 \quad (i)'$$

$$3y_1 + y_2 \geq 4 \quad (ii)'$$

$$4y_1 + 3y_2 \geq 2 \quad (iii)'$$

$$y_1, y_2 \geq 0$$

- **Example 1:** Consider the following primal problem (P):

$$\text{Max } 4x_1 + 4x_2 + 2x_3$$

subject to

$$2x_1 + 3x_2 + 4x_3 \leq 10 \quad (i)$$

$$2x_1 + x_2 + 3x_3 \leq 4 \quad (ii)$$

$$x_1, x_2, x_3 \geq 0$$

The dual (D) of the above problem is given by:

$$\text{Min } 10y_1 + 4y_2$$

subject to

$$2y_1 + 2y_2 \geq 4 \quad (i)'$$

$$3y_1 + y_2 \geq 4 \quad (ii)'$$

$$4y_1 + 3y_2 \geq 2 \quad (iii)'$$

$$y_1, y_2 \geq 0$$

- Check that $[\frac{1}{2}, 3, 0]^T \in \text{Opt}(P)$ and $[1, 1]^T \in \text{Opt}(D)$