

Introduction of a new variable

- Consider the problem (**P**),
Min $\mathbf{c}^T \mathbf{x}$
subject to
 $A\mathbf{x} = \mathbf{b}, \mathbf{x} \geq \mathbf{0}$.
Let $\mathbf{x}_0$ be an optimal BFS of this problem.
- A new variable x_{n+1} is added to the LPP given above.

Introduction of a new variable

- Consider the problem (P),

$$\text{Min } \mathbf{c}^T \mathbf{x}$$

subject to

$$\mathbf{Ax} = \mathbf{b}, \mathbf{x} \geq \mathbf{0}.$$

Let $\mathbf{x}_0$ be an optimal BFS of this problem.

- A new variable x_{n+1} is added to the LPP given above.
- A column is added to the matrix A call it $\tilde{\mathbf{a}}_{n+1}$ and a component c_{n+1} is added to the cost vector $\mathbf{c}$, which is the cost associated with the new variable x_{n+1} .

Introduction of a new variable

- Consider the problem (**P**),

$$\text{Min } \mathbf{c}^T \mathbf{x}$$

subject to

$$\mathbf{Ax} = \mathbf{b}, \mathbf{x} \geq \mathbf{0}.$$

Let $\mathbf{x}_0$ be an **optimal BFS** of this problem.

- A new variable x_{n+1} is added to the LPP given above.
- A column is added to the matrix A call it $\tilde{\mathbf{a}}_{n+1}$ and a component c_{n+1} is added to the cost vector $\mathbf{c}$, which is the cost associated with the new variable x_{n+1} .
- Since $\mathbf{x}_0$ is optimal for **P**
 $\mathbf{x}'_0 = [\mathbf{x}_0^T, 0]^T$ is feasible for the new problem (P'), given by:

Introduction of a new variable

- Consider the problem (**P**),

$$\text{Min } \mathbf{c}^T \mathbf{x}$$

subject to

$$\mathbf{Ax} = \mathbf{b}, \mathbf{x} \geq \mathbf{0}.$$

Let $\mathbf{x}_0$ be an **optimal BFS** of this problem.

- A new variable x_{n+1} is added to the LPP given above.
- A column is added to the matrix A call it $\tilde{\mathbf{a}}_{n+1}$ and a component c_{n+1} is added to the cost vector $\mathbf{c}$, which is the cost associated with the new variable x_{n+1} .
- Since $\mathbf{x}_0$ is optimal for **P**
 $\mathbf{x}'_0 = [\mathbf{x}_0^T, 0]^T$ is feasible for the new problem (P'), given by:
- Min $[\mathbf{c}, c_{n+1}]^T \mathbf{x}_{(n+1) \times 1}$
subject to
 $[\mathbf{A} : \tilde{\mathbf{a}}_{n+1}] \mathbf{x}_{(n+1) \times 1} = \mathbf{b}, \quad \mathbf{x}_{(n+1) \times 1} \geq \mathbf{0}.$

- To check whether $\mathbf{x}'_0$ is optimal for (P') , add a new column to the optimal table and calculate $C_{n+1} - Z_{n+1}$.

- To check whether $\mathbf{x}'_0$ is optimal for (P'), add a new column to the optimal table and calculate $c_{n+1} - z_{n+1}$.
- If $c_{n+1} - z_{n+1} \geq 0$, then $\mathbf{x}'_0$ is optimal for the new problem.

- To check whether $\mathbf{x}'_0$ is optimal for (P'), add a new column to the optimal table and calculate $c_{n+1} - z_{n+1}$.
- If $c_{n+1} - z_{n+1} \geq 0$, then $\mathbf{x}'_0$ is optimal for the new problem.
- If **not**, then enter $\tilde{\mathbf{a}}_{n+1}$ in the basis and use the **simplex** to either obtain an optimal solution of (P') or to conclude that (P') **does not** have an optimal solution.

- To check whether $\mathbf{x}'_0$ is optimal for (P'), add a new column to the optimal table and calculate $c_{n+1} - z_{n+1}$.
- If $c_{n+1} - z_{n+1} \geq 0$, then $\mathbf{x}'_0$ is optimal for the new problem.
- If **not**, then enter $\tilde{\mathbf{a}}_{n+1}$ in the basis and use the **simplex** to either obtain an optimal solution of (P') or to conclude that (P') **does not** have an optimal solution.
- Adding a **variable** to **P** results in adding a **constraint** to the dual,

- To check whether $\mathbf{x}'_0$ is optimal for (P'), add a new column to the optimal table and calculate $c_{n+1} - z_{n+1}$.
- If $c_{n+1} - z_{n+1} \geq 0$, then $\mathbf{x}'_0$ is optimal for the new problem.
- If **not**, then enter $\tilde{\mathbf{a}}_{n+1}$ in the basis and use the **simplex** to either obtain an optimal solution of (P') or to conclude that (P') **does not** have an optimal solution.
- Adding a **variable** to **P** results in adding a **constraint** to the dual, If the feasible region of the new dual becomes empty,

- To check whether $\mathbf{x}'_0$ is optimal for (P'), add a new column to the optimal table and calculate $c_{n+1} - z_{n+1}$.
- If $c_{n+1} - z_{n+1} \geq 0$, then $\mathbf{x}'_0$ is optimal for the new problem.
- If **not**, then enter $\tilde{a}_{n+1}$ in the basis and use the **simplex** to either obtain an optimal solution of (P') or to conclude that (P') **does not** have an optimal solution.
- Adding a **variable** to **P** results in adding a **constraint** to the dual, If the feasible region of the new dual becomes empty, then **P'** will **not** have an optimal solution.

Introduction of a new constraint

- Addition of a new **constraint** makes the feasible region of **P** smaller.

Introduction of a new constraint

- Addition of a new **constraint** makes the feasible region of **P** smaller.
- Let us assume that the new constraint added is of the form $\mathbf{a}_{m+1}^T \mathbf{x} \leq b_{m+1}$, where $\mathbf{a}_{m+1}^T$ is a row vector.

Introduction of a new constraint

- Addition of a new **constraint** makes the feasible region of **P** smaller.
- Let us assume that the new constraint added is of the form $\mathbf{a}_{m+1}^T \mathbf{x} \leq b_{m+1}$, where $\mathbf{a}_{m+1}^T$ is a row vector.
- **Case 1:** $\mathbf{x}_0$ satisfies $\mathbf{a}_{m+1}^T \mathbf{x} \leq b_{m+1}$.

Introduction of a new constraint

- Addition of a new **constraint** makes the feasible region of **P** smaller.
- Let us assume that the new constraint added is of the form $\mathbf{a}_{m+1}^T \mathbf{x} \leq b_{m+1}$, where $\mathbf{a}_{m+1}^T$ is a row vector.
- **Case 1:** $\mathbf{x}_0$ satisfies $\mathbf{a}_{m+1}^T \mathbf{x} \leq b_{m+1}$.
Then $\mathbf{x}_0$ is optimal for the changed problem also.

Introduction of a new constraint

- Addition of a new **constraint** makes the feasible region of **P** smaller.
- Let us assume that the new constraint added is of the form $\mathbf{a}_{m+1}^T \mathbf{x} \leq b_{m+1}$, where $\mathbf{a}_{m+1}^T$ is a row vector.
- **Case 1:** $\mathbf{x}_0$ satisfies $\mathbf{a}_{m+1}^T \mathbf{x} \leq b_{m+1}$.
Then $\mathbf{x}_0$ is optimal for the changed problem also.
- **Case 2:** $\mathbf{x}_0$ does not satisfy the new constraint, that is $\mathbf{a}_{m+1}^T \mathbf{x}_0 > b_{m+1}$,

Introduction of a new constraint

- Addition of a new **constraint** makes the feasible region of **P** smaller.
- Let us assume that the new constraint added is of the form $\mathbf{a}_{m+1}^T \mathbf{x} \leq b_{m+1}$, where $\mathbf{a}_{m+1}^T$ is a row vector.
- **Case 1:** $\mathbf{x}_0$ satisfies $\mathbf{a}_{m+1}^T \mathbf{x} \leq b_{m+1}$.
Then $\mathbf{x}_0$ is optimal for the changed problem also.
- **Case 2:** $\mathbf{x}_0$ does not satisfy the new constraint, that is $\mathbf{a}_{m+1}^T \mathbf{x}_0 > b_{m+1}$,
or $\mathbf{a}_{B,m+1}^T B^{-1} \mathbf{b} > b_{m+1}$,

Introduction of a new constraint

- Addition of a new **constraint** makes the feasible region of **P** smaller.
- Let us assume that the new constraint added is of the form $\mathbf{a}_{m+1}^T \mathbf{x} \leq b_{m+1}$, where $\mathbf{a}_{m+1}^T$ is a row vector.
- **Case 1:** $\mathbf{x}_0$ satisfies $\mathbf{a}_{m+1}^T \mathbf{x} \leq b_{m+1}$.
Then $\mathbf{x}_0$ is optimal for the changed problem also.
- **Case 2:** $\mathbf{x}_0$ does not satisfy the new constraint, that is $\mathbf{a}_{m+1}^T \mathbf{x}_0 > b_{m+1}$,
or $\mathbf{a}_{B,m+1}^T B^{-1} \mathbf{b} > b_{m+1}$,
where $\mathbf{a}_{B,m+1}^T$ and $\mathbf{a}_{N,m+1}^T$ are the components of $\mathbf{a}_{m+1}^T$ corresponding to the basic variables and nonbasic variables of $\mathbf{x}_0$, respectively.

- The new constraint can be written as $\mathbf{a}_{m+1}^T \mathbf{x} + s_{m+1} = b_{m+1}$, where $s_{m+1} \geq 0$.

- The new constraint can be written as $\mathbf{a}_{m+1}^T \mathbf{x} + s_{m+1} = b_{m+1}$, where $s_{m+1} \geq 0$.
- If B denotes the basis matrix corresponding to $\mathbf{x}_0$, then

- The new constraint can be written as $\mathbf{a}_{m+1}^T \mathbf{x} + s_{m+1} = b_{m+1}$, where $s_{m+1} \geq 0$.
- If B denotes the basis matrix corresponding to $\mathbf{x}_0$, then
$$\begin{bmatrix} B & \mathbf{0} \\ \mathbf{a}_{B,m+1}^T & 1 \end{bmatrix}$$
 is the new basis matrix after the new constraint is added, the newly added **basic variable** being s_{m+1} .

- The new constraint can be written as

$$\mathbf{a}_{m+1}^T \mathbf{x} + s_{m+1} = b_{m+1}, \text{ where } s_{m+1} \geq 0.$$

- If B denotes the basis matrix corresponding to $\mathbf{x}_0$, then

$$\begin{bmatrix} B & \mathbf{0} \\ \mathbf{a}_{B,m+1}^T & 1 \end{bmatrix}$$

is the new basis matrix after the new constraint is added, the newly added **basic variable** being s_{m+1} .

- The **inverse** of this new basis matrix is given by

$$\begin{bmatrix} B^{-1} & \mathbf{0} \\ -\mathbf{a}_{B,m+1}^T B^{-1} & 1 \end{bmatrix}.$$

- The new row added to the (main) simplex table will be of the form

$$\begin{aligned} & [-\mathbf{a}_{B,m+1}^T B^{-1}, 1] \begin{bmatrix} \mathbf{a}_{B,m+1}^T & \mathbf{a}_{N,m+1}^T & \mathbf{0} \\ 1 & & \end{bmatrix} = \\ & [\mathbf{0}, -\mathbf{a}_{B,m+1}^T B^{-1} N + \mathbf{a}_{N,m+1}^T, 1] \end{aligned}$$

- The new row added to the (main) simplex table will be of the form

$$\begin{bmatrix} -\mathbf{a}_{B,m+1}^T B^{-1}, 1 \end{bmatrix} \begin{bmatrix} \mathbf{a}_{B,m+1}^T & \mathbf{a}_{N,m+1}^T & 0 \\ \mathbf{0}, -\mathbf{a}_{B,m+1}^T B^{-1} N + \mathbf{a}_{N,m+1}^T, 1 \end{bmatrix} =$$

- The new RHS becomes

$$\begin{bmatrix} B^{-1} & 0 \\ -\mathbf{a}_{B,m+1}^T B^{-1} & 1 \end{bmatrix} \begin{bmatrix} \mathbf{b} \\ b_{m+1} \end{bmatrix} = \begin{bmatrix} B^{-1} \mathbf{b} \\ b_{m+1} - \mathbf{a}_{B,m+1}^T B^{-1} \mathbf{b} \end{bmatrix}$$

- The new row added to the (main) simplex table will be of the form

$$\begin{bmatrix} -\mathbf{a}_{B,m+1}^T B^{-1}, 1 \end{bmatrix} \begin{bmatrix} \mathbf{a}_{B,m+1}^T & \mathbf{a}_{N,m+1}^T & \mathbf{0} \\ \mathbf{0}, -\mathbf{a}_{B,m+1}^T B^{-1} N + \mathbf{a}_{N,m+1}^T, 1 \end{bmatrix} =$$

- The new RHS becomes

$$\begin{bmatrix} B^{-1} & \mathbf{0} \\ -\mathbf{a}_{B,m+1}^T B^{-1} & 1 \end{bmatrix} \begin{bmatrix} \mathbf{b} \\ b_{m+1} \end{bmatrix} = \begin{bmatrix} B^{-1} \mathbf{b} \\ b_{m+1} - \mathbf{a}_{B,m+1}^T B^{-1} \mathbf{b} \end{bmatrix}$$

- If $\mathbf{x}_0$ does not satisfy the newly added constraint then the $(m+1)$ th entry of the RHS will be **strictly** less than zero.

- The new row added to the (main) simplex table will be of the form

$$\begin{bmatrix} -\mathbf{a}_{B,m+1}^T B^{-1}, 1 \end{bmatrix} \begin{bmatrix} \mathbf{a}_{B,m+1}^T & \mathbf{a}_{N,m+1}^T & \mathbf{0} \\ \mathbf{0}, -\mathbf{a}_{B,m+1}^T B^{-1} N + \mathbf{a}_{N,m+1}^T, 1 \end{bmatrix} =$$

- The new RHS becomes

$$\begin{bmatrix} B^{-1} & \mathbf{0} \\ -\mathbf{a}_{B,m+1}^T B^{-1} & 1 \end{bmatrix} \begin{bmatrix} \mathbf{b} \\ b_{m+1} \end{bmatrix} = \begin{bmatrix} B^{-1} \mathbf{b} \\ b_{m+1} - \mathbf{a}_{B,m+1}^T B^{-1} \mathbf{b} \end{bmatrix}$$

- If $\mathbf{x}_0$ does not satisfy the newly added constraint then the $(m+1)$ th entry of the RHS will be **strictly** less than zero.
- Since the cost associated with s_{m+1} is equal to zero, all the $c_j - z_j$ entries remain unchanged and are non negative.

- The new row added to the (main) simplex table will be of the form

$$\begin{bmatrix} -\mathbf{a}_{B,m+1}^T B^{-1}, 1 \end{bmatrix} \begin{bmatrix} \mathbf{a}_{B,m+1}^T & \mathbf{a}_{N,m+1}^T & \mathbf{0} \\ \mathbf{0} & -\mathbf{a}_{B,m+1}^T B^{-1} N + \mathbf{a}_{N,m+1}^T & 1 \end{bmatrix} =$$

- The new RHS becomes

$$\begin{bmatrix} B^{-1} & \mathbf{0} \\ -\mathbf{a}_{B,m+1}^T B^{-1} & 1 \end{bmatrix} \begin{bmatrix} \mathbf{b} \\ b_{m+1} \end{bmatrix} = \begin{bmatrix} B^{-1} \mathbf{b} \\ b_{m+1} - \mathbf{a}_{B,m+1}^T B^{-1} \mathbf{b} \end{bmatrix}$$

- If $\mathbf{x}_0$ does not satisfy the newly added constraint then the $(m+1)$ th entry of the RHS will be **strictly** less than zero.
- Since the cost associated with s_{m+1} is equal to zero, all the $c_j - z_j$ entries remain unchanged and are non negative.
- Then **Dual Simplex** can be used either to obtain the new optimal solution or to conclude that new (P) is **infeasible**.

Changing entries in the coefficient matrix A :

- **Case 1:** The column corresponding to a **nonbasic variable** say $\tilde{a}_j$ of the optimal solution is changed to $\tilde{a}'_j$.

Changing entries in the coefficient matrix A :

- **Case 1:** The column corresponding to a **nonbasic variable** say $\tilde{\mathbf{a}}_j$ of the optimal solution is changed to $\tilde{\mathbf{a}}'_j$.
- The column corresponding to that variable in the optimal simplex table $B^{-1}\tilde{\mathbf{a}}_j$ is changed to $B^{-1}\tilde{\mathbf{a}}'_j$.

Changing entries in the coefficient matrix A :

- **Case 1:** The column corresponding to a **nonbasic variable** say $\tilde{\mathbf{a}}_j$ of the optimal solution is changed to $\tilde{\mathbf{a}}'_j$.
- The column corresponding to that variable in the optimal simplex table $B^{-1}\tilde{\mathbf{a}}_j$ is changed to $B^{-1}\tilde{\mathbf{a}}'_j$.
- If the new $c_j - z'_j$ value satisfies the **optimality condition**, then the previous optimal solution will be **optimal** for the new problem.

Changing entries in the coefficient matrix A :

- **Case 1:** The column corresponding to a **nonbasic variable** say $\tilde{\mathbf{a}}_j$ of the optimal solution is changed to $\tilde{\mathbf{a}}'_j$.
- The column corresponding to that variable in the optimal simplex table $B^{-1}\tilde{\mathbf{a}}_j$ is changed to $B^{-1}\tilde{\mathbf{a}}'_j$.
- If the new $c_j - z'_j$ value satisfies the **optimality condition**, then the previous optimal solution will be **optimal** for the new problem.
- If **not**, then use **Simplex** either to obtain the new **optimal solution** or to conclude that new (P) does **not** have an optimal solution.

Changing entries in the coefficient matrix A :

- **Case 2:** If the column corresponding to the j th **basic variable** say $\tilde{\mathbf{a}}_j$ is changed to $\tilde{\mathbf{a}}'_j$.

Changing entries in the coefficient matrix A :

- **Case 2:** If the column corresponding to the j th **basic variable** say $\tilde{\mathbf{a}}_j$ is changed to $\tilde{\mathbf{a}}'_j$.
- Then treat this case as the case when a variable (say a variable x_{n+1}) is added to the problem with column $\tilde{\mathbf{a}}'_j$ and cost c_j .

Changing entries in the coefficient matrix A :

- **Case 2:** If the column corresponding to the j th **basic variable** say $\tilde{\mathbf{a}}_j$ is changed to $\tilde{\mathbf{a}}'_j$.
- Then treat this case as the case when a variable (say a variable x_{n+1}) is added to the problem with column $\tilde{\mathbf{a}}'_j$ and cost c_j .
- Add a column $B^{-1}\tilde{\mathbf{a}}'_j$ and the corresponding $c_{n+1} - z_{n+1}$ value to the previous simplex table.

Changing entries in the coefficient matrix A :

- **Case 2:** If the column corresponding to the j th **basic variable** say $\tilde{\mathbf{a}}_j$ is changed to $\tilde{\mathbf{a}}'_j$.
- Then treat this case as the case when a variable (say a variable x_{n+1}) is added to the problem with column $\tilde{\mathbf{a}}'_j$ and cost c_j .
- Add a column $B^{-1}\tilde{\mathbf{a}}'_j$ and the corresponding $c_{n+1} - z_{n+1}$ value to the previous simplex table.
- If $u_{j,(n+1)} = 0$, then it implies that $\{\tilde{\mathbf{a}}_1, \dots, \tilde{\mathbf{a}}_{j-1}, \tilde{\mathbf{a}}'_j, \tilde{\mathbf{a}}_{j+1}, \dots, \tilde{\mathbf{a}}_m\}$ is **LD** and we have to find an **initial BFS** for the changed problem.

Changing entries in the coefficient matrix A :

- **Case 2:** If the column corresponding to the j th **basic variable** say $\tilde{\mathbf{a}}_j$ is changed to $\tilde{\mathbf{a}}'_j$.
- Then treat this case as the case when a variable (say a variable x_{n+1}) is added to the problem with column $\tilde{\mathbf{a}}'_j$ and cost c_j .
- Add a column $B^{-1}\tilde{\mathbf{a}}'_j$ and the corresponding $c_{n+1} - z_{n+1}$ value to the previous simplex table.
- If $u_{j,(n+1)} = 0$, then it implies that $\{\tilde{\mathbf{a}}_1, \dots, \tilde{\mathbf{a}}_{j-1}, \tilde{\mathbf{a}}'_j, \tilde{\mathbf{a}}_{j+1}, \dots, \tilde{\mathbf{a}}_m\}$ is **LD** and we have to find an **initial BFS** for the changed problem.
- If $u_{j,(n+1)} \neq 0$, then pivot on this element and make x_{n+1} enter the basis and x_j leave the basis.

- Perform the necessary elementary row operations to make the $(n+1)$ th column as the j **th column** of I and $C_{n+1} - Z_{n+1}$ equal to 0.

- Perform the necessary elementary row operations to make the $(n+1)$ th column as the j **th column** of I and $c_{n+1} - z_{n+1}$ equal to 0.
- Delete the column corresponding to $\tilde{a}_j$ from all subsequent calculations, and the variable x_{n+1} is now indexed as x_j .

- Perform the necessary elementary row operations to make the $(n+1)$ th column as the j **th column** of I and $C_{n+1} - Z_{n+1}$ equal to 0.
- Delete the column corresponding to $\tilde{a}_j$ from all subsequent calculations, and the variable x_{n+1} is now indexed as x_j .
- The necessary calculations mentioned above might disturb the optimality as well as the feasibility of the current **BFS**.

- Perform the necessary elementary row operations to make the $(n+1)$ th column as the j **th column** of I and $c_{n+1} - z_{n+1}$ equal to 0.
- Delete the column corresponding to $\tilde{a}_j$ from all subsequent calculations, and the variable x_{n+1} is now indexed as x_j .
- The necessary calculations mentioned above might disturb the optimality as well as the feasibility of the current **BFS**.
- If however the new $c_j - z_j$ values are all **nonnegative** and all the RHS entries remain **non negative**, then the table is **optimal** and the corresponding BFS is **optimal** for the new (P).

- If the new table has all RHS entries **nonnegative**, but atleast one of the $c_j - z_j$ values is **negative** then use **Simplex** either to obtain the new optimal solution, or to conclude that the new **(P)** **does not** have an optimal solution.

- If the new table has all RHS entries **nonnegative**, but atleast one of the $c_j - z_j$ values is **negative** then use **Simplex** either to obtain the new optimal solution, or to conclude that the new **(P)** **does not** have an optimal solution.
- If all the $c_j - z_j$ values are **nonnegative** but atleast one of the RHS entries in the simplex table is **negative** then use **Dual Simplex** to obtain the new optimal solution or to conclude that the new **(P)** has **no** feasible solution.

Artificial Variable Method to find an initial BFS of an LPP: Big- M method:

- Consider the problem **P**:
Min $\mathbf{c}^T \mathbf{x}$
subject to $\mathbf{Ax} = \mathbf{b}, \mathbf{x} \geq \mathbf{0}$.

Artificial Variable Method to find an initial BFS of an LPP: Big- M method:

- Consider the problem **P**:
Min $\mathbf{c}^T \mathbf{x}$
subject to $\mathbf{Ax} = \mathbf{b}, \mathbf{x} \geq \mathbf{0}$.
- Assume $\mathbf{b} \geq \mathbf{0}$.

Artificial Variable Method to find an initial BFS of an LPP: Big- M method:

- Consider the problem **P**:
Min $\mathbf{c}^T \mathbf{x}$
subject to $\mathbf{Ax} = \mathbf{b}, \mathbf{x} \geq \mathbf{0}$.
- Assume $\mathbf{b} \geq \mathbf{0}$.
- Add artificial variables $w_1, \dots, w_m$ with cost M (M large).

Artificial Variable Method to find an initial BFS of an LPP: Big- M method:

- Consider the problem **P**:
Min $\mathbf{c}^T \mathbf{x}$
subject to $\mathbf{Ax} = \mathbf{b}, \mathbf{x} \geq \mathbf{0}$.
- Assume $\mathbf{b} \geq \mathbf{0}$.
- Add artificial variables $w_1, \dots, w_m$ with cost M (M large).
- The new problem **LP(M)** is given by,

Artificial Variable Method to find an initial BFS of an LPP: Big- M method:

- Consider the problem **P**:
Min $\mathbf{c}^T \mathbf{x}$
subject to $\mathbf{Ax} = \mathbf{b}, \mathbf{x} \geq \mathbf{0}$.
- Assume $\mathbf{b} \geq \mathbf{0}$.
- Add artificial variables $w_1, \dots, w_m$ with cost M (M large).
- The new problem **LP(M)** is given by,
Min $\mathbf{c}^T \mathbf{x} + [M, \dots, M] \mathbf{w}$
subject to

$$\mathbf{Ax} + \mathbf{w} = [\mathbf{A} : \mathbf{I}] \begin{bmatrix} \mathbf{x} \\ \mathbf{w} \end{bmatrix} = \mathbf{b}, \mathbf{x} \geq \mathbf{0}, \mathbf{w} \geq \mathbf{0},$$

where $\mathbf{w} = [w_1, \dots, w_m]^T$.

Artificial Variable Method to find an initial BFS of an LPP: Big- M method:

- Consider the problem **P**:
Min $\mathbf{c}^T \mathbf{x}$
subject to $\mathbf{Ax} = \mathbf{b}, \mathbf{x} \geq \mathbf{0}$.
- Assume $\mathbf{b} \geq \mathbf{0}$.
- Add artificial variables $w_1, \dots, w_m$ with cost M (M large).
- The new problem **LP(M)** is given by,
Min $\mathbf{c}^T \mathbf{x} + [M, \dots, M] \mathbf{w}$
subject to
$$\mathbf{Ax} + \mathbf{w} = [\mathbf{A} : \mathbf{I}] \begin{bmatrix} \mathbf{x} \\ \mathbf{w} \end{bmatrix} = \mathbf{b}, \mathbf{x} \geq \mathbf{0}, \mathbf{w} \geq \mathbf{0},$$

where $\mathbf{w} = [w_1, \dots, w_m]^T$.
- An initial BFS is $[\mathbf{0}_{1 \times n}, b_1, \dots, b_m]^T$

Artificial Variable Method to find an initial BFS of an LPP: Big- M method:

- Consider the problem **P**:
Min $\mathbf{c}^T \mathbf{x}$
subject to $\mathbf{Ax} = \mathbf{b}, \mathbf{x} \geq \mathbf{0}$.
- Assume $\mathbf{b} \geq \mathbf{0}$.
- Add artificial variables $w_1, \dots, w_m$ with cost M (M large).
- The new problem **LP(M)** is given by,
Min $\mathbf{c}^T \mathbf{x} + [M, \dots, M] \mathbf{w}$
subject to
$$\mathbf{Ax} + \mathbf{w} = [\mathbf{A} : \mathbf{I}] \begin{bmatrix} \mathbf{x} \\ \mathbf{w} \end{bmatrix} = \mathbf{b}, \mathbf{x} \geq \mathbf{0}, \mathbf{w} \geq \mathbf{0},$$
where $\mathbf{w} = [w_1, \dots, w_m]^T$.
- An initial BFS is $[\mathbf{0}_{1 \times n}, b_1, \dots, b_m]^T$ with basic variables $w_1, \dots, w_m$.

Artificial Variable Method to find an initial BFS of an LPP: Big- M method:

- Consider the problem **P**:
Min $\mathbf{c}^T \mathbf{x}$
subject to $\mathbf{Ax} = \mathbf{b}, \mathbf{x} \geq \mathbf{0}$.
- Assume $\mathbf{b} \geq \mathbf{0}$.
- Add artificial variables $w_1, \dots, w_m$ with cost M (M large).
- The new problem **LP(M)** is given by,
Min $\mathbf{c}^T \mathbf{x} + [M, \dots, M] \mathbf{w}$
subject to
$$\mathbf{Ax} + \mathbf{w} = [\mathbf{A} : \mathbf{I}] \begin{bmatrix} \mathbf{x} \\ \mathbf{w} \end{bmatrix} = \mathbf{b}, \mathbf{x} \geq \mathbf{0}, \mathbf{w} \geq \mathbf{0},$$
where $\mathbf{w} = [w_1, \dots, w_m]^T$.
- An initial BFS is $[\mathbf{0}_{1 \times n}, b_1, \dots, b_m]^T$ with basic variables $w_1, \dots, w_m$.
- $\mathbf{x} \in \text{Fea}(\mathbf{P}) \Leftrightarrow [\mathbf{x}^T, \mathbf{0}_{1 \times m}]^T \in \text{Fea}(\text{LP}(\mathbf{M}))$.

- **Case 1:** $LP(M)$ has an optimal solution.

- **Case 1:** $\text{LP}(\mathbf{M})$ has an **optimal** solution.
- **Case 1a:** $[\mathbf{x}_*^T, \mathbf{O}_{1 \times m}]^T$ is an **optimal** BFS of $\text{LP}(\mathbf{M})$ with all the artificial variables as nonbasic variables.

- **Case 1:** $\text{LP}(\mathbf{M})$ has an **optimal** solution.
- **Case 1a:** $[\mathbf{x}_*^T, \mathbf{0}_{1 \times m}]^T$ is an **optimal** BFS of $\text{LP}(\mathbf{M})$ with all the artificial variables as nonbasic variables.
Then $\mathbf{x}_*$ is an **optimal** BFS for $\mathbf{P}$.

- **Case 1:** $\text{LP}(\mathbf{M})$ has an **optimal** solution.
- **Case 1a:** $[\mathbf{x}_*^T, \mathbf{O}_{1 \times m}]^T$ is an **optimal** BFS of $\text{LP}(\mathbf{M})$ with all the artificial variables as nonbasic variables.
Then $\mathbf{x}_*$ is an **optimal** BFS for $\mathbf{P}$.
- **Case 1b:** $[\mathbf{x}_*^T, \mathbf{w}_*^T]^T$ is optimal for $\text{LP}(\mathbf{M})$ and $\mathbf{w}_* \neq \mathbf{O}$.

- **Case 1:** $\text{LP}(\mathbf{M})$ has an **optimal** solution.
- **Case 1a:** $[\mathbf{x}_*^T, \mathbf{O}_{1 \times m}]^T$ is an **optimal** BFS of $\text{LP}(\mathbf{M})$ with all the artificial variables as nonbasic variables.
Then $\mathbf{x}_*$ is an **optimal** BFS for $\mathbf{P}$.
- **Case 1b:** $[\mathbf{x}_*^T, \mathbf{w}_*^T]^T$ is optimal for $\text{LP}(\mathbf{M})$ and $\mathbf{w}_* \neq \mathbf{O}$.
Then $\mathbf{P}$ **does not** have a **feasible solution**.

- **Case 1:** $\text{LP}(\mathbf{M})$ has an **optimal** solution.
- **Case 1a:** $[\mathbf{x}_*^T, \mathbf{0}_{1 \times m}]^T$ is an **optimal** BFS of $\text{LP}(\mathbf{M})$ with all the artificial variables as nonbasic variables.
Then $\mathbf{x}_*$ is an **optimal** BFS for $\mathbf{P}$.
- **Case 1b:** $[\mathbf{x}_*^T, \mathbf{w}_*^T]^T$ is optimal for $\text{LP}(\mathbf{M})$ and $\mathbf{w}_* \neq \mathbf{0}$.
Then $\mathbf{P}$ **does not** have a **feasible solution**.
- **Case 2:** $\text{LP}(\mathbf{M})$ **does not** have an **optimal** solution.

- **Case 2a:** In some iteration (or simplex table), there exists a k such that $c_k - z_k < 0$, the corresponding column $B^{-1}\tilde{\mathbf{a}}_k \leq \mathbf{0}$ and $\mathbf{w} = \mathbf{0}$ in the corresponding BFS for $\text{LP}(\mathbf{M})$.

- **Case 2a:** In some iteration (or simplex table), there exists a k such that $c_k - z_k < 0$, the corresponding column $B^{-1}\tilde{\mathbf{a}}_k \leq \mathbf{0}$ and $\mathbf{w} = \mathbf{0}$ in the corresponding BFS for $\mathbf{LP}(\mathbf{M})$.
- Then there exists a **direction** $\mathbf{d} = \begin{bmatrix} \mathbf{d}_1 \\ \mathbf{d}_2 \end{bmatrix}$ of $\text{Fea}(\mathbf{LP}(\mathbf{M}))$ such that

- **Case 2a:** In some iteration (or simplex table), there exists a k such that $c_k - z_k < 0$, the corresponding column $B^{-1}\tilde{\mathbf{a}}_k \leq \mathbf{0}$ and $\mathbf{w} = \mathbf{0}$ in the corresponding BFS for $\mathbf{LP}(\mathbf{M})$.
- Then there exists a **direction** $\mathbf{d} = \begin{bmatrix} \mathbf{d}_1 \\ \mathbf{d}_2 \end{bmatrix}$ of $\text{Fea}(\mathbf{LP}(\mathbf{M}))$ such that $\mathbf{c}^T \mathbf{d}_1 + [M, \dots, M] \mathbf{d}_2 < 0$.

- **Case 2a:** In some iteration (or simplex table), there exists a k such that $c_k - z_k < 0$, the corresponding column $B^{-1}\tilde{\mathbf{a}}_k \leq \mathbf{0}$ and $\mathbf{w} = \mathbf{0}$ in the corresponding BFS for $\mathbf{LP}(\mathbf{M})$.
- Then there exists a **direction** $\mathbf{d} = \begin{bmatrix} \mathbf{d}_1 \\ \mathbf{d}_2 \end{bmatrix}$ of $\text{Fea}(\mathbf{LP}(\mathbf{M}))$ such that $\mathbf{c}^T \mathbf{d}_1 + [M, \dots, M] \mathbf{d}_2 < 0$.
- Then $\mathbf{d}_1$ is a **direction** of $\text{Fea}(\mathbf{P})$ and $\mathbf{c}^T \mathbf{d}_1 < 0$.

- **Case 2a:** In some iteration (or simplex table), there exists a k such that $c_k - z_k < 0$, the corresponding column $B^{-1}\tilde{\mathbf{a}}_k \leq \mathbf{0}$ and $\mathbf{w} = \mathbf{0}$ in the corresponding BFS for $\mathbf{LP}(\mathbf{M})$.
- Then there exists a **direction** $\mathbf{d} = \begin{bmatrix} \mathbf{d}_1 \\ \mathbf{d}_2 \end{bmatrix}$ of $\text{Fea}(\mathbf{LP}(\mathbf{M}))$ such that $\mathbf{c}^T \mathbf{d}_1 + [M, \dots, M] \mathbf{d}_2 < 0$.
- Then $\mathbf{d}_1$ is a **direction** of $\text{Fea}(\mathbf{P})$ and $\mathbf{c}^T \mathbf{d}_1 < 0$.
Hence $\mathbf{P}$ does **not** have an **optimal** solution but $\mathbf{P}$ has a **feasible** solution.

- **Case 2a:** In some iteration (or simplex table), there exists a k such that $c_k - z_k < 0$, the corresponding column $B^{-1}\tilde{\mathbf{a}}_k \leq \mathbf{0}$ and $\mathbf{w} = \mathbf{0}$ in the corresponding BFS for $\mathbf{LP}(\mathbf{M})$.
- Then there exists a **direction** $\mathbf{d} = \begin{bmatrix} \mathbf{d}_1 \\ \mathbf{d}_2 \end{bmatrix}$ of $\text{Fea}(\mathbf{LP}(\mathbf{M}))$ such that $\mathbf{c}^T \mathbf{d}_1 + [M, \dots, M] \mathbf{d}_2 < 0$.
- Then $\mathbf{d}_1$ is a **direction** of $\text{Fea}(\mathbf{P})$ and $\mathbf{c}^T \mathbf{d}_1 < 0$.
Hence $\mathbf{P}$ does **not** have an **optimal** solution but $\mathbf{P}$ has a **feasible** solution.
- **Case 2b:** In some iteration (or simplex table), $c_k - z_k = \min\{c_j - z_j : c_j - z_j < 0\}$, $B^{-1}\tilde{\mathbf{a}}_k \leq \mathbf{0}$ and $\mathbf{w} \neq \mathbf{0}$ in the corresponding BFS for $\mathbf{LP}(\mathbf{M})$.

- **Case 2a:** In some iteration (or simplex table), there exists a k such that $c_k - z_k < 0$, the corresponding column $B^{-1}\tilde{\mathbf{a}}_k \leq \mathbf{0}$ and $\mathbf{w} = \mathbf{0}$ in the corresponding BFS for $\text{LP}(\mathbf{M})$.
- Then there exists a **direction** $\mathbf{d} = \begin{bmatrix} \mathbf{d}_1 \\ \mathbf{d}_2 \end{bmatrix}$ of $\text{Fea}(\text{LP}(\mathbf{M}))$ such that $\mathbf{c}^T \mathbf{d}_1 + [M, \dots, M] \mathbf{d}_2 < 0$.
- Then $\mathbf{d}_1$ is a **direction** of $\text{Fea}(\mathbf{P})$ and $\mathbf{c}^T \mathbf{d}_1 < 0$.
Hence $\mathbf{P}$ does **not** have an **optimal** solution but $\mathbf{P}$ has a **feasible** solution.
- **Case 2b:** In some iteration (or simplex table), $c_k - z_k = \min\{c_j - z_j : c_j - z_j < 0\}$, $B^{-1}\tilde{\mathbf{a}}_k \leq \mathbf{0}$ and $\mathbf{w} \neq \mathbf{0}$ in the corresponding BFS for $\text{LP}(\mathbf{M})$.
Then $\mathbf{P}$ does **not** have a **feasible** solution.

- **Case 2a:** In some iteration (or simplex table), there exists a k such that $c_k - z_k < 0$, the corresponding column $B^{-1}\tilde{\mathbf{a}}_k \leq \mathbf{0}$ and $\mathbf{w} = \mathbf{0}$ in the corresponding BFS for $\text{LP}(\mathbf{M})$.
- Then there exists a **direction** $\mathbf{d} = \begin{bmatrix} \mathbf{d}_1 \\ \mathbf{d}_2 \end{bmatrix}$ of $\text{Fea}(\text{LP}(\mathbf{M}))$ such that $\mathbf{c}^T \mathbf{d}_1 + [M, \dots, M] \mathbf{d}_2 < 0$.
- Then $\mathbf{d}_1$ is a **direction** of $\text{Fea}(\mathbf{P})$ and $\mathbf{c}^T \mathbf{d}_1 < 0$.
Hence $\mathbf{P}$ does **not** have an **optimal** solution but $\mathbf{P}$ has a **feasible** solution.
- **Case 2b:** In some iteration (or simplex table), $c_k - z_k = \min\{c_j - z_j : c_j - z_j < 0\}$, $B^{-1}\tilde{\mathbf{a}}_k \leq \mathbf{0}$ and $\mathbf{w} \neq \mathbf{0}$ in the corresponding BFS for $\text{LP}(\mathbf{M})$.
Then $\mathbf{P}$ does **not** have a **feasible** solution.
- The above conclusion of **Case 2b** that $\mathbf{P}$ is **infeasible** may not be true if $c_k - z_k$ is **not** the **most negative** among the $c_j - z_j$ values.

- **Case 2(continued):** In some iteration (or simplex table), there exists a k such that $c_k - z_k < 0$, the corresponding column $B^{-1}\tilde{a}_k \leq 0$ for **LP(M)**.

- **Case 2(continued):** In some iteration (or simplex table), there exists a k such that $c_k - z_k < 0$, the corresponding column $B^{-1}\tilde{a}_k \leq \mathbf{0}$ for **LP(M)**.
Then **P** does **not** have an **optimal** solution (may be **infeasible**, may not be **infeasible**).

- **Case 2(continued):** In some iteration (or simplex table), there exists a k such that $c_k - z_k < 0$, the corresponding column $B^{-1}\tilde{a}_k \leq \mathbf{0}$ for **LP(M)**.
Then **P** does **not** have an **optimal** solution (may be **infeasible**, may not be **infeasible**).
- **Remark 1:** If at some stage of solving the artificial variables **leave** the basis, then the corresponding **BFS** is a **BFS** of **P**.

- **Case 2(continued):** In some iteration (or simplex table), there exists a k such that $c_k - z_k < 0$, the corresponding column $B^{-1}\tilde{a}_k \leq \mathbf{0}$ for **LP(M)**.
Then **P** does **not** have an **optimal** solution (may be **infeasible**, may not be **infeasible**).
- **Remark 1:** If at some stage of solving the artificial variables **leave** the basis, then the corresponding **BFS** is a **BFS** of **P**.
Then **delete** all columns corresponding to the **artificial variables** and continue.

- **Example 1** : Consider the problem **P**,

Minimize $x_1 - x_2$

subject to

$$2x_1 + x_2 \geq 4$$

$$x_1 - x_2 \leq 1$$

$$x_1 \geq 0, x_2 \geq 0.$$

- **Example 1** : Consider the problem **P**,

Minimize $x_1 - x_2$

subject to

$$2x_1 + x_2 \geq 4$$

$$x_1 - x_2 \leq 1$$

$$x_1 \geq 0, x_2 \geq 0.$$

- By adding variables we get the following problem,

Minimize $x_1 - x_2$

subject to

$$2x_1 + x_2 - s_1 = 4$$

$$x_1 - x_2 + s_2 = 1$$

$$x_1 \geq 0, x_2 \geq 0, s_1 \geq 0, s_2 \geq 0.$$

- **Example 1** : Consider the problem **P**,

Minimize $x_1 - x_2$

subject to

$$2x_1 + x_2 \geq 4$$

$$x_1 - x_2 \leq 1$$

$$x_1 \geq 0, x_2 \geq 0.$$

- By adding variables we get the following problem,

Minimize $x_1 - x_2$

subject to

$$2x_1 + x_2 - s_1 = 4$$

$$x_1 - x_2 + s_2 = 1$$

$$x_1 \geq 0, x_2 \geq 0, s_1 \geq 0, s_2 \geq 0.$$

- If we consider the initial basic solution with basic variables s_1 and s_2 , then the $c_j - z_j$ values are not ≥ 0 for all j .

- **Example 1** : Consider the problem **P**,

Minimize $x_1 - x_2$

subject to

$$2x_1 + x_2 \geq 4$$

$$x_1 - x_2 \leq 1$$

$$x_1 \geq 0, x_2 \geq 0.$$

- By adding variables we get the following problem,

Minimize $x_1 - x_2$

subject to

$$2x_1 + x_2 - s_1 = 4$$

$$x_1 - x_2 + s_2 = 1$$

$$x_1 \geq 0, x_2 \geq 0, s_1 \geq 0, s_2 \geq 0.$$

- If we consider the initial basic solution with basic variables s_1 and s_2 , then the $c_j - z_j$ values are not ≥ 0 for all j .
We do **not** have a **feasible solution** of the **dual** of **P** at hand.

- **Example 1** : Consider the problem **P**,

Minimize $x_1 - x_2$

subject to

$$2x_1 + x_2 \geq 4$$

$$x_1 - x_2 \leq 1$$

$$x_1 \geq 0, x_2 \geq 0.$$

- By adding variables we get the following problem,

Minimize $x_1 - x_2$

subject to

$$2x_1 + x_2 - s_1 = 4$$

$$x_1 - x_2 + s_2 = 1$$

$$x_1 \geq 0, x_2 \geq 0, s_1 \geq 0, s_2 \geq 0.$$

- If we consider the initial basic solution with basic variables s_1 and s_2 , then the $c_j - z_j$ values are not ≥ 0 for all j .
We do **not** have a **feasible solution** of the **dual** of **P** at hand.
- The (**Big-M**) is used which provides an **initial BFS** of **P** for Simplex algorithm.

- Consider the modified problem

Minimize $x_1 - x_2 + Mw$

subject to

$$2x_1 + x_2 - s_1 + w = 4$$

$$x_1 - x_2 + s_2 = 1$$

$$x_1, x_2, s_1, s_2, w \geq 0.$$

- Consider the modified problem

Minimize $x_1 - x_2 + Mw$

subject to

$$2x_1 + x_2 - s_1 + w = 4$$

$$x_1 - x_2 + s_2 = 1$$

$$x_1, x_2, s_1, s_2, w \geq 0.$$

Here w is called the artificial variable and cost M associated with it is **very large**.

- Consider the modified problem

Minimize $x_1 - x_2 + Mw$

subject to

$$2x_1 + x_2 - s_1 + w = 4$$

$$x_1 - x_2 + s_2 = 1$$

$$x_1, x_2, s_1, s_2, w \geq 0.$$

Here w is called the artificial variable and cost M associated with it is **very large**.

- The initial table corresponding to the basic variables w and s_2 is given below.

$c_j - z_j$	$1 - 2M$	$-M - 1$	M	0	0	
	$B^{-1}\tilde{\mathbf{a}}_1$	$B^{-1}\tilde{\mathbf{a}}_2$	$B^{-1}\mathbf{s}_1$	$B^{-1}\mathbf{s}_2$	$B^{-1}\mathbf{w}$	$B^{-1}\mathbf{b}$
w	2	1	-1	0	1	4
s_2	1	-1	0	1	0	1

- Consider the modified problem

Minimize $x_1 - x_2 + Mw$

subject to

$$2x_1 + x_2 - s_1 + w = 4$$

$$x_1 - x_2 + s_2 = 1$$

$$x_1, x_2, s_1, s_2, w \geq 0.$$

Here w is called the artificial variable and cost M associated with it is **very large**.

- The initial table corresponding to the basic variables w and s_2 is given below.

$c_j - z_j$	$1 - 2M$	$-M - 1$	M	0	0	
	$B^{-1}\tilde{a}_1$	$B^{-1}\tilde{a}_2$	$B^{-1}s_1$	$B^{-1}s_2$	$B^{-1}w$	$B^{-1}b$
w	2	1	-1	0	1	4
s_2	1	-1	0	1	0	1

- x_1 will be the **entering variable** and s_2 will be the **leaving variable** for the next table.



$c_j - z_j$	0	$-3M$	M	$2M - 1$	0	
	$B^{-1}\tilde{\mathbf{a}}_1$	$B^{-1}\tilde{\mathbf{a}}_2$	$B^{-1}\mathbf{s}_1$	$B^{-1}\mathbf{s}_2$	$B^{-1}\mathbf{w}$	$B^{-1}\mathbf{b}$
w	0	3	-1	-2	1	2
x_1	1	-1	0	1	0	1



$c_j - z_j$	0	$-3M$	M	$2M - 1$	0	
	$B^{-1}\tilde{\mathbf{a}}_1$	$B^{-1}\tilde{\mathbf{a}}_2$	$B^{-1}\mathbf{s}_1$	$B^{-1}\mathbf{s}_2$	$B^{-1}\mathbf{w}$	$B^{-1}\mathbf{b}$
w	0	3	-1	-2	1	2
x_1	1	-1	0	1	0	1

- x_2 will be the **entering variable** and the artificial variable w will be the **leaving variable**.



$c_j - z_j$	0	$-3M$	M	$2M - 1$	0	
	$B^{-1}\tilde{\mathbf{a}}_1$	$B^{-1}\tilde{\mathbf{a}}_2$	$B^{-1}\mathbf{s}_1$	$B^{-1}\mathbf{s}_2$	$B^{-1}\mathbf{w}$	$B^{-1}\mathbf{b}$
w	0	3	-1	-2	1	2
x_1	1	-1	0	1	0	1

- x_2 will be the **entering variable** and the artificial variable w will be the **leaving variable**.



$c_j - z_j$	0	0	0	-1	M	
	$B^{-1}\tilde{\mathbf{a}}_1$	$B^{-1}\tilde{\mathbf{a}}_2$	$B^{-1}\mathbf{s}_1$	$B^{-1}\mathbf{s}_2$	$B^{-1}\mathbf{w}$	$B^{-1}\mathbf{b}$
x_2	0	1				$\frac{2}{3}$
x_1	1	0				$1 + \frac{2}{3}$

- Continue with the BFS corresponding to the basic vectors x_1 and x_2 and drop the column corresponding to the artificial variable w from all future calculations.

- Continue with the BFS corresponding to the basic vectors x_1 and x_2 and drop the column corresponding to the artificial variable w from all future calculations.
Now proceed with **simplex method**.

- Continue with the BFS corresponding to the basic vectors x_1 and x_2 and drop the column corresponding to the artificial variable w from all future calculations.
Now proceed with **simplex method**.

- Example 2:** Consider the problem,

Minimize $x_1 + x_2$

subject to

$$x_1 + 2x_2 \leq 2$$

$$3x_1 + 5x_2 \geq 15$$

$$x_1 \geq 0, x_2 \geq 0.$$

- Continue with the BFS corresponding to the basic vectors x_1 and x_2 and drop the column corresponding to the artificial variable w from all future calculations.
Now proceed with **simplex method**.

- Example 2:** Consider the problem,
Minimize $x_1 + x_2$
subject to

$$x_1 + 2x_2 \leq 2$$

$$3x_1 + 5x_2 \geq 15$$

$$x_1 \geq 0, x_2 \geq 0.$$

We consider the corresponding modified problem

- Minimize $x_1 + x_2 + Mw$
subject to

$$x_1 + 2x_2 + s_1 = 2$$

$$3x_1 + 5x_2 - s_2 + w = 15$$

$$x_1, x_2, s_1, s_2, w \geq 0.$$

- **Note** that this problem can also be solved by **dual simplex method**.

We will solve it by using the **Big-M** method.

- **Note** that this problem can also be solved by **dual simplex method**.

We will solve it by using the **Big-M** method.

- The initial table is given by

$c_j - z_j$	$1 - 3M$	$1 - 5M$	0	M	0	
	$B^{-1}\tilde{\mathbf{a}}_1$	$B^{-1}\tilde{\mathbf{a}}_2$	$B^{-1}\mathbf{s}_1$	$B^{-1}\mathbf{s}_2$	$B^{-1}\mathbf{w}$	$B^{-1}\mathbf{b}$
s_1	1	2	1	0	0	2
w	3	5	0	-1	1	15

- **Note** that this problem can also be solved by **dual simplex method**.

We will solve it by using the **Big-M** method.

- The initial table is given by

$c_j - z_j$	$1 - 3M$	$1 - 5M$	0	M	0	
	$B^{-1}\tilde{\mathbf{a}}_1$	$B^{-1}\tilde{\mathbf{a}}_2$	$B^{-1}\mathbf{s}_1$	$B^{-1}\mathbf{s}_2$	$B^{-1}\mathbf{w}$	$B^{-1}\mathbf{b}$
s_1	1	2	1	0	0	2
w	3	5	0	-1	1	15

- Here x_2 is the **entering variable** and s_1 the **leaving variable**, hence the next table is given by,

- **Note** that this problem can also be solved by **dual simplex method**.

We will solve it by using the **Big-M** method.

- The initial table is given by

$c_j - z_j$	$1 - 3M$	$1 - 5M$	0	M	0	
	$B^{-1}\tilde{\mathbf{a}}_1$	$B^{-1}\tilde{\mathbf{a}}_2$	$B^{-1}\mathbf{s}_1$	$B^{-1}\mathbf{s}_2$	$B^{-1}\mathbf{w}$	$B^{-1}\mathbf{b}$
s_1	1	2	1	0	0	2
w	3	5	0	-1	1	15

- Here x_2 is the **entering variable** and s_1 the **leaving variable**, hence the next table is given by,



$c_j - z_j$	$\frac{1}{2}(1 - M)$	0	$\frac{1}{2}(5M - 1)$	M	0	
	$B^{-1}\tilde{\mathbf{a}}_1$	$B^{-1}\tilde{\mathbf{a}}_2$	$B^{-1}\mathbf{s}_1$	$B^{-1}\mathbf{s}_2$	$B^{-1}\mathbf{w}$	$B^{-1}\mathbf{b}$
x_2	$\frac{1}{2}$	1	$\frac{1}{2}$	0	0	1
w	$\frac{1}{2}$	0	$-\frac{5}{2}$	-1	1	10

- Here x_1 is the entering variable and x_2 the leaving variable and the next table is given by

- Here x_1 is the entering variable and x_2 the leaving variable and the next table is given by

$c_j - z_j$	0	$M - 1$	$3M - 1$	M	0	
	$B^{-1}\tilde{\mathbf{a}}_1$	$B^{-1}\tilde{\mathbf{a}}_2$	$B^{-1}\mathbf{s}_1$	$B^{-1}\mathbf{s}_2$	$B^{-1}\mathbf{w}$	$B^{-1}\mathbf{b}$
x_1	1	2	1	0	0	2
w	0	-1	-3	-1	1	9

- Here x_1 is the entering variable and x_2 the leaving variable and the next table is given by

$c_j - z_j$	0	$M - 1$	$3M - 1$	M	0	
	$B^{-1}\tilde{\mathbf{a}}_1$	$B^{-1}\tilde{\mathbf{a}}_2$	$B^{-1}\mathbf{s}_1$	$B^{-1}\mathbf{s}_2$	$B^{-1}\mathbf{w}$	$B^{-1}\mathbf{b}$
x_1	1	2	1	0	0	2
w	0	-1	-3	-1	1	9

- Since the above optimal table has an artificial variable taking positive value hence the original problem (without the artificial variable) has no feasible solution.

- If we solve the above problem by the **Dual simplex method** then the problem is written as:

- If we solve the above problem by the **Dual simplex method** then the problem is written as:

Minimize $x_1 + x_2$

subject to

$$x_1 + 2x_2 + s_1 = 2$$

$$-3x_1 - 5x_2 + s_2 = -15$$

$$x_1, x_2, s_1, s_2 \geq 0.$$

- If we solve the above problem by the **Dual simplex method** then the problem is written as:

Minimize $x_1 + x_2$

subject to

$$x_1 + 2x_2 + s_1 = 2$$

$$-3x_1 - 5x_2 + s_2 = -15$$

$$x_1, x_2, s_1, s_2 \geq 0.$$

The initial table is given by:

$c_j - z_j$	1	1	0	0	
	$B^{-1}\tilde{\mathbf{a}}_1$	$B^{-1}\tilde{\mathbf{a}}_2$	$B^{-1}\mathbf{e}_1$	$B^{-1}\mathbf{e}_2$	$B^{-1}\mathbf{b}$
s_1	1	2	1	0	2
s_2	-3	-5	0	1	-15

- If we solve the above problem by the **Dual simplex method** then the problem is written as:

Minimize $x_1 + x_2$

subject to

$$x_1 + 2x_2 + s_1 = 2$$

$$-3x_1 - 5x_2 + s_2 = -15$$

$$x_1, x_2, s_1, s_2 \geq 0.$$

The initial table is given by:

$c_j - z_j$	1	1	0	0	
	$B^{-1}\tilde{\mathbf{a}}_1$	$B^{-1}\tilde{\mathbf{a}}_2$	$B^{-1}\mathbf{e}_1$	$B^{-1}\mathbf{e}_2$	$B^{-1}\mathbf{b}$
s_1	1	2	1	0	2
s_2	-3	-5	0	1	-15

Here s_2 is the **leaving variable** and x_2 the **entering variable**.

- The next table is given by,

- The next table is given by,

$c_j - z_j$	$\frac{2}{5}$	0	0	$\frac{1}{5}$	
	$B^{-1}\tilde{\mathbf{a}}_1$	$B^{-1}\tilde{\mathbf{a}}_2$	$B^{-1}\mathbf{e}_1$	$B^{-1}\mathbf{e}_2$	$B^{-1}\mathbf{b}$
s_1	$-\frac{1}{5}$	0	1	$\frac{2}{5}$	-4
x_2	$\frac{3}{5}$	1	0	$-\frac{1}{5}$	3

- The next table is given by,

$c_j - z_j$	$\frac{2}{5}$	0	0	$\frac{1}{5}$	
	$B^{-1}\tilde{\mathbf{a}}_1$	$B^{-1}\tilde{\mathbf{a}}_2$	$B^{-1}\mathbf{e}_1$	$B^{-1}\mathbf{e}_2$	$B^{-1}\mathbf{b}$
s_1	$-\frac{1}{5}$	0	1	$\frac{2}{5}$	-4
x_2	$\frac{3}{5}$	1	0	$-\frac{1}{5}$	3

- Here s_1 is the leaving variable and x_1 the entering variable. The next table is given by:

$c_j - z_j$	0	0			
	$B^{-1}\tilde{\mathbf{a}}_1$	$B^{-1}\tilde{\mathbf{a}}_2$	$B^{-1}\mathbf{e}_1$	$B^{-1}\mathbf{e}_2$	$B^{-1}\mathbf{b}$
x_1	1	0	-5	-2	20
x_2	0	1	3	1	-9

- The next table is given by,

$c_j - z_j$	$\frac{2}{5}$	0	0	$\frac{1}{5}$	
	$B^{-1}\tilde{\mathbf{a}}_1$	$B^{-1}\tilde{\mathbf{a}}_2$	$B^{-1}\mathbf{e}_1$	$B^{-1}\mathbf{e}_2$	$B^{-1}\mathbf{b}$
s_1	$-\frac{1}{5}$	0	1	$\frac{2}{5}$	-4
x_2	$\frac{3}{5}$	1	0	$-\frac{1}{5}$	3

- Here s_1 is the leaving variable and x_1 the entering variable. The next table is given by:

$c_j - z_j$	0	0			
	$B^{-1}\tilde{\mathbf{a}}_1$	$B^{-1}\tilde{\mathbf{a}}_2$	$B^{-1}\mathbf{e}_1$	$B^{-1}\mathbf{e}_2$	$B^{-1}\mathbf{b}$
x_1	1	0	-5	-2	20
x_2	0	1	3	1	-9

Hence clearly the problem is infeasible.