
Annual groundwater discharge ("baseflow") during water−resources year
before 1978
after 1978
10
15
20
m3/s

Annual groundwater discharge ("baseflow") during water−resources year

Student: t = −8.565, p = 0, m1 = 9.086, m2 = 13.783
Fisher: F = 0.62, p = 0.25127, cv1 = 0.206, cv2 = 0.172



Maximum annual discharge during seasonal flood wave

Student: t = 1.526, p = 0.14949, m1 = 304.748, m2 = 121.514
Fisher: F = 4.379, p = 0.06852, cv1 = 0.533, cv2 = 0.572



Annual groundwater discharge ("baseflow") during water−resources year

Student: t = 3.738, p = 0.00043, m1 = 10.445, m2 = 13.829
Fisher: F = 1.416, p = 0.33843, cv1 = 0.262, cv2 = 0.186



Annual groundwater discharge ("b
resources year

Student: t = 1.526, p = 0.14949, m1 = 1
Fisher: F = 4.379, p = 0.06852, cv1 = 0.



First date of 10−day window disch

Student: t = −2.665, p = 0.01341, m1 =
Fisher: F = 2.464, p = 0.01576, cv1 = 0.



Maximum annual discharge during

Student: t = 3.738, p = 0.00043, m1 = 3
Fisher: F = 1.416, p = 0.33843, cv1 = 0.



Seasonal flood runoff (with ground

Student: t = 3.469, p = 0.00101, m1 = 12
Fisher: F = 1.516, p = 0.25254, cv1 = 0.