
Tolerant − sensitive parental phenotype within tissue: estimate ± 95% CI from a mixed model with family random intercept
(fish-level OLS where the family variance is at its boundary); raw p shown, * BH-FDR < 0.05. Δ = phenotype × treatment interaction
Global TE axis (SD)
logit strict-TE fraction
logit all-repeat fraction
brain ctrl
brain dev
brain intergen
brain mean
brain Δdev effect
brain Δintergen effect
gill ctrl
gill dev
gill intergen
gill mean
gill Δdev effect
gill Δintergen effect
liver ctrl
liver dev
liver intergen
liver mean
liver Δdev effect
liver Δintergen effect
p=0.011
p=0.024
p=0.78
p=0.024
p=0.82
p=0.034
p=0.11
p=0.5
p=0.61
p=0.22
p=0.34
p=0.36
p=0.33
p=0.33
p=0.15
p=0.4
p=0.17
p=0.74
p=0.025
p=0.038
p=0.79
p=0.019
p=0.9
p=0.074
p=0.095
p=0.44
p=0.16
p=0.029
p=0.51
p=0.84
p=0.11
p=0.034
p=0.96
p=0.74
p=0.0085
p=0.25
p=0.02
p=0.14
p=0.78
p=0.075
p=0.47
p=0.048
p=0.32
p=0.46
p=0.2
p=0.086
p=0.86
p=0.83
p=0.45
p=0.022
p=0.93
p=0.33
p=0.031
p=0.55
−2
−1
0
1
2
3
−0.2
0.0
0.2
0.4
−0.2
0.0
0.2
0.4