

Paired tumour-vs-normal T cells from 43 patients: which genes does a foundation model read as T-cell state?

Kaisar Dauyey · Shinji Nakaoka

Laboratory of Mathematical Biology, Faculty of Advanced Life Science, Hokkaido University

43 donors, both tissues,
100 cells each

Delete AND overexpress,
donor's own normal as goal

Registered before any number;
all 54 rows reported

Geneformer V2-316M (bf16) · in silico perturbation is a hypothesis generator, not a knockout screen

Two registered questions, one cleaner design

Panel A — do the July hits replicate?

The 15 LUAD→normal top genes of the July screen.
July compared LUAD with normal tissue from other studies (study-confounded).

Panel B — does the model read T-cell genes?


The lab's 39 curated T-cell genes (36 perturbable).
July never ranked them in its top 120.

What 'a hit' means here

Per donor: shift of tumour T cells toward the SAME donor's normal T cells, minus the median of 20 matched control genes.
Exact Wilcoxon over donors, Holm per panel, for deletion and overexpression separately.
A signal needs BOTH arms significant with opposite signs (dose concordance).

Pairing within a donor cancels study, chemistry and person — not ambient RNA

Design: every donor contributes both tissues; after the cap each donor weighs exactly 1/43




What pairing cannot fix

Tumour and normal tissue differ in ambient contamination within each donor. 20 of 43 donors overlap July's LUAD pool. Selection favours T-cell-rich samples; 17 unpaired donors excluded.

The model tells a donor's tumour from normal T cells: 43/43 donors above chance

Classifier gate PASS: pooled 0.825; 43/43 donors above chance (sign test $p = 2.3e-13$)



Gate: PASS


Pooled balanced accuracy 0.825 (gate 0.60).

Sign test $p = 2.3e-13$.

Every donor scored by a model that never saw it.

Fine-tune not bit-reproducible (band 0.045 per donor); cannot flip the gate.

13 of 15 July hits are not expressed in T cells — not testable, not refuted



Reading


Keratins, mucin, a secretoglobin, a haemoglobin chain: epithelial and blood markers, present in ≤ 8 of 43 donors' T cells.

The one testable gene, POLR2J3, is OPEN ($p = 0.10 / 0.25$).

The design did not refute the July panel; it showed the panel is mostly not T-cell biology.

Confound: 316M replaced 104M (no 104M arm), so a non-replication could not be attributed — moot, none occurred.

13 of 34 tested T-cell genes give a dose-concordant signal



Toward normal (11)

CD3D, CD3G, CD247, CD7, LCK, LAT, CD8A, CCR7, GZMA, CD27, ICOS

Away from normal (2)

ITK, CTSW


Stable — and what cannot be judged

None of the 13 moves under any registered sensitivity.

Cannot attain significance at any effect size: CD28, CTLA4, PDCD1 ($n = 9, 6, 5$) — no evidence either way.

TRAC/TRBC1/TRBC2: not in the model's vocabulary.

Opposite-signed arms are common even for control genes — effect size separates few



Confound

360 matched control entries are anti-correlated too ($\rho = -0.62$); 35% sit in the toward-normal quadrant.


Post hoc, descriptive: only CCR7, CD3D, GZMA, CD7 (and CD27 on overexpression) lie beyond ~95% of controls.

Measured on the same cells (Amendment 3h)

Geneformer overexpresses into all cells; the token-positive subset was reconstructed and checked.


15,142 / 15,142 count checks; 8 / 8 GPU identity pairs bit-identical.

The medians are not carried by a few donors



The coherent set does not move under any registered sensitivity

Sensitivities: reported alongside, never used to change a registered status



Three fragile rows, outside the coherent set

GZMK: deletion-only → away under all-cells overexpress. PRF1: open → deletion-only if any control counts. CD69: deletion-only → dose-incoherent under Holm-over-tested and under the global goal.

Panel-level: 21 toward vs 13 away, sign test $p = 0.23$ — genes share controls, so signs are not independent.

What each status licenses — and what it does not

T-cell signal (13)

Licenses: Model-internal, donor-consistent, dose-concordant response beyond matched controls.

Does not: Not a causal driver; mostly not unusual among genes.

Deletion only (2)

Licenses: A consistent deletion effect.

Does not: No direction; no signal claim.

Open (20)

Licenses: The registered test did not call it.

Does not: Not 'no effect'; for CD28/CTLA4/PDCD1 no evidence either way.

Not run (18) • no controls (1)

Licenses: Not expressed in enough donors' T cells, or not in the vocabulary.

Does not: Not non-replication; not no effect.

What this design still cannot rule out

Ambient RNA

Tumour vs normal tissue differ in contamination within a donor; pairing cannot remove it.

Model change

316M bf16 replaced July's 104M fp32; no 104M arm.

Not independent

20 of 43 donors overlap July's LUAD pool.

Selection

T-cell-rich samples only; 17 unpaired donors excluded.

Nondeterminism

Fine-tuning is not bit-reproducible; the ISP inherits these fold models.

Process

No-op and wrapper checks ran after Phase 6 started; the overexpress cell set was fixed by Amendment 3h after Phase 6.

For the July screen: its LUAD → normal list was mostly not T-cell biology

What this changes

13 of 15 July hits cannot be tested in T cells under a paired, balanced design.

The model does read canonical T-cell genes (CD3 complex, LCK, LAT, CD8A, CCR7, GZMA) in a donor-consistent, dose-concordant way.

Effect size, not significance, separates a handful from arbitrary genes.

Next

Genome-wide control distribution, to rank effect size formally (registered first).

Checkpoint genes (CTLA4, PDCD1, CD28) need more token-positive donors: a larger cohort or a looser cap.

A 104M arm, if attribution to model vs design is wanted.

Independent tissue validation for the top four (CCR7, CD3D, GZMA, CD7).