

# MASTEQ Manual

## [Abstract]

This code is a numerical estimator of atomic diffusivity in crystals using a master equation approach. The diffusion tensor is calculated under the independent particle approximation only by specifying the initial and final sites, jump vector, and jump frequency of every atomic jump in unitcell. See the following references for the detailed theoretical background. Note that interaction between diffusion carriers cannot be treated in the current version. This code was written by Kazuaki Toyoura. Mr. Takeo Fujii has great contribution to the code development with fruitful discussions.

- [1] K. Toyoura, T. Fujii, N. Hatada, D. Han, T. Uda, Carrier–Carrier Interaction in Proton-Conducting Perovskites: Carrier Blocking vs Trap-Site Filling, *The Journal of Physical Chemistry C* **123**, 26823–26830 (2019).  
 [2] K. Toyoura, T. Fujii, K. Kanamori, I. Takeuchi, Sampling strategy in efficient potential energy surface mapping for predicting atomic diffusivity in crystals by machine learning, *Phys. Rev. B* **101**, 184117/1–11 (2020).  
 [3] T. Fujii, K. Toyoura, T. Uda, S. Kasamatsu, Theoretical study on proton diffusivity in Y-doped BaZrO<sub>3</sub> with realistic dopant configurations, *Phys. Chem. Chem. Phys.* **23**, 5908–5918 (2021).

## [Programming language]

Python 3.x (installed through anaconda, <https://www.anaconda.com/products/individual>)

Imported modules in this code: numpy, scipy, argparse, copy, datetime, os

\*mkemig.py requires *pymatgen*, <https://pymatgen.org/>.

## [Input file]

Only an input file, *jmpdata.csv*, is required, which includes initial and final site IDs, jump vectors, and jump frequencies for every atomic jump in unitcell. Note that all atomic jumps with initial sites in the unitcell have to be specified including the atomic jumps across the periodic boundary, and that both jumps in the opposite directions have to be specified separately. New line for different atomic jumps. “#” denotes a comment line. Put the following items for every atomic jump separated by comma. (Site IDs have to be sequential numbers starting from 1.)

- Initial site ID [integer]: ID of the initial site for an atomic jump
- Final site ID [integer]: ID of the final site for an atomic jump
- *x* component of the jump vector [float]: The *x* component of jump vector in Å for an atomic jump
- *y* component of the jump vector [float]: The *y* component of jump vector in Å for an atomic jump
- *z* component of the jump vector [float]: The *z* component of jump vector in Å for an atomic jump
- Jump frequency [float]: Jump frequency in Hz for an atomic jump

(jmpdata.csv example)

```
#InitialSiteID,FinalSiteID,jmpVec_x[Ang.], jmpVec_y[Ang.], jmpVec_z[Ang.],frequency[Hz]
```

```
1,7,-1.2067005467,1.2067005467,0.0000000000,5.4961104742e+11
```

```
1,9,0.0000000000,-0.9114043880,-0.9114043880,1.3907136217e+12
```

**[Options]**

- h, --help Help information. List of options in this code.
- jump File name of atomic jump data in a given system. If it is the default name (jmpdata.csv), the argument is not necessary.
- nonzero Non-zero elements in diffusion tensor. Specified non-zero elements are estimated in this program. Specify element indexes separated by comma. Default is 1,2,3,4,5,6. Note that the independent elements are six in diffusion tensor, where the index is as follows:

$$\text{Diffusion tensor: } \begin{pmatrix} D_1 & D_6 & D_5 \\ D_6 & D_2 & D_4 \\ D_5 & D_4 & D_3 \end{pmatrix}$$

For example, “--nonzero 1,2,3” means estimating only the diagonal elements.

- factor Scaling factor  $f$  for wave vector  $\mathbf{Q}$ . This factor have to be enough small to correspond to large scale for atomic diffusion. The length of the shortest jump vector  $l_{\min}$  is used for the standardized scale of reciprocal space, given by  $2\pi/l_{\min}$ . The magnitude of wave vector  $|\mathbf{Q}|$  for estimating diffusion tensor is then  $(2\pi/l_{\min}) \times f$ . Default value is 1.0e-3.

**[Usage]**

Type the following command at the directory with jmpdata.csv. The estimated diffusion tensor is printed on the standard output. Please redirect it to a file, e.g., stdout.

```
python [MASTEQ_DIR]/masteq.py --jump jmpdata.csv --nonzero 1,2,3 --factor 1.0e-3 > stdout
```


**[Note]**

This program assumes a single diffusion network without any other competing network in a given system. For example, if several parallel two-dimensional (2D) networks coexist in a given system, the estimated diffusivity reflects only the diffusivity of the fastest one. In such a case, the jump matrix has several eigenvalues close to zero, and this program prints a **warning** on the standard output. This warning is often output for anisotropic diffusivity at low temperatures, because a single network at high temperatures can substantially be separated into several networks at low temperatures. Generally, the differences in jump frequency between atomic jumps become larger rapidly with decreasing temperatures, which creates negligible paths for atomic jumps at low temperatures, leading to the separated networks.

The accuracy of estimated diffusivity corresponds to the accuracy of the minimum-magnitude eigenvalue in the jump matrix. The smaller scaling factor  $f$  for wave vector  $\mathbf{Q}$  is better in terms of larger scale for atomic diffusion, but worse in terms of the accuracy of eigenvalues in the jump matrix. Please carefully check the temperature dependence of estimated diffusivity, which should be a smooth curve (approximately a straight line) in the Arrhenius plot. In the future, this program has a function for checking the diffusion network (site connectivity) in a given system.

**[Example]**

In example directory, the proton diffusivity in the perfect crystal of BaZrO<sub>3</sub> [3] is estimated in the range of 200-1000 K. In the perfect crystal, a proton migrates two types of proton jumps, i.e., *rotation* around single oxide ions and *hopping* between adjacent oxide ions. The calculated migration energies of proton rotation and hopping are 0.17 and 0.25 eV, respectively. The input files (jmpdata.csv) at all temperatures can be generated all at once by **mkjmpdata.py** in the next section.

**[Useful tools]****mkjmpdata.py**

This program makes multiple **jmpdata.csv** all at once in a specified temperature range, using the migration energy  $\Delta E^{\text{mig}}$  and vibrational prefactor  $\nu_0$  for atomic jump. The jump frequency  $\nu$  is estimated on the basis of the classical transition state theory,  $\nu = \nu_0 \exp(-\Delta E^{\text{mig}}/k_B T)$ , where  $k_B$  and  $T$  are the Boltzmann constant and temperature. The input file (default name: emig.csv) have to be prepared in the csv format. The file format is almost same as jmpdata.csv, where  $\nu$  [Hz], is replaced by  $\Delta E^{\text{mig}}$  [eV] and  $\nu_0$  [Hz] separated by comma. Emig.csv have to contain the information on all atomic jumps in the unitcell as well as jmpdata.csv. The file example is as follows:

(emig.csv example)

```
#InitialSiteID,FinalSiteID,jmpVec_x[Ang.],jmpVec_y[Ang.],jmpVec_z[Ang.],DEmig[eV],v0[Hz]
1,7,-1.2067005467,1.2067005467,0.0000000000,0.25,1.0e+13
1,9,0.0000000000,-0.9114043880,-0.9114043880,0.17,1.0e+13
```

The temperature range [K] is specified by three parameters, i.e., lb\_T (lower bound of temperature), ub\_T (upper bound of temperature), and int\_T (interval of temperature), which are specified as arguments.

```
-h, --help          Help information. List of options in this code.
--emig             File name of migration energy [eV] and vibrational prefactor [Hz] for every atomic jump. If it is
                   the default name (emig.csv), this argument is not necessary.
--lb_T            Lower bound of the temperature range [K]. Default value is 300.
--ub_T            Upper bound of the temperature range [K]. Default value is 1000.
--int_T           Interval of the temperature step [K]. Default value is 100.
```

When typing the following command, jmpdata\_300K.csv, jmpdata\_400K.csv, and jmpdata\_500K.csv are generated.

```
python [MASTEQ_DIR]/tools/mkjmpdata.py --emig emig.csv --lb_T 300 --ub_T 500 --int_T 100
```




Fig. 1. Migration paths and estimated diffusivity of a proton in BaZrO<sub>3</sub>.

**mkemig.py (pymatgen is required!)**

This program makes *emig.csv* from only the information on crystallographically non-equivalent atomic jumps. Specifically, two files are required, *site.cif* with site information of diffusion carriers and *emig\_noneq.csv* with jump information including the initial and final site types, jump distance [ang.], migration energy [eV], and vibrational prefactor [Hz]. All equivalent sites in the unitcell are generated using symmetry operations in *site.cif*, and the equivalent atomic jumps are then searched by the jump distance and the initial and final site types specified in *emig\_noneq.csv*.

For reading *site.cif* by *pymatgen*, non-equivalent site types of diffusion carriers have to be distinguished by *element symbols* (H, He, Li, Be, ...), whatever the diffusion carriers are. The example of site information line is shown below.

(Site information example in *site.cif*)

```
loop_
  _atom_site_label
  _atom_site_occupancy
  _atom_site_fract_x
  _atom_site_fract_y
  _atom_site_fract_z
  _atom_site_type_symbol
H      1.0    0.000000  0.000000  0.000000    H
He     1.0    0.500000  0.500000  0.500000   He
Li     1.0    0.000000  0.500000  0.500000    Li
```

Concerning the file format of *emig\_noneq.csv*, non-equivalent atomic jumps are specified by the initial and final site types (element symbols in *site.cif*), the jump distance [ang.],  $\Delta E^{\text{mig}}$  [eV], and  $\nu_0$  [Hz], separated by comma. New line for different atomic jumps. Both jumps in the opposite directions have to be specified separately, if they are non-equivalent. “#” denotes a comment line. The file example is as follows:

(*emig\_noneq.csv* example)

```
#InitialSiteType,FinalSiteType,jumpDistance[Ang.],DEmig[eV],v0[Hz]
H,Li,1.41421356,0.30,1.0e+13
Li,H,1.41421356,0.20,1.0e+13
He,Li,1.00000000,0.35,1.0e+13
Li,He,1.00000000,0.30,1.0e+13
```

Type the following command at the directory with *site.cif* and *emig\_noneq.csv*, *emig.csv* is generated.

```
python [MASTEQ_DIR]/tools/mkemig.py -cif site.cif -emig_noneq emig_noneq.csv --supercell 1,1,1 --prec 1.0e-4
```

The option of this program is as follows:

- h, --help            Help information. List of options in this code.
- cif                File name of the cif file with site information. If it is the default name (site.cif), this argument is not necessary.
- emig\_noneq        File name with information on non-equivalent atomic jumps. Initial and final site type, jump vector, Emig, and vibrational prefactor in the csv format. New line for non-equivalent atomic jumps. Both jumps in the opposite directions have to be specified if they are non-equivalent. Site types have to be specified by ELEMENT SYMBOL! e.g., H, He, Li, Be, B, etc. If it is the default name (emig\_noneq.cif), this argument is not necessary.
- supercell        Supercell size for expanding unitcell. Since the inter-site distance is defined as the 1NN distance with periodicity, longer jump vectors than the half of lattice vectors are not accepted in this program. '3,3,3' is enough even for a small cell with few sites. Default value: 1,1,1
- prec               Precision used in symmetry operation and atomic jump search. Default value: 1.0e-4

## [Theoretical background]

### Master equation

Under the independent-particle approximation, the master equation corresponding to the balance of the existence probability  $p_i$  of a single particle at site  $i$  is given by

$$\frac{\partial p_i(t)}{\partial t} = \sum_j [\Gamma_{ji} p_j(t) - \Gamma_{ij} p_i(t)],$$

where  $t$  is the time, and  $\Gamma_{ij}$  is the jump frequency between sites  $i$  and  $j$ . In the case of no path for atomic jump between sites  $i$  and  $j$ ,  $\Gamma_{ij}$  is zero. The first and second terms on the right side correspond to the inflow and outflow of the existence probability, respectively. The jump frequency matrix  $\mathbf{\Gamma}$  is defined as the negative of the Laplacian matrix for a weighted directed graph, in which the off-diagonal elements are  $\Gamma_{ij}$  and the diagonal elements are  $-\sum_{j \neq i} \Gamma_{ij}$ . Using the matrix  $\mathbf{\Gamma}$ , the above master equation can be expressed as

$$\frac{\partial \mathbf{p}}{\partial t} = \mathbf{\Gamma}^T \mathbf{p},$$

where  $\mathbf{p}$  is the vector of the existence probabilities of the single particle at all sites,  $\mathbf{p} = [p_1(t), \dots, p_N(t)]^T$  ( $N$ : number of sites in the system). The solution of the master equation is expressed using a given initial condition  $\mathbf{p}_0$  at  $t = 0$ ,

$$\mathbf{p} = \exp(\mathbf{\Gamma}^T t) \mathbf{p}_0.$$

In a crystal, the dimensions of the matrix  $\mathbf{\Gamma}$  and the vector  $\mathbf{p}$  can be reduced by the translational symmetry. The master equation is rewritten as

$$\frac{\partial p_i(\mathbf{r}, t)}{\partial t} = \sum_{(j, \alpha) \in A_i} [\Gamma_{ji}^\alpha p_j(\mathbf{r} + \mathbf{s}_{ij}^\alpha, t) - \Gamma_{ij}^\alpha p_i(\mathbf{r}, t)],$$

where  $i$  is the site index in the unitcell ( $i = 1, \dots, n$ ),  $p_i(\mathbf{r}, t)$  is the existence probability of the single particle at site  $i$  as a function of position  $\mathbf{r}$  and time  $t$ , and  $A_i$  is the set of all adjacent sites to site  $i$ . The set  $A_i$  includes adjacent sites in different unitcells from the focused unitcell beyond the periodic boundaries, if any. Therefore, when site  $i$  has several adjacent sites  $j$  in different unitcells, they are distinguished by the unitcell index  $\alpha$ .  $\Gamma_{ij}^\alpha$  and  $\mathbf{s}_{ij}^\alpha$  are the jump frequency and jump vector from site  $i$  in the focused unitcell to site  $j$  in unitcell  $\alpha$ , respectively.

This master equation is generally solved in Fourier space. With the Fourier transform,  $p_i(\mathbf{r}, t)$  is transformed into  $P_i(\mathbf{Q}, t)$  ( $= \int \exp(i\mathbf{Q}\mathbf{r}) p_i(\mathbf{r}, t) d\mathbf{r}$ ), where  $\mathbf{Q}$  is the Fourier variable for position  $\mathbf{r}$ , resulting in the following master equation in Fourier space,

$$\frac{\partial P_i(\mathbf{Q}, t)}{\partial t} = \sum_{(j, \alpha) \in A_i} [\Gamma_{ji}^\alpha P_j(\mathbf{Q}, t) \exp(-i\mathbf{Q}\mathbf{s}_{ij}^\alpha) - \Gamma_{ij}^\alpha P_i(\mathbf{Q}, t)].$$

Defining the jump matrix  $\mathbf{\Lambda}$  as the  $n \times n$  matrix with the elements  $\Lambda_{ij}$ ,

$$\Lambda_{ij} = \sum_\alpha \Gamma_{ij}^\alpha \exp(i\mathbf{Q}\mathbf{s}_{ij}^\alpha) - \delta_{ij} \sum_{j', \alpha} \Gamma_{ij'}^\alpha \quad (\delta_{ij}: \text{Kronecker delta}),$$

the master equation in Fourier space is simply expressed as

$$\frac{\partial \mathbf{P}}{\partial t} = \mathbf{\Lambda}^T \mathbf{P},$$

where  $\mathbf{P}$  is the vector of the existence probabilities of the single particle at all  $n$  sites in Fourier space. The solution of the master equation is also expressed as  $\exp(\mathbf{\Lambda}^T t) \mathbf{P}_0$  ( $\mathbf{P}_0$ : existence probability vector in Fourier space at  $t = 0$ ). When  $\mathbf{\Lambda}^T$  is eigendecomposed into  $\mathbf{X}\mathbf{Y}\mathbf{X}^{-1}$  ( $\mathbf{Y}$ : diagonal matrix with eigenvalues  $\lambda_i$ ,  $\mathbf{X}$ : transformation matrix), the

solution can finally be transformed as follows:

$$\mathbf{P} = \exp(\mathbf{X}\mathbf{Y}\mathbf{X}^{-1}t)\mathbf{P}_0 = \mathbf{X}\exp(\mathbf{Y}t)\mathbf{X}^{-1}\mathbf{P}_0 = \mathbf{X} \begin{pmatrix} e^{\lambda_1 t} & & 0 \\ & \ddots & \\ 0 & & e^{\lambda_n t} \end{pmatrix} \mathbf{X}^{-1}\mathbf{P}_0.$$

To connect the above solution with the diffusion coefficient tensor  $\mathbf{D}$ , the Fick's second law,  $\frac{\partial p(\mathbf{r},t)}{\partial t} = \nabla[\mathbf{D}\nabla p(\mathbf{r},t)]$ , is solved also in Fourier space, where  $p(\mathbf{r},t)$  is the existence probability distribution of a particle at position  $\mathbf{r}$  and time  $t$ . Under the initial condition,  $p(\mathbf{r},0) = \delta(\mathbf{r})$ , the solution is given by

$$P(\mathbf{Q},t) = \exp\left[-\sum_{m,n=x,y,z} D_{mn} Q_m Q_n t\right],$$

where  $D_{mn}$  and  $Q_m$  are the elements of the  $\mathbf{D}$  tensor and the  $\mathbf{Q}$  vector, respectively. Considering the time and spatial scales of atomic diffusion ( $t \rightarrow \infty$ ,  $|Q| \rightarrow 0$ ),  $-\sum_{m,n=x,y,z} D_{mn} Q_m Q_n$  coincides with the minimum-magnitude eigenvalue of matrix  $\mathbf{\Lambda}^T$ . Note that all eigenvalues are negative real numbers due to the mathematical property of matrix  $\mathbf{\Lambda}^T$  relevant to the Laplacian matrix in graph theory.