

演習 6-1

$$\int_0^x \frac{\pi-t}{2} dt = \int_0^x \sum_{n=1}^{\infty} \frac{\sin(nt)}{n} dt.$$

$$\left[\frac{\pi}{2}t - \frac{t^2}{4} \right]_0^x = \sum_{n=1}^{\infty} \int_0^x \frac{1}{n} \sin(nt) dt$$

$$\frac{\pi}{2}x - \frac{x^2}{4} = \sum_{n=1}^{\infty} \frac{1}{n} \left[-\frac{\cos(nt)}{n} \right]_0^x$$

$$\frac{\pi}{2}x - \frac{x^2}{4} = \sum_{n=1}^{\infty} \frac{1 - \cos(nx)}{n^2}$$

$$= \sum_{n=1}^{\infty} \frac{1}{n^2} - \sum_{n=1}^{\infty} \frac{\cos(nx)}{n^2}$$

演習 6-2

$$\sum_{n=1}^{\infty} \frac{\cos(nx)}{n^2} = \frac{\pi^2}{6} - \frac{\pi}{2}x + \frac{x^2}{4} = \frac{3x^2 - 6\pi x + 2\pi^2}{12}$$

$$x = \frac{\pi}{2} \text{ とおく。}$$

$$-\frac{1}{2^2} + \frac{1}{4^2} - \frac{1}{6^2} + \dots = \frac{3\left(\frac{\pi}{2}\right)^2 - 6\pi\left(\frac{\pi}{2}\right) + 2\pi^2}{12} = -\frac{\pi^2}{48}$$

$$-\left[\frac{1}{4 \cdot 1^2} - \frac{1}{4 \cdot 2^2} + \frac{1}{4 \cdot 3^2} + \dots \right] = -\frac{\pi^2}{48}$$

$$\frac{1}{1^2} - \frac{1}{2^2} + \frac{1}{3^2} + \dots = \frac{\pi^2}{12}$$