

演習 9

$$f(t) = \begin{cases} E & 0 \leq t < 2d \\ 0 & t < 0, t \geq 2d \end{cases}$$

$$F(\omega) = \int_{-\infty}^{\infty} f(t) e^{-i\omega t} dt$$

$$= \int_0^{2d} E \cdot e^{-i\omega t} dt$$

$$= \frac{E}{-i\omega} [e^{-i\omega t}]_0^{2d}$$

$$= -\frac{E}{i\omega} (e^{-i2\omega d} - 1)$$

$$= -\frac{E}{i\omega} e^{-i\omega d} (e^{-i\omega d} - e^{+i\omega d})$$

$$= 2E \cdot (2d - 0) \cdot \frac{\sin \omega d}{\omega d} \exp[-i\omega d]$$

$$= 4Ed \cdot \frac{\sin \omega d}{\omega d} \cdot e^{-i\omega d}$$

$$|G(w)| = Ed \left| \frac{\sin wd}{wd} \right| \text{ の零点は、}$$

$$\sin wd = 0$$

を無限可周級数に表す。つまり、

$$w_0 d = \pm \pi, \pm 2\pi, \pm 3\pi, \dots$$

$$w_0 = \pm \frac{\pi}{d}, \pm \frac{2\pi}{d}, \pm \frac{3\pi}{d}$$

$$\theta(w) = -wd.$$

$$\theta_0\left(\frac{\pi}{d}\right) = -\pi \left(\frac{d+\beta}{d-\beta} \right)$$

$$= -\pi \left(\frac{2d}{2d} \right)$$

$$= -\pi.$$