

# Bounding the rank of SDP

Lijun Ding

January 29, 2019

## 1 SDP and Why Low Rank Solutions

- What is SDP?
- Why Low Rank Solutions?

## 2 Bounding the rank via number of constraints ( $m$ )

- Bound of an exact solution
- Bound of an approximate solution

## 3 Questions

# Semidefinite Programming (SDP)

A semidefinite program is an optimization problem of the form:

$$\begin{aligned} & \underset{X \in \mathbb{S}^n}{\text{minimize}} && \mathbf{trace}(A_0 X) \\ & \text{subject to} && \mathbf{trace}(A_i X) = b_i, \quad i = 1, \dots, m, \\ & && X \succeq 0, \end{aligned} \tag{P}$$

where  $A_i \in \mathbb{S}^n \subset \mathbb{R}^{n \times n}$ ,  $b \in \mathbb{R}^m$  are problem data.

# Semidefinite Programming (SDP)

A semidefinite program is an optimization problem of the form:

$$\begin{aligned} & \underset{X \in \mathbb{S}^n}{\text{minimize}} && \mathbf{trace}(A_0 X) \\ & \text{subject to} && \mathbf{trace}(A_i X) = b_i, \quad i = 1, \dots, m, \\ & && X \succeq 0, \end{aligned} \tag{P}$$

where  $A_i \in \mathbb{S}^n \subset \mathbb{R}^{n \times n}$ ,  $b \in \mathbb{R}^m$  are problem data.

Here  $X \succeq 0$  means  $X$  is positive semidefinite and symmetric, i.e., exists some  $k \in \mathbb{N}$  and  $F \in \mathbb{R}^{n \times k}$  such that

$$X = FF^\top.$$

# Applications of SDP

SDP is applicable to a variety of problems: meaning

- analyzing problem,
- providing useful solutions computationally.

An incomplete list of application includes

- Combinatorial optimization: Max-Cut, TSP
- Optics: Phase Retrieval
- Recommendation System: Matrix Completion
- Power Systems: Power flow
- Machine Learning: Community Detection
- Statistics: Experiment Design

Also see [Vandenberghe and Boyd 96] for more applications.

# A detailed example: EDMC Set-up

Euclidean Distance Matrix Completion (EDMC):

- ①  $n$  many vectors  $\bar{v}_i \in \mathbb{R}^r$
- ② An index set  $\Omega \subset \{1, \dots, n\} \times \{1, \dots, n\}$
- ③ Pairwise distance  $d_{ij}^2 = \|\bar{v}_i - \bar{v}_j\|^2, (i, j) \in \Omega$

# A detailed example: EDMC Set-up

Euclidean Distance Matrix Completion (EDMC):

- 1  $n$  many vectors  $\bar{v}_i \in \mathbb{R}^r$
- 2 An index set  $\Omega \subset \{1, \dots, n\} \times \{1, \dots, n\}$
- 3 Pairwise distance  $d_{ij}^2 = \|\bar{v}_i - \bar{v}_j\|^2, (i, j) \in \Omega$

Q: Can we recover the matrix  $D = [d_{ij}^2]$  from  $d_{ij}, (i, j) \in \Omega$  with

$$|\Omega| \ll \frac{n(n+1)}{2}, \text{ e.g., } |\Omega| = \mathcal{O}(n)?$$

# A detailed example: EDMC Set-up

## Euclidean Distance Matrix Completion (EDMC):

- 1  $n$  many vectors  $\bar{v}_i \in \mathbb{R}^r$
- 2 An index set  $\Omega \subset \{1, \dots, n\} \times \{1, \dots, n\}$
- 3 Pairwise distance  $d_{ij}^2 = \|\bar{v}_i - \bar{v}_j\|^2, (i, j) \in \Omega$

Q: Can we recover the matrix  $D = [d_{ij}^2]$  from  $d_{ij}, (i, j) \in \Omega$  with

$$|\Omega| \ll \frac{n(n+1)}{2}, \text{ e.g., } |\Omega| = \mathcal{O}(n)?$$

- Important in sensor network
- Infer all pairwise distance from a few pairwise distance
- $r = 3$  usually



# A detail example : Convexification to SDP

Consider the variable matrix:

$$V = [v_1, \dots, v_n] \in \mathbb{R}^{r \times n},$$

and

$$X = V^T V \in \mathbb{S}^n, X \succeq 0, \quad \text{and} \quad \mathbf{rank}(\bar{X}) \leq r$$

In particular, when  $V = \bar{V} = [\bar{v}_1, \dots, \bar{v}_n]$ ,  $\bar{X} = \bar{V}^T \bar{V}$ .

# A detail example : Convexification to SDP

Consider the variable matrix:

$$V = [v_1, \dots, v_n] \in \mathbb{R}^{r \times n},$$

and

$$X = V^T V \in \mathbb{S}^n, X \succeq 0, \quad \text{and} \quad \mathbf{rank}(\bar{X}) \leq r$$

In particular, when  $V = \bar{V} = [\bar{v}_1, \dots, \bar{v}_n]$ ,  $\bar{X} = \bar{V}^T \bar{V}$ .

The condition based on  $d_{ij}^2$  is

$$d_{ij}^2 = \|v_i - v_j\|_2^2 = \|v_i\|_2^2 + \|v_j\|_2^2 - 2v_i^T v_j = x_{ii} + x_{jj} - 2x_{ij}.$$

# A detail example : Convexification to SDP

Consider the variable matrix:

$$V = [v_1, \dots, v_n] \in \mathbb{R}^{r \times n},$$

and

$$X = V^T V \in \mathbb{S}^n, X \succeq 0, \quad \text{and} \quad \text{rank}(\bar{X}) \leq r$$

In particular, when  $V = \bar{V} = [\bar{v}_1, \dots, \bar{v}_n]$ ,  $\bar{X} = \bar{V}^T \bar{V}$ .

The condition based on  $d_{ij}^2$  is

$$d_{ij}^2 = \|v_i - v_j\|_2^2 = \|v_i\|_2^2 + \|v_j\|_2^2 - 2v_i^T v_j = x_{ii} + x_{jj} - 2x_{ij}.$$

Convexification to SDP: expect  $\bar{X}$  to be a solution (up to a shift in  $\bar{V}$ )

Find	$X, V$	minimize	$\text{trace}(X)$
s.t.	$d_{ij}^2 = x_{ii} + x_{jj} - 2x_{ij}$ $(i, j) \in \Omega$ $X = V^T V$	subject to	$d_{ij}^2 = x_{ii} + x_{jj} - 2x_{ij}$ $(i, j) \in \Omega$ $X \succeq 0$

## 1 SDP and Why Low Rank Solutions

- What is SDP?
- Why Low Rank Solutions?

## 2 Bounding the rank via number of constraints ( $m$ )

- Bound of an exact solution
- Bound of an approximate solution

## 3 Questions

# Why Low Rank? Interpretability

*Low Rank* solutions  $X^*$  to SDP usually are more interpretable.

- ① Euclidean distance matrix completion:  $\text{rank}(X^*) > \text{rank}(\bar{X})$ , hard to interpret. Expect  $\text{rank}(X^*) = \text{rank}(\bar{X})$ .

# Why Low Rank? Interpretability

*Low Rank* solutions  $X^*$  to SDP usually are more interpretable.

- 1 Euclidean distance matrix completion:  $\text{rank}(X^*) > \text{rank}(\bar{X})$ , hard to interpret. Expect  $\text{rank}(X^*) = \text{rank}(\bar{X})$ .
- 2 Combinatorial optimization:  $X^* = xx^\top$  and  $x = \{\pm 1\}^n$ , binary assignment.
- 3 Phase Retrieval:  $X^* = xx^\top$ ,  $x \in \mathbb{R}^n$  where  $x$  stands for an approximation of the underlying phase/object.

# Why Low Rank? Computational opportunity

Recall  $X^* \in \mathbb{S}^n$ . Let  $r^* = \mathbf{rank}(X^*)$  with  $r^* \ll n$ . Efficient Algorithm opportunity:

# Why Low Rank? Computational opportunity

Recall  $X^* \in \mathbb{S}^n$ . Let  $r^* = \mathbf{rank}(X^*)$  with  $r^* \ll n$ . Efficient Algorithm opportunity:

- Space:

	$X^* \in \mathbb{S}^n$	$X^* = FF^\top$
Space	$\mathcal{O}(n^2)$	$\mathcal{O}(nr^*)$



# Why Low Rank? Computational opportunity

Recall  $X^* \in \mathbb{S}^n$ . Let  $r^* = \mathbf{rank}(X^*)$  with  $r^* \ll n$ . Efficient Algorithm opportunity:

- Space:

	$X^* \in \mathbb{S}^n$	$X^* = FF^\top$
Space	$\mathcal{O}(n^2)$	$\mathcal{O}(nr^*)$

- Time:

	$X \in \mathbb{S}^n$	$X = FF^\top$
Matrix-vector product	$\mathcal{O}(n^2)$	$\mathcal{O}(nr^*)$

## 1 SDP and Why Low Rank Solutions

- What is SDP?
- Why Low Rank Solutions?

## 2 Bounding the rank via number of constraints ( $m$ )

- Bound of an exact solution
- Bound of an approximate solution

## 3 Questions

# Exact Bound: $\frac{r^*(r^*+1)}{2} \leq m + 1$

Recall the SDP (P):

$$\begin{array}{ll} \underset{X}{\text{minimize}} & \mathbf{trace}(A_0 X) \\ \text{subject to} & \mathbf{trace}(A_i X) = b_i, \ i = 1, \dots, m, \\ & X \succeq 0. \end{array}$$

Exact Bound:  $\frac{r^*(r^*+1)}{2} \leq m + 1$

Recall the SDP (P):

$$\begin{array}{ll} \underset{X}{\text{minimize}} & \mathbf{trace}(A_0 X) \\ \text{subject to} & \mathbf{trace}(A_i X) = b_i, \quad i = 1, \dots, m, \\ & X \succeq 0. \end{array}$$

Theorem 1 [Barvinok 95, Pataki 98]

If there is a solution to (P), then there is a solution  $X^*$  with rank  $r^*$  satisfying

$$\frac{r^*(r^* + 1)}{2} \leq m + 1$$

## Theorem 1 [Barvinok 95, Pataki 98]

If there is a solution to (P), then there is a solution  $X^*$  with rank  $r^*$  satisfying

$$\frac{r^*(r^* + 1)}{2} \leq m + 1$$

- 1 The bound on  $r^*$  depend only on  $m$ .

## Theorem 1 [Barvinok 95, Pataki 98]

If there is a solution to (P), then there is a solution  $X^*$  with rank  $r^*$  satisfying

$$\frac{r^*(r^* + 1)}{2} \leq m + 1$$

- 1 The bound on  $r^*$  depend only on  $m$ .
- 2 If  $m = \mathcal{O}(n)$ , e.g., EDMC  $|\Omega| = m = \mathcal{O}(n)$ , then  $r^* \ll n$ .

## Theorem 1 [Barvinok 95, Pataki 98]

If there is a solution to (P), then there is a solution  $X^*$  with rank  $r^*$  satisfying

$$\frac{r^*(r^* + 1)}{2} \leq m + 1$$

- ① The bound on  $r^*$  depend only on  $m$ .
- ② If  $m = \mathcal{O}(n)$ , e.g., EDMC  $|\Omega| = m = \mathcal{O}(n)$ , then  $r^* \ll n$ .
- ③ Rank grows  $\mathcal{O}(\sqrt{m})$  though actual  $r^*$  can be much lower.

## Theorem 1 [Barvinok 95, Pataki 98]

If there is a solution to (P), then there is a solution  $X^*$  with rank  $r^*$  satisfying


$$\frac{r^*(r^* + 1)}{2} \leq m + 1$$

- 1 The bound on  $r^*$  depend only on  $m$ .
- 2 If  $m = \mathcal{O}(n)$ , e.g., EDMC  $|\Omega| = m = \mathcal{O}(n)$ , then  $r^* \ll n$ .
- 3 Rank grows  $\mathcal{O}(\sqrt{m})$  though actual  $r^*$  can be much lower.
- 4 Existence of a “low” rank solution but not every solution is “low rank”.



# Exact bound: Remarks

- Existence of a “low” rank solution but not every solution is “low rank”.



## [Barvinok 95]

- ① Motivation: a graph problem which now are considered as *Euclidean distance matrix completion*
- ② Proof strategy: relies on convex duality theory
- ③ Technique: Need results from algebraic geometry and semialgebraic set.

## [Barvinok 95]

- ① Motivation: a graph problem which now are considered as *Euclidean distance matrix completion*
- ② Proof strategy: relies on convex duality theory
- ③ Technique: Need results from algebraic geometry and semialgebraic set.

## [Pataki 98]

- ① Motivation: eigenvalue optimization
- ② Proof strategy: direct bound the rank of  $X^*$ .
- ③ Technique: Linear algebra (only)
- ④ Generality: More general, deals with all solutions of (P)

# Rank-Nullity Theorem

Given a linear map  $\Psi : \mathbb{R}^l \rightarrow \mathbb{R}^h$ , define the null space and image space as

$$\mathbf{nullspace}(\Psi) := \{x \in \mathbb{R}^l \mid \Psi(x) = 0\}, \quad \mathbf{im}(\Psi) := \{\Psi(x) \in \mathbb{R}^h \mid x \in \mathbb{R}^l\}.$$

Recall the rank-nullity theorem says that

$$\dim(\mathbf{nullspace}(\Psi)) + \dim(\mathbf{im}(\Psi)) = l.$$

In particular, if  $h < l$ , then  $\mathbf{nullspace}(\Psi) \neq \emptyset$ .

# Exact Bound: Proof (inspired by [Pataki 98])

- 1 Given a solution  $\bar{X} = FF^\top$ , where  $F \in \mathbb{R}^{n \times k}$

# Exact Bound: Proof (inspired by [Pataki 98])

- ① Given a solution  $\bar{X} = FF^\top$ , where  $F \in \mathbb{R}^{n \times k}$
- ② If  $\frac{k(k+1)}{2} \leq m+1$ , we are done. Otherwise, consider the linear map

$$\psi_F : S \in \mathbb{S}^k \rightarrow \begin{bmatrix} \text{trace}(A_0 F S F^\top) \\ \text{trace}(A_1 F S F^\top) \\ \vdots \\ \text{trace}(A_m F S F^\top) \end{bmatrix} \in \mathbb{R}^{m+1}.$$

# Exact Bound: Proof (inspired by [Pataki 98])

- ① Given a solution  $\bar{X} = FF^\top$ , where  $F \in \mathbb{R}^{n \times k}$
- ② If  $\frac{k(k+1)}{2} \leq m+1$ , we are done. Otherwise, consider the linear map

$$\psi_F : S \in \mathbb{S}^k \rightarrow \begin{bmatrix} \text{trace}(A_0 F S F^\top) \\ \text{trace}(A_1 F S F^\top) \\ \vdots \\ \text{trace}(A_m F S F^\top) \end{bmatrix} \in \mathbb{R}^{m+1}.$$

- ③ Since  $\frac{k(k+1)}{2} > m+1$ , rank-nullity theorem tells us that exists  $S \neq 0$ , such that  $\psi_F(S) = 0$

# Exact Bound: Proof (inspired by [Pataki 98])

- ① Given a solution  $\bar{X} = FF^\top$ , where  $F \in \mathbb{R}^{n \times k}$
- ② If  $\frac{k(k+1)}{2} \leq m+1$ , we are done. Otherwise, consider the linear map

$$\Psi_F : S \in \mathbb{S}^k \rightarrow \begin{bmatrix} \text{trace}(A_0 F S F^\top) \\ \text{trace}(A_1 F S F^\top) \\ \vdots \\ \text{trace}(A_m F S F^\top) \end{bmatrix} \in \mathbb{R}^{m+1}.$$

- ③ Since  $\frac{k(k+1)}{2} > m+1$ , rank-nullity theorem tells us that exists  $S \neq 0$ , such that  $\Psi_F(S) = 0$
- ④ Consider  $X_\alpha = F(I + \alpha S)F^\top$  for  $\alpha \in \mathbb{R}$ , which is feasible and optimal for all small  $\alpha$  (as  $I \succ 0$ )



# Exact Bound: Proof (inspired by [Pataki 98])

- ① Given a solution  $\bar{X} = FF^\top$ , where  $F \in \mathbb{R}^{n \times k}$
- ② If  $\frac{k(k+1)}{2} \leq m+1$ , we are done. Otherwise, consider the linear map

$$\psi_F : S \in \mathbb{S}^k \rightarrow \begin{bmatrix} \text{trace}(A_0 F S F^\top) \\ \text{trace}(A_1 F S F^\top) \\ \vdots \\ \text{trace}(A_m F S F^\top) \end{bmatrix} \in \mathbb{R}^{m+1}.$$


- ③ Since  $\frac{k(k+1)}{2} > m+1$ , rank-nullity theorem tells us that exists  $S \neq 0$ , such that  $\psi_F(S) = 0$
- ④ Consider  $X_\alpha = F(I + \alpha S)F^\top$  for  $\alpha \in \mathbb{R}$ , which is feasible and optimal for all small  $\alpha$  (as  $I \succ 0$ )
- ⑤ Set  $-\frac{1}{\alpha}$  be a certain eigenvalue of  $S$  to reduce the rank and repeat step 1-4 until  $\frac{k(k+1)}{2} \leq m+1$ .

# Exact bound: Illustration of Step 5

- Set  $\frac{1}{\alpha}$  be a certain eigenvalue of  $S$  to reduce the rank


Eigenvalue of  
 $S \in \mathbb{S}^k$

Case  
1:



$$\text{Set } \alpha = -\frac{1}{\lambda_{\max}(S)},$$
$$\text{rank}(I + \alpha S) < k,$$
$$I + \alpha S \geq 0$$

Case  
2:



$$\text{Set } \alpha = -\frac{1}{\lambda_{\min}(S)},$$
$$\text{rank}(I + \alpha S) < k,$$
$$I + \alpha S \geq 0$$

## 1 SDP and Why Low Rank Solutions

- What is SDP?
- Why Low Rank Solutions?

## 2 Bounding the rank via number of constraints ( $m$ )

- Bound of an exact solution
- Bound of an approximate solution

## 3 Questions

Approximation bound:  $\mathcal{O}\left(\frac{\ln(m)}{\epsilon^2}\right)$

What if we look for an approximate solution?

# Approximation bound: $\mathcal{O}\left(\frac{\ln(m)}{\epsilon^2}\right)$

What if we look for an approximate solution?

## Theorem 2 [Barvinok 2002]

Fix an approximation level  $\epsilon \in (0, 1)$ . If  $A_i$ s are positive semidefinite and (P) admits a solution  $X^*$  with optimal value  $b_0$ , then there is a matrix  $X_0 \succeq 0$  such that for all  $0 \leq i \leq m$

$$\text{trace}(A_i X_0) \in b_i[1 - \epsilon, 1 + \epsilon]$$

and

$$\text{rank}(X_0) \leq \frac{8}{\epsilon^2} \ln(4(m + 1))$$

*Fact:* for any  $A, B \succeq 0$ ,  $\text{trace}(AB) \geq 0$ .

# Approximation: Remarks

- ① Dependence comparison:

Theorem 1	Theorem 2
$\mathcal{O}(\sqrt{m})$	$\mathcal{O}(\frac{\ln m}{\epsilon^2})$

# Approximation: Remarks

- ① Dependence comparison:

Theorem 1	Theorem 2
$\mathcal{O}(\sqrt{m})$	$\mathcal{O}(\frac{\ln m}{\epsilon^2})$

- ② Genreal version: Exists but more complicated

# Approximation: Remarks

- ① Dependence comparison:

Theorem 1	Theorem 2
$\mathcal{O}(\sqrt{m})$	$\mathcal{O}(\frac{\ln m}{\epsilon^2})$

- ② Genreal version: Exists but more complicated
- ③  $\mathcal{O}(\frac{1}{\epsilon^2})$  Dependence: Not great. Similar dependence as Johnson-Lindenstrauss Lemma



# Approximation: Remarks

- ① Dependence comparison:

Theorem 1	Theorem 2
$\mathcal{O}(\sqrt{m})$	$\mathcal{O}(\frac{\ln m}{\epsilon^2})$

- ② Genreal version: Exists but more complicated
- ③  $\mathcal{O}(\frac{1}{\epsilon^2})$  Dependence: Not great. Similar dependence as Johnson-Lindenstrauss Lemma
- ④ Proof technique in [Barvinok 2002]: Semidefinite quadratic form and Gaussian measures.

# Approximation: Key Lemma

## (Generalized) Johnson Lindenstrauss Lemma

Fix an approximation level  $\epsilon \in (0, 1)$ , given  $m$  matrices  $F_i \in \mathbb{R}^{n \times d_i}$ ,  $i = 1, \dots, m$ ,  $d_i \leq n$ , there is a matrix  $U \in \mathbb{R}^{r \times n}$  with  $r = \frac{8 \ln(4m)}{\epsilon^2}$  such that for all  $i$ ,

$$\|UF_i\|_F^2 \in \|F_i\|_F^2 [1 - \epsilon, 1 + \epsilon].$$

- Take  $d = 1$  reduces to normal J-L Lemma.
- Normal J-L will make the  $r = \mathcal{O}(\frac{\ln(m + \sum d_i)}{\epsilon^2})$ .

# Approximation: Proof based on (generalized) J-L Lemma

① First suppose the solution  $X^* = I$ . Then

$$b_i = \mathbf{trace}(A_i X^*) = \mathbf{trace}(A_i), \quad i = 0, 1, \dots, m.$$

# Approximation: Proof based on (generalized) J-L Lemma

- ① First suppose the solution  $X^* = I$ . Then

$$b_i = \mathbf{trace}(A_i X^*) = \mathbf{trace}(A_i), \quad i = 0, 1, \dots, m.$$

- ② Since  $A_i \succeq 0$ , exists  $F_i \in \mathbb{R}^{n \times d_i}$  such that

$$A_i = F_i F_i^\top, \quad i = 0, \dots, m.$$

# Approximation: Proof based on (generalized) J-L Lemma

- ① First suppose the solution  $X^* = I$ . Then

$$b_i = \mathbf{trace}(A_i X^*) = \mathbf{trace}(A_i), \quad i = 0, 1, \dots, m.$$

- ② Since  $A_i \succeq 0$ , exists  $F_i \in \mathbb{R}^{n \times d_i}$  such that

$$A_i = F_i F_i^\top, \quad i = 0, \dots, m.$$

- ③ Thus

$$b_i = \mathbf{trace}(F_i F_i^\top) = \|F_i\|_F^2, \quad i = 0, \dots, m.$$

# Approximation: Proof based on (generalized) J-L Lemma

- ① First suppose the solution  $X^* = I$ . Then

$$b_i = \mathbf{trace}(A_i X^*) = \mathbf{trace}(A_i), \quad i = 0, 1, \dots, m.$$

- ② Since  $A_i \succeq 0$ , exists  $F_i \in \mathbb{R}^{n \times d_i}$  such that

$$A_i = F_i F_i^\top, \quad i = 0, \dots, m.$$

- ③ Thus

$$b_i = \mathbf{trace}(F_i F_i^\top) = \|F_i\|_F^2, \quad i = 0, \dots, m.$$

- ④ Using generalized J-L to conclude there exists  $U \in \mathbb{R}^{r \times n}$  with  $r = \frac{8 \ln(4(m+1))}{\epsilon^2}$  such that

$$\|UF_i\|_F^2 \in \|F_i\|_F^2[1 - \epsilon, 1 + \epsilon] = b_i[1 - \epsilon, 1 + \epsilon].$$

# Approximation: Proof based on J-L Lemma conti.

① We have  $U \in \mathbb{R}^{r \times n}$  with  $r = \frac{8 \ln(4(m+1))}{\epsilon^2}$  and

$$\|UF_i\|_F^2 \in \|F_i\|_F^2[1 - \epsilon, 1 + \epsilon] = b_i[1 - \epsilon, 1 + \epsilon]$$

# Approximation: Proof based on J-L Lemma conti.

- ① We have  $U \in \mathbb{R}^{r \times n}$  with  $r = \frac{8 \ln(4(m+1))}{\epsilon^2}$  and

$$\|UF_i\|_F^2 \in \|F_i\|_F^2[1 - \epsilon, 1 + \epsilon] = b_i[1 - \epsilon, 1 + \epsilon]$$

- ② Now set  $X_0 = U^\top U$  with rank no larger than  $\frac{8 \ln(4(m+1))}{\epsilon^2}$ , and for all  $i$

$$\mathbf{trace}(A_i X_0) = \mathbf{trace}(F_i F_i^\top U^\top U) = \mathbf{trace}((UF_i)^\top UF_i) = \|UF_i\|_F^2.$$



# Approximation: Proof based on J-L Lemma conti.

- ① We have  $U \in \mathbb{R}^{r \times n}$  with  $r = \frac{8 \ln(4(m+1))}{\epsilon^2}$  and

$$\|UF_i\|_F^2 \in \|F_i\|_F^2[1 - \epsilon, 1 + \epsilon] = b_i[1 - \epsilon, 1 + \epsilon]$$

- ② Now set  $X_0 = U^\top U$  with rank no larger than  $\frac{8 \ln(4(m+1))}{\epsilon^2}$ , and for all  $i$

$$\mathbf{trace}(A_i X_0) = \mathbf{trace}(F_i F_i^\top U^\top U) = \mathbf{trace}((UF_i)^\top UF_i) = \|UF_i\|_F^2.$$

- ③ For general  $X^*$ , consider  $X^* = FF^\top$  and

$$A'_i = F^\top A_i F, \quad i = 1, \dots, m.$$

# Approximation: Proof based on J-L Lemma conti.

- ① We have  $U \in \mathbb{R}^{r \times n}$  with  $r = \frac{8 \ln(4(m+1))}{\epsilon^2}$  and

$$\|UF_i\|_F^2 \in \|F_i\|_F^2[1 - \epsilon, 1 + \epsilon] = b_i[1 - \epsilon, 1 + \epsilon]$$

- ② Now set  $X_0 = U^\top U$  with rank no larger than  $\frac{8 \ln(4(m+1))}{\epsilon^2}$ , and for all  $i$

$$\text{trace}(A_i X_0) = \text{trace}(F_i F_i^\top U^\top U) = \text{trace}((UF_i)^\top UF_i) = \|UF_i\|_F^2.$$

- ③ For general  $X^*$ , consider  $X^* = FF^\top$  and

$$A'_i = F^\top A_i F, \quad i = 1, \dots, m.$$

- ④ Repeat previous step for  $A'_i$  and get  $X'_0$ . Set  $X_0 = FX'_0F^\top$ .

## 1 SDP and Why Low Rank Solutions

- What is SDP?
- Why Low Rank Solutions?

## 2 Bounding the rank via number of constraints ( $m$ )

- Bound of an exact solution
- Bound of an approximate solution

## 3 Questions

- If a solution to (P) exists, there is a solution with rank  $r^*$  must satisfy

$$(a)r^* > \frac{1}{2}m \quad (b)r^* = \mathcal{O}(\sqrt{m}) \quad (c)r^* > \frac{1}{2}n \quad (d)r^* = \mathcal{O}(m^{\frac{1}{3}}).$$

- If a solution to (P) exists and all  $A_i \succeq 0$ , there is a matrix  $X_0$  with  $\epsilon$ -approximation level, i.e.,  $\text{trace}(A_i X_0) \in b_i[1 - \epsilon, 1 + \epsilon]$ , and has rank  $r$  no more than

$$(a)\mathcal{O}\left(\frac{\ln(m)}{\epsilon^2}\right) \quad (b)\mathcal{O}\left(\frac{\ln(\ln(m))}{\epsilon^2}\right) \quad (c)\mathcal{O}\left(\frac{\ln(m)}{\epsilon}\right) \quad (d)\mathcal{O}\left(\frac{\ln(n)}{\epsilon}\right)$$

# References



Vandenberghe, Lieven, and Stephen Boyd. "Semidefinite programming." SIAM review 38, no. 1 (1996): 49-95.



Barvinok, Alexander I. "Problems of distance geometry and convex properties of quadratic maps." Discrete & Computational Geometry 13, no. 2 (1995): 189-202.



Pataki, Gabor. "On the rank of extreme matrices in semidefinite programs and the multiplicity of optimal eigenvalues." Mathematics of operations research 23, no. 2 (1998): 339-358.



Barvinok, Alexander. A course in convexity. Vol. 54. American Mathematical Soc., 2002.