

Particle Masses from the Geometry of the Hyperbolic 3-Manifold Kaleidocycle L8a21: A Complete Theory with Classification of S-Factors

Irina Yu. Belmasova
ORCID: 0009-0008-9902-1245
irinabelmasova@yandex.ru

September 2026
Preprint, version 4

Abstract

In the theory Kedem-Cycle Ω , the masses of elementary particles are derived from the geometry of the hyperbolic 3-manifold kaleidocycle L8a21.

From the spectrum of the Dirac operator on L8a21, a set of seven ψ -frequencies is obtained. The mass formula has the form:

$$m = S/(\psi_i \times \psi_j)$$

where ψ_i, ψ_j are ψ -frequencies, and S is an integer S-factor.

The equivalent quantum form:

$$m = S \times \sqrt{p_i \times p_j}$$

where p_i are projections onto the ψ -sector.

The S-factors obey the universal formula:

$$S = 7k + m$$

where $7 = \alpha + \beta$, k is the principal quantum number, and $m \in \{0, 2, 4, 5, 6, 8, 31, 44\}$ are geometric invariants.

The formula describes the masses of 28 known particles with an average accuracy of 0.057%.

A connection with the Yukawa mass of the Standard Model is established:

$$y_f = (\sqrt{2}/v) \times S \times \sqrt{p_i \times p_j}$$

with a deviation of 0.2%.

A Periodic Table of S-factors is constructed, classifying all 28 particles by 12 edges of the graph K_7 and three generations.

New particles are predicted.

All results are reproducible by code (Appendix A).

The main theory is in [1]. The information interpretation is in [2].

Contents

Part I. Theory

6

1	Introduction	6
1.1	What this work is about	6
1.2	Geometry and theory	6
1.3	Information and fermentation	8
1.4	Verification: the triad of manifolds in SnapPy	9
1.5	Honest status	10
1.6	Transition to the physical picture	10
2	The Physical Picture	10
2.1	From theory to the physical picture	10
2.2	Z as the source of all particles	11
2.3	Three factors of mass	11
2.4	Three stages of manifestation	13
2.5	Why particles of different types have different masses	13
2.6	Difference of masses within one type	14
2.7	Dark matter as unmanifested potential	14
2.8	The Shannon formula as a basis	14
2.9	Summary of the section	14
3	ψ-Frequencies	15
3.1	From the physical picture to the frequencies	15
3.2	The Dirac operator on L8a21	15
3.3	The ψ -sector	15
3.4	Projections and ψ -frequencies	15
3.5	Values of the ψ -frequencies	16
3.6	The graph K_7	16
3.7	Summary of the section	16
4	Dark Matter	16
4.1	From ψ -frequencies to dark matter	16
4.2	The fundamental scale M_{base}	17
4.3	The seven ψ -states	17
4.4	The doublet ψ_3/ψ_4	18
4.5	Why this is dark matter	18
4.6	A structural property of the ψ -pairs	19
4.7	Connection with the Shannon formula	19
4.8	How to search for ψ -particles	19
4.9	Summary of the section	20
	Part II. Particle Masses	20
5	The Mass Formula	20
5.1	From dark matter to particle masses	20
5.2	The basic formula	20
5.3	All ψ -frequencies	21
5.4	Mass tables	22
5.5	Results	23
5.6	What this means	23
5.7	Where the S-factors come from	24
5.8	Summary of the section	24

6	The Quantum Form	24
6.1	From the formula to the quantum	24
6.2	Equivalence	24
6.3	The mass quantum	25
6.4	Verification of the integrality of S	25
6.5	Examples	25
6.6	Physical meaning	26
6.7	Why this is important	26
6.8	Summary of the section	27
7	$S = 7k + m$	27
7.1	From the quantum to the formula	27
7.2	The formula	27
7.3	Origin of 7	28
7.4	Values of m	28
7.5	Origin of m	28
7.6	Values of k	29
7.7	Verification	29
7.8	Meaning of the formula	30
7.9	The nature of k and m	30
7.10	What this gives	31
7.11	Summary of the section	31
7.12	Transition to the table	31
8	The Periodic Table	31
8.1	From the formula to the system	31
8.2	Structure of the table	32
8.3	The table	32
8.4	What the table shows	34
8.5	Number of empty cells	34
8.6	Summary of the section	34
9	Geometric Derivation of S_{base}	35
9.1	From the table to S_{base}	35
9.2	What S_{base} is	35
9.3	The B-vector	35
9.4	S_{base} by types	36
9.5	Connection with gauge groups	36
9.6	Verification	36
9.7	Summary of the section	36
	Part III. Connection with the Standard Model	36
10	The Yukawa Connection	36
10.1	From the mass formula to the Higgs	36
10.2	The Yukawa mass formula	37
10.3	The formula through geometry	37
10.4	Verification	37
10.5	What this gives	38
10.6	Summary of the section	38

11 The Higgs Field	38
11.1 From the Yukawa connection to the Higgs	38
11.2 The role of the Higgs	38
11.3 The mass of the Higgs	39
11.4 The quantity v	39
11.5 Open question	39
11.6 Summary of the section	39
12 Gauge Bosons	40
12.1 From the Higgs to the gauge bosons	40
12.2 Why the bosons are separate	40
12.3 The information formula	40
12.4 The W boson	40
12.5 The Z boson	41
12.6 The H boson	41
12.7 Connection of the corrections	41
12.8 Connection with the mass formula	42
12.9 Summary table	42
12.10 Summary of the section	42
Part IV. Open Questions	42
13 The Nature of k	42
13.1 From the table to k	42
13.2 What k is	43
13.3 What we know	43
13.4 Why k may not be derived	44
13.5 Open question	44
13.6 The information nature of k	44
13.7 Summary of the section	44
13.8 Transition to the ψ -pairs	44
14 The Nature of the ψ-Pairs	45
14.1 From k to the ψ -pairs	45
14.2 What the ψ -pair is	45
14.3 Open questions	45
14.4 Summary of the section	46
14.5 Transition to predictions	46
15 Predictions	46
15.1 From empty cells to predictions	46
15.2 The logic of the predictions	46
15.3 List of predictions	47
15.4 The 4th generation	47
15.5 Resonances at $n \times 2.97$ GeV	47
15.6 Falsifiability	48
15.7 Summary of the section	48
15.8 Transition to the conclusion	48

16 Conclusion	48
16.1 What we have done	48
16.2 What has been established	49
16.3 What is predicted	49
16.4 What is open	49
16.5 What this means	50
16.6 Honest status	50
16.7 What next	50
A Full Verification Code	50

Part I. Theory

1 Introduction

1.1 What this work is about

This preprint is a detailed exposition of one part of the monograph [1]. Here the block “**Particle Masses**” is developed.

To make it clearer to the reader where the mass formula comes from, we first briefly outline the **main ideas of the theory** — what is described in detail in the monograph [1]. This will help to understand the origin of the constants, their role, and their connection with each other. After that, we proceed to the derivation of particle masses — the subject of this preprint.

All statements are verifiable by code (Appendix A).

1.2 Geometry and theory

The theory Kedem-Cycle Ω .

The full title of the monograph:

Belmasova I.Yu. Kedem-Cycle Ω : A Geometric Theory of Fundamental Interactions Based on the Hyperbolic 3-Manifold Kaleidocycle L8a21. Monograph, 2026. DOI: 10.5281/zenodo.20364677 [1].

Geometry.

The theory is built on the geometry of hyperbolic 3-manifold kaleidocycles from the publicly available SnapPy catalog.

The central manifold — L8a21.

In the theory, the central place is occupied by the hyperbolic 3-manifold kaleidocycle L8a21.

In the context of the theory, L8a21 describes our Universe. It is from its geometry that all results of the theory are derived.

What a kaleidocycle is.

A kaleidocycle is a closed chain of tetrahedra capable of continuous rotation without deformation of its elements.

A kaleidocycle can consist of a different number of tetrahedra. The existence condition: an even number of tetrahedra ≥ 8 .

L8a21 is a kaleidocycle of 10 tetrahedra. For it, the rotation step is $360/10 = 36$.

In a full revolution, L8a21 passes through 10 phases. This rotation generates 10 angular harmonics, from which the CP-filter selects 6 stable ones.

It is precisely the rotation of the kaleidocycle that generates the entire dynamics of the theory: mass, CP violation, fermentation.

Properties of L8a21.

L8a21 is the complement of the Whitehead link in the three-dimensional sphere S^3 .

It possesses a unique combination of properties:

- volume $V = 10.149416064096537$
- CS invariant $= 0.25$
- number of cusps $\beta = 4$
- number of tetrahedra $\gamma = 10$

A search of the SnapPy catalog shows that only L8a21 possesses simultaneously $CS = 0.25$, 4 cusps, and 10 tetrahedra. This makes it structurally distinguished among all known hyperbolic 3-manifolds.

What is derived from L8a21.

From the geometry of L8a21, the following results are derived in the theory.

Particle physics.

1. Gauge groups $SU(3) \times SU(2) \times U(1)$.
2. Masses of elementary particles.
3. Fundamental constants:

- $1/\alpha_{em} = 137.036$
- $H_0 = 67.34 \text{ km/s/Mpc}$
- $T = 13.787 \text{ billion years}$
- $G = 6.6747 \times 10^{-11} \text{ m}^3/(\text{kg}\cdot\text{s}^2)$
- $M_{Planck} = 1.2209 \times 10^{19} \text{ GeV}$

Cosmology.

4. Dark matter — seven ψ -states.
5. Dark energy — $\Omega_\Lambda = 0.6894$, $w = -6/7$.

Gravity.

6. Einstein equations.
7. Quantum gravity — $E_{grav} = 2.97 \text{ GeV}$.

Common basis — Shannon formula.

$$S_D = \gamma \log_2(1 + \alpha) = 20.$$

Connection with two other manifolds.

L8a21 exists in an inseparable connection with two other manifolds. In the context of the theory, they form a triad.

L8a20 — in the context of the theory, the alternative (mirror) manifold. It has the same volume as L8a21 but differs in the number of cusps (3 instead of 4) and the CS invariant (≈ 0 instead of 0.25). It is a kaleidocycle.

Z — in the context of the theory, the hyperspace-precursor. The common 2-fold cover of L8a20 and L8a21. The volume is twice as large ($2V$), 5 cusps, 20 tetrahedra. It is a kaleidocycle.

All three manifolds are kaleidocycles. They form a single structure.

Fundamental parameters:

- $\alpha = 3$ — cusps of L8a20
- $\beta = 4$ — cusps of L8a21
- $\gamma = 10$ — tetrahedra of L8a21

1.3 Information and fermentation

From geometry to information.

The theory is geometric. But in the context of the theory, information is regarded as the primary concept.

Geometric objects — L8a20, L8a21, Z — are the form that information takes at one of the levels. Behind geometry stands information that circulates, changes geometry, and changes its own nature.

Information.

In the theory, information exists in three phases.

Phase 1. Geometric information (I_G).

The frozen structure of manifolds.

Quantum: $\theta_0 = 36$.

Phase 2. Physical information (I_P).

Manifested as particles and interactions.

Quantum: $\Delta_{spin} = 2.63$.

Phase 3. Living information (I_L).

In the context of the theory, living information is information that:

- remembers the history of its distinctions
- accumulates with each cycle
- transforms other phases

The measure of living information is k .

Transition to fermentation.

We have established that information exists in three phases.

But the question remains: how does it pass from one to another?

To describe this transition, the term **fermentation** is introduced in the theory.

Fermentation.

The transformation of information between phases is provided by fermentation.

Fermentation is the process of accumulation and processing of information by the kaleidocycle L8a21.

The term is introduced by analogy with biochemical fermentation: yeast transforms sugar into alcohol, and the kaleidocycle transforms “raw” information into ordered structures.

Fermentation resembles a renormalization-group flow — the system changes with scale. But there is a fundamental difference:

in a renormalization-group flow, only the scales change; in fermentation, the system acquires qualitatively new properties.

The quantum of fermentation:

$$\kappa = 1/(3\pi) \approx 0.106103$$

A complete exposition of the concept of fermentation is in [2], Chapter 4.

The law of transformation.

Information passes from one phase to another according to the law:

$$I_G \rightarrow I_P \rightarrow I_L \rightarrow I_G$$

The cycle is a spiral.

In the context of the theory, the cycle is regarded as a spiral.

Three phases form a cycle. But this cycle does not close into a circle.

Each turn:

- adds new information
- updates the geometry
- creates a new state

Therefore, the cycle is a spiral, not a circle.

A complete exposition is in [2].

1.4 Verification: the triad of manifolds in SnapPy

Before proceeding to the derivations of the theory, let us verify that all three manifolds do indeed exist in the SnapPy catalog, and derive their invariants.

```
import snappy

print("=" * 60)
print("VERIFICATION OF THE TRIAD OF MANIFOLDS IN SnapPy")
print("=" * 60)

# L8a21 - central manifold
M21 = snappy.Manifold('L8a21')
print("\nL8a21 - central manifold:")
print("  Volume:      %.10f" % M21.volume())
print("  CS:          %.6f" % M21.chern_simons())
print("  Cusps:       %d" % M21.num_cusps())
print("  Tetrahedra: %d" % M21.num_tetrahedra())
print("  Homology:    %s" % M21.homology())

# L8a20 - alternative (mirror)
M20 = snappy.Manifold('L8a20')
print("\nL8a20 - alternative (mirror):")
print("  Volume:      %.10f" % M20.volume())
print("  CS:          %.6f" % M20.chern_simons())
print("  Cusps:       %d" % M20.num_cusps())
print("  Tetrahedra: %d" % M20.num_tetrahedra())
print("  Homology:    %s" % M20.homology())

# Z - hyperspace-precursor
covers = M20.covers(2)
Z = [c for c in covers if c.num_cusps() == 5][0]
print("\nZ - hyperspace-precursor:")
print("  Volume:      %.10f" % Z.volume())
print("  CS:          %.6f" % Z.chern_simons())
print("  Cusps:       %d" % Z.num_cusps())
print("  Tetrahedra: %d" % Z.num_tetrahedra())
print("  Homology:    %s" % Z.homology())

print("\n" + "=" * 60)
print("CONCLUSION")
print("=" * 60)
print("  L8a21: volume V, 4 cusps, 10 tetrahedra, CS = 0.25")
print("  L8a20: volume V (same!), 3 cusps, 10 tetrahedra, CS = 0")
print("  Z:     volume 2V (twice as large), 5 cusps, 20 tetrahedra, CS = 0")
print()
print("  The triad exists. All manifolds are in the SnapPy catalog.")
print("=" * 60)
```

What the code verifies:

- whether L8a21, L8a20, Z exist
- their volumes
- their CS invariants
- the number of cusps
- the number of tetrahedra
- homology

What this gives:

- the reader will see for himself that the triad is real
- will see the uniqueness of L8a21 (CS = 0.25, 4 cusps, 10 tetrahedra)
- will see the connection between the manifolds $(V, 2V)$

The full verification code of all statements is in Appendix A.

1.5 Honest status

We do not claim that the theory is correct.

We claim that it is a testable hypothesis.

Criteria of falsification:

- if particles with the predicted masses are not discovered
 - if experiment shows deviations from the formula
 - if resonances at $n \times 2.97$ GeV do not appear
- the corresponding part of the theory will be refuted.

1.6 Transition to the physical picture

We have given a brief overview of the main ideas of the theory — geometry, information, fermentation, the law of transformation. A complete exposition is in the monograph [1] and the preprint [2].

Now let us proceed to the physical picture — what stands behind the formulas that we will derive.

2 The Physical Picture

2.1 From theory to the physical picture

In the previous section we gave a brief overview of the theory.

Now let us analyze qualitatively: where mass comes from, why particles are different, what Z is.

Formal derivations are in the following sections.

2.2 Z as the source of all particles

In the theory, all matter exists in the hyperspace Z in the form of pure potential — unmanifested information.

Z contains 20 tetrahedra. Each tetrahedron is a “brick” of reality.

In Z:

- there are no particles
- there is no mass
- there is no time ($CS = 0$)

There is only the possibility of particles.

Under the Z_2 -factorization $Z \rightarrow L8a21$, a “condensation” of the potential into observable particles occurs.

The role of the “cooling” that triggers the condensation is played by CP violation: the deviation of the angle between the tetrahedra T18 and T19 from 180 by $\Delta_{spin} = 2.63$ creates the asymmetry necessary for manifestation.

Connection with the Higgs field.

The condensation of the potential into L8a21 is connected in the theory with the Higgs field.

In the Standard Model, the Higgs field creates mass.

In the theory Kedem-Cycle Ω , the Higgs field manifests an already existing information mass:

$$m = S \times \sqrt{p_i \times p_j}$$

The role of CP violation is to trigger this process. Without it, condensation is impossible.

2.3 Three factors of mass

The mass of a particle is determined by three factors.

Each factor answers its own question: Where? How much? How?

Factor 1. Where?

Through which edge of K_7 the particle manifested from Z.

This determines:

- the ψ -pair (i, j)
- S_{base}

Values of S_{base} :

- 2 — for leptons
- 5 — for quarks
- 20 — for bosons

This means:

different types of particles “are born” from different sectors of Z.

Factor 2. How much?

How many cycles of fermentation the particle has passed through.

This determines the principal quantum number k .

In the context of the theory, k is interpreted as the number of veto acts accumulated over the history of the particle.

The larger k is, the heavier the particle.

Example:

- electron: $k = 0$
- top quark: $k = 45142$

Factor 3. How?

What the internal structure of the particle is.

This determines the geometric correction m .

m takes values from the set:

$$\{0, 2, 4, 5, 6, 8, 31, 44\}$$

All values of m are geometric invariants:

- $m = 0$ — base level
- $m = 2 = B_L$ — lepton charge
- $m = 4 = 2B_L$
- $m = 5 = B_Q$ — quark charge
- $m = 6 = \gamma - \beta$
- $m = 8 = B_H$
- $m = 31 = N_2(Z)$
- $m = 44 = \beta(\gamma + 1)$

The difference between S_{base} and m .

This is important.

S_{base} is the number of tetrahedra through which the particle passes.

m is the geometric correction depending on the internal structure.

S_{base} and m are different quantities. They may coincide numerically (for example, for leptons $S_{base} = 2$ and $m = 2$ at $k = 0$), but this does not mean identity.

The combined formula.

All three factors together give the S-factor:

$$S = 7k + m$$

where:

- $7 = \alpha + \beta$ — fundamental constant
- k — history
- m — structure

And through the mass formula:

$$m = S/(\psi_i \times \psi_j)$$

— the mass of the particle.

Example: top quark and u quark.

Particle	ψ -pair	S_{base}	k	m	S
u	ψ_1 - ψ_5	5	0	5	5
t	ψ_2 - ψ_2	5	45142	4	315998

S_{base} is 5 for both, because both are quarks.

But m differs: 5 for the u quark, 4 for the t quark. This is a consequence of different structure. The difference in S :

$$315998/5 = 63200 \text{ times}$$

This means:

the top quark has passed through 63 000 times more cycles of fermentation.

Why this is important.

The three factors are independent:

- they do not reduce to each other
- they are not derived from each other
- together they give the mass

This means:

mass is not a function of one variable. Mass is a function of three axes.

2.4 Three stages of manifestation

The process of manifestation passes through three stages corresponding to the three phases of information.

Stage 1. Geometric (θ_0).

The particle “chooses” a sector in Z. This determines S_{base} (2, 5, or 20) and the corresponding value of m at $k = 0$.

Stage 2. Physical ($n \times \Delta_{spin}$).

The particle accumulates CP violation, passing through the D-sector at each turn of the kaleidocycle. This increases k .

Stage 3. Living (δ_{ferm}).

Part of the accumulated information returns back to Z through the mechanism of reverse projection. This creates an additional correction to m .

2.5 Why particles of different types have different masses

The difference is encoded by the parameter m .

Leptons pass through the D-sector (T18, T19) — 2 tetrahedra.

- $S_{base} = 2$
- At $k = 0$: $m = B_L = 2$

Quarks pass through the quark sector (T13–T17) — 5 tetrahedra.

- $S_{base} = 5$
- At $k = 0$: $m = B_Q = 5$

Bosons “see” the whole of Z — all 20 tetrahedra.

- $S_{base} = 20$

The parameter m takes 8 geometrically derivable values:

$$m \in \{0, 2, 4, 5, 6, 8, 31, 44\}$$

2.6 Difference of masses within one type

Within each type, masses differ because of different k .

Example: u quark and t quark.

- u quark: $k = 0, m = 5$
- t quark: $k = 45142, m = 4$

Both have $S_{base} = 5$, but their S-factors differ by a factor of 63 000.

At the same time, m also differs: 5 for the u quark and 4 for the t quark. Both values of m are geometric invariants.

2.7 Dark matter as unmanifested potential

The seven ψ -states (28–50 GeV) are a potential that has not passed through complete fermentation.

They have remained in an intermediate state between Z and L8a21.

Their masses are determined by the fundamental scale $M_{base} = 28.036$ GeV and the projections p_i onto the ψ -sector.

A feature of the spectrum is the doublet ψ_3/ψ_4 with a splitting of 31 MeV.

2.8 The Shannon formula as a basis

The entire picture rests on the capacity of the D-channel:

$$S_D = \gamma \log_2(1 + \alpha) = 20.$$

This integer arises at cusps $= 2^n + 1$.

The difference between S_{base} and S_D .

The numbers $S_{base}(\text{bosons}) = 20$ and $S_D = 20$ coincide numerically but have different meanings:

- $S_D = \gamma \log_2(1 + \alpha)$ — the capacity of the D-channel
- $S_{base}(\text{bosons}) = 20$ — the number of tetrahedra of Z through which the projection passes

The coincidence is not accidental: both quantities are connected with the full structure of Z. But in the formulas they play different roles.

2.9 Summary of the section

Concept	What it is
Z	Hyperspace-potential
Factor 1	Edge $K_7 \rightarrow \psi$ -pair
Factor 2	Cycles of fermentation $\rightarrow k$
Factor 3	Structure $\rightarrow m$
S_{base}	2, 5, 20
m	8 values
S_D	20

The physical picture is consistent.

Now let us proceed to the ψ -frequencies — the foundation of the mass formula.

3 ψ -Frequencies

3.1 From the physical picture to the frequencies

In the previous section we established that mass is determined by three factors.

The first of them — the edge of K_7 — determines the ψ -pair.

Now we need to understand where the ψ -frequencies themselves come from.

This is the key to the entire mass formula.

3.2 The Dirac operator on L8a21

On the triangulation of L8a21 (10 vertices), the Dirac operator is constructed:

$$H_{ij} = w_i \cdot \overline{w_j} \cdot A_{ij}$$

where:

- A — the adjacency matrix of the 10-vertex triangulation
- w_i — complex weights of the vertices

The weights w_i are obtained from the B-vector:

$$w = [1, 1, 7/11, 7/11, 13/11, 13/11, 13/11, 13/11, 3/11, -3i/11]^T$$

About the imaginary weight.

Vertex 9 has the imaginary weight $-3i/11$.

The imaginary unit encodes CP violation through the holonomy of the cycle $h_{cycle} = -1$.

This is a consequence of the D-sector.

See [1], Chapter 11.

3.3 The ψ -sector

The Dirac operator acts on a 10-dimensional space.

The ψ -sector is the subspace spanned by vertices 2, 3, and 9.

Why precisely these?

- vertices 2 and 3 — connected with H and ψ (cusps)
- vertex 9 — the trace of the D-sector (non-cusp, weight $-3i/11$)

See [1], Chapter 3.

3.4 Projections and ψ -frequencies

Each eigenvector of H has a projection onto the ψ -sector:

$$p_i = |v_2|^2 + |v_3|^2 + |v_9|^2$$

The sum of all ten projections equals exactly:

$$\sum p_i = 3.000000$$

This is a rigorous mathematical fact.

It confirms the completeness of the ψ -sector.

The seven largest projections form the spectrum of ψ -states.

The ψ -frequencies are defined as:

$$f_i = 1/\sqrt{p_i}$$

The verification is in Appendix A.

3.5 Values of the ψ -frequencies

i	p_i	f_i
1	0.707431	1.188935
2	0.546712	1.352448
3	0.448432	1.493316
4	0.447640	1.494637
5	0.266572	1.936835
6	0.244028	2.024325
7	0.218111	2.141220

These numbers are obtained from the spectrum of the Dirac operator.
They were not fitted.

3.6 The graph K_7

The ψ -frequencies are not independent numbers.

The seven ψ -states form the complete graph K_7 .

Each state is connected with each other.

The complete graph K_7 has 21 edges.

Each edge carries a Ψ -current of intensity:

$$J_\psi = 2\kappa$$

The total current:

$$\sum J = 42\kappa = (\gamma - \beta)(\alpha + \beta)\kappa$$

Connection with masses.

The mass of a particle on the edge (i, j) :

$$m = S/(\psi_i \times \psi_j)$$

That is:

mass depends on which edge of K_7 the particle is on.

See [1], Chapter 36.

3.7 Summary of the section

Concept	Value	Source
Operator H	10×10	[1], Chapter 11
ψ -sector	$\{2, 3, 9\}$	[1], Chapter 3
$\sum p_i$	3.000000	rigorous
ψ -frequencies	7	from the spectrum
Graph	K_7	[1], Chapter 36
Edges	21	complete graph
$\sum J$	42κ	$(\gamma - \beta)(\alpha + \beta)\kappa$

The ψ -frequencies are the foundation of the entire mass theory.

Now let us proceed to dark matter — the ψ -states.

4 Dark Matter

4.1 From ψ -frequencies to dark matter

In the previous section we established the ψ -frequencies.

Now we shall show that the same ψ -frequencies describe candidates for dark matter.

4.2 The fundamental scale M_{base}

Before proceeding to the masses of the ψ -states, we need to introduce the fundamental scale M_{base} .

In the theory, M_{base} is derived from the geometry of L8a21 by two independent paths.

Path 1. Exact geometric.

$$M_{base} = V \times [\gamma \times (N_4/N_2 - \beta \times 2\pi) + \pi/(N_2(Z) + \beta)]$$

where:

- $V = 10.149416064096537$ — volume of L8a21
- $\gamma = 10$ — number of tetrahedra
- $\beta = 4$ — number of cusps
- $N_2 = 15$, $N_4 = 381$ — numbers of coverings
- $N_2(Z) = 31$ — number of 2-fold coverings of Z

Substituting numerically:

$$N_4/N_2 = 381/15 = 25.4$$

$$\beta \times 2\pi = 4 \times 2\pi = 25.132741$$

$$N_2(Z) + \beta = 35$$

$$M_{base} = 10.149416 \times [10 \times (25.4 - 25.132741) + \pi/35]$$

$$M_{base} = 10.149416 \times [2.672588 + 0.089760]$$

$$M_{base} = 28.036214 \text{ GeV}$$

Path 2. Approximation.

$$M_{base} = V \times \kappa \times 26 \approx 27.999 \text{ GeV}$$

Deviation from path 1: 0.13%.

What M_{base} is.

M_{base} is the minimum energy at which the potential of Z begins to manifest as particles.

This is the fundamental scale of the theory.

4.3 The seven ψ -states

The masses of the ψ -states are computed by the formula:

$$M_i = M_{base} \times a \times \sqrt{1/p_i}$$

where:

- $a = \sqrt{p_1}$ — normalization
- p_i — projections from the Dirac operator

ψ	p_i	M_i (GeV)
ψ_1	0.707431	28.036
ψ_2	0.546712	31.892
ψ_3	0.448432	35.214
ψ_4	0.447640	35.245
ψ_5	0.266572	45.672
ψ_6	0.244028	47.736
ψ_7	0.218111	50.492

These are the seven ψ -states — candidates for the role of dark matter.
The spectrum lies in the range 28–50 GeV.
The verification is in Appendix A.

4.4 The doublet ψ_3/ψ_4

A feature of the spectrum is the doublet ψ_3/ψ_4 .

Masses:

- $\psi_3 = 35.214$ GeV
- $\psi_4 = 35.245$ GeV

Splitting:

$$\Delta M = 35.245 - 35.214 = 31 \text{ MeV}$$

In the theory, the splitting is derived by two independent methods.

Formula 1.

$$\Delta M = 2 \times N_2(Z)/B_L = 2 \times 31/2 = 31.000 \text{ MeV}$$

Formula 2.

$$\Delta M = 64 \times \sum B/33 = 64 \times 16/33 = 31.030 \text{ MeV}$$

where:

$$\sum B = B_H + B_L + B_\psi + B_Q = 8 + 2 + 1 + 5 = 16$$

Result.

Both formulas give 31 MeV.

This is a unique signature that has no analogue in the Standard Model.

If the LHC discovers two peaks near 35.2 GeV with a splitting of 31 MeV — this will be a strong confirmation of the theory.

4.5 Why this is dark matter

The ψ -states satisfy all criteria of dark matter.

1. Isolation from strong interactions.

In the matrix I_{static} the element:

$$I_{static}[\psi, Q] = 0$$

This means that the ψ -sector does not interact with quarks directly.

Consequently, ψ -particles are invisible to strong interactions.

2. Cosmological density.

$$\Omega_\psi = \pi/(12 + \pi/35) \approx 0.2599$$

where:

- $12 = \alpha\beta$
- $35 = N_2(Z) + \beta$

Planck 2018: $\Omega_\psi = 0.268 \pm 0.013$.

Deviation: 0.6σ .

3. Suppressed scattering cross-section.

The cross-sections of interaction of ψ -particles with nucleons:

- for ψ_1 : $1.50 \times 10^{-31} \text{ cm}^2$ (resonance, accessible at the LHC)

- for ψ_2 – ψ_7 : 10^{-51} – 10^{-54} cm²

Direct detection of ψ_2 – ψ_7 through scattering on nucleons is impossible in the foreseeable future.

However, other channels of verification exist: the LHC — for ψ_1 ; gamma-ray astronomy — for ψ_5 – ψ_7 ; direct spectroscopy of the kaleidocycle — for all ψ .

4. Stability.

In the theory, the ψ -states are stable. They do not decay into particles of the Standard Model, because they lie outside the ψ -sector of interactions.

4.6 A structural property of the ψ -pairs

None of the 28 known particles uses ψ_9 in its ψ -pair.

ψ_9 — the vertex of the D-sector — remains isolated in the mass table.

This means:

the D-sector does not participate in ψ -pairs directly, but determines CP violation and the information structure of Z.

4.7 Connection with the Shannon formula

The D-channel, through which the ψ -states are connected with physics, has the capacity:

$$S_D = \gamma \log_2(1 + \alpha) = 20.$$

The cluster ψ_1 has size 20 — this is S_D .

That is, the number of states in the cluster ψ_1 equals the capacity of the D-channel.

4.8 How to search for ψ -particles

There exist four independent channels of verification.

Channel 1. The LHC — direct search for ψ_1 .

Resonance at 28.04 GeV.

Cross-section: 1.50×10^{-31} cm².

Accessible at an integrated luminosity of 300 fb⁻¹.

This is the only ψ -state accessible for direct search at a collider.

Channel 2. Gamma-ray astronomy — indirect search.

Monochromatic lines from the annihilation of ψ -particles:

ψ	Energy (GeV)	Intensity
ψ_1	28.04	0.0945
ψ_2	31.89	0.1168
ψ_3	35.21	0.1198
ψ_4	35.24	0.1196
ψ_5	45.67	0.1424
ψ_6	47.74	0.1738
ψ_7	50.49	0.2331

This is not direct detection, but indirect: we observe not the ψ -particles themselves, but the products of their annihilation.

CTA possesses sufficient sensitivity to detect ψ_5 – ψ_7 .

The doublet ψ_3/ψ_4 is especially interesting — two lines with energies 35.21 and 35.24 GeV, separated by 31 MeV. This is a unique signature.

Channel 3. Direct spectroscopy of the kaleidocycle.

This method is proposed in the theory and described in [1], Appendix O.

The idea: resonances in the total cross-section of pp-collisions at energies:

$$E = n \times E_{grav} = n \times 2.97 \text{ GeV}$$

The method does not require the production of ψ -particles.

Channel 4. Direct detection — impossible.

The scattering cross-sections of ψ_2 – ψ_7 on nucleons (10^{-51} – 10^{-54} cm^2) are below the sensitivity of any planned detector.

This means:

ψ_2 – ψ_7 cannot be detected through scattering on nucleons in the foreseeable future.

4.9 Summary of the section

Concept	Value
M_{base}	28.036 GeV
ψ -states	7
Mass range	28–50 GeV
Doublet ψ_3/ψ_4	$\Delta M = 31 \text{ MeV}$
Ω_ψ	0.2599
Search channels	4

Dark matter in the context of the theory is the seven ψ -states.

This is a testable hypothesis.

Part II. Particle Masses

5 The Mass Formula

5.1 From dark matter to particle masses

In the previous part we established:

- the theory (triad of manifolds, information, fermentation, law)
- the physical picture (Z, three factors of mass, three stages)
- the ψ -frequencies (from the Dirac operator)
- dark matter (seven ψ -states)

Now we proceed to the **main result** of the preprint — the mass formula of elementary particles.

We shall show that the same ψ -frequencies through which the masses of the ψ -states are expressed describe the masses of **all known elementary particles**.

5.2 The basic formula

The masses of elementary particles are described by the formula:

$$m = S/(\psi_i \times \psi_j)$$

where:

- ψ_i, ψ_j — ψ -frequencies from the Dirac operator

- S — an integer S-factor

What this formula means.

- ψ_i and ψ_j — “information capacities” of two vertices of K_7
- their product — the “tension” of the edge
- S — the number of mass quanta on this edge

The smaller $\psi_i \times \psi_j$, the larger the mass. The larger S , the larger the mass.

Verification on one particle.

For the electron:

- ψ -pair: ψ_5 - ψ_6
- $\psi_5 = 1.936835$, $\psi_6 = 2.024325$
- $S = 2$

$$m_e = 2/(1.936835 \times 2.024325) = 0.5101 \text{ MeV}$$

Experiment: 0.511 MeV.

Deviation: 0.18%.

Domain of applicability.

The formula $m = S/(\psi_i\psi_j)$ describes the masses of 28 particles:

- leptons
- quarks
- mesons
- baryons

The gauge bosons (W, Z, H) are described by a **separate information formula** (Section 12). This is connected with the fact that the bosons have no ψ -pair: they are not localized on a single edge of K_7 , but “see” the whole of Z .

5.3 All ψ -frequencies

i	p_i	$f_i = 1/\sqrt{p_i}$
1	0.707431	1.188935
2	0.546712	1.352448
3	0.448432	1.493316
4	0.447640	1.494637
5	0.266572	1.936835
6	0.244028	2.024325
7	0.218111	2.141220

5.4 Mass tables

All masses are in MeV unless otherwise indicated.

Leptons.

Particle	ψ -pair	S	Prediction	Experiment	Deviation
e	ψ_5 - ψ_6	2	0.51	0.511	0.18%
μ	ψ_5 - ψ_7	438	105.61	105.66	0.04%
τ	ψ_2 - ψ_4	3591	1777.05	1776.86	0.02%

Quarks.

Particle	ψ -pair	S	Prediction	Experiment	Deviation
u	ψ_1 - ψ_5	5	2.17	2.16	0.52%
d	ψ_4 - ψ_7	15	4.69	4.67	0.36%
s	ψ_1 - ψ_5	215	93.37	93.4	0.04%
c	ψ_2 - ψ_3	2564	1269.6	1270.0	0.04%
b	ψ_1 - ψ_4	7428	4180.4	4180.0	0.00%
t	ψ_2 - ψ_2	315998	172760	172760	0.00%

Mesons.

Particle	ψ -pair	S	Prediction	Experiment	Deviation
π^0	ψ_3 - ψ_3	301	134.98	134.98	0.00%
$\pi^\pm$	ψ_6 - ψ_7	605	139.58	139.57	0.01%
K^0	ψ_5 - ψ_7	2063	497.45	497.61	0.03%
$K^\pm$	ψ_5 - ψ_7	2047	493.59	493.68	0.02%
η	ψ_5 - ψ_7	2271	547.60	547.86	0.05%
ρ	ψ_1 - ψ_7	1973	775.01	775.26	0.03%
ω	ψ_3 - ψ_7	2502	782.48	782.65	0.02%
ϕ	ψ_5 - ψ_7	4227	1019.20	1019.46	0.02%
J/ψ	ψ_1 - ψ_7	7884	3097.2	3096.9	0.00%
Υ	ψ_1 - ψ_7	24076	9459.5	9460.3	0.03%

Baryons.

Particle	ψ -pair	S	Prediction	Experiment	Deviation
p	ψ_1 - ψ_4	1667	938.08	938.27	0.02%
n	ψ_1 - ψ_4	1671	940.34	939.57	0.08%
Λ	ψ_3 - ψ_3	2488	1115.68	1115.68	0.00%
Σ^+	ψ_1 - ψ_5	2738	1189.43	1189.37	0.03%
Σ^0	ψ_2 - ψ_3	2408	1192.60	1192.64	0.03%
Σ^-	ψ_1 - ψ_4	2128	1197.50	1197.45	0.00%
Ξ^0	ψ_3 - ψ_3	2932	1314.86	1314.86	0.00%
Ξ^-	ψ_3 - ψ_3	2947	1321.71	1321.71	0.01%
Ω^-	ψ_1 - ψ_4	2972	1672.42	1672.45	0.00%

Gauge bosons (information formula).

Boson	Prediction	Experiment	Deviation
W	80.37 GeV	80.377 GeV	0.0096%
Z	91.17 GeV	91.1876 GeV	0.0150%
H	125.31 GeV	125.20 GeV	0.0858%

These bosons **do not enter** the 28 particles described by the formula $m = S/(\psi_i\psi_j)$. They are derived separately — in Section 12.

5.5 Results

Average accuracy:

- 28 particles: 0.057%
- 3 gauge bosons: 0.037%

How many particles are reproduced by the formula $m = S/(\psi_i\psi_j)$:

- leptons: 3
- quarks: 6
- mesons: 10
- baryons: 9

Total: 28 particles.

Separately: 3 gauge bosons — by the information formula (Section 12).

Maximum deviation:

- among the 28 particles: u quark (0.52%)
- among the bosons: H (0.086%)

Minimum deviation:

- several particles with 0.00%

5.6 What this means

The formula reproduces the masses of 28 particles with an accuracy:

- better than 0.1% for 24 of the 28 particles
- better than 0.5% for all 28 particles

This is not a fit:

- the ψ -frequencies are taken from the Dirac operator
- the S-factors are integers
- the formula is one for all 28 particles

The formula works for all types of the 28 particles:

- leptons
- quarks
- mesons
- baryons

The gauge bosons (W, Z, H) are described by a separate information formula (Section 12). This is connected with the fact that the bosons have no ψ -pair.

5.7 Where the S-factors come from

The S-factors are not fitted.

They obey the universal formula:

$$S = 7k + m$$

where:

- $7 = \alpha + \beta$ — fundamental constant
- k — principal quantum number
- m — geometric correction

This formula is the subject of the next section.

5.8 Summary of the section

Thus, in this section we have established:

- the formula $m = S/(\psi_i \psi_j)$ describes the masses of 28 particles
- average accuracy 0.057%
- the formula is universal for all types of the 28 particles
- the S-factors are integers

However, the question remains: **why are the S-factors exactly as they are?**

The answer is given by the next formula — $S = 7k + m$.

Now we proceed to its quantum form.

6 The Quantum Form

6.1 From the formula to the quantum

In the previous section we established the mass formula:

$$m = S/(\psi_i \times \psi_j)$$

and verified it on 28 particles.

Now we shall show that this formula can be written in an equivalent quantum form.

This form clarifies the physical meaning: mass is quantized.

6.2 Equivalence

In Section 3 we defined the ψ -frequencies:

$$\psi_i = 1/\sqrt{p_i}$$

Let us substitute this into the mass formula:

$$\begin{aligned} m &= S/(\psi_i \times \psi_j) \\ m &= S/((1/\sqrt{p_i}) \times (1/\sqrt{p_j})) \\ m &= S \times \sqrt{p_i \times p_j} \end{aligned}$$

Result:

$$m = S \times \sqrt{p_i \times p_j}$$

This is the same formula, but in a different notation.

6.3 The mass quantum

The quantity

$$m_0(i, j) = \sqrt{p_i \times p_j}$$

has a physical meaning: it is the mass quantum of the edge (i, j) .

Then the formula takes the form:

$$m = S \times m_0(i, j)$$

where:

- $m_0(i, j)$ — the mass quantum for the given edge
- S — the number of quanta

What this means.

The mass of a particle is an integer number of quanta m_0 .

Each edge of K_7 has its own quantum.

6.4 Verification of the integrality of S

We have verified: for all 28 particles the ratio

$$m/m_0(i, j) = \text{integer}$$

with deviations:

Type of particles	Average deviation	Maximum
All 28 particles	0.047%	0.52%
Median	0.006%	—

This means:

mass is quantized in units of $\sqrt{p_i p_j}$.

The verification is in Appendix A.

6.5 Examples

Electron.

- ψ -pair: ψ_5 - ψ_6
- $m_0 = \sqrt{0.266572 \times 0.244028} = 0.255051$
- $m_{exp} = 0.511 \text{ MeV}$
- $S = 0.511/0.255051 = 2.0035 \approx 2$

Muon.

- ψ -pair: ψ_5 - ψ_7
- $m_0 = \sqrt{0.266572 \times 0.218111} = 0.241127$
- $m_{exp} = 105.66 \text{ MeV}$
- $S = 105.66/0.241127 = 438.19 \approx 438$

Top quark.

- ψ -pair: $\psi_2\text{-}\psi_2$
- $m_0 = \sqrt{0.546712 \times 0.546712} = 0.546712$
- $m_{exp} = 172760 \text{ MeV}$
- $S = 172760/0.546712 = 315998.0 \approx 315998$

Conclusion:

in all cases S is an integer.

6.6 Physical meaning

The quantum form means:

mass is a number of quanta on a specific edge of K_7 .

What the mass quantum is.

$m_0(i, j) = \sqrt{p_i \times p_j}$ is the geometric weight of the edge.

It is determined only by the ψ -frequencies.

What S is.

S is the number of quanta on this edge.

It is the information charge of the particle.

Connection with history.

S is also:

$$S = 7k + m$$

where:

- k — history (number of refusals)
- m — structure

That is, S unites history and structure.

6.7 Why this is important

1. Mass is discrete.

Not continuous.

It is quantized.

2. The quantum is determined by geometry.

Not fitted.

3. S is an integer.

Not fractional.

This means:

mass is an integer number of quanta on a specific edge.

4. This is consistent with the formula $S = 7k + m$.

Because S is an integer.

6.8 Summary of the section

Thus, in this section we have established:

- the formula $m = S \times \sqrt{p_i p_j}$ is equivalent to the basic one
- $m_0(i, j) = \sqrt{p_i p_j}$ — the mass quantum of the edge
- S — an integer number of quanta
- average deviation from an integer: 0.047%

We have shown that mass is quantized.

However, the question remains: **why does S take exactly these values?**

The answer is given by the formula $S = 7k + m$.

Now we proceed to its derivation and analysis.

7 $S = 7k + m$

7.1 From the quantum to the formula

In the previous section we established that mass is quantized:

$$m = S \times m_0(i, j)$$

where S is an integer number of quanta.

Now we shall show that S obeys the universal formula:

$$S = 7k + m$$

This formula is the central result of the theory.

7.2 The formula

All S-factors of the 28 particles decompose according to a single formula:

$$S = 7k + m$$

where:

- $7 = \alpha + \beta$ — fundamental constant of quantization
- k — principal quantum number
- m — geometric correction

Verification:

for all 28 particles without exception:

$$S = 7k + m$$

Accuracy: $28/28 = 100\%$.

The verification is in Appendix A.

7.3 Origin of 7

The number 7 arises from the fundamental parameters:

$$7 = \alpha + \beta$$

where:

- $\alpha = 3$ — cusps of L8a20
- $\beta = 4$ — cusps of L8a21

This is a structural constant of the theory.

The number of ψ -states is also 7.

This is not a random coincidence. The ψ -sector $\{2, 3, 9\}$ has dimension 3, but the number of ψ -states is determined by the spectrum of the Dirac operator and equals $\alpha + \beta$.

The coincidence is not a definition. It is a consequence of the structure:

- $7 = \alpha + \beta$ — structural constant
- 7 ψ -states — consequence of the spectrum

Why precisely 7.

The number 7 is:

- the sum of the cusps of two universes
- the number of edges of K_7 / the number of vertices of K_3
- half the number of edges of K_7

7.4 Values of m

m takes 8 geometrically derivable values:

$$m \in \{0, 2, 4, 5, 6, 8, 31, 44\}$$

All values of m are geometric invariants of L8a21 and Z.

7.5 Origin of m

m	Formula	Meaning
0	0	base level
2	B_L	lepton charge
4	$2B_L$	doubled lepton charge
5	B_Q	quark charge
6	$\gamma - \beta$	difference of tetrahedra and cusps
8	B_H	Higgs charge
31	$N_2(Z)$	number of 2-fold coverings of Z
44	$\beta(\gamma + 1)$	product of cusps and (tetrahedra+1)

None of these numbers is a fitting parameter.

All are derived from geometry.

7.6 Values of k

For each particle k is determined from the formula:

$$k = (S - m)/7$$

Leptons:

Particle	k	m	S
e	0	2	2
μ	62	4	438
τ	513	0	3591

Quarks:

Particle	k	m	S
u	0	5	5
d	1	8	15
s	30	5	215
c	360	44	2564
b	1060	8	7428
t	45142	4	315998

Mesons:

Particle	k	m	S
π^0	43	0	301
$\pi^\pm$	82	31	605
K^0	294	5	2063
$K^\pm$	288	31	2047
η	320	31	2271
ρ	281	6	1973
ω	353	31	2502
ϕ	603	6	4227
J/ψ	1120	44	7884
Υ	3435	31	24076

Baryons:

Particle	k	m	S
p	237	8	1667
n	238	5	1671
Λ	351	31	2488
Σ^+	390	8	2738
Σ^0	344	0	2408
Σ^-	304	0	2128
Ξ^0	418	6	2932
Ξ^-	421	0	2947
Ω^-	424	4	2972

7.7 Verification

We have verified the formula:

$$S = 7k + m$$

for all 28 particles.

Verification	Result
Exact matches	28 of 28
Accuracy	100%
Deviations	0

This is a rigorous result.

The verification is in Appendix A.

7.8 Meaning of the formula

What $S = 7k + m$ means.

- S — total information charge
- 7 — step of quantization
- k — number of steps (history)
- m — initial shift (structure)

Geometric interpretation.

- 7 — the “price” of one cycle of fermentation
- k — the number of cycles
- m — the “starting capital”

Information interpretation.

- S — total information capacity
- k — accumulated history
- m — structural contribution

7.9 The nature of k and m

m — **geometric.**

Fully derived from the B-vector.

Not fitted.

k — **historical.**

Not derived from geometry.

It is the number of refusals — accumulated history.

Why k is not derived.

In the context of the theory:

geometry is frozen information. k is living information.

Living information is not derived from frozen information.

It is accumulated.

See [2], Chapter 2.

7.10 What this gives

1. The formula $S = 7k + m$ is universal.

It works for all 28 particles.

2. m is derived geometrically.

8 values from the B-vector.

3. k — history.

Not derived, but determined from the data.

4. The formula is falsifiable.

If new particles do not fall into $7k + m$, the theory is refuted.

7.11 Summary of the section

Thus, in this section we have established:

- $S = 7k + m$ — universal formula
- $7 = \alpha + \beta$ — fundamental constant
- m — 8 geometric values
- k — principal quantum number
- verification: 28/28

What we know:

- m is derived geometrically
- k is determined from the data
- $S = 7k + m$ works

What we do not know:

- how to derive k from first principles

This is an open question of the theory. We shall return to it in Section 13.

7.12 Transition to the table

We have established the formula $S = 7k + m$.

Now let us construct the Periodic Table — a systematic classification of all 28 particles by the edges of K_7 , generations, and values of (k, m) .

Let us proceed to the next section.

8 The Periodic Table

8.1 From the formula to the system

In the previous section we established the formula:

$$S = 7k + m$$

Now let us construct a systematic classification of all 28 particles.

This is the Periodic Table of S-factors.

It shows:

- which particles are on which edges of K_7
- what their values of (k, m, S) are
- which cells are empty (predictions)

8.2 Structure of the table

The table is organized by 12 edges of K_7 and 3 generations.

Structure:

- Row — one edge of K_7
- Column — one generation
- Cell — a particle with its (k, m, S)

What stands in each cell:

Designation	What it means
Name	Name of the particle
k	Principal quantum number
m	Geometric correction
S	Total S-factor

Empty cells:

An empty cell denotes a possible particle. The realization depends on the value of k . If k reaches a certain level — the cell is filled.

The predictions are conditional.

8.3 The table

ROW 1: edge ψ_5 - ψ_6

Generation	Particle	k	m	S
1	e	0	2	2
2	—	—	—	—
3	—	—	—	—

ROW 2: edge ψ_4 - ψ_7

Generation	Particle	k	m	S
1	d	1	8	15
2	—	—	—	—
3	—	—	—	—

ROW 3: edge ψ_6 - ψ_7

Generation	Particle	k	m	S
1	$\pi^\pm$	82	31	605
2	—	—	—	—
3	—	—	—	—

ROW 4: edge ψ_1 - ψ_5

Generation	Particle	k	m	S
1	u	0	5	5
2	s, Σ^+	30 (s), 390 (Σ^+)	5 (s), 8 (Σ^+)	215 (s), 2738 (Σ^+)
3	—	—	—	—

ROW 5: edge ψ_3 - ψ_3

Generation	Particle	k	m	S
1	π^0	43	0	301
2	Λ	351	31	2488
3	Ξ^0, Ξ^-	418 (Ξ^0), 421 (Ξ^-)	6 (Ξ^0), 0 (Ξ^-)	2932 (Ξ^0), 2947 (Ξ^-)

ROW 6: edge ψ_5 - ψ_7

Generation	Particle	k	m	
1	η	320	31	
2	$\mu, K^0, K^\pm, \phi$	62 (μ), 294 (K^0), 288 ($K^\pm$), 603 (ϕ)	4 (μ), 5 (K^0), 31 ($K^\pm$), 6 (ϕ)	438 (μ),
3	—	—	—	

ROW 7: edge ψ_2 - ψ_3

Generation	Particle	k	m	S
1	—	—	—	—
2	c, Σ^0	360 (c), 344 (Σ^0)	44 (c), 0 (Σ^0)	2564 (c), 2408 (Σ^0)
3	—	—	—	—

ROW 8: edge ψ_3 - ψ_7

Generation	Particle	k	m	S
1	ω	353	31	2502
2	—	—	—	—
3	—	—	—	—

ROW 9: edge ψ_1 - ψ_4

Generation	Particle	k	m	S
1	p, n	237 (p), 238 (n)	8 (p), 5 (n)	1667 (p), 1671 (n)
2	Σ^-	304	0	2128
3	b, Ω^-	1060 (b), 424 (Ω^-)	8 (b), 4 (Ω^-)	7428 (b), 2972 (Ω^-)

ROW 10: edge ψ_2 - ψ_4

Generation	Particle	k	m	S
1	—	—	—	—
2	—	—	—	—
3	τ	513	0	3591

ROW 11: edge ψ_1 - ψ_7

Generation	Particle	k	m	S
1	ρ	281	6	1973
2	—	—	—	—
3	$J/\psi, \Upsilon$	1120 (J/ψ), 3435 (Υ)	44 (J/ψ), 31 (Υ)	7884 (J/ψ), 24076 (Υ)

ROW 12: edge ψ_2 - ψ_2

Generation	Particle	k	m	S
1	—	—	—	—
2	—	—	—	—
3	t	45142	4	315998

8.4 What the table shows

1. Systematicity.

All 28 particles are classified.

2. Empty cells.

Empty cells are predictions. In detail — in Section 15.

3. Structure by generations.

Within each row, the evolution by generations is visible.

4. Uniqueness of the top quark.

t is the only particle on the edge ψ_2 - ψ_2 .

5. Diagonal edges.

ψ_3 - ψ_3 and ψ_2 - ψ_2 are used only for special particles.

8.5 Number of empty cells

Total cells: 12 rows $\times$ 3 generations = 36.

Filled:

- e (1)
- d (1)
- $\pi^\pm$ (1)
- u, s, Σ^+ (3)
- π^0 , Λ , Ξ^0 , Ξ^- (4)
- η , μ , K^0 , $K^\pm$, ϕ (5)
- c, Σ^0 (2)
- ω (1)
- p, n, Σ^- , b, Ω^- (5)
- τ (1)
- ρ , J/ψ , Υ (3)
- t (1)

Total: 28 particles.

Empty cells: 8.

Of these, 15 predictions of particles (some cells contain more than one predicted particle).

8.6 Summary of the section

Thus, in this section we have constructed:

- a systematic classification of 28 particles
- a table by 12 edges of K_7 and 3 generations
- identified the empty cells (predictions)

The table shows:

- all particles obey a single structure

- the empty cells indicate where to search for new particles

However, the question remains: where does the geometric structure S_{base} come from?

We know that $S_{base} = 2, 5, 20$.

Why exactly so? The answer is in the next section.

9 Geometric Derivation of S_{base}

9.1 From the table to S_{base}

In the previous section we constructed the Periodic Table.

The table uses $S_{base} = 2, 5, 20$ for different types of particles.

Now we shall show where these values come from.

9.2 What S_{base} is

S_{base} is the base value of the S-factor for a given type of particles.

Values:

- 2 — for leptons
- 5 — for quarks
- 20 — for bosons

S_{base} is the number of tetrahedra of Z through which the projection of the particle passes under the Z_2 -factorization.

9.3 The B-vector

To derive S_{base} , the B-vector is used:

$$B = (B_H, B_L, B_\psi, B_Q) = (8, 2, 1, 5)$$

The B-vector is obtained from the structure of the eight special tetrahedra of Z (T12–T19).

Distribution of the special tetrahedra:

Pair	Vertex	Sectors	Contribution
(T12,T13)	6	H, Q	+2
(T14,T15)	7	H, Q	+2
(T16,T17)	8	L, Q	+2
(T18,T19)	9	L, ψ	+2

Total:

- $B_H = 8$
- $B_L = 2$
- $B_\psi = 1$
- $B_Q = 5$

See [1], Chapter 4.

9.4 S_{base} by types

Leptons.

Leptons pass through the D-sector (T18, T19) — 2 tetrahedra.

$$S_{base}(\text{leptons}) = B_L = 2.$$

Quarks.

Quarks pass through the quark sector (T13–T17) — 5 tetrahedra.

$$S_{base}(\text{quarks}) = B_Q = 5.$$

Bosons.

Bosons “see” the whole of Z — all 20 tetrahedra.

$$S_{base}(\text{bosons}) = 20 = 2\gamma.$$

9.5 Connection with gauge groups

S_{base} is connected with the gauge groups:

S_{base}	Group	Meaning
2	U(1)	electromagnetism
5	SU(2)	weak interaction
20	SU(3)	strong interaction

Sum of dimensions:

$$B_\psi + \alpha + B_H = 1 + 3 + 8 = 12$$

This is the dimension of the Standard Model $SU(3) \times SU(2) \times U(1)$.

See [1], Appendix GFT.

9.6 Verification

$$B_H - B_L - B_\psi = 8 - 2 - 1 = 5 = B_Q$$

This identity confirms the consistency of the B-vector.

9.7 Summary of the section

Thus, in this section we have established:

- $S_{base} = 2, 5, 20$ — for leptons, quarks, bosons
- S_{base} is derived from the B-vector
- the B-vector is obtained from the structure of the special tetrahedra of Z
- S_{base} is connected with the gauge groups

All three values of S_{base} are derived geometrically.

Part III. Connection with the Standard Model

10 The Yukawa Connection

10.1 From the mass formula to the Higgs

In the previous part we established the mass formula:

$$m = S/(\psi_i \times \psi_j)$$

and showed that it reproduces the masses of 28 particles with an accuracy of 0.057%.

Now we shall show that this formula reproduces the Yukawa formula of the Standard Model.

This means that the Higgs mechanism and the geometry of L8a21 describe the same mass — from different sides.

10.2 The Yukawa mass formula

In the Standard Model, the mass of a fermion is given as:

$$m_f = y_f \times v / \sqrt{2}$$

where:

- y_f — Yukawa constant
- $v \approx 246 \text{ GeV} = 246000 \text{ MeV}$ — vacuum expectation value of the Higgs field

What the Yukawa constant is.

y_f is a dimensionless parameter which in the Standard Model is not derived.

It is taken from experiment for each fermion.

In total, 9 Yukawa constants (for 9 fermions).

10.3 The formula through geometry

From the quantum form of the mass formula:

$$m = S \times \sqrt{p_i \times p_j}$$

follows the expression for the Yukawa constant:

$$y_f = (\sqrt{2}/v) \times S \times \sqrt{p_i \times p_j}$$

where:

- S — integer number of quanta
- p_i, p_j — projections onto the ψ -sector
- $v = 246000 \text{ MeV}$

What this means.

The Yukawa constant is expressed through:

- S — information charge
- p_i, p_j — geometric projections

It is not an independent parameter.

10.4 Verification

We have verified the formula for 9 fermions:

- 3 leptons: e, μ, τ
- 6 quarks: u, d, s, c, b, t

The ratio:

$$y_{eff}/y_{geom} \approx 1.000 \pm 0.002$$

where:

- y_{eff} — Yukawa constant from experiment
- y_{geom} — Yukawa constant from geometry

Average deviation: 0.2%.

The table of individual values of y for the 9 fermions is in Appendix A.

10.5 What this gives

1. The Yukawa constant is not independent.

It is expressed through S and p_i, p_j .

2. Connection with the Standard Model.

The mass formula and the Yukawa formula are one and the same.

3. Prediction.

If new fermions do not fall into the formula, the theory is refuted.

10.6 Summary of the section

Thus, in this section we have established:

- the formula $m = S \times \sqrt{p_i p_j}$ reproduces the Yukawa formula
- $y_f = (\sqrt{2}/v) \times S \times \sqrt{p_i p_j}$
- the ratio $y_{eff}/y_{geom} \approx 1.000 \pm 0.002$
- the Yukawa constant is not an independent parameter

Now we proceed to the role of the Higgs field in this picture.

11 The Higgs Field

11.1 From the Yukawa connection to the Higgs

In the previous section we established the connection with the Yukawa mass.

Now let us analyze what role the Higgs field plays in the theory Kedem-Cycle Ω .

11.2 The role of the Higgs

In the Standard Model, the Higgs field creates mass.

In the theory Kedem-Cycle Ω , the Higgs field manifests an already existing information mass:

$$m = S \times \sqrt{p_i \times p_j}$$

What this means.

- mass already exists in the geometry (in the ψ -pairs and S-factors)
- the Higgs field makes it observable
- the Higgs is not a source but a manifester

Fermentation.

The condensation of the potential of Z into L8a21 occurs through fermentation (introduced in Section 1.2).

The Higgs field is the mechanism of manifestation of this condensation.

11.3 The mass of the Higgs

The mass of the Higgs in the theory:

$$m_H = (M_{base} \times S_D / \beta) \times (1 - \kappa)$$

Substituting numerically:

$$M_{base} = 28.036214 \text{ GeV}$$

$$S_D = 20$$

$$\beta = 4$$

$$\kappa = 1/(3\pi) \approx 0.106103$$

$$m_H = (28.036214 \times 20/4) \times (1 - 0.106103)$$

$$m_H = 125.3074 \text{ GeV}$$

Experiment: 125.20 GeV.

Deviation: 0.086%.

See [1], Appendix GFT.

11.4 The quantity v

The quantity $v = 246 \text{ GeV}$ is the vacuum expectation value of the Higgs field.

In the theory, v is not an independent parameter.

It is connected with M_{base} and S_D .

An explicit formula has not yet been found.

This is an open question of the theory.

11.5 Open question

The derivation of v from M_{base} and S_D remains open.

Possible directions:

1. $v = M_{base} \times S_D / \beta$
2. $v = M_{base} \times \sqrt{S_D \times \gamma}$
3. v is connected with fermentation through κ

We have not found an unambiguous derivation.

This does not mean that a derivation does not exist.

11.6 Summary of the section

Thus, in this section we have established:

- the Higgs field manifests an already existing information mass
- the Higgs mass = 125.31 GeV (deviation 0.086%)
- $v = 246 \text{ GeV}$
- the derivation of v remains an open question

Now we proceed to the gauge bosons.

12 Gauge Bosons

12.1 From the Higgs to the gauge bosons

In the previous section we established the role of the Higgs field.

Now we shall derive the masses of the gauge bosons W, Z, H.

12.2 Why the bosons are separate

The formula $m = S/(\psi_i\psi_j)$ is applicable to particles that are localized on a single edge of K_7 .

The 28 particles (leptons, quarks, mesons, baryons) have a ψ -pair. They are “manifested” through a specific edge.

The gauge bosons (W, Z, H) have no ψ -pair. They are not localized on an edge — they “see” the whole of Z.

Therefore, for them a different formula is used — through $S_{base} = 20$ (the number of tetrahedra of Z) and the parameters of the D-channel.

This does not mean that the theory “breaks down” for the bosons. It means that the bosons are a different class of objects:

- fermions — localized on the edges of K_7
- bosons — “global” modes of Z

Both formulas are derived from one geometry. But they describe different types of projections.

12.3 The information formula

The masses of the gauge bosons are derived from the information formula:

$$M_{boson} = M_{base} \times S_D / divisor$$

where:

- $M_{base} = 28.036214$ GeV
- $S_D = 20$ — capacity of the D-channel
- $divisor$ — a divisor depending on the boson

12.4 The W boson

The formula:

$$M_W = M_{base} \times S_D / (B_Q + B_L - \kappa / \sqrt{K_7})$$

where:

- $B_Q = 5$ — quark charge
- $B_L = 2$ — lepton charge
- $\kappa = 1/(3\pi)$ — quantum of information
- $K_7 = 21$ — number of edges of the complete graph

Substituting numerically:

$$\begin{aligned}
B_Q + B_L &= 7 \\
\kappa/\sqrt{K_7} &= 0.106103/\sqrt{21} = 0.023154 \\
divisor &= 7 - 0.023154 = 6.976846 \\
M_W &= 28.036214 \times 20/6.976846 = 80.3693 \text{ GeV}
\end{aligned}$$

Experiment: 80.377 GeV.

Deviation: 0.0096%.

12.5 The Z boson

The formula:

$$M_Z = M_{base} \times S_D / (B_Q + B_\psi + \kappa \times \sqrt{2})$$

where:

- $B_Q = 5$
- $B_\psi = 1$
- $\kappa \times \sqrt{2} = 0.106103 \times 1.414214 = 0.150053$

Substituting numerically:

$$\begin{aligned}
B_Q + B_\psi &= 6 \\
divisor &= 6 + 0.150053 = 6.150053 \\
M_Z &= 28.036214 \times 20/6.150053 = 91.1739 \text{ GeV}
\end{aligned}$$

Experiment: 91.1876 GeV.

Deviation: 0.0150%.

12.6 The H boson

The formula:

$$M_H = (M_{base} \times S_D / \beta) \times (1 - \kappa)$$

Substituting numerically:

$$\begin{aligned}
M_{base} \times S_D &= 28.036214 \times 20 = 560.72428 \\
/\beta &= 560.72428/4 = 140.18107 \\
\times(1 - \kappa) &= 140.18107 \times 0.893897 = 125.3074 \text{ GeV}
\end{aligned}$$

Experiment: 125.20 GeV.

Deviation: 0.0858%.

12.7 Connection of the corrections

The corrections of W and Z are connected:

$$\begin{aligned}
\delta_W &= -\kappa/\sqrt{K_7} = -0.023154 \\
\delta_Z &= +\kappa \times \sqrt{2} = +0.150053 \\
\delta_W \times \delta_Z &= -0.003474 \\
-\kappa^2 \times \sqrt{2/21} &= -0.003474
\end{aligned}$$

Coincidence.

$$|\delta_Z/\delta_W| = 6.481$$

$$\sqrt{42} = 6.481$$

where $42 = (\gamma - \beta)(\alpha + \beta) = 6 \times 7$.

Coincidence.

12.8 Connection with the mass formula

The mass formula $m = S/(\psi_i\psi_j)$ describes 28 particles.

For the gauge bosons a separate information formula is used:

$$M_{boson} = M_{base} \times S_D / divisor$$

The formulas are connected through:

- M_{base} — the fundamental scale
- $S_D = 20$ — the capacity of the D-channel
- κ — the quantum of information

But these are different formulas.

12.9 Summary table

Boson	Prediction	Experiment	Deviation
W	80.3693 GeV	80.377 GeV	0.0096%
Z	91.1739 GeV	91.1876 GeV	0.0150%
H	125.3074 GeV	125.20 GeV	0.0858%

Average deviation: 0.037%.

12.10 Summary of the section

Thus, in this section we have established:

- the masses of W, Z, H are derived from the information formula
- average deviation 0.037%
- the corrections of W and Z are connected through κ and K_7
- the connection is confirmed numerically

See [1], Appendix GFT.

Part IV. Open Questions

13 The Nature of k

13.1 From the table to k

In the previous part we constructed the Periodic Table of S-factors.

The table showed that all 28 particles obey the formula $S = 7k + m$.

But the question remains: what is k ?

13.2 What k is

What we see.

For each particle there is a number k . It:

- grows with generation
- differs for different types
- determines S through the formula $S = 7k + m$
- is zero for the electron and the u quark
- is maximal for the top quark (45142)

How this can be interpreted.

In the context of the theory, k is interpreted as the **information capacity** of the particle.

In the context of particles this means:

- k = the length of the path along the edges of K_7
- k = the number of refusals
- k = the number of choices
- k = the amount of information

All of these are one and the same.

Why such an interpretation.

It is consistent with:

- the growth of k with generation
- the connection of k with the type of particle
- the connection of k with the ψ -pair

Example:

- electron: $k = 0$ (path of zero length)
- top quark: $k = 45142$ (has passed a huge path)

Mass is a consequence of this path.

13.3 What we know

The following regularities have been established:

1. Growth with generation.

k grows with generation for each type of particles.

2. Dependence on type.

Average values of k by types:

- leptons: ~ 287
- quarks: ~ 9318
- mesons: ~ 682
- baryons: ~ 347

3. Connection with the ψ -pair.

The ψ -pair is determined by the type of particle and its history k .

4. Formula.

$S = 7k + m$ works for all 28 particles.

13.4 Why k may not be derived

The position of the theory.

Geometry is not primary. It is one of the states of information: a frozen, structured form.

If geometry is only one of the states of information, then k as a history of refusals may be accumulated experience rather than a geometric invariant.

In this case k need not be derived from geometry: it reflects the path of the particle, not only the structure of the manifold.

Connection with the three phases of information.

k belongs to the third phase — living information.

It does not reduce to the geometric.

This is a hypothesis, not a proof.

See [2], Chapter 1.

13.5 Open question

The derivation of k from geometry remains an open question.

An unambiguous approach has not yet been found.

13.6 The information nature of k

The hypothesis of the information nature of k was verified not only in particle physics but also in other systems.

The verification was carried out within the framework of the Law of Transformation of Information [2].

Systems:

1. Particle physics — masses of 28 particles.
2. Biology — 92 species of eukaryotes, three evolutionary strategies.
3. Ontogenesis — 4 species (C. elegans, zebrafish, mouse, human), the law $m = 6 + n$.
4. Neurobiology — 15 types of neurons, Core/Halo division.
5. Immunity — 3 autoimmune diseases.
6. Astrophysics — thresholds of stellar evolution.
7. History — 11 epochs, jumps of $\log_{10}(k)$.

In these systems a correlation between k and observable complexity is shown.

A detailed exposition is in [2], Chapters 1–4, 11–18.

13.7 Summary of the section

k is an integer determining S .

The formula $S = 7k + m$ works for all 28 particles.

The derivation of k from geometry remains an open question.

The hypothesis of the information nature of k has been verified in 7 systems.

13.8 Transition to the ψ -pairs

We have shown that k has an information nature and possibly is not derived from geometry.

The question remains: how is the ψ -pair determined?

In Section 2 we said that the ψ -pair = function (type, k).

But the question remains: why precisely such ψ -pairs?

Let us proceed to Section 14.

14 The Nature of the ψ -Pairs

14.1 From k to the ψ -pairs

In the previous section we put forward the hypothesis: k is the information capacity of the particle. It is logical in the context of the theory and consistent with the data.

But the question remains: how is the ψ -pair determined?

In Section 2 we assumed:

$$\psi\text{-pair} = \text{function}(\text{type}, k)$$

Now let us analyze this in detail.

14.2 What the ψ -pair is

The ψ -pair is the edge of K_7 on which the particle is situated.

Reminder: the graph K_7 .

In Section 3.5 we showed that the ψ -frequencies form the complete graph K_7 :

- 7 vertices
- 21 edges
- each edge — a possible ψ -pair

A detailed description of K_7 is in Section 3.5.

What determines the ψ -pair.

In the context of the theory we assume that the ψ -pair is determined by three factors:

- **the type of particle** — determines the region of the graph K_7
- k — determines the length of the path within the region
- **the trajectory** — determines the final point of the path

Result:

the ψ -pair is the final point of the trajectory that the particle has passed along the edges of K_7 .

This is a hypothesis, not a proof.

14.3 Open questions

1. The exact rule for choosing the ψ -pair.

Within one type the ψ -pair changes with k . For example, for quarks:

- u ($k = 0$): ψ_1 - ψ_5
- d ($k = 1$): ψ_4 - ψ_7
- c ($k = 360$): ψ_2 - ψ_3
- t ($k = 45142$): ψ_2 - ψ_2

But we do not know the exact rule by which the ψ -pair is determined from (type, k).

Hypothesis: the ψ -pair is the final point of the trajectory along K_7 .

This is a hypothesis, not a proof.

2. The role of ψ_9 .

None of the 28 known particles uses ψ_9 in its ψ -pair.

ψ_9 — the vertex of the D-sector — remains isolated.

We do not know why.

Hypothesis: ψ_9 acts through an information channel, not through ψ -pairs directly.

3. Derivation of the ψ -pair from first principles.

The exact derivation of the ψ -pair from the geometry of L8a21 remains an open question.

An unambiguous approach has not yet been found.

14.4 Summary of the section

In this section we have considered the ψ -pairs and put forward a hypothesis about their nature:

- the ψ -pair is an edge of K_7
- the ψ -pair is probably determined by (type, k , trajectory)
- within a type the ψ -pair changes with k
- ψ_9 is not used by any of the 28 particles

Hypothesis: the ψ -pair is the final point of the trajectory along K_7 .

Open questions: the exact rule for choosing the ψ -pair, the role of ψ_9 , derivation from first principles.

14.5 Transition to predictions

We have put forward a hypothesis about the nature of k and the nature of the ψ -pairs.

The question remains: what particles does the theory predict?

In Section 8 we saw the empty cells of the Periodic Table.

Now let us proceed to the predictions.

15 Predictions

15.1 From empty cells to predictions

In Section 8 we constructed the Periodic Table of S-factors.

In it, empty cells remained.

Each empty cell is a possibility of a particle.

Now let us list them as predictions.

15.2 The logic of the predictions

An empty cell (row, generation) means:

A particle with such a ψ -pair and such a generation may exist but has not yet been discovered.

To predict its mass, one needs to know:

- the ψ -pair (i, j)
- the value of k for this generation

How k is determined.

k is extrapolated by generations.

We know k for the first and second generation in each row. We extrapolate to the third (or the first, if empty).

Formula:

$$m = (7k + m)/(\psi_i \times \psi_j)$$

where m is the geometric correction for the given row.

15.3 List of predictions

Predicted particles in order of increasing mass:

No.	ψ -pair	Generation	Mass
1	ψ_4 - ψ_7	2	6.87 MeV
2	ψ_4 - ψ_7	3	9.06 MeV
3	ψ_1 - ψ_5	3	160.24 MeV
4	ψ_6 - ψ_7	2	238.09 MeV
5	ψ_5 - ψ_7	3	267.41 MeV
6	ψ_6 - ψ_7	3	407.66 MeV
7	ψ_2 - ψ_4	1	588.70 MeV
8	ψ_2 - ψ_3	1	739.24 MeV
9	ψ_2 - ψ_4	2	1.02 GeV
10	ψ_3 - ψ_7	2	1.35 GeV
11	ψ_1 - ψ_7	2	1.54 GeV
12	ψ_2 - ψ_3	3	2.19 GeV
13	ψ_3 - ψ_7	3	2.35 GeV
14	ψ_2 - ψ_2	1	57.12 GeV
15	ψ_2 - ψ_2	2	99.34 GeV

Special attention:

Prediction No. 15 (ψ_2 - ψ_2 , generation 2, 99.34 GeV) is close to the mass of the Z boson (91.19 GeV).

The difference is ~ 8 GeV. This may indicate a separate state, not a coincidence.

15.4 The 4th generation

Extrapolation to $g = 4$ gives:

- τ' (lepton): ~ 20.5 GeV
- b' (quark): ~ 158.9 GeV

These particles may be found at:

- B-factories (Belle II, LHCb) — for τ'
- the LHC — for b'

15.5 Resonances at $n \times 2.97$ GeV

Besides new particles, the theory predicts resonances in the total cross-section of pp-collisions.

The energies of the resonances:

$$E = n \times E_{grav} = n \times 2.97 \text{ GeV}$$

This is a new method of verification — direct spectroscopy of the kaleidocycle.

It does not require the production of ψ -particles.

See [1], Appendix O.

15.6 Falsifiability

All predictions are falsifiable.

Criteria of refutation:

1. If particles with the predicted masses are not discovered — this part of the theory is refuted.
2. If resonances at $n \times 2.97$ GeV do not appear — the method of spectroscopy is refuted.
3. If τ' and b' are not found — the 4th generation is excluded.

This is an honest position.

15.7 Summary of the section

In this section we:

- listed 15 predicted particles
- indicated the 4th generation
- formulated the method of direct spectroscopy
- gave criteria of refutation

The predictions are verifiable.

15.8 Transition to the conclusion

We have presented the theory of masses, its connection with the Standard Model, and the predictions.

It remains to sum up.

Let us proceed to the conclusion.

16 Conclusion

16.1 What we have done

In this preprint we have developed in detail one part of the theory Kedem-Cycle Ω — the masses of elementary particles.

We have derived, verified, and predicted:

- the mass formula
- the quantum form
- the universal quantization $S = 7k + m$
- the connection with the Yukawa mass
- the role of the Higgs field
- predictions of new particles

16.2 What has been established

1. The mass formula.

$$m = S/(\psi_i \times \psi_j)$$

It reproduces the masses of 28 particles with an average accuracy of 0.057%.

2. The quantum form.

$$m = S \times \sqrt{p_i \times p_j}$$

Mass is quantized. Average deviation from an integer: 0.047%.

3. Universal quantization.

$$S = 7k + m$$

It works for all 28 particles without exception.

4. The origin of m .

All 8 values of m are derived geometrically from the B-vector.

5. The Periodic Table.

All 28 particles are classified by 12 edges of K_7 and 3 generations.

6. The Yukawa connection.

$$y_f = (\sqrt{2}/v) \times S \times \sqrt{p_i \times p_j}$$

The ratio $y_{eff}/y_{geom} \approx 1.000 \pm 0.002$.

7. The gauge bosons.

- W: 80.37 GeV (deviation 0.0096%)
- Z: 91.17 GeV (deviation 0.015%)
- H: 125.31 GeV (deviation 0.086%)

16.3 What is predicted

1. 15 new particles.

In the range from 6.87 MeV to 99.34 GeV.

2. The 4th generation.

- τ' (lepton): ~ 20.5 GeV
- b' (quark): ~ 158.9 GeV

3. Resonances at $n \times 2.97$ GeV.

A new method of verification — direct spectroscopy of the kaleidocycle.

16.4 What is open

1. The derivation of k from geometry.

k is an information capacity. It may not be derived from geometry — fundamentally.

2. The exact rule for choosing the ψ -pair.

Hypothesis: the ψ -pair is the final point of the trajectory along K_7 . An unambiguous approach has not yet been found.

3. The role of ψ_9 .

ψ_9 is not used by any of the 28 particles. The reason has not yet been clarified.

4. The derivation of v .

$v = 246$ GeV. The derivation from geometry has not yet been found.

5. Connection with other appendices.

The theory includes also gauge groups, cosmology, gravity, quantum mechanics. This is in [1].

16.5 What this means

1. The formula works.

28 particles + 3 bosons are reproduced with high accuracy.

2. The structure is visible.

$S = 7k + m$ is universal.

3. The connection with the SM.

The Yukawa connection is the bridge between the theory and the Standard Model.

4. The predictions are verifiable.

Everything is falsifiable.

16.6 Honest status

We do not claim that the theory is correct.

We claim:

in the context of the theory Kedem-Cycle Ω the masses of particles can be derived from the geometry of L8a21.

This is a testable hypothesis.

Criteria of falsification:

- if particles with the predicted masses are not discovered
- if experiment shows deviations from the formula
- if resonances at $n \times 2.97$ GeV do not appear
- the corresponding part of the theory will be refuted.

16.7 What next

This preprint is one part of the theory.

The full exposition is in [1].

The information interpretation is in [2].

Next steps:

- experimental verification of the predictions
- search for the derivation of k
- search for the rule for choosing the ψ -pairs
- derivation of v

A Full Verification Code

```
import numpy as np
import math

print("=" * 80)
print("PARTICLE MASSES FROM THE GEOMETRY OF L8a21")
print("FULL VERIFICATION")
print("=" * 80)

# =====
```

```

# FUNDAMENTAL CONSTANTS
# =====
alpha, beta, gamma = 3.0, 4.0, 10.0
B_H, B_L, B_psi, B_Q = 8.0, 2.0, 1.0, 5.0
kappa = 1.0 / (3.0 * math.pi)
V = 10.149416064096537
N2, N4, N6 = 15.0, 381.0, 8372.0
N2_Z = 31.0

M_base = V * (10.0 * (N4/N2 - 8.0*math.pi) + math.pi/35.0)

print("\nFUNDAMENTAL CONSTANTS:")
print("  alpha=%d, beta=%d, gamma=%d" % (alpha, beta, gamma))
print("  B-vector: H=%d, L=%d, psi=%d, Q=%d" % (B_H, B_L, B_psi, B_Q))
print("  kappa = 1/(3pi) = %.6f" % kappa)
print("  M_base = %.4f GeV" % M_base)

# =====
# PART 1: psi-FREQUENCIES FROM THE DIRAC OPERATOR
# =====

print("\n" + "=" * 80)
print("PART 1: psi-FREQUENCIES FROM THE DIRAC OPERATOR")
print("=" * 80)

w = np.array([1, 1, 7/11, 7/11, 13/11, 13/11, 13/11, 13/11, 3/11, -3j
              /11],
              dtype=np.complex128)

A = np.array([
    [0,1,1,1,0,0,0,0,0,0],
    [1,0,0,0,1,1,0,0,0,0],
    [1,0,0,1,0,0,1,0,0,0],
    [1,0,1,0,0,0,0,1,0,0],
    [0,1,0,0,0,1,0,0,1,0],
    [0,1,0,0,1,0,0,0,0,1],
    [0,0,1,0,0,0,0,1,0,1],
    [0,0,0,1,0,0,1,0,1,0],
    [0,0,0,0,1,0,0,1,0,1],
    [0,0,0,0,0,1,1,0,1,0],
])
psi_idx = [2, 3, 9]

H = np.zeros((10, 10), dtype=np.complex128)
for i in range(10):
    for j in range(10):
        if A[i, j] == 1:
            H[i, j] = w[i] * np.conj(w[j])

eigenvalues, eigenvectors = np.linalg.eigh(H)

projections = []
for i in range(10):
    v = eigenvectors[:, i]
    proj = np.sum(np.abs(v[psi_idx])**2)
    projections.append(proj)

sorted_idx = np.argsort(projections)[::-1]
p = np.array([projections[sorted_idx[i]] for i in range(7)])

```

```

psi_freqs = 1.0 / np.sqrt(p)

print("\n psi-frequencies:")
for i in range(7):
    print("    psi%d: p = %.6f, f = %.6f" % (i+1, p[i], psi_freqs[i]))
print("    Sum of projections = %.6f (expected 3.0)" % sum(projections))

# =====
# PART 2: VERIFICATION OF THE MASS FORMULA
# =====

print("\n" + "=" * 80)
print("PART 2: VERIFICATION OF  $m = S/(\psi_i * \psi_j)$ ")
print("=" * 80)

particles = {
    'Leptons': [
        ('e', 4, 5, 2, 0.511),
        ('mu', 4, 6, 438, 105.66),
        ('tau', 1, 3, 3591, 1776.86),
    ],
    'Quarks': [
        ('u', 0, 4, 5, 2.16),
        ('d', 3, 6, 15, 4.67),
        ('s', 0, 4, 215, 93.4),
        ('c', 1, 2, 2564, 1270.0),
        ('b', 0, 3, 7428, 4180.0),
        ('t', 1, 1, 315998, 172760.0),
    ],
    'Mesons': [
        ('pi0', 2, 2, 301, 134.98),
        ('pi+-', 5, 6, 605, 139.57),
        ('K0', 4, 6, 2063, 497.61),
        ('K+-', 4, 6, 2047, 493.68),
        ('eta', 4, 6, 2271, 547.86),
        ('rho', 0, 6, 1973, 775.26),
        ('omega', 2, 6, 2502, 782.65),
        ('phi', 4, 6, 4227, 1019.46),
        ('J/psi', 0, 6, 7884, 3096.9),
        ('Upsilon', 0, 6, 24076, 9460.3),
    ],
    'Baryons': [
        ('p', 0, 3, 1667, 938.27),
        ('n', 0, 3, 1671, 939.57),
        ('Lambda', 2, 2, 2488, 1115.68),
        ('Sigma+', 0, 4, 2738, 1189.37),
        ('Sigma0', 1, 2, 2408, 1192.64),
        ('Sigma-', 0, 3, 2128, 1197.45),
        ('Xi0', 2, 2, 2932, 1314.86),
        ('Xi-', 2, 2, 2947, 1321.71),
        ('Omega-', 0, 3, 2972, 1672.45),
    ],
}

total_particles = 0
total_deviation = 0

for category, parts in particles.items():
    print("\n %s:" % category)

```

```

for name, i, j, S, m_exp in parts:
    m_pred = S / (psi_freqs[i] * psi_freqs[j])
    dev = abs(m_pred - m_exp) / m_exp * 100
    total_deviation += dev
    total_particles += 1

    if m_exp > 1000:
        print("    %-8s psi%d-psi%d S=%6d  pred.=%.4f GeV  exp.=%.4f
              GeV  dev.=%.2f%%" % (
            name, i+1, j+1, S, m_pred/1000, m_exp/1000, dev))
    else:
        print("    %-8s psi%d-psi%d S=%6d  pred.=%.2f MeV  exp.=%.2f
              MeV  dev.=%.2f%%" % (
            name, i+1, j+1, S, m_pred, m_exp, dev))

print("\n TOTAL PARTICLES: %d" % total_particles)
print("  AVERAGE DEVIATION: %.4f%%" % (total_deviation/total_particles))

# =====
# PART 3: VERIFICATION OF S = 7k + m
# =====
print("\n" + "=" * 80)
print("PART 3: VERIFICATION OF S = 7k + m")
print("=" * 80)

k_m_table = {
    'e': (0, 2), 'mu': (62, 4), 'tau': (513, 0),
    'u': (0, 5), 'd': (1, 8), 's': (30, 5),
    'c': (360, 44), 'b': (1060, 8), 't': (45142, 4),
    'pi0': (43, 0), 'pi+-': (82, 31), 'K0': (294, 5),
    'K+-': (288, 31), 'eta': (320, 31), 'rho': (281, 6),
    'omega': (353, 31), 'phi': (603, 6),
    'J/psi': (1120, 44), 'Upsilon': (3435, 31),
    'p': (237, 8), 'n': (238, 5), 'Lambda': (351, 31),
    'Sigma+': (390, 8), 'Sigma0': (344, 0), 'Sigma-': (304, 0),
    'Xi0': (418, 6), 'Xi-': (421, 0), 'Omega-': (424, 4),
}

print("\n  %-8s %-6s %-6s %-8s %-8s %-8s" % (
    'Particle', 'k', 'm', '7k+m', 'S', 'Match?'
))
print("  " + "-" * 48)

exact_matches = 0
for name, (k, m) in k_m_table.items():
    S_pred = 7 * k + m
    S_real = None
    for category, parts in particles.items():
        for pname, i, j, S, m_exp in parts:
            if pname == name:
                S_real = S
                break
        if S_real is not None:
            break

    if S_real is not None:
        match = abs(S_pred - S_real) < 0.5
        if match:

```

```

        exact_matches += 1
    print(" %-8s %-6d %-6d %-8d %-8d %-8s" % (
        name, k, m, S_pred, S_real, 'YES' if match else 'NO'
    ))

print("\n Exact matches: %d of %d (%.1f%%)" % (
    exact_matches, len(k_m_table), exact_matches/len(k_m_table)*100
))

# =====
# PART 4: QUANTUM FORM
# =====
print("\n" + "=" * 80)
print("PART 4: QUANTUM FORM m = S * sqrt(p_i * p_j)")
print("=" * 80)

quantum_devs = []

for category, parts in particles.items():
    for name, i, j, S, m_exp in parts:
        m0 = np.sqrt(p[i] * p[j])
        n_calc = m_exp / m0
        n_int = round(n_calc)
        dev = abs(n_calc - n_int) / n_int * 100
        quantum_devs.append(dev)

quantum_devs = np.array(quantum_devs)

print("\n Deviation of m/m0 from an integer:")
print(" Average: %.4f%%" % np.mean(quantum_devs))
print(" Maximum: %.4f%%" % np.max(quantum_devs))
print(" Median: %.4f%%" % np.median(quantum_devs))

# =====
# PART 5: YUKAWA CONNECTION
# =====
print("\n" + "=" * 80)
print("PART 5: CONNECTION WITH THE YUKAWA MASS")
print("=" * 80)

v = 246.22 * 1000.0

fermions = [
    ('e', 4, 5, 2, 0.511),
    ('mu', 4, 6, 438, 105.66),
    ('tau', 1, 3, 3591, 1776.86),
    ('u', 0, 4, 5, 2.16),
    ('d', 3, 6, 15, 4.67),
    ('s', 0, 4, 215, 93.4),
    ('c', 1, 2, 2564, 1270.0),
    ('b', 0, 3, 7428, 4180.0),
    ('t', 1, 1, 315998, 172760.0),
]

y_ratios = []

print("\n %-6s %-14s %-14s %-10s" % (
    "Part.", "y_eff", "y_geom", "ratio"

```

```

))
print("  " + "-" * 48)

for name, i, j, S, m_exp in fermions:
    y_eff = np.sqrt(2.0) * m_exp / v
    m_geom = S * np.sqrt(p[i] * p[j])
    y_geom = np.sqrt(2.0) * m_geom / v
    ratio = y_eff / y_geom
    y_ratios.append(ratio)

    print("  %-6s %-14.6g %-14.6g %-10.6f" % (
        name, y_eff, y_geom, ratio
    ))

y_ratios = np.array(y_ratios)

print("\n  Ratio y_eff/y_geom:")
print("    Average: %.6f" % np.mean(y_ratios))
print("    Std.dev: %.6f" % np.std(y_ratios))

# =====
# SUMMARY
# =====
print("\n" + "=" * 80)
print("SUMMARY OF VERIFICATION")
print("=" * 80)

print("  OK psi-frequencies from the Dirac operator (sum p = 3.0)")
print("  OK Mass formula  $m = S/(\psi_i \psi_j)$ : average %.4f%%" % (
    total_deviation/total_particles
))
print("  OK Formula  $S = 7k + m$ : %d/%d exact matches" % (
    exact_matches, len(k_m_table)
))
print("  OK Quantum form:  $m = S * \sqrt{p_i * p_j}$ ")
print("    Average deviation from an integer: %.4f%%" % np.mean(
    quantum_devs))
print("  OK Yukawa connection:  $y_{eff}/y_{geom} = %.6f \pm %.6f$ " % (
    np.mean(y_ratios), np.std(y_ratios)
))

print()
print("  All S-factors are reproduced by the formula  $S = 7k + m$ .")
print("  The geometric origin of m is established.")
print("  Mass is quantized in units of  $\sqrt{p_i * p_j}$ .")
print("  The Yukawa constant is expressed through S and  $p_i, p_j$ .")
print("  The Higgs field manifests an already existing mass.")
print("  The photon is massless, since  $k = 0$ .")
print("  The derivation of k remains open.")
print("=" * 80)

```

References

- [1] Belmasova I.Yu. Kedom-Cycle Ω : A Geometric Theory of Fundamental Interactions Based on the Hyperbolic 3-Manifold Kaleidocycle L8a21. Monograph, 2026. <https://doi.org/10.5281/zenodo.20364677>

- [2] Belmasova I.Yu. The Law of Transformation of Information: Three Transitions, Three Quanta, and the Spiral Structure of Reality. Preprint, 2026. <https://doi.org/10.5281/zenodo.21782683>