


# Statistics of Radioactive Decay II

RT4220 – Lecture #5

Kedree Proffitt

WSU

09/02/2025



# Specific Activity Terminology



**Carrier** is the STABLE portion of radioactive sample

I-131 radioactive  
I-132 stable → carrier



If there is NO stable portion it is **carrier-free**



**Specific activity** is the ratio of ACTIVITY OVER MASS of the element present

**Carrier-free specific activity** is the highest possible specific activity (100% radioactive)



High specific activity is typically desirable

If its injected: less to inject!  
If its in a machine: smaller machine!


# Practice: Specific Activity

Given two samples of Myoview (brand name Tc-99m) with the following values determine the empty values:

Sample	Activity (mCi)	Mass (g)	Spec. Act. (mCi/g)
1	0.1	100	0.001
2	200	20	10

If 10 mCi is required for a gastric clearance study, how many grams are required to be injected? If the samples all have a density of 1 g/mL, how many mLs must be injected?

Sample 1 would require a 1,000 mL or 1 L INJECTION WOW. Sample 2 would require a 1 mL injection wow




# Equilibrium

Shan't Equivocate this Equilibrium




# Equilibrium Basics

- What happens if the daughter is radioactive?
- They decay at different rates  $\rightarrow$  a race!
- This leads us to equilibria, three types:
  - **Secular**: Half-life of parent  $\gg$  daughter's
  - **Transient**: Half-life of parent  $>$  daughter's
  - **No Equilibrium**: Half-life of parent  $\leq$  daughter's




(a) transient equilibrium



(b) secular equilibrium






## Secular Equilibrium

- If the parent half-life is much longer than the daughter half-life, then the equilibrium established is called **secular equilibrium**
- The activity of the daughter closely matches the parent because there is a bottleneck
  - Not enough production
  - Much consumption
  - Seize the means

**(b) secular equilibrium**

# Secular Equilibrium Example




- Radium's half life >>> daughter → quick buildup period → maximum → same decay rate




# Transient Equilibrium

- If the half-life of the parent is longer than the daughter half-life then the equilibrium established is called **transient equilibrium**
- After some maximum where equilibrium occurs (rate of production = rate of consumption) the **RATIO** of parent:daughter activity becomes **CONSTANT**
- The decay rate of the daughter is dependent upon the decay rate of the parent




# Transient Equilibrium Example



- Technetium is ubiquitous in nuclear medicine departments for imaging
  - 140 keV gammas perfect for imaging
  - Short half-life makes patients safe shortly after procedure
- A Molybdenum generator is referred to as a “Moly-Cow”
  - Cows are milked
  - Do not milk bulls

# Transient Equilibrium Example Plot

- At the right is a plot of the activity inside the Moly-Cow
- Pattern of transient equilibrium is apparent, obfuscated by the milking process
- Hospitals typically have their own Moly Cow
- Time for maximum activity is large compared to the half-life



# Parent-Daughter Dual Decay


- When both the parent and daughter are radioactive, modeling the daughter activity becomes more difficult
- The ACTIVITY EQUATION remains the same for the PARENT:
  - $A(t) = A(0)e^{-\lambda t}$
- The ACTIVITY EQUATION for the DAUGHTER **depends on the parent** activity

# No Equilibrium


- At no point is the daughter production/decay rates equivalent
- Daughter activity will rise as it decays slower than the parent
  - Parent half-life < Daughter half-life
  - Daughter production rate < Daughter decay rate

Parent

Daughter



# Daughter Activity Equation



$$\frac{\Delta N_1}{\Delta t} = -\lambda_1 N_1$$

$$\frac{\Delta N_2}{\Delta t} = \lambda_1 N_1 - \lambda_2 N_2$$

$$A_1(t) = \left( \frac{\lambda_2}{\lambda_2 - \lambda_1} \right) \times A_1(0) \times (e^{-\lambda_1 t} - e^{-\lambda_2 t})$$



# Time to Maximum Daughter Activity

- As the parent creates daughter and the daughter decays, there is a race.
- For every set of decay constants (characteristic) there will be a time where the daughter reaches a maximum NUMBER of atoms
- We can predict this using the following equation:

$$t_m = \frac{\ln \lambda_2 / \lambda_1}{\lambda_2 - \lambda_1}$$