



Please print Name in BLOCK LETTER in accordance to NRIC/Passport (To Circle: Dr. / Mr / Ms. / Mrs / Madam)

[illegible]

Last updated: 6 Jan 2026

(Please ☒ or mark x as performed)

(Please ☒ or mark x as performed)

Please turn over to next page ...



The Singapore Life Saving Society

Adult Foreign Body Airway Obstruction - Conscious who becomes unconscious

(Please ✓ or mark x as performed)

Critical Performance		Attempt	
		1st	2nd
1	Ask: "Are you choking? I can help." Rescuer to encourage coughing and upon ascertain that the person is UNABLE to Speak, Breath and/or Cough		
2	Position - Stand behind the casualty, wrap arms around the casualty's abdomen. Landmark- Locate the navel using the Ring Finger. Place 2 finger above the navel and well below the tip of xiphoid process.		
3	Make a fist with the other hand with the thumb in the palm. Place thumbside of the fist, Against the casualty's abdomen, in the midline (i.e. 2 fingers just above navel)		
4	Lean the casualty forward with one hand, while maintaining the fist against the abdomen		
5	With your other hand, Grasp the fist and start a quick inward and upward abdominal thrusts in one motion into the casualty's abdomen		
6	Repeat thrusts until the foreign body is expelled or the casualty becomes unconscious		
7	If casualty becomes unconscious. Land person safely onto the ground		
8	Shout for help Rescuer: "Help!, Call 995 for ambulance. Get AED!"		
9	Perform 30 chest compressions (same as 1-Rescuer CPR method)		
10	Assess Airway - Open airway with Head-Tilt-Chin-Lift maneuver. Rescuer looks for foreign body in the mouth and if found, remove it with the use of hooking action with index finger.		
11	if not visible continue 30 chest compressions, then proceed back to Step 10		
12	if visible, remove it with the use of hooking action index finger (finger sweep), Attempts one ventilation (with the use of resuscitation mask) while maintaining an open airway (first attempt)		
13	If the chest does not rise, reposition with Head-Tilt-Chin-Lift maneuver, attempt second ventilation		
14	If airway is still blocked, repeat step 9		
15	Continue as instructed by Examiner and be able to state that in reality, continue until (1) casualty show sign of life (2) AED arrives & analysing heart rhythm (3) EMS takes over CPR		
16	Upon Examiner remarked : "Victim regains consciousness", place casualty in Recovery Position (The victim should be placed in the "recovery position" if the victim is (a) unresponsive, has no evidence of trauma, and is obviously breathing adequately, or (b) successfully resuscitated and has resumed breathing.)		
Result:		Please Circle	C/NYC
Comments (if any):			

Pregnant or Obese Person - Conscious

(Please ✓ or mark x as performed)

Critical Performance		Attempt	
		1st	2nd
1	Ask: "Are you choking? I can help. "Rescuer to encourage coughing and upon ascertain that the person is UNABLE to speak, Breath and/or Cough		
2	Position - Stand behind the casualty, place both arms around the casualty's armpits encircling the chest.		
3	Landmark - Make a fist with one hand, grasp the fist with the other hand and place thumb-side of the fist on the middle of the casualty's sternum (breastbone). backward thrusts on the middle of the person's chest wall/breastbone) (i.e. mid of sternum)		
4	Chest thrusts - Grasp your fist with your other hand and bring yourself close to the casualty Deliver each backward thrust firmly and distinctly. Perform continuous backward chest thrusts		
5	Continues successive inward-backward chest thrusts.		
6	Repeat thrusts until the foreign body is expelled of the person becomes unconscious		
Results		Please Circle	C/NYC
Comments (if any):			

Please turn over to next page ...



The Singapore Life Saving Society

One-Rescuer Infant CPR

(Please ✓ or mark x as performed)

Critical Performance		Attempt	
		1st	2nd
1	Check for Danger		
2	Check Responsiveness		
	• Tap the infant's shoulders and the sole of the feet		
	• Avoid unnecessary movements/prevent injuries on the head and neck		
	• Infant must be lying on his/her back on a firm, flat surface		
3	Shout to get help		
	• Call 995 for EMS		
	• Activate the emergency response system		
	• Someone to get AED		
4	Check breathing and brachial pulse (for healthcare providers)		
	• Look for rise and fall of the chest (10 seconds)		
	• If no breathing or only gasping, start CPR		
5	Locate landmark for chest compression		
	• Place index finger below nipple line		
	• Place middle and ring fingers on sternum and lift up index finger		
6	Proper chest compression technique, compress chest 3cm to 4cm vertically. Counting: 1 and 2 and 3 and 4 and 5 and 1 and 2 and 3 and 4 and 10 and 1 and 2 and 3 and 4 and 15 and 1 and 2 and 3 and 4 and 20 and 1 and 2 and 3 and 4 and 25 and 1 and 2 and 3 and 4 and 30 Chest compression rate if at 100 to 120 per minute		
7	Mouth-to-mouth and nose breathing		
	• Give 2 breaths (1 sec and 30ml of air) and watch chest rise		
8	Perform 5 cycles of 30 compression 2 ventilations		
9	Check breathing and brachial pulse (for healthcare providers)		
	• Present: Check for injury. No injury, Recovery position (lateral)		
	• Absent: Continue CPR till help arrives and take over or infant shows sign of life		
Result:		Please Circle	C/NYC
Comments (if any):			

Initiative Test

(Please ✓ or mark x as performed)

Demonstrate initiative in dealing with a simulated live casualty		Attempt	
		1st	2nd
Please specify details:			
Result:		Please Circle	C/NYC
Comments (if any):			

For Official Use Only:

Overall Result (Please circle): C (Competent) / NYC (Not Yet Competent)

Name of Examiner: _____

Date: _____

Signature of Examiner: _____