

Essentials of Analytical Geometry and Linear Algebra. Lecture 9.

Vladimir Ivanov

Innopolis University

October 22, 2021

Lecture 9. Outline

- Part 1. Quadratic curves
- Part 2. Ellipse
- Part 3. Hyperbola
- Part 4. Parabola

Curves



Part 1. Quadratic curves

In general any quadratic curve is a set of points satisfying the equation:

$$Ax^2 + Bxy + Cy^2 + Dx + Ey + F = 0$$

Without proof...

$$Ax^2 + Bxy + Cy^2 + Dx + Ey + F = 0$$

Any five (5) points on a plane uniquely define a quadratic curve.

Goals

- Understand similarities
- Understand differences
- Solve some basic problems

Part 2. Ellipse


Definition

Ellipse. Canonical equation of an ellipse

An ellipse is a plane curve, which is represented by the equation

$$\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$$

in some Cartesian coordinate system.



Parametric form of the equation of an ellipse

$$\begin{array}{l} \text{Given} \\ \left\{ \begin{array}{l} x = a \cos t \\ y = b \sin t \end{array} \right. \end{array}$$

Parametric form of the equation of an ellipse

Given

$$\begin{cases} x = a \cos t \\ y = b \sin t \end{cases}$$

and eliminating the parameter t , we get

$$\begin{cases} \frac{x^2}{a^2} = \cos^2 t \\ \frac{y^2}{b^2} = \sin^2 t \end{cases}$$

This gives you a nice way to plot a point $M(x,y)$ on an ellipse.

Question

Is this an equation of an ellipse?

$$\frac{(x - x_0)^2}{a^2} + \frac{(y - y_0)^2}{b^2} = 1$$

Question


Is this an equation of an ellipse?

$$\frac{(x - x_0)^2}{a^2} + \frac{(y - y_0)^2}{b^2} = 1$$


Center of the ellipse is at $M(x_0, y_0)$.

Ellipse: foci, eccentricity and focal distances

Foci and eccentricity



Foci and eccentricity




Foci (aka focuses)


Given an ellipse with major axis $2a$. Foci (plural from *focus*) are points $F_1(-c, 0)$ and $F_2(c, 0)$ that satisfy:

$$c^2 = a^2 - b^2$$

Foci and eccentricity




Foci and eccentricity



Eccentricity

Given an ellipse with major axis $2a$ and foci $F_1(-c, 0)$, $F_2(c, 0)$, the eccentricity of ellipse is denoted as ε : $\varepsilon = \frac{c}{a}$

Foci and eccentricity




Eccentricity

Given an ellipse with major axis $2a$ and foci $F_1(-c, 0)$, $F_2(c, 0)$, the eccentricity of ellipse is denoted as ε : $\varepsilon = \frac{c}{a}$


What is the range for eccentricity?

"Eccentric Ellipse"


THE ELLIPSE BECAME INCREASINGLY ECCENTRIC.



Focal distances



Focal distances



Focal distances

Distance from a point $M(x, y)$ on an ellipse to each of foci.

$$r_1 = a + x\varepsilon$$

$$r_2 = a - x\varepsilon$$

Proof

We need to show that $r_1 = a + x\varepsilon$.

$$r_1 = \sqrt{(x + c)^2 + y^2}$$

Proof

We need to show that $r_1 = a + x\varepsilon$.

$$r_1 = \sqrt{(x + c)^2 + y^2}$$

$$y^2 = (a^2 - x^2) \frac{b^2}{a^2}$$

Proof

We need to show that $r_1 = a + x\varepsilon$.

$$r_1 = \sqrt{(x + c)^2 + y^2}$$

$$y^2 = (a^2 - x^2) \frac{b^2}{a^2}$$

$$c = a\varepsilon; \text{ Note also: } b^2 = a^2 - c^2 = a^2(1 - \varepsilon^2)$$

Proof

We need to show that $r_1 = a + x\varepsilon$.

$$r_1 = \sqrt{(x + c)^2 + y^2}$$

$$y^2 = (a^2 - x^2) \frac{b^2}{a^2}$$

$$c = a\varepsilon; \text{ Note also: } b^2 = a^2 - c^2 = a^2(1 - \varepsilon^2)$$

$$y^2 = (a^2 - x^2) \frac{b^2}{a^2} = a^2 - a^2\varepsilon^2 - x^2 + x^2\varepsilon^2$$

Proof

We need to show that $r_1 = a + x\varepsilon$.

$$r_1 = \sqrt{(x+c)^2 + y^2}$$

$$y^2 = (a^2 - x^2) \frac{b^2}{a^2}$$


$$c = a\varepsilon; \text{ Note also: } b^2 = a^2 - c^2 = a^2(1 - \varepsilon^2)$$

$$y^2 = (a^2 - x^2) \frac{b^2}{a^2} = a^2 - a^2\varepsilon^2 - x^2 + x^2\varepsilon^2$$


$$r_1^2 = (x+c)^2 + y^2 = x^2 + 2xc + c^2 + a^2 - a^2\varepsilon^2 - x^2 + x^2\varepsilon^2 = (a + x\varepsilon)^2$$

Note, that $r_2 = a - x\varepsilon$ and hence $r_1 + r_2 = 2a$


Directrices



What are the equations of the directrices?




Why $\frac{r_1}{d_1} = \frac{r_2}{d_2} = \varepsilon$?



Why $\frac{r_1}{d_1} = \frac{r_2}{d_2} = \varepsilon$?

$$d_1 = \frac{a}{\varepsilon} + x = \frac{1}{\varepsilon}r_1; d_2 = \frac{a}{\varepsilon} - x = \frac{1}{\varepsilon}r_2$$

Tangent lines



Example

Check whether this equation is an equation of ellipse?

$$2x^2 + 4x + 3y^2 - 12 = 1$$

Break, 5 min.

Interesting question to study:

Propose a formula for the length (perimeter) of an ellipse.

or

Write a program to calculate it.

Part 3. Hyperbola

Hyperbola

Definition. Canonical equation

A hyperbola is a plane curve, which can be represented in some Cartesian coordinate system by one of the below equations


$$\frac{x^2}{a^2} - \frac{y^2}{b^2} = 1 \quad \text{or} \quad \frac{x^2}{a^2} - \frac{y^2}{b^2} = -1$$

Hyperbola

Definition. Canonical equation

A hyperbola is a plane curve, which can be represented in some Cartesian coordinate system by one of the below equations

$$\frac{x^2}{a^2} - \frac{y^2}{b^2} = 1 \quad \text{or} \quad \frac{y^2}{b^2} - \frac{x^2}{a^2} = 1$$




Hyperbola

Definition. Canonical equation

A hyperbola is a plane curve, which can be represented in some Cartesian coordinate system by one of the below equations

$$\frac{x^2}{a^2} - \frac{y^2}{b^2} = 1 \quad \text{or} \quad \frac{y^2}{b^2} - \frac{x^2}{a^2} = 1$$



If $a = b$, then it is a **equilateral hyperbola**.

Question

Is this an equation of a hyperbola?

$$\frac{(x - x_0)^2}{a^2} - \frac{(y - y_0)^2}{b^2} = \pm 1$$

Question


Is this an equation of a hyperbola?

$$\frac{(x - x_0)^2}{a^2} - \frac{(y - y_0)^2}{b^2} = \pm 1$$


Center of the hyperbola is at $M(x_0, y_0)$.

Hyperbola: foci, eccentricity and focal distances

Foci and eccentricity



Foci and eccentricity




Foci (aka focuses)


Given a hyperbola $\frac{x^2}{a^2} - \frac{y^2}{b^2} = 1$. Foci are points $F_1(-c, 0)$ and $F_2(c, 0)$ that satisfy:

$$c^2 = a^2 + b^2$$

Foci and eccentricity



Foci and eccentricity




Eccentricity

Given $\frac{x^2}{a^2} - \frac{y^2}{b^2} = 1$. Eccentricity of the hyperbola ε :


$$\varepsilon = \frac{c}{a}$$

Note, $\varepsilon > 1$

Focal distances



Focal distances




Focal distances

Distance from a point $M(x, y)$ on a hyperbola to each of foci.

$$r_1 = \pm(x\varepsilon + a)$$

$$r_2 = \pm(x\varepsilon - a)$$

Focal distances



Focal distances

Distance from a point $M(x, y)$ on a hyperbola to each of foci.


$$r_1 = \pm(x\varepsilon + a)$$

$$r_2 = \pm(x\varepsilon - a)$$

For any point of hyperbola:

$$r_1 - r_2 = \pm 2a$$

Focal distances



$$r_1 = \pm(x\varepsilon + a)$$

$$r_2 = \pm(x\varepsilon - a)$$

$$r_1 - r_2 = \pm 2a \text{ (or, just } |r_1 - r_2| = 2a)$$

Why do we use \pm here?


Hint: we have 2 branches (for one x is positive for another one $x < 0$)

Assignment

Define foci, eccentricity and focal distances if a hyperbola has the equation:

$$\frac{x^2}{a^2} - \frac{y^2}{b^2} = -1$$

Directrices



Directrices (plural from *directrix*) are two vertical lines $x = \pm \frac{a}{\varepsilon}$


$$\frac{r_1}{d_1} = \frac{r_2}{d_2} = \varepsilon$$

Directrix is not DirectX

Directrix a fixed line used in describing a curve or surface.

Thus, Ellipse, Hyperbola (and also a parabola) can be defined using a **directrix** and a **point**.

Asymptotes



Given a hyperbola $\frac{x^2}{a^2} - \frac{y^2}{b^2} = 1$

Two straight lines $y = \pm \frac{b}{a}x$ are the asymptotes of the hyperbola.

Use limits to prove it (as $x \rightarrow \pm\infty$).


Example

Reduce the equation


$$x^2 - 6x + y^2 + 8y = 0$$

to the canonical form

Part 4. Parabola




Parabola. Definitions



Parabola

- A parabola is the locus of points, which are equidistant from a given point **F** and line **L**.


Parabola. Definitions



Parabola

- A parabola is the locus of points, which are equidistant from a given point **F** and line **L**.
- The point **F** is called the **focus**.

Parabola. Definitions




Parabola

- A parabola is the locus of points, which are equidistant from a given point **F** and line **L**.
- The point **F** is called the **focus**.
- The line **L** is called the **directrix** of the parabola.


Sometimes, p is denoted as $2a$.

Parabola. Canonical equation



$$d = r,$$


Parabola. Canonical equation



$$d = r,$$

$$x + \frac{p}{2} = \sqrt{\left(x - \frac{p}{2}\right)^2 + y^2}$$

Parabola. Canonical equation






$$d = r,$$

$$x + \frac{p}{2} = \sqrt{(x - \frac{p}{2})^2 + y^2}$$

$$y^2 = 2px$$

Parabola. Other cases



Write the canonical equations.

Parabola. Eccentricity

Recall how we defined eccentricity for ellipse and hyperbola.

Parabola. Eccentricity

Recall how we defined eccentricity for ellipse and hyperbola.
What is the eccentricity of a parabola?

Parabola. Eccentricity

Recall how we defined eccentricity for ellipse and hyperbola.
What is the eccentricity of a parabola?

$$\varepsilon = \frac{r}{d} = 1$$


Reduce the equation

$$x^2 + 4x - 3y = -5$$

to the canonical form.



Interesting case 1

- $x^2 - y^2 = 0$ (a pair of intersecting lines)




Interesting case 2

- $2x^2 + 3y^2 = -1$ (an imaginary ellipse)




Summary

Summary.




Quadratic curves as sections of a circle cone

Conics!



Degenerate conics



Relation to Quadratic forms and Matrices

Conic sections are the sets of points whose coordinates satisfy a second-degree polynomial equation (A, B, C, D, E, F are numbers):

$$Q(x, y) = Ax^2 + Bxy + Cy^2 + Dx + Ey + F = 0.$$

Relation to Quadratic forms and Matrices

Conic sections are the sets of points whose coordinates satisfy a second-degree polynomial equation (A, B, C, D, E, F are numbers):

$$Q(x, y) = Ax^2 + Bxy + Cy^2 + Dx + Ey + F = 0.$$

In matrix form (it is the **same** equation):

$$\begin{bmatrix} x & y \end{bmatrix} \begin{bmatrix} A & B/2 \\ B/2 & C \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} + \begin{bmatrix} D & E \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} + F = 0.$$

Relation to Quadratic forms and Matrices

Conic sections are the sets of points whose coordinates satisfy a second-degree polynomial equation (A, B, C, D, E, F are numbers):

$$Q(x, y) = Ax^2 + Bxy + Cy^2 + Dx + Ey + F = 0.$$

In matrix form (it is the **same** equation):

$$\begin{bmatrix} x & y \end{bmatrix} \begin{bmatrix} A & B/2 \\ B/2 & C \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} + \begin{bmatrix} D & E \end{bmatrix} \begin{bmatrix} x \\ y \end{bmatrix} + F = 0.$$

This following expression is called the **quadratic form**: $Ax^2 + Bxy + Cy^2$.

Matrix of the **quadratic form** : $\begin{bmatrix} A & B/2 \\ B/2 & C \end{bmatrix}$

**WHAT IF THIS IS NOT
USELESS**

**AND CONIC SECTIONS ARE
EVERYWHERE**

memegenerator.net

Useful links

- <https://www.geogebra.org>
- https://youtu.be/fNk_zzaMoSs
- <http://immersivemath.com/ila>