

# Theoretical Computer Science

## Tutorial Week 5

Prof. Andrey Frolov



## Regular languages

- **Myhill-Nerode criteria**
  - Positive Examples
  - Negative Examples
- Pumping Lemma

# Regular Languages

## Definition

A language is called regular, if it is recognized by a FSA.

## Problem

Which languages are regular?

# Myhill-Nerode criteria


For a language  $L$  over an alphabet  $A$ ,

$$s_1 \equiv_L s_2 \Leftrightarrow (\forall t \in A^*) (s_1 t \in L \leftrightarrow s_2 t \in L)$$

$\equiv_L$  is an equivalence relation

# Myhill-Nerode criteria

What are equivalence relations in general?



# Myhill-Nerode criteria

For a language  $L$  over an alphabet  $A$ ,

$$s_1 \not\equiv_L s_2 \Leftrightarrow (\exists t \in A^*) [(s_1 t \notin L \& s_2 t \in L) \vee (s_1 t \in L \& s_2 t \notin L)]$$

$t$  is called a **distinguishing extension**.

## Myhill-Nerode theorem

A language  $L$  is regular iff  $\equiv_L$  has a finite number of equivalent classes.

## Regular languages

- Myhill-Nerode criteria
  - **Positive Examples**
  - Negative Examples
- Pumping Lemma



# Myhill-Nerode method. Examples

$$s_1 \not\equiv_L s_2 \Leftrightarrow (\exists t \in A^*) [(s_1 t \notin L \& s_2 t \in L) \vee (s_1 t \in L \& s_2 t \notin L)]$$

Example 1:  $L_1 = \{0x \mid x \in \Sigma^*\}$ , where  $\Sigma = \{0, 1\}$

# Myhill-Nerode method. Examples

$$s_1 \not\equiv_L s_2 \Leftrightarrow (\exists t \in A^*) [(s_1 t \notin L \& s_2 t \in L) \vee (s_1 t \in L \& s_2 t \notin L)]$$

Example 1:  $L_1 = \{0x \mid x \in \Sigma^*\}$ , where  $\Sigma = \{0, 1\}$   
 $\epsilon$ ,

# Myhill-Nerode method. Examples

$$s_1 \not\equiv_L s_2 \Leftrightarrow (\exists t \in A^*) [(s_1 t \notin L \& s_2 t \in L) \vee (s_1 t \in L \& s_2 t \notin L)]$$

Example 1:  $L_1 = \{0x \mid x \in \Sigma^*\}$ , where  $\Sigma = \{0, 1\}$   
 $\epsilon, 0,$

# Myhill-Nerode method. Examples

$$s_1 \not\equiv_L s_2 \Leftrightarrow (\exists t \in A^*) [(s_1 t \notin L \& s_2 t \in L) \vee (s_1 t \in L \& s_2 t \notin L)]$$

Example 1:  $L_1 = \{0x \mid x \in \Sigma^*\}$ , where  $\Sigma = \{0, 1\}$   
 $\epsilon, 0, 1,$

# Myhill-Nerode method. Examples

$$s_1 \not\equiv_L s_2 \Leftrightarrow (\exists t \in A^*) [(s_1 t \notin L \& s_2 t \in L) \vee (s_1 t \in L \& s_2 t \notin L)]$$

Example 1:  $L_1 = \{0x \mid x \in \Sigma^*\}$ , where  $\Sigma = \{0, 1\}$   
 $\epsilon, 0, 1, 00,$

# Myhill-Nerode method. Examples

$$s_1 \not\equiv_L s_2 \Leftrightarrow (\exists t \in A^*) [(s_1 t \notin L \& s_2 t \in L) \vee (s_1 t \in L \& s_2 t \notin L)]$$

Example 1:  $L_1 = \{0x \mid x \in \Sigma^*\}$ , where  $\Sigma = \{0, 1\}$   
 $\epsilon, 0, 1, 00, 01,$

# Myhill-Nerode method. Examples

$$s_1 \not\equiv_L s_2 \Leftrightarrow (\exists t \in A^*) [(s_1 t \notin L \& s_2 t \in L) \vee (s_1 t \in L \& s_2 t \notin L)]$$

Example 1:  $L_1 = \{0x \mid x \in \Sigma^*\}$ , where  $\Sigma = \{0, 1\}$   
 $\epsilon, 0, 1, 00, 01, 10, 11, \dots$

1.  $\epsilon$

# Myhill-Nerode method. Examples

$$s_1 \not\equiv_L s_2 \Leftrightarrow (\exists t \in A^*) [(s_1 t \notin L \& s_2 t \in L) \vee (s_1 t \in L \& s_2 t \notin L)]$$

Example 1:  $L_1 = \{0x \mid x \in \Sigma^*\}$ , where  $\Sigma = \{0, 1\}$   
 $\epsilon, 0, 1, 00, 01, 10, 11, \dots$

1.  $\epsilon$
2. 0:  $0 \not\equiv_{L_1} \epsilon$ , since  $0 \cdot \epsilon \notin L_1 \& \epsilon \cdot \epsilon \in L_1$  (a disting. ext. is  $\epsilon$ )



# Myhill-Nerode method. Examples

$$s_1 \not\equiv_L s_2 \Leftrightarrow (\exists t \in A^*) [(s_1 t \notin L \& s_2 t \in L) \vee (s_1 t \in L \& s_2 t \notin L)]$$

Example 1:  $L_1 = \{0x \mid x \in \Sigma^*\}$ , where  $\Sigma = \{0, 1\}$   
 $\epsilon, 0, 1, 00, 01, 10, 11, \dots$

1.  $\epsilon$
2. 0:  $0 \not\equiv_{L_1} \epsilon$ , since  $0 \cdot \epsilon \notin L_1 \& \epsilon \cdot \epsilon \in L_1$  (a disting. ext. is  $\epsilon$ )
3. 1:

# Myhill-Nerode method. Examples

$$s_1 \not\equiv_L s_2 \Leftrightarrow (\exists t \in A^*) [(s_1 t \notin L \& s_2 t \in L) \vee (s_1 t \in L \& s_2 t \notin L)]$$

Example 1:  $L_1 = \{0x \mid x \in \Sigma^*\}$ , where  $\Sigma = \{0, 1\}$   
 $\epsilon, 0, 1, 00, 01, 10, 11, \dots$

1.  $\epsilon$
2. 0:  $0 \not\equiv_{L_1} \epsilon$ , since  $0 \cdot \epsilon \notin L_1 \& \epsilon \cdot \epsilon \in L_1$  (a disting. ext. is  $\epsilon$ )
3. 1:  
 $1 \not\equiv_{L_1} 0$ , since  $1 \notin L_1, 0 \in L_1$

# Myhill-Nerode method. Examples

$$s_1 \not\equiv_L s_2 \Leftrightarrow (\exists t \in A^*) [(s_1 t \notin L \& s_2 t \in L) \vee (s_1 t \in L \& s_2 t \notin L)]$$

Example 1:  $L_1 = \{0x \mid x \in \Sigma^*\}$ , where  $\Sigma = \{0, 1\}$   
 $\epsilon, 0, 1, 00, 01, 10, 11, \dots$

1.  $\epsilon$
2. 0:  $0 \not\equiv_{L_1} \epsilon$ , since  $0 \cdot \epsilon \notin L_1 \& \epsilon \cdot \epsilon \in L_1$  (a disting. ext. is  $\epsilon$ )
3. 1:
  - $1 \not\equiv_{L_1} 0$ , since  $1 \notin L_1, 0 \in L_1$
  - $1 \not\equiv_{L_1} \epsilon$ , since  $1 \cdot 0 \notin L_1, \epsilon \cdot 0 \in L_1$  (a distinguishing ext. is 0)

# Myhill-Nerode method. Examples

$$s_1 \not\equiv_L s_2 \Leftrightarrow (\exists t \in A^*) [(s_1 t \notin L \& s_2 t \in L) \vee (s_1 t \in L \& s_2 t \notin L)]$$

Example 1:  $L_1 = \{0x \mid x \in \Sigma^*\}$ , where  $\Sigma = \{0, 1\}$   
 $\epsilon, 0, 1, 00, 01, 10, 11, \dots$

1.  $\epsilon$
2.  $0: 0 \not\equiv_{L_1} \epsilon$ , since  $0 \cdot \epsilon \notin L_1 \& \epsilon \cdot \epsilon \in L_1$  (a disting. ext. is  $\epsilon$ )
3.  $1:$   
 $1 \not\equiv_{L_1} 0$ , since  $1 \notin L_1, 0 \in L_1$   
 $1 \not\equiv_{L_1} \epsilon$ , since  $1 \cdot 0 \notin L_1, \epsilon \cdot 0 \in L_1$  (a distinguishing ext. is  $0$ )
4.  $0t \equiv_{L_1} 0$

# Myhill-Nerode method. Examples

$$s_1 \not\equiv_L s_2 \Leftrightarrow (\exists t \in A^*) [(s_1 t \notin L \& s_2 t \in L) \vee (s_1 t \in L \& s_2 t \notin L)]$$

Example 1:  $L_1 = \{0x \mid x \in \Sigma^*\}$ , where  $\Sigma = \{0, 1\}$   
 $\epsilon, 0, 1, 00, 01, 10, 11, \dots$

1.  $\epsilon$
2.  $0: 0 \not\equiv_{L_1} \epsilon$ , since  $0 \cdot \epsilon \notin L_1 \& \epsilon \cdot \epsilon \in L_1$  (a disting. ext. is  $\epsilon$ )
3.  $1:$   
 $1 \not\equiv_{L_1} 0$ , since  $1 \notin L_1, 0 \in L_1$   
 $1 \not\equiv_{L_1} \epsilon$ , since  $1 \cdot 0 \notin L_1, \epsilon \cdot 0 \in L_1$  (a distinguishing ext. is  $0$ )
4.  $0t \equiv_{L_1} 0$
5.  $1t \equiv_{L_1} 1$

# Myhill-Nerode method. Examples

Example 1:  $L_1 = \{0x \mid x \in \Sigma^*\}$

$[\epsilon] = \{\epsilon\}$ ,  $[0] = \{0x \mid x \in \Sigma^*\} = L_1$ ,  $[1] = \{1x \mid x \in \Sigma^*\}$

$\delta$	0	1
$\rightarrow [\epsilon]$		
$*[0]$		
$[1]$		

# Myhill-Nerode method. Examples

Example 1:  $L_1 = \{0x \mid x \in \Sigma^*\}$

$[\epsilon] = \{\epsilon\}$ ,  $[0] = \{0x \mid x \in \Sigma^*\} = L_1$ ,  $[1] = \{1x \mid x \in \Sigma^*\}$

$\delta$	0	1
$\rightarrow [\epsilon]$	$[0]$	
$^*[0]$		
$[1]$		

# Myhill-Nerode method. Examples

Example 1:  $L_1 = \{0x \mid x \in \Sigma^*\}$

$[\epsilon] = \{\epsilon\}$ ,  $[0] = \{0x \mid x \in \Sigma^*\} = L_1$ ,  $[1] = \{1x \mid x \in \Sigma^*\}$

$\delta$	0	1
$\rightarrow [\epsilon]$	$[0]$	$[1]$
$^*[0]$		
$[1]$		



# Myhill-Nerode method. Examples

Example 1:  $L_1 = \{0x \mid x \in \Sigma^*\}$

$[\epsilon] = \{\epsilon\}$ ,  $[0] = \{0x \mid x \in \Sigma^*\} = L_1$ ,  $[1] = \{1x \mid x \in \Sigma^*\}$

$\delta$	0	1
$\rightarrow [\epsilon]$	$[0]$	$[1]$
$*[0]$	$[0]$	
$[1]$		

# Myhill-Nerode method. Examples

Example 1:  $L_1 = \{0x \mid x \in \Sigma^*\}$

$[\epsilon] = \{\epsilon\}$ ,  $[0] = \{0x \mid x \in \Sigma^*\} = L_1$ ,  $[1] = \{1x \mid x \in \Sigma^*\}$

$\delta$	0	1
$\rightarrow [\epsilon]$	$[0]$	$[1]$
$*[0]$	$[0]$	$[0]$
$[1]$		

# Myhill-Nerode method. Examples

Example 1:  $L_1 = \{0x \mid x \in \Sigma^*\}$

$[\epsilon] = \{\epsilon\}$ ,  $[0] = \{0x \mid x \in \Sigma^*\} = L_1$ ,  $[1] = \{1x \mid x \in \Sigma^*\}$

$\delta$	0	1
$\rightarrow [\epsilon]$	$[0]$	$[1]$
$*[0]$	$[0]$	$[0]$
$[1]$	$[1]$	

# Myhill-Nerode method. Examples


Example 1:  $L_1 = \{0x \mid x \in \Sigma^*\}$

$[\epsilon] = \{\epsilon\}$ ,  $[0] = \{0x \mid x \in \Sigma^*\} = L_1$ ,  $[1] = \{1x \mid x \in \Sigma^*\}$

$\delta$	0	1
$\rightarrow [\epsilon]$	$[0]$	$[1]$
$^*[0]$	$[0]$	$[0]$
$[1]$	$[1]$	$[1]$


# Myhill-Nerode method. Examples

Example 1:  $L_1 = \{0x \mid x \in \Sigma^*\}$



# Myhill-Nerode method. Examples

Example 1:  $L_1 = \{0x \mid x \in \Sigma^*\}$



# Myhill-Nerode method. Examples

$$s_1 \not\equiv_L s_2 \Leftrightarrow (\exists t \in A^*) [(s_1 t \notin L \& s_2 t \in L) \vee (s_1 t \in L \& s_2 t \notin L)]$$

Example 2:  $L_2 = \{x00 \mid x \in \Sigma^*\}$

1.  $\epsilon$

# Myhill-Nerode method. Examples

$$s_1 \not\equiv_L s_2 \Leftrightarrow (\exists t \in A^*) [(s_1 t \notin L \& s_2 t \in L) \vee (s_1 t \in L \& s_2 t \notin L)]$$

Example 2:  $L_2 = \{x00 \mid x \in \Sigma^*\}$

1.  $\epsilon$
2. 0:  $0 \not\equiv_{L_2} \epsilon$ , since  $00 \in L_2$  &  $\epsilon 0 = 0 \notin L_2$  (a disting. ext. is 0)



# Myhill-Nerode method. Examples

$$s_1 \not\equiv_L s_2 \Leftrightarrow (\exists t \in A^*) [(s_1 t \notin L \& s_2 t \in L) \vee (s_1 t \in L \& s_2 t \notin L)]$$

Example 2:  $L_2 = \{x00 \mid x \in \Sigma^*\}$

1.  $\epsilon$
2.  $0: 0 \not\equiv_{L_2} \epsilon$ , since  $00 \in L_2$  &  $\epsilon 0 = 0 \notin L_2$  (a disting. ext. is 0)
3.  $1: 1 \equiv_{L_2} \epsilon$ ,  $t1 \equiv_{L_2} \epsilon$

# Myhill-Nerode method. Examples

$$s_1 \not\equiv_L s_2 \Leftrightarrow (\exists t \in A^*) [(s_1 t \notin L \& s_2 t \in L) \vee (s_1 t \in L \& s_2 t \notin L)]$$

Example 2:  $L_2 = \{x00 \mid x \in \Sigma^*\}$

1.  $\epsilon$
2.  $0: 0 \not\equiv_{L_2} \epsilon$ , since  $00 \in L_2$  &  $\epsilon 0 = 0 \notin L_2$  (a disting. ext. is 0)
3.  $1: 1 \equiv_{L_2} \epsilon$ ,  $t1 \equiv_{L_2} \epsilon$
4.  $00:$

# Myhill-Nerode method. Examples

$$s_1 \not\equiv_L s_2 \Leftrightarrow (\exists t \in A^*) [(s_1 t \notin L \& s_2 t \in L) \vee (s_1 t \in L \& s_2 t \notin L)]$$

Example 2:  $L_2 = \{x00 \mid x \in \Sigma^*\}$

1.  $\epsilon$
2.  $0: 0 \not\equiv_{L_2} \epsilon$ , since  $00 \in L_2$  &  $\epsilon 0 = 0 \notin L_2$  (a disting. ext. is 0)
3.  $1: 1 \equiv_{L_2} \epsilon$ ,  $t1 \equiv_{L_2} \epsilon$
4.  $00$ :  
 $00 \not\equiv_{L_2} \epsilon$ , since  $00 \in L_2$  &  $\epsilon \notin L_2$  (a distinguishing ext. is  $\epsilon$ )

# Myhill-Nerode method. Examples

$$s_1 \not\equiv_L s_2 \Leftrightarrow (\exists t \in A^*) [(s_1 t \notin L \& s_2 t \in L) \vee (s_1 t \in L \& s_2 t \notin L)]$$

Example 2:  $L_2 = \{x00 \mid x \in \Sigma^*\}$

1.  $\epsilon$
2.  $0: 0 \not\equiv_{L_2} \epsilon$ , since  $00 \in L_2 \& \epsilon 0 = 0 \notin L_2$  (a disting. ext. is 0)
3.  $1: 1 \equiv_{L_2} \epsilon$ ,  $t1 \equiv_{L_2} \epsilon$
4.  $00$ :  
 $00 \not\equiv_{L_2} \epsilon$ , since  $00 \in L_2 \& \epsilon \notin L_2$  (a distinguishing ext. is  $\epsilon$ )  
 $00 \not\equiv_{L_2} 0$ , since  $00 \in L_2 \& 0 \notin L_2$  (a distinguishing ext. is  $\epsilon$ )

# Myhill-Nerode method. Examples

Example 2:  $L_2 = \{x00 \mid x \in \Sigma^*\}$

$$[\epsilon] = \{\epsilon\} \cup \{x1 \mid x \in \Sigma^*\}, [0] = \{x10 \mid x \in \Sigma^*\}, \\ [00] = \{x00 \mid x \in \Sigma^*\} = L_2$$

$\delta$	0	1
$\rightarrow [\epsilon]$		
$[0]$		
$*[00]$		

# Myhill-Nerode method. Examples

Example 2:  $L_2 = \{x00 \mid x \in \Sigma^*\}$

$$[\epsilon] = \{\epsilon\} \cup \{x1 \mid x \in \Sigma^*\}, [0] = \{x10 \mid x \in \Sigma^*\}, \\ [00] = \{x00 \mid x \in \Sigma^*\} = L_2$$

$\delta$	0	1
$\rightarrow [\epsilon]$	$[0]$	
$[0]$		
$*[00]$		

# Myhill-Nerode method. Examples

Example 2:  $L_2 = \{x00 \mid x \in \Sigma^*\}$

$$[\epsilon] = \{\epsilon\} \cup \{x1 \mid x \in \Sigma^*\}, [0] = \{x10 \mid x \in \Sigma^*\}, \\ [00] = \{x00 \mid x \in \Sigma^*\} = L_2$$

$\delta$	0	1
$\rightarrow [\epsilon]$	$[0]$	$[\epsilon]$
$[0]$		
$*[00]$		

# Myhill-Nerode method. Examples

Example 2:  $L_2 = \{x00 \mid x \in \Sigma^*\}$

$$[\epsilon] = \{\epsilon\} \cup \{x1 \mid x \in \Sigma^*\}, [0] = \{x10 \mid x \in \Sigma^*\}, \\ [00] = \{x00 \mid x \in \Sigma^*\} = L_2$$

$\delta$	0	1
$\rightarrow [\epsilon]$	$[0]$	$[\epsilon]$
$[0]$	$[00]$	
$*[00]$		



# Myhill-Nerode method. Examples

Example 2:  $L_2 = \{x00 \mid x \in \Sigma^*\}$

$$[\epsilon] = \{\epsilon\} \cup \{x1 \mid x \in \Sigma^*\}, [0] = \{x10 \mid x \in \Sigma^*\}, \\ [00] = \{x00 \mid x \in \Sigma^*\} = L_2$$

$\delta$	0	1
$\rightarrow [\epsilon]$	$[0]$	$[\epsilon]$
$[0]$	$[00]$	$[\epsilon]$
$*[00]$		

# Myhill-Nerode method. Examples

Example 2:  $L_2 = \{x00 \mid x \in \Sigma^*\}$

$$[\epsilon] = \{\epsilon\} \cup \{x1 \mid x \in \Sigma^*\}, [0] = \{x10 \mid x \in \Sigma^*\}, \\ [00] = \{x00 \mid x \in \Sigma^*\} = L_2$$

$\delta$	0	1
$\rightarrow [\epsilon]$	$[0]$	$[\epsilon]$
$[0]$	$[00]$	$[\epsilon]$
$*[00]$	$[00]$	

# Myhill-Nerode method. Examples


Example 2:  $L_2 = \{x00 \mid x \in \Sigma^*\}$

$$[\epsilon] = \{\epsilon\} \cup \{x1 \mid x \in \Sigma^*\}, [0] = \{x10 \mid x \in \Sigma^*\}, \\ [00] = \{x00 \mid x \in \Sigma^*\} = L_2$$

$\delta$	0	1
$\rightarrow [\epsilon]$	$[0]$	$[\epsilon]$
$[0]$	$[00]$	$[\epsilon]$
$*[00]$	$[00]$	$[\epsilon]$

# Myhill-Nerode method. Examples

Example 2:  $L_2 = \{x00 \mid x \in \Sigma^*\}$



# Myhill-Nerode method. Examples

Example 3:  $L_3 = \{x \in \Sigma^* \mid x \text{ is a binary representation of an integer divisible by 5 and it begins with 1}\}$ , where  $\Sigma = \{0, 1\}$

1.  $[\epsilon]$
2.  $[0] = \{0x \mid x \in \Sigma^*\}$
3.  $[1] = \{x \mid \text{the remainder after dividing } x \text{ by 5 is 1}\}$
4.  $[10] = \{x \mid \text{the remainder after dividing } x \text{ by 5 is 2}\}$
5.  $[11] = \{x \mid \text{the remainder after dividing } x \text{ by 5 is 3}\}$
6.  $[100] = \{x \mid \text{the remainder after dividing } x \text{ by 5 is 4}\}$
7.  $[101] = \{x \mid x \text{ is divisible by 5}\}$

# Myhill-Nerode method. Examples

Example 3:  $L_3 = \{x \in \Sigma^* \mid x \text{ is a binary representation of an integer divisible by 5 and it begins with 1}\}$

$\delta$	0	1
$\rightarrow [\epsilon]$	[0]	[1]
[0]	[0]	[0]
[1]		
[10]		
[11]		
[100]		
*[101]		

# Myhill-Nerode method. Examples

Example 3:  $L_3 = \{x \in \Sigma^* \mid x \text{ is a binary representation of an integer divisible by 5 and it begins with 1}\}$

$\delta$	0	1
$\rightarrow [\epsilon]$	[0]	[1]
[0]	[0]	[0]
[1]	[10]	
[10]		
[11]		
[100]		
*[101]		

# Myhill-Nerode method. Examples

Example 3:  $L_3 = \{x \in \Sigma^* \mid x \text{ is a binary representation of an integer divisible by 5 and it begins with 1}\}$

$\delta$	0	1
$\rightarrow [\epsilon]$	[0]	[1]
[0]	[0]	[0]
[1]	[10]	[11]
[10]		
[11]		
[100]		
*[101]		



# Myhill-Nerode method. Examples

Example 3:  $L_3 = \{x \in \Sigma^* \mid x \text{ is a binary representation of an integer divisible by 5 and it begins with 1}\}$

$\delta$	0	1
$\rightarrow [\epsilon]$	[0]	[1]
[0]	[0]	[0]
[1]	[10]	[11]
[10]	[100]	
[11]		
[100]		
*[101]		

# Myhill-Nerode method. Examples

Example 3:  $L_3 = \{x \in \Sigma^* \mid x \text{ is a binary representation of an integer divisible by 5 and it begins with 1}\}$

$\delta$	0	1
$\rightarrow [\epsilon]$	[0]	[1]
[0]	[0]	[0]
[1]	[10]	[11]
[10]	[100]	[101]
[11]		
[100]		
*[101]		

# Myhill-Nerode method. Examples

Example 3:  $L_3 = \{x \in \Sigma^* \mid x \text{ is a binary representation of an integer divisible by 5 and it begins with 1}\}$

$\delta$	0	1
$\rightarrow [\epsilon]$	[0]	[1]
[0]	[0]	[0]
[1]	[10]	[11]
[10]	[100]	[101]
[11]	[1]	
[100]		
*[101]		

# Myhill-Nerode method. Examples

Example 3:  $L_3 = \{x \in \Sigma^* \mid x \text{ is a binary representation of an integer divisible by 5 and it begins with 1}\}$

$\delta$	0	1
$\rightarrow [\epsilon]$	[0]	[1]
[0]	[0]	[0]
[1]	[10]	[11]
[10]	[100]	[101]
[11]	[1]	[10]
[100]		
*[101]		

# Myhill-Nerode method. Examples

Example 3:  $L_3 = \{x \in \Sigma^* \mid x \text{ is a binary representation of an integer divisible by 5 and it begins with 1}\}$

$\delta$	0	1
$\rightarrow [\epsilon]$	[0]	[1]
[0]	[0]	[0]
[1]	[10]	[11]
[10]	[100]	[101]
[11]	[1]	[10]
[100]	[11]	
*[101]		

# Myhill-Nerode method. Examples

Example 3:  $L_3 = \{x \in \Sigma^* \mid x \text{ is a binary representation of an integer divisible by 5 and it begins with 1}\}$

$\delta$	0	1
$\rightarrow [\epsilon]$	[0]	[1]
[0]	[0]	[0]
[1]	[10]	[11]
[10]	[100]	[101]
[11]	[1]	[10]
[100]	[11]	[100]
*[101]		

# Myhill-Nerode method. Examples

Example 3:  $L_3 = \{x \in \Sigma^* \mid x \text{ is a binary representation of an integer divisible by 5 and it begins with 1}\}$

$\delta$	0	1
$\rightarrow [\epsilon]$	[0]	[1]
[0]	[0]	[0]
[1]	[10]	[11]
[10]	[100]	[101]
[11]	[1]	[10]
[100]	[11]	[100]
*[101]	[101]	

# Myhill-Nerode method. Examples

Example 3:  $L_3 = \{x \in \Sigma^* \mid x \text{ is a binary representation of an integer divisible by 5 and it begins with 1}\}$

$\delta$	0	1
$\rightarrow [\epsilon]$	[0]	[1]
[0]	[0]	[0]
[1]	[10]	[11]
[10]	[100]	[101]
[11]	[1]	[10]
[100]	[11]	[100]
*[101]	[101]	[1]



## Regular languages

- Myhill-Nerode criteria
  - Positive Examples
  - **Negative Examples**
- Pumping Lemma

# Myhill-Nerode method. Examples

## Negative Example 1

$L_1 = \{a^n b^n \mid n \in \mathbb{N}\}$  is not regular.

## Proof

For  $m \neq k$ ,

$$a^m \not\equiv_{L_1} a^k,$$

since  $a^m b^k \notin L_1, a^k b^k \in L_1$  (a distinguishing ext. is  $b^k$ ).

Therefore, there are **infinity many equivalence classes**!

So,  $L_1$  is not regular.

# Myhill-Nerode method. Examples

## Negative Example 2

$L_2 = \{a^n ba^n \mid n \in \mathbb{N}\}$  is not regular.

## Proof

For  $m \neq k$ ,

$$a^m b \not\equiv_{L_2} a^k b,$$

since  $a^m ba^k \notin L_2$ ,  $a^k ba^k \in L_2$  (a distinguishing ext. is  $a^k$ ).

Therefore, there are **infinitely many equivalence classes!**

So,  $L_2$  is not regular.

## Regular languages

- Myhill-Nerode criteria
  - Positive Examples
  - Negative Examples
- **Pumping Lemma**

# Pumping lemma

## Pumping lemma

If  $L \subseteq \Sigma^*$  is a regular language then there exists  $m \geq 1$  such that any  $w \in L$  with  $|w| \geq m$  can be represented as  $w = xyz$  such that

- $y \neq \epsilon$ ,
- $xy^iz \in L$  for any  $i \geq 1$ .

# How Pumping lemma is useful?

- **Can we use this theorem to prove that a set is regular?**  
No, because it gives only a necessary condition for a language to be regular (and not a sufficient condition).
- **We can use it to prove that a language is not regular.**  
**How?**

# Pumping lemma

## Pumping lemma

If  $L \subseteq \Sigma^*$  is a regular language then there exists  $m \geq 1$  such that any  $w \in L$  with  $|w| \geq m$  can be represented as  $w = xyz$  such that

- $y \neq \epsilon$ ,
- $xy^iz \in L$  for any  $i \geq 1$ .

## Corollary

If for any  $m \geq 1$  there is  $w \in L$  such that  $|w| \geq m$  and for any representation  $w = xyz$  with  $y \neq \epsilon$

$$xy^iz \notin L \text{ for some } i \geq 1.$$

Then  $L$  is **not** a regular language.

# Pumping lemma. Examples

## Example 1

$L_1 = \{a^n b^n \mid n \in \mathbb{N}\}$  is not regular.

## Proof

$xyz \in L_1$  with  $y \neq \epsilon$



# Pumping lemma. Examples

## Example 1

$L_1 = \{a^n b^n \mid n \in \mathbb{N}\}$  is not regular.

## Proof

$xyz \in L_1$  with  $y \neq \epsilon$

$$1) \underbrace{a^{n-p_1-p_2}}_x \underbrace{(a^{p_1})}_y \underbrace{a^{p_2} b^n}_z \text{ and } p_1 \neq 0$$

# Pumping lemma. Examples

## Example 1

$L_1 = \{a^n b^n \mid n \in \mathbb{N}\}$  is not regular.

## Proof

$xyz \in L_1$  with  $y \neq \epsilon$

$$1) \underbrace{a^{n-p_1-p_2}}_x \underbrace{(a^{p_1})}_y \underbrace{a^{p_2} b^n}_z \text{ and } p_1 \neq 0$$

$$2) \underbrace{a^n b^{p_1}}_x \underbrace{(b^{p_2})}_y \underbrace{b^{n-p_1-p_2}}_z \text{ and } p_2 \neq 0$$

# Pumping lemma. Examples

## Example 1

$L_1 = \{a^n b^n \mid n \in \mathbb{N}\}$  is not regular.

## Proof

$xyz \in L_1$  with  $y \neq \epsilon$

- 1)  $\underbrace{a^{n-p_1-p_2}}_x \underbrace{(a^{p_1})}_y \underbrace{a^{p_2} b^n}_z$  and  $p_1 \neq 0$
- 2)  $\underbrace{a^n b^{p_1}}_x \underbrace{(b^{p_2})}_y \underbrace{b^{n-p_1-p_2}}_z$  and  $p_2 \neq 0$
- 3)  $\underbrace{a^{n-p_1}}_x \underbrace{(a^{p_1} b^{p_2})}_y \underbrace{b^{n-p_2}}_z$  and  $p_1, p_2 \neq 0$

# Pumping lemma. Examples

## Example 1

$L_1 = \{a^n b^n \mid n \in \mathbb{N}\}$  is not regular.

## Proof

1) for  $\underbrace{a^{n-p_1-p_2}}_x \underbrace{(a^{p_1})}_y \underbrace{a^{p_2} b^n}_z$  and  $p_1 \neq 0$ , we have

$$xy^2z =$$

# Pumping lemma. Examples

## Example 1

$L_1 = \{a^n b^n \mid n \in \mathbb{N}\}$  is not regular.

## Proof

1) for  $\underbrace{a^{n-p_1-p_2}}_x \underbrace{(a^{p_1})}_y \underbrace{a^{p_2} b^n}_z$  and  $p_1 \neq 0$ , we have

$$xy^2z = a^{n-p_1-p_2}(a^{p_1})^2 a^{p_2} b^n$$

# Pumping lemma. Examples

## Example 1

$L_1 = \{a^n b^n \mid n \in \mathbb{N}\}$  is not regular.

## Proof

1) for  $\underbrace{a^{n-p_1-p_2}}_x \underbrace{(a^{p_1})}_y \underbrace{a^{p_2} b^n}_z$  and  $p_1 \neq 0$ , we have

$$xy^2z = a^{n-p_1-p_2}(a^{p_1})^2 a^{p_2} b^n = a^{n+p_1} b^n$$

# Pumping lemma. Examples

## Example 1

$L_1 = \{a^n b^n \mid n \in \mathbb{N}\}$  is not regular.

## Proof

1) for  $\underbrace{a^{n-p_1-p_2}}_x \underbrace{(a^{p_1})}_y \underbrace{a^{p_2} b^n}_z$  and  $p_1 \neq 0$ , we have

$$xy^2z = a^{n-p_1-p_2}(a^{p_1})^2 a^{p_2} b^n = a^{n+p_1} b^n \notin L,$$

# Pumping lemma. Examples

## Example 1

$L_1 = \{a^n b^n \mid n \in \mathbb{N}\}$  is not regular.

## Proof

1) for  $\underbrace{a^{n-p_1-p_2}}_x \underbrace{(a^{p_1})}_y \underbrace{a^{p_2} b^n}_z$  and  $p_1 \neq 0$ , we have

$$xy^2z = a^{n-p_1-p_2}(a^{p_1})^2 a^{p_2} b^n = a^{n+p_1} b^n \notin L,$$

2) for  $\underbrace{a^n b^{p_1}}_x \underbrace{(b^{p_2})}_y \underbrace{b^{n-p_1-p_2}}_z$  and  $p_2 \neq 0$ , we have

$$xy^2z =$$



# Pumping lemma. Examples

## Example 1

$L_1 = \{a^n b^n \mid n \in \mathbb{N}\}$  is not regular.

## Proof

1) for  $\underbrace{a^{n-p_1-p_2}}_x \underbrace{(a^{p_1})}_y \underbrace{a^{p_2} b^n}_z$  and  $p_1 \neq 0$ , we have

$$xy^2z = a^{n-p_1-p_2} (a^{p_1})^2 a^{p_2} b^n = a^{n+p_1} b^n \notin L,$$

2) for  $\underbrace{a^n b^{p_1}}_x \underbrace{(b^{p_2})}_y \underbrace{b^{n-p_1-p_2}}_z$  and  $p_2 \neq 0$ , we have

$$xy^2z = a^n b^{n+p_2} \notin L,$$

# Pumping lemma. Examples

## Example 1

$L_1 = \{a^n b^n \mid n \in \mathbb{N}\}$  is not regular.

## Proof

1) for  $\underbrace{a^{n-p_1-p_2}}_x \underbrace{(a^{p_1})}_y \underbrace{a^{p_2} b^n}_z$  and  $p_1 \neq 0$ , we have

$$xy^2z = a^{n-p_1-p_2}(a^{p_1})^2 a^{p_2} b^n = a^{n+p_1} b^n \notin L,$$

2) for  $\underbrace{a^n b^{p_1}}_x \underbrace{(b^{p_2})}_y \underbrace{b^{n-p_1-p_2}}_z$  and  $p_2 \neq 0$ , we have

$$xy^2z = a^n b^{n+p_2} \notin L,$$

3) for  $\underbrace{a^{n-p_1}}_x \underbrace{(a^{p_1} b^{p_2})}_y \underbrace{b^{n-p_2}}_z$  and  $p_1, p_2 \neq 0$ , we have  $xy^2z =$

# Pumping lemma. Examples

## Example 1

$L_1 = \{a^n b^n \mid n \in \mathbb{N}\}$  is not regular.

## Proof

1) for  $\underbrace{a^{n-p_1-p_2}}_x \underbrace{(a^{p_1})}_y \underbrace{a^{p_2} b^n}_z$  and  $p_1 \neq 0$ , we have

$$xy^2z = a^{n-p_1-p_2}(a^{p_1})^2 a^{p_2} b^n = a^{n+p_1} b^n \notin L,$$

2) for  $\underbrace{a^n b^{p_1}}_x \underbrace{(b^{p_2})}_y \underbrace{b^{n-p_1-p_2}}_z$  and  $p_2 \neq 0$ , we have

$$xy^2z = a^n b^{n+p_2} \notin L,$$

3) for  $\underbrace{a^{n-p_1}}_x \underbrace{(a^{p_1} b^{p_2})}_y \underbrace{b^{n-p_2}}_z$  and  $p_1, p_2 \neq 0$ , we have  $xy^2z =$   
 $a^{n-p_1}(a^{p_1} b^{p_2})^2 b^{n-p_2} =$

# Pumping lemma. Examples

## Example 1

$L_1 = \{a^n b^n \mid n \in \mathbb{N}\}$  is not regular.

## Proof

1) for  $\underbrace{a^{n-p_1-p_2}}_x \underbrace{(a^{p_1})}_y \underbrace{a^{p_2} b^n}_z$  and  $p_1 \neq 0$ , we have

$$xy^2z = a^{n-p_1-p_2} (a^{p_1})^2 a^{p_2} b^n = a^{n+p_1} b^n \notin L,$$

2) for  $\underbrace{a^n b^{p_1}}_x \underbrace{(b^{p_2})}_y \underbrace{b^{n-p_1-p_2}}_z$  and  $p_2 \neq 0$ , we have

$$xy^2z = a^n b^{n+p_2} \notin L,$$

3) for  $\underbrace{a^{n-p_1}}_x \underbrace{(a^{p_1} b^{p_2})}_y \underbrace{b^{n-p_2}}_z$  and  $p_1, p_2 \neq 0$ , we have  $xy^2z =$

$$a^{n-p_1} (a^{p_1} b^{p_2})^2 b^{n-p_2} = a^{n-p_1} a^{p_1} b^{p_2} a^{p_1} b^{p_2} b^{n-p_2} =$$

# Pumping lemma. Examples

## Example 1

$L_1 = \{a^n b^n \mid n \in \mathbb{N}\}$  is not regular.

## Proof

1) for  $\underbrace{a^{n-p_1-p_2}}_x \underbrace{(a^{p_1})}_y \underbrace{a^{p_2} b^n}_z$  and  $p_1 \neq 0$ , we have

$$xy^2z = a^{n-p_1-p_2}(a^{p_1})^2 a^{p_2} b^n = a^{n+p_1} b^n \notin L,$$

2) for  $\underbrace{a^n b^{p_1}}_x \underbrace{(b^{p_2})}_y \underbrace{b^{n-p_1-p_2}}_z$  and  $p_2 \neq 0$ , we have

$$xy^2z = a^n b^{n+p_2} \notin L,$$

3) for  $\underbrace{a^{n-p_1}}_x \underbrace{(a^{p_1} b^{p_2})}_y \underbrace{b^{n-p_2}}_z$  and  $p_1, p_2 \neq 0$ , we have  $xy^2z =$

$$a^{n-p_1}(a^{p_1} b^{p_2})^2 b^{n-p_2} = a^{n-p_1} a^{p_1} b^{p_2} a^{p_1} b^{p_2} b^{n-p_2} = a^n b^{p_2} a^{p_1} b^n$$

# Pumping lemma. Examples

## Example 1

$L_1 = \{a^n b^n \mid n \in \mathbb{N}\}$  is not regular.

## Proof

1) for  $\underbrace{a^{n-p_1-p_2}}_x \underbrace{(a^{p_1})}_y \underbrace{a^{p_2} b^n}_z$  and  $p_1 \neq 0$ , we have

$$xy^2z = a^{n-p_1-p_2}(a^{p_1})^2 a^{p_2} b^n = a^{n+p_1} b^n \notin L,$$

2) for  $\underbrace{a^n b^{p_1}}_x \underbrace{(b^{p_2})}_y \underbrace{b^{n-p_1-p_2}}_z$  and  $p_2 \neq 0$ , we have

$$xy^2z = a^n b^{n+p_2} \notin L,$$

3) for  $\underbrace{a^{n-p_1}}_x \underbrace{(a^{p_1} b^{p_2})}_y \underbrace{b^{n-p_2}}_z$  and  $p_1, p_2 \neq 0$ , we have  $xy^2z =$

$$a^{n-p_1}(a^{p_1} b^{p_2})^2 b^{n-p_2} = a^{n-p_1} a^{p_1} b^{p_2} a^{p_1} b^{p_2} b^{n-p_2} = a^n b^{p_2} a^{p_1} b^n \notin L$$

# Pumping lemma. Examples

## Example 2

$L_2 = \{a^n ba^n \mid n \in \mathbb{N}\}$  is not regular.

## Proof

Let  $w = xyz \in L_2$  and  $y \neq \epsilon$

# Pumping lemma. Examples

## Example 2

$L_2 = \{a^n b a^n \mid n \in \mathbb{N}\}$  is not regular.

## Proof

Let  $w = xyz \in L_2$  and  $y \neq \epsilon$

1) if  $y$  contains  $b$  (that is  $w = \underbrace{a^{n-p_1}}_x \underbrace{a^{p_1} b a^{p_2}}_y \underbrace{a^{n-p_2}}_z$ )



# Pumping lemma. Examples

## Example 2

$L_2 = \{a^n b a^n \mid n \in \mathbb{N}\}$  is not regular.

## Proof

Let  $w = xyz \in L_2$  and  $y \neq \epsilon$

1) if  $y$  contains  $b$  (that is  $w = \underbrace{a^{n-p_1}}_x \underbrace{a^{p_1} b a^{p_2}}_y \underbrace{a^{n-p_2}}_z$ )

then  $xy^2z$  contains  $b$  twice and hence  $xy^2z \notin L_2$

# Pumping lemma. Examples

## Example 2

$L_2 = \{a^n b a^n \mid n \in \mathbb{N}\}$  is not regular.

## Proof

Let  $w = xyz \in L_2$  and  $y \neq \epsilon$

1) if  $y$  contains  $b$  (that is  $w = \underbrace{a^{n-p_1}}_x \underbrace{a^{p_1} b a^{p_2}}_y \underbrace{a^{n-p_2}}_z$ )

then  $xy^2z$  contains  $b$  twice and hence  $xy^2z \notin L_2$

2) if  $y$  does not contain  $b$ , that is

$$w = \underbrace{a^{n-p_1}}_x \underbrace{a^{p_1}}_y \underbrace{b a^n}_z, \text{ or } w = \underbrace{a^n}_x \underbrace{b a^{p_2}}_y \underbrace{a^{n-p_2}}_z$$

# Pumping lemma. Examples

## Example 2

$L_2 = \{a^n ba^n \mid n \in \mathbb{N}\}$  is not regular.

## Proof

Let  $w = xyz \in L_2$  and  $y \neq \epsilon$

1) if  $y$  contains  $b$  (that is  $w = \underbrace{a^{n-p_1}}_x \underbrace{a^{p_1} ba^{p_2}}_y \underbrace{a^{n-p_2}}_z$ )

then  $xy^2z$  contains  $b$  twice and hence  $xy^2z \notin L_2$

2) if  $y$  does not contain  $b$ , that is

$$w = \underbrace{a^{n-p_1}}_x \underbrace{a^{p_1}}_y \underbrace{ba^n}_z, \text{ or } w = \underbrace{a^n}_x \underbrace{b}_y \underbrace{a^{p_2}}_y \underbrace{a^{n-p_2}}_z$$

then  $xy^2z = a^k ba^m$  where  $k \neq m$  and hence  $xy^2z \notin L_2$

# Pumping lemma. For practice

## Pumping lemma

If  $L \subseteq \Sigma^*$  is a regular language then there exists  $m \geq 1$  such that any  $w \in L$  with  $|w| \geq m$  can be represented as  $w = xyz$  such that

- $y \neq \epsilon$ ,
- $|xy| \leq m$ ,
- $xy^iz \in L$  for any  $i \geq 1$ .

## Corollary

If for any  $m \geq 1$  there is  $w \in L$  such that  $|w| \geq m$  and for any representation  $w = xyz$  with  $y \neq \epsilon$  and  $|xy| \leq m$

$$xy^iz \notin L \text{ for some } i \geq 1.$$

Then  $L$  is **not** a regular language.

# Pumping lemma. For practice

## Corollary

If for any  $m \geq 1$  there is  $w \in L$  such that  $|w| \geq m$  and for any representation  $w = xyz$  with  $y \neq \epsilon$  and  $|xy| \leq m$

$$xy^iz \notin L \text{ for some } i \geq 1.$$

Then  $L$  is **not** a regular language.

## Negative Example 1

$L_1 = \{a^n b^n \mid n \in \mathbb{N}\}$  is not regular.

## Proof

For each  $m$ , let  $w = a^k b^k$ , where  $k \geq m$ . If  $w = xyz$  and  $|xy| \leq m$ , then  $w = \underbrace{a^{n-p_1-p_2}}_x \underbrace{(a^{p_1})}_y \underbrace{a^{p_2} b^n}_z$  and hence  $xy^2z \notin L$

# Pumping lemma. For practice

## Corollary

If for any  $m \geq 1$  there is  $w \in L$  such that  $|w| \geq m$  and for any representation  $w = xyz$  with  $y \neq \epsilon$  and  $|xy| \leq m$

$$xy^iz \notin L \text{ for some } i \geq 1.$$

Then  $L$  is **not** a regular language.

## Negative Example 2

$L_2 = \{a^n ba^n \mid n \in \mathbb{N}\}$  is not regular.

## Proof

For each  $m$ , let  $w = a^k ba^k$ , where  $k \geq m + 1$ . If  $w = xyz$  and  $|xy| \leq m$ , then  $w = \underbrace{a^{n-p_1}}_x \underbrace{a^{p_1}}_y \underbrace{ba^n}_z$  and hence  $xy^2z \notin L_2$

Thank you for your attention!