# MECP2 duplication syndrome

MECP2 duplication syndrome is a condition that occurs almost exclusively in males and is characterized by moderate to severe intellectual disability. Most people with this condition also have weak muscle tone in infancy, feeding difficulties, poor or absent speech, or muscle stiffness (rigidity). Individuals with MECP2 duplication syndrome have delayed development of motor skills such as sitting and walking. About half of individuals have seizures, often of the tonic-clonic type. This type of seizure involves a loss of consciousness, muscle rigidity, and convulsions and may not respond to medication. Some affected individuals experience the loss of previously acquired skills (developmental regression). Approximately half of individuals learn to walk, and about one-third of people with this condition require assistance when walking. Many individuals with MECP2 duplication syndrome have recurrent respiratory tract infections. These respiratory infections are a major cause of death in affected individuals, with only half surviving past age 25.

# **Frequency**

The prevalence of *MECP2* duplication syndrome is unknown; more than 200 affected individuals have been described in the scientific literature. It is estimated that this condition is responsible for 1 to 2 percent of all cases of intellectual disability caused by changes in the X chromosome.

#### Causes

MECP2 duplication syndrome is caused by a genetic change in which there is an extra copy of the MECP2 gene in each cell. This extra copy of the MECP2 gene is caused by a duplication of genetic material on the long (q) arm of the X chromosome. The size of the duplication varies from 100,000 to a few million DNA building blocks (base pairs). The MECP2 gene is always included in this duplication, and other genes may also be involved, depending on the size of the duplicated segment. It is unclear whether extra copies of these other genes affect the severity of the condition.

The *MECP2* gene provides instructions for making a protein called MeCP2 that is critical for normal brain function. Researchers believe that this protein has several functions, including regulating other genes in the brain by controlling when they are able to participate in protein production. An extra copy of the *MECP2* gene leads to the production of excess MeCP2 protein and an increase in protein function. The resulting changes in gene regulation and protein production in the brain lead to abnormal nerve cell (neuronal) function. These neuronal changes disrupt normal brain activity, causing the signs and symptoms of *MECP2* duplication syndrome.

#### **Inheritance Pattern**

*MECP2* duplication syndrome is inherited in an X-linked pattern. The gene associated with this condition is located on the X chromosome, which is one of the two sex chromosomes in each cell. In males (who have only one X chromosome), a duplication of the only copy of the *MECP2* gene in each cell is sufficient to cause the condition. In females (who have two X chromosomes), a duplication of one of the two copies of the gene typically does not cause the disorder, but can be associated with behavioral and psychiatric symptoms such as depression, anxiety, and features of autism spectrum disorder that affect communication and social interaction.

Females with a *MECP2* gene duplication tend to be unaffected or less severely affected than males because the X chromosome that contains the duplication may be turned off (inactive) in many of their cells due to a process called X-inactivation. Early in embryonic development in females, one of the two X chromosomes is permanently inactivated in somatic cells (cells other than egg and sperm cells). X-inactivation ensures that females, like males, have only one active copy of the X chromosome in each body cell. Usually X-inactivation occurs randomly, such that each X chromosome is active in about half of the body's cells. Sometimes X-inactivation is not random, and one X chromosome is active in more than half of cells. When X-inactivation does not occur randomly, it is called skewed X-inactivation.

Females with a *MECP2* gene duplication often have skewed X-inactivation, which results in the inactivation of the X chromosome containing the duplication in most cells of the body. Although this skewed X-inactivation ensures that the chromosome with the normal *MECP2* gene is active most often, some of these females develop behavioral and psychiatric symptoms thought to be related to the additional genetic material. It is unclear why these features develop in a small number of females with skewed X-inactivation. Researchers speculate that in these females some cells in the brain may have a different pattern of X-inactivation than the cells in the rest of the body so that the X chromosome with the duplicated *MECP2* gene is active, resulting in behavioral and psychiatric symptoms.

#### Other Names for This Condition

- Lubs X-linked mental retardation syndrome
- trisomy Xq28

## **Diagnosis & Management**

**Genetic Testing Information** 

- What is genetic testing?
   /primer/testing/genetictesting
- Genetic Testing Registry: Syndromic X-linked intellectual disability Lubs type https://www.ncbi.nlm.nih.gov/gtr/conditions/C1846058/

# Research Studies from ClinicalTrials.gov

 ClinicalTrials.gov https://clinicaltrials.gov/ct2/results?cond=%22MECP2+duplication+syndrome %22+OR+%22Trisomy+Xq28%22+OR+%22RTT-related+conditions%22+OR+ %22Rett-related+disorder%22

# Other Diagnosis and Management Resources

- GeneReview: MECP2 Duplication Syndrome https://www.ncbi.nlm.nih.gov/books/NBK1284
- RettSyndrome.org: Rett Syndrome, MECP2 Duplication, and Rett-related Disorders Natural History Study https://www.rettsyndrome.org/for-families/natural-history-study/

#### **Additional Information & Resources**

# Health Information from MedlinePlus

- Encyclopedia: Hypotonia https://medlineplus.gov/ency/article/003297.htm
- Encyclopedia: Spasticity https://medlineplus.gov/ency/article/003297.htm
- Health Topic: Developmental Disabilities https://medlineplus.gov/developmentaldisabilities.html

#### Genetic and Rare Diseases Information Center

 MECP2 duplication syndrome https://rarediseases.info.nih.gov/diseases/9781/mecp2-duplication-syndrome

#### Additional NIH Resources

- National Institute of Neurological Disorders and Stroke: Seizures and Epilepsy https://www.ninds.nih.gov/Disorders/Patient-Caregiver-Education/Hope-Through-Research/Epilepsies-and-Seizures-Hope-Through
- National Institute of Neurological Disorders and Stroke: Spasticity Information Page https://www.ninds.nih.gov/Disorders/All-Disorders/Spasticity-Information-Page

### **Educational Resources**

- Boston Children's Hospital: Respiratory Distress https://www.childrenshospital.org/conditions-and-treatments/conditions/r/respiratory-distress
- Boston Children's Hospital: Seizures http://www.childrenshospital.org/conditions-and-treatments/conditions/s/seizures

- Centers for Disease Control and Prevention: Intellectual Disability https://www.cdc.gov/ncbddd/actearly/pdf/parents\_pdfs/IntellectualDisability.pdf
- March of Dimes: Chromosomal Conditions https://www.marchofdimes.org/baby/chromosomal-conditions.aspx
- Orphanet: Trisomy Xq28
   https://www.orpha.net/consor/cgi-bin/OC\_Exp.php?Lng=EN&Expert=1762

# Patient Support and Advocacy Resources

- American Association on Intellectual and Developmental Disabilities (AAIDD) https://www.aaidd.org/
- Rare Diseases Clinical Research Network: Rett Consortium https://www.rarediseasesnetwork.org/cms/rett/Learn-More/Disorder-Definitions#MECP2
- RettSyndrome.org
   https://www.rettsyndrome.org/
- RettSyndrome.org: Rett Syndrome, MECP2 Duplication, and Rett-related Disorders Natural History Study https://www.rettsyndrome.org/for-families/natural-history-study/

## Clinical Information from GeneReviews

 MECP2 Duplication Syndrome https://www.ncbi.nlm.nih.gov/books/NBK1284

#### Scientific Articles on PubMed

PubMed

https://www.ncbi.nlm.nih.gov/pubmed?term=%28mecp2+duplication+syndrome %5BTIAB%5D%29+AND+english%5Bla%5D+AND+human%5Bmh%5D+AND+%22last+3600+days%22%5Bdp%5D

# Catalog of Genes and Diseases from OMIM

 LUBS X-LINKED MENTAL RETARDATION SYNDROME http://omim.org/entry/300260

#### Medical Genetics Database from MedGen

 Trisomy Xq28 syndrome https://www.ncbi.nlm.nih.gov/medgen/930014

# **Sources for This Summary**

- El Chehadeh S, Touraine R, Prieur F, Reardon W, Bienvenu T, Chantot-Bastaraud S, Doco-Fenzy M, Landais E, Philippe C, Marle N, Callier P, Mosca-Boidron AL, Mugneret F, Le Meur N, Goldenberg A, Guerrot AM, Chambon P, Satre V, Coutton C, Jouk PS, Devillard F, Dieterich K, Afenjar A, Burglen L, Moutard ML, Addor MC, Lebon S, Martinet D, Alessandri JL, Doray B, Miguet M, Devys D, Saugier-Veber P, Drunat S, Aral B, Kremer V, Rondeau S, Tabet AC, Thevenon J, Thauvin-Robinet C, Perreton N, Des Portes V, Faivre L. Xq28 duplication including MECP2 in six unreported affected females: what can we learn for diagnosis and genetic counselling? Clin Genet. 2016 Oct 19. doi: 10.1111/cge.12898. [Epub ahead of print] Citation on PubMed: https://www.ncbi.nlm.nih.gov/pubmed/27761913
- Gonzales ML, LaSalle JM. The role of MeCP2 in brain development and neurodevelopmental disorders. Curr Psychiatry Rep. 2010 Apr;12(2):127-34. doi: 10.1007/s11920-010-0097-7. Review. Citation on PubMed: https://www.ncbi.nlm.nih.gov/pubmed/20425298
   Free article on PubMed Central: https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2847695/
- Lombardi LM, Baker SA, Zoghbi HY. MECP2 disorders: from the clinic to mice and back. J Clin Invest. 2015 Aug 3;125(8):2914-23. doi: 10.1172/JCI78167. Review.
   Citation on PubMed: https://www.ncbi.nlm.nih.gov/pubmed/26237041
   Free article on PubMed Central: https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4563741/
- Lugtenberg D, Kleefstra T, Oudakker AR, Nillesen WM, Yntema HG, Tzschach A, Raynaud M, Rating D, Journel H, Chelly J, Goizet C, Lacombe D, Pedespan JM, Echenne B, Tariverdian G, O'Rourke D, King MD, Green A, van Kogelenberg M, Van Esch H, Gecz J, Hamel BC, van Bokhoven H, de Brouwer AP. Structural variation in Xq28: MECP2 duplications in 1% of patients with unexplained XLMR and in 2% of male patients with severe encephalopathy. Eur J Hum Genet. 2009 Apr;17(4):444-53. doi: 10.1038/ejhg.2008.208. Epub 2008 Nov 5. Erratum in: Eur J Hum Genet. 2009 May;17(5):697. Citation on PubMed: https://www.ncbi.nlm.nih.gov/pubmed/18985075
- Ramocki MB, Peters SU, Tavyev YJ, Zhang F, Carvalho CM, Schaaf CP, Richman R, Fang P, Glaze DG, Lupski JR, Zoghbi HY. Autism and other neuropsychiatric symptoms are prevalent in individuals with MeCP2 duplication syndrome. Ann Neurol. 2009 Dec;66(6):771-82. doi: 10.1002/

Free article on PubMed Central: https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2986218/

- Citation on PubMed: https://www.ncbi.nlm.nih.gov/pubmed/20035514

  Free article on PubMed Central: https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2801873/
- Yi Z, Pan H, Li L, Wu H, Wang S, Ma Y, Qi Y. Chromosome Xq28 duplication encompassing MECP2: Clinical and molecular analysis of 16 new patients from 10 families in China. Eur J Med Genet. 2016 Jun;59(6-7):347-53. doi: 10.1016/j.ejmg.2016.05.004.
   Citation on PubMed: https://www.ncbi.nlm.nih.gov/pubmed/27180140

#### Reprinted from Genetics Home Reference:

https://ghr.nlm.nih.gov/condition/mecp2-duplication-syndrome

Reviewed: March 2017 Published: May 26, 2020

ana.21715.

Lister Hill National Center for Biomedical Communications U.S. National Library of Medicine National Institutes of Health Department of Health & Human Services