

Contents lists available at ScienceDirect

Pediatric Neurology

journal homepage: www.elsevier.com/locate/pnu



Original Article

Leveraging a Sturge-Weber Gene Discovery: An Agenda for Future Research



Anne M. Comi MD ^{a,b,*}, Mustafa Sahin MD, PhD ^c, Adrienne Hammill MD, PhD ^d, Emma H. Kaplan BA ^a, Csaba Juhász MD, PhD ^{e,f}, Paula North MD, PhD ^g, Karen L. Ball BS ^h, Alex V. Levin MD, MHSc, FRCSC ^{i,j}, Bernard Cohen MD ^{b,k}, Jill Morris PhD ^l, Warren Lo MD ^m, E. Steve Roach MD ^m, and participants of the 2015 Sturge-Weber Syndrome Research Workshop

- ^a Department of Neurology, Kennedy Krieger Institute and Johns Hopkins University School of Medicine, Baltimore, Maryland
- ^b Department of Pediatrics, Johns Hopkins University School of Medicine, Baltimore, Maryland
- ^c Department of Radiology, Boston Children's Hospital and Harvard Medical School, Boston, Massachusetts
- d Cancer and Blood Diseases Institute, Cincinnati Children's Hospital Medical Center and University of Cincinnati, Cincinnati, Ohio
- ^e Department of Pediatrics, Wayne State University School of Medicine, Detroit, Michigan
- ^f Department of Neurology, Wayne State University School of Medicine, Detroit, Michigan
- ^g Department of Pathology, Medical College of Wisconsin, Milwaukee, Wisconsin
- ^h The Sturge-Weber Foundation, Randolph, New Jersey
- Department of Ophthalmology, Wills Eye Hospital, Sidney Kimmel Medical College, Thomas University, Philadelphia, Pennsylvania
- ^j Department of Pediatrics, Wills Eye Hospital, Sidney Kimmel Medical College, Thomas Jefferson University, Philadelphia, Pennsylvania
- ^k Department of Dermatology, Johns Hopkins University School of Medicine, Baltimore, Maryland
- ¹National Institute of Neurologic Disorders and Stroke, National Institutes of Health, Bethesda, Maryland
- ^m Departments of Pediatrics and Neurology, The Ohio State University College of Medicine and Nationwide Children's Hospital, Columbus, Ohio

ABSTRACT

Sturge-Weber syndrome (SWS) is a vascular neurocutaneous disorder that results from a somatic mosaic mutation in GNAQ, which is also responsible for isolated port-wine birthmarks. Infants with SWS are born with a cutaneous capillary malformation (port-wine birthmark) of the forehead or upper eyelid which can signal an increased risk of brain and/or eye involvement prior to the onset of specific symptoms. This symptom-free interval represents a time when a targeted intervention could help to minimize the neurological and ophthalmologic manifestations of the disorder. This paper summarizes a 2015 SWS workshop in Bethesda, Maryland that was sponsored by the National Institutes of Health. Meeting attendees included a diverse group of clinical and translational researchers with a goal of establishing research priorities for the next few years. The initial portion of the meeting included a thorough review of the recent genetic discovery and what is known of the pathogenesis of SWS. Breakout sessions related to neurology, dermatology, and ophthalmology aimed to establish SWS research priorities in each field. Key priorities for future development include the need for clinical consensus guidelines, further work to develop a clinical trial network, improvement of tissue banking for research purposes, and the need for multiple animal and cell culture models of SWS.

Keywords: Sturge-Weber syndrome, GNAQ, somatic mutation, port-wine birthmark, glaucoma, seizures

Pediatr Neurol 2016; 58: 12-24 © 2016 Published by Elsevier Inc.

Article History:

Received November 16, 2015; Accepted in final form November 21, 2015

* Communications should be addressed to: Dr. Comi; Department of Neurology; Kennedy Krieger Institute and Johns Hopkins University School of Medicine; 801 North Broadway; Baltimore, MD 21205.

E-mail address: comi@kennedykrieger.org