Review Article

Are There Differences in the Prevalence of Palliative Care-Related Problems in People Living With Advanced Cancer and Eight Non-Cancer Conditions? A Systematic Review

Katrien Moens, MSc, Irene J. Higginson, PhD, and Richard Harding, PhD, on behalf of EURO IMPACT

Department of Palliative Care, Policy & Rehabilitation, Cicely Saunders Institute, King's College London, London, United Kingdom

Abstract

Context. If access to effective palliative care is to extend beyond cancer patients, an understanding of the comparative prevalence of palliative care problems among cancer and non-cancer patients is necessary.

Objectives. This systematic review aimed to describe and compare the prevalence of seventeen palliative care-related problems across the four palliative care domains among adults with advanced cancer, acquired immune deficiency syndrome, chronic heart failure, end-stage renal disease (ESRD), chronic obstructive pulmonary disease, multiple sclerosis, motor neuron disease, Parkinson's disease, and dementia.

Methods. Three databases were searched using three groups of keywords. The results of the extraction of the prevalence figures were summarized.

Results. The electronic searches yielded 4697 hits after the removal of 1784 duplicates. Of these hits, 143 met the review criteria. The greatest number of studies were found for advanced cancer (n=57) and ESRD patients (n=47), and 75 of the 143 studies used validated scales. Few data were available for people living with multiple sclerosis (n=2) and motor neuron disease (n=3). The problems with a prevalence of 50% or more found across most of the nine studied diagnostic groups were: pain, fatigue, anorexia, dyspnea, and worry.

Conclusion. There are commonalities in the prevalence of problems across cancer and non-cancer patients, highlighting the need for palliative care to be provided irrespective of diagnosis. The methodological heterogeneity across the studies and the lack of non-cancer studies need to be addressed in future research. J Pain Symptom Manage 2014;48:660–677. © 2014 American Academy of Hospice and Palliative Medicine. Published by Elsevier Inc. Open access under CC BY-NC-ND license.

Address correspondence to: Katrien Moens, MSc, Department of Palliative Care, Policy & Rehabilitation, Cicely Saunders Institute, King's College London, Bessemer Road, Denmark Hill, London SE5 9PJ, United Kingdom. E-mail: Katrien.Moens@kcl.ac.uk

Accepted for publication: November 21, 2013.

Key Words

Prevalence, palliative care, cancer, acquired immune deficiency syndrome, heart diseases, chronic obstructive pulmonary disease, renal disease, neurodegenerative diseases, dementia

Introduction

The number of deaths from chronic rather than acute diseases is expected to increase worldwide. 1-3 A consequence of the aging population is that the time before death is differentiated by longer dying trajectories, with a higher prevalence of symptoms, resulting in longer and varied care needs at the end of life.⁴ To effectively anticipate patients' needs, and plan palliative care services for non-cancer patients, we must understand how prevalent palliative care-related problems are across different diagnostic groups.^{5,6} Although individual studies have considered this, only the Solano et al⁷ review in 2006 attempted to bring data together across studies. However, only five conditions were appraised, and a considerable amount of data have been published since then. In addition, the review of Solano et al' review focused on physical and psychological symptoms, rather than wider palliative care related problems. Therefore, this systematic review aimed to compare the prevalence of 17 palliative care-related problems within the four World Health Organization (WHO) palliative care domains (the physical, psychological, social, and spiritual domains)⁸ among adults with advanced stage cancer, acquired immune deficiency syndrome (AIDS), chronic heart failure (CHF), end-stage renal disease (ESRD), obstructive pulmonary chronic (COPD), multiple sclerosis (MS), motor neuron disease (MND), Parkinson's disease, and dementia. We chose to include people living with MS, MND, Parkinson's disease, and dementia in our search because these neurodegenerative diseases show a different disease trajectory in comparison with the other diseases.⁹ People living with dementia were included because of the increasing aging population.¹⁰

Two main objectives were set: 1) to describe and compare the prevalence of palliative care—related problems among advanced cancer and these eight advanced non-cancer diagnostic groups of patients, and 2) to describe and compare the characteristics of the prevalence studies.

Methods

Study Design

This systematic review was conducted in line with the PRISMA 2009 checklist, the PRISMA Flow Diagram, and the Center for Reviews and Dissemination procedures. 11,12 We used the Assessment of Multiple SysTemAtic Reviews (AMSTAR) measurement tool to ensure the methodological quality of our review.¹³ The AMSTAR uses 11 questions each of which needs to be answered using the following possibilities: yes, no, cannot answer, or not applicable. To have the highest quality ranking, all applicable questions should be answered with "yes." K. M. conducted the search and the initial screening; R. H. reviewed any articles where there was a question as to whether they met study criteria. I. J. H. adjudicated. The reference lists within the full texts also were searched.

Search Strategy

We searched three electronic medical databases, namely Medline (1966 to February 2013), Embase (1988 to February 2013), and PsycINFO (1985 to February 2013), using three groups of keywords. Within each group of keywords, the words were combined using OR. The three groups of keywords were then combined using AND.

Keyword Group 1: worry; depression, low mood, sadness, mood; anxiety, adjustment disorders (psychological palliative care domain); wellbeing; peace; spiritual wellbeing; meaning of life; religion; spirituality (spiritual palliative care domain); support; information; family (social palliative care domain); pain; confusion, delirium, cognitive failure; breathlessness, dyspnoea, dyspnea; fatigue, weakness; insomnia, poor sleeping; anorexia; nausea; diarrhea, diarrhea; constipation (physical palliative care domain); symptoms. These words were selected as the most appropriate words to investigate the 17 palliative care-related problems. Because different terms are used to define problems, we used more than one keyword for the following problems: poor sleeping for insomnia; weakness for fatigue; low mood, mood and sadness for depression; dyspnoea or dyspnea for breathlessness; and delirium and cognitive failure for confusion.

Keyword Group 2: dying, end of life, terminally ill, hospice\$, palliative care, terminal care, advanced, terminal disease\$. These words were selected to capture samples of patients in advanced disease stage.

Keyword Group 3: terminal disease\$, advanced cancer\$, metastatic cancer\$, AIDS, end stage heart disease\$, end stage heart failure, end stage respiratory disease\$, end stage chronic obstructive pulmonary disease\$, end stage COPD, end stage renal disease\$, end stage renal failure, MS, MND, Parkinson disease\$, dementia. These words were selected to identify the specified diagnostic groups.

These keywords were derived from those that were successfully used in the reviews of Solano et al⁷ and Simms et al.¹⁴ Simms et al¹⁴ used the WHO definition of palliative care⁸ to extract keywords identifying each palliative care domain. We refined our keywords using the synonyms for problems reported in the literature on symptom guidance and systematic reviews in palliative care.¹⁵

Inclusion/Exclusion Criteria

Fig. 1 presents the inclusion and exclusion criteria that were applied in the literature search.

Data Extraction

The data from each study were extracted to a common table with predesigned headings, namely reference, year of publication, country of origin; sample description; study design (measurement methods/tools, point or period

Inclusion criteria

- Studies on adults (aged 18 years or older), with advanced or terminal illness suffering from either cancer, AIDS, HD, COPD, ESRD, MS, MND, Parkinson's disease, or dementia
- Patients in palliative care who deteriorated despite treatment or with poor prognosis (i.e., less than one year)
- Studies reporting the prevalence of specified symptoms
- Original studies (quantitative)
- Articles that included data from at least one of the four palliative care domains

Exclusion criteria

- · Case reports of single patients
- · Articles not written in English, Dutch, French, or German
- Studies that did not disaggregate the palliative care–related problems within the reported sample
- Randomized controlled trials and other intervention studies (because of the application of selection criteria to the population of study)

Fig. 1. Inclusion and exclusion criteria of search. AIDS = acquired immune deficiency syndrome; HD = heart disease; COPD = chronic obstructive pulmonary disease; ESRD = end-stage renal disease; MS = multiple sclerosis; MND = motor neuron disease.

prevalence); and findings (% prevalence of problems per palliative care domain).

Describing and Comparing the Studies' Characteristics

The characteristics of all the studies are described in Table 1, which presents the total number of studies retained, total combined sample size, the nature of the prevalence measured (period/point), the period prevalence time ranges (minimum-maximum), the countries in which the studies were conducted, and the measurement methods used.

Data Analysis: Describing and Comparing the Prevalence of Problems

The results of the extraction of the prevalence figures of all included studies are summarized using minimum and maximum percentage ranges, and the problems are categorized by palliative care domain in line with the WHO definition⁸ (Table 2). Meta-analytic techniques were not appropriate to analyze the prevalence data because of the heterogeneity between studies in terms of measurement tools and study designs. Nevertheless, we reviewed and present the number of studies that showed the prevalence of a problem in a particular diagnostic group of 50% or more.

Results

Retrieval Data

The electronic searches of the three data-bases yielded 4697 hits after removal of 1784 duplicates. A total of 143 studies^{1,5,16–156} met the inclusion/exclusion criteria for the synthesis. The PRISMA flow diagram reports the search strategy and the results yielded (Fig. 2).

The assessment of the methodological quality of our systematic review using the AMSTAR measurement tool indicated that 10 of the 11 questions could be answered positively, that is, that the following actions are executed and reported in the review: an *a priori* design is provided, a duplicate study selection and data extraction are executed, a comprehensive literature search is performed, the status of publication is used as an inclusion criterion, a list of studies that were included and excluded is provided, the characteristics of the included studies are provided, the scientific quality of the included studies is used to formulate

conclusions, the method used to combine the findings of the studies is appropriate, and the conflict of interest is stated at the end of this article. The 11th item or question of the AMSTAR reviews if the likelihood of publication bias is assessed. This was not applicable for our study as we could not use statistical tests to analyze our results because of the heterogeneity across all included studies.

Characteristics of Included Studies

The greatest number of studies by diagnostic group were found for advanced cancer (n=57) and ESRD patients (n=47; Table 1). The lowest number of studies were found for MND (n=3) and MS patients (n=2). The greatest number of studies originated from the U.S. (n=57), the U.K. (n=20), and Canada (n = 13). In the advanced cancer diagnostic group, 37 of the 57 studies used a point prevalence to collect data, and in the ESRD patient disease group, 28 of the 47 studies used a period prevalence with a minimum-maximum time range of "past two days" to "the last two months." For all the other diagnostic groups, a period prevalence was primarily used. Considering the measurement tools used across the advanced cancer and non-cancer diagnostic groups, 75 of the 143 studies used validated scales. For data collection among patients with advanced dementia, four of the seven studies used chart reviews.

Describing and Comparing Prevalence Data

The prevalence figures reported in the 143 studies are summarized using minimum and maximum prevalence ranges (Table 2). In this table, the different terms used for each palliative care—related problem across all the studies are presented to highlight the varying definitions. For example, we found studies investigating "shortness of breath" or "respiratory distress" to indicate the study of dyspnea or breathlessness.

Among people living with MND and MS, there were many problems for which we did not find prevalence data, for example, anorexia, delirium, diarrhea, and worry, as well as for all the spiritual and social palliative care—related problems. For certain diseases, we only found one study on the prevalence of a problem (e.g., fatigue in people living with dementia, Parkinson's disease, and MS; Table 2).

Characteristics	Cancer	COPD	CHF	ESRD	Dementia	AIDS	Parkinson	MND	MS
Total number of studies	57	10	8	47	7	8	9	3	2
Total sample (patients), all studies	34,866	2045	1310	11,140	868	1864	25,941	221	214
Total number of point prevalence studies ^a	37, of which two studies presented both a point and a period prevalence	1	1	11, of which one study presented both a point and a period prevalence	2	_	_	_	_
Total number of period prevalence studies	17, of which two studies presented both a point and a period prevalence	8	7, of which one study presented both a point and a period prevalence	28, of which one study presented both a point and a period prevalence	_	5	6	1	1
Period prevalence time ranges (minimum- maximum)	Three Days-past week	Last three days- final year of life	Last year of life	Past two days-last two months	_	Last two days-last two weeks	Past two weeks-last month	Last month of life	Last three days
Total number of studies of which the point or period of prevalence was not specified	7	1	1	10	5	3	3	2	1
	idies conducted in th	ne following countrie							
The Netherlands, $n = 6$	3	1	1	1					
Germany, $n = 5$ Austria, $n = 1$ France, $n = 3$	4	1		1 2		1			
Italy, $n = 7$	5			1	1	1			
Spain, $n=5$	3		1	1	1				
Switzerland, $n=3$	1		1		1				
United Kingdom, $n = 20$	7	3	2	5			1	1	1
Ireland, $n = 1$ China, $n = 6$ Korea, $n = 1$	4			1 2 1					

Sweden, $n=3$	1		1				1		
Norway, $n=1$	1								
Denmark, $n=1$	1								
Romania, $n=1$	1								
Hungary, $n=1$				1					
South Africa,	1					1			
n=2									
Israel, $n=3$				2	1				
Africa, $n=1$				1					
U.S., $n = 57$	16	4	3	18	3	6	6	1	
Number of studies using	the following sy	mptom/problem mea	asurement methods:						
Validated	24	7	2	29	1	4	5	1	2
assessment									
scales									
Self-developed	22	2	4	13	1	2	2	1	
items list or									
checklist									
Medical records	10		2	5	4	2		1	
or patient									
reports									
Minimum data					1		1		
set review									
Method not	1	1					1		
specified									

10

Japan, n=3India, n=1Australia, n = 5New Zealand, n = 2Canada, n = 13

sclerosis.

^aPoint prevalence time points include measurements at the moment and those over the last 24 hours.

 ${\it Table~2} \\ {\bf Prevalence~Summary~Grid~Derived~From~143~Studies}$

	Problems and Number of Different Studies for Each	Minimum-Maximum Prevalence % of Each Problem References Total Sample of Patients (N) for Each Problem Total Number of Studies (n) for Each Problem Number of Studies That Report Prevalence for a Problem as $\geq 50\%$ $(n_{\geq 50\%})$										
Domain	Problem for All Diagnostic Groups (n)	Cancer	COPD	CHF	ESRD	Dementia	AIDS	Parkinson	MND	MS		
Physical:	Fatigue/ tiredness/ weakness n = 58	23-100% 5,17,18, 21,23-26,28, 31,34,37, 38,40,43, 45,47-50, 52-54,56-59, 64,65,67-71	$32 - 96\%^{1,74},$ $75,77,79,a$ $N = 500$ $n = 5$ $n \ge 50\% = 3$	$42-82\%^{79,83,84}$ $N=212$ $n=3$ $n_{\geq 50\%}=2$	$13-100\% ^{87,88,90}, \\ 91,93,100,101, \\ 106,117,128,129$ $N=1403$ $n=11$ $n \ge 50\% = 7$	$22\%^{137} \\ N = 50 \\ n = 1$	$43 - 95\%^{141,143},$ $N = 528$ $n = 3$ $n \ge 50\% = 2$	$42\%^{149} \\ N = 101 \\ n = 1$	-	$80\%^{156}$ $N = 52$ $n = 1$		
	Anorexia $n = 10$	$N = 23,825$ $n = 34$ $n \ge 50\% = 25$ $76 - 95\%^{63,67,70}$ $N = 1098$ $n = 3$	$64-67\%^{1,a}$ $N=87$ $n=1$	-	$38-64\%^{95,103},$ $106,128$ $N = 788$ $n = 4$	-	$82\%^{145} N = 52 n = 1$	$13\%^{154}$ $N = 123$ $n = 1$	-	-		
	Pain (including different origins) $n = 86$	$\begin{array}{c} 30 - 94\%^{5,16-19,} \\ 21 - 25,27 - 29,31,\\ 32,34,35,37 - 40,\\ 42 - 52,54 - 60,\\ 63 - 65,67 - 71 \end{array}$ $\begin{array}{c} N = 29,966 \\ n = 47 \end{array}$	$21-77\%^{1,72-75},$ $77,79,a$ $N = 1730$ $n = 7$ $n \ge 50\% = 3$	$14-78\%^{73,79},$ $81-85$ $N=1230$ $n=7$ $n \ge 50\% = 2$	$\begin{array}{l} n-1 \\ n \ge 50\% = 3 \\ 11-83\% & 86-88,93, \\ 94,100,101, \\ 103,131,b \\ N=814 \\ n=9 \\ n \ge 50\% = 6 \end{array}$	$14-63\%^{85},$ $132-135,137$ $N = 745$ $n = 6$ $n \ge_{50\%} = 1$	$30-98\%^{138,139},$ $142-145,\epsilon$ $N=1325$ $n=6$ $n \ge 50\% = 5$	$42-85\% \frac{147,148}{152,154}$ $N = 327$ $n = 4$ $n \ge 50\% = 2$	$52 - 76\%^{152,155,d}$ $N = 102$ $n = 2$	$68\%^{156}$ $N = 52$ $n = 1$		
	Nausea-vomiting $n = 67$	$n = 17$ $n \ge 50\% = 4$ $2 - 78\% $	$4\%^{79}$ $N = 81$ $n = 1$	$2-48\%^{79,81},$ $83,84$ $N=812$ $n=4$	$8-52\%^{86,87,90,}_{91,101,103,}_{106,117,128,b}$ $N=1176$ $n=9$	$8\%^{137}$ $N = 50$ $n = 1$	$41-57\%^{138,139},$ $143,145,c$ $N=914$ $n=4$ $n \ge 50\% = 1$	-	-	$26\%^{156}$ $N = 52$ $n = 1$		
	Breathlessness/ dyspnea/ shortness of breath/respiratory distress n = 73	$n \ge 50\% = 3$ $16 - 77\%^{5,16-21,23-35},$ $37 - 54,56 - 60,63,65,$ $67 - 71$	$56-98\%^{1,72-75}_{77,79,80,a}$ $N=1835$ $n=8$	$18-88\%^{73,79},$ $81-85$ $N = 1230$ $n = 7$ $n \ge 50\% = 5$	$n = 50\% = 1$ $n \ge 50\% = 1$ $11 - 82\% 86 - 88.90$ $.91,93,95,100$, $101,117,131,b$ $N = 1031$ $n = 11$ $n \ge 50\% = 4$	$12-52\%^{85},$ $132-134,137$ $N = 674$ $n = 5$ $n \ge 50\% = 1$	$43-62\%^{138},$ $139,145,\epsilon$ $N = 811$ $n = 3$ $n \ge 50\% = 3$	-	$81 - 88\%^{155,d}$ $N = 52$ $n = 1$	$26\%^{156}$ $N = 52$ $n = 1$		

	Insomnia/ poor sleeping/ difficulty sleeping/ sleep disturbance n = 65	$\begin{array}{l} 3-67\% 5.16-18.20.21, \\ 23-25.27.29.31,32.34, \\ 35.37.38.41, \\ 43.45.47-50, \\ 52.56-59, \\ 63.65.67.70 \\ N=21.725 \\ n=33 \end{array}$	$15-77\%^{1,74,75,77,a}$ $N=419$ $n=4$ $n \ge 50\% = 3$	$36{-}48\%^{81,83,84}$ $N{=}746$ $n{=}3$	$\begin{aligned} &1 - 83\% ^{87,88,91,93,}_{&98,101,103,}_{&117 - 120,}_{&122 - 127,131} \\ &N = 2265\\ &n = 18\\ &n \geq 50\% = 7 \end{aligned}$	$ \begin{array}{l} 14\%^{136} \\ N = 123 \\ n = 1 \end{array} $	$40-74\%^{138,139},$ $145,c$ $N=811$ $n=3$ $n_{\geq 50\%}=3$	$43-50\%^{149,154}$ $N=224$ $n=2$ $n_{\geq 50\%}=1$	$24-33\%^{155,d}$ $N=52$ $n=1$	-
	Confusion/ delirium/ cognitive failure/ cognitive symptoms	$\begin{array}{l} n \geq 50\% = 11 \\ 2-68\% & 18,22,23,27, \\ 29,31-34, \\ 38,40,41,44, \\ 46-52,56, \\ 57,62,65,71 \\ N = 18,768 \end{array}$	$14 - 33\%^{1,73,a}$ $N = 309$ $n = 2$	$15-48\%^{73,82,83}$ $N=433$ $n=3$	$35-70\% {}^{86,b}$ N=32 n=1	-	-	$24\%^{154}$ $N=123$ $n=1$	-	-
	n = 31 Constipation $n = 60$	$n = 25$ $n \le 50\% = 3$ $4 - 64\% 5.18 - 25.29.30,$ $32 - 38.40 - 50.52.54,$ $56 - 60.63.65,$ $67.69 - 71$ $N = 30.841$	$12-44\%^{1,75,a}$ $N=150$ $n=2$	$12 - 42\% {}^{81,83,84}$ $N = 746$ $n = 3$	$8-65\%^{87,88,92,}_{101,106,130,131}$ $N=1136$ $n=7$ $n \ge 50\%=3$	$40\%^{137}$ $N = 50$ $n = 1$	$19 - 35\% {}^{138,139,\epsilon}$ $N = 759$ $n = 2$	$24\%^{153} \\ N = 123 \\ n = 1$	$52-56\%^{155,d}$ $N=52$ $n=1$	$46\%^{156} \\ N = 52 \\ n = 1$
	Diarrhea $n = 33$	$n = 42$ $n \ge 50\% = 10$ $1 - 25\% 5.20.21.23 - 25.27,$ $29.32.34.37.40 - 42.44,$ $46 - 48.56 - 60.69$ $N = 17.813$	-	$12\%^{83,84} \\ N = 146 \\ n = 2$	$8-36\%^{86-88,131,b}$ $N=105$ $n=4$	-	$29-53\%^{138,139,143,c}$ $N=862$ $n=3$ $n_{\geq 50\%}=2$	-	-	-
Psychological:	Depression/ depressive mood/sadness/ low mood n = 71	$\begin{array}{l} n=24\\ 4-80\%, 5,16,\\ 18=20,23,\\ 27-29,32,\\ 34,37,38,\\ 41,47,48,50,\\ 52,56,57,63,\\ 65,68-71\\ N=16,200\\ n=26 \end{array}$	$17-77\%^{1,74,76-80,a}$ $N = 752$ $n = 7$ $n \ge 50\% = 4$	$6-59\%^{79-81,83,84} \\ N=892 \\ n=5 \\ n \geq 50\% = 1$	$\begin{array}{l} 2-61\%^{87-89,91},\\ 96-100,102,\\ 104-116 \end{array}$ $\begin{array}{l} N=7644\\ n=23\\ n_{\geq 50\%}=3 \end{array}$	$46\%^{136} \\ N = 123 \\ n = 1$	$17-82\% {}^{138-140,142},$ $143,145$ $N=1211$ $n=6$ $n \ge 50\% = 4$	$15-50\%^{146},_{149-151,153}$ $N=24,759$ $n=5$ $n \ge 50\%=1$	$23\%^{153}$ $N = 119$ $n = 1$	$ 15\%^{153} $ $N = 162$ $n = 1$
	Anxiety/ nervousness/ agitation n = 50	$n = 250\% = 10$ $n \ge 50\% = 10$ $3 - 74\% 5.16.20.28.32.38,$ $47.48.50.52.56.57.63,$ $65.66.68 - 71$ $N = 14.724$ $n = 19$	$23-53\%^{74,77,79,80} \\ N=455 \\ n=4 \\ n \ge 50\% = 1$	$2-49\%^{79-81,83,84}$ $N=892$ $n=5$	$7-52\%^{86-88,91},$ $96,99,100,$ $106,107,109,b$ $N=818$ $n=10$	$8-72\%^{133-137}$ $N=708$ $n=5$ $n_{\geq 50\%}=2$	$\begin{array}{l} 13 - 76\%^{138 - 140,145} \\ N = 923 \\ n = 4 \\ n \ge_{50\%} = 2 \end{array}$	$5-62\%^{146,149},$ $151,153,154$ $N=24,842$ $n=5$ $n \ge 50\% = 2$	$ 19\%^{153} \\ N = 119 \\ n = 1 $	$24\%^{153} \\ N = 162 \\ n = 1$
	Worry $n = 12$	$n \ge 50\% = 7$ $3-71\%^{16,23},$ $27,29,32,57,69$ $N=1500$ $n=8$ $n \ge 50\% = 3$	$65\%^{77}$ $N = 60$ $n = 1$	-	$\begin{array}{l} n \geq 50\% = 1 \\ 32 - 55\% ^{87,88} \\ N = 30 \\ n = 2 \\ n \geq 50\% = 1 \end{array}$	-	$51-86\%^{138,139,c}$ $N=759$ $n=2$	-	-	-

Table 2
Continued

				COL	iiiiiiaca							
	Problems and Number of Different Studies	Minimum-Maximum Prevalence % of Each Problem References Total Sample of Patients (N) for Each Problem Total Number of Studies (n) for Each Problem Number of Studies That Report Prevalence for a Problem as $\geq 50\%$ $(n_{\geq 50\%})$										
Domain	for Each Problem for All Diagnostic Groups (n)	Cancer	COPD	CHF	ESRD	Dementia	AIDS	Parkinson	MND	MS		
Spiritual:	Well-being $n=2$	$91\%^{70}$ $N = 406$ $n = 1$	$14\%^{79}$ $N = 81$ $n = 1$	$5\%^{79}$ $N = 66$ $n = 1$	-	-	-	-	-	-		
	Spiritual pain $n=1$	$44\%^{61}$ $N = 91$ $n = 1$	-	-	-	-	-	-	-	-		
	Spiritual wellbeing $n=1$	<i>n</i> – 1	-	$6\%^{84}$ $N = 66$ $n = 1$	-	-	-	-	-	-		
	Peace $n=0$	-	-	-	-	-	-	-	-	-		
Social:	Support $n = 1$	$49\%^{71}$ $N = 4035$ $n = 1$	-	-	-	-	-	-	-	-		
	Information $n = 0$	-	-	-	-	-	-	-	-	-		
	Psychosocial distress $n = 0$	-	-	-	-	-	-	-	-	-		

COPD = chronic obstructive pulmonary disease; CHF = chronic heart failure; ESRD = end-stage renal disease; AIDS = acquired immune deficiency syndrome; MND = motor neuron disease; MS = multiple sclerosis.

A dash indicates that no data were found for this specific problem in this specific diagnostic group.

[&]quot;This study included a period prevalence measurement on two time points: in the final year of life and in the final week of life.

^bThis study included a point and a period prevalence measurement in the same sample of patients.

^oThis study took place in three different sites and contains data for three different samples of patients.

^dThis study contains the data of two groups of patients: one group who died at home and one group who died in a hospice.




Fig. 2. PRISMA flow diagram of search process.

We found only four studies on spiritual, ^{61,70,79,84} and one study on social palliative care—related⁷¹ problems. In people living with advanced cancer, lack of well-being and support were the problems that were considerably prevalent (91% and 49%, respectively).

The problems for which we found the widest minimum (<10%) and maximum $(\ge50\%)$ prevalence ranges were: nausea-vomiting in advanced cancer and ESRD; insomnia in advanced cancer and ESRD; delirium in advanced cancer; constipation in advanced cancer and

ESRD; depression in advanced cancer, CHF and ESRD; anxiety in advanced cancer, ESRD, dementia, and Parkinson's disease; and worry in advanced cancer.

Looking at most studies that presented a prevalence of 50% and more in comparison with the total number of studies for each problem per diagnostic group, the following problems showed a prevalence of 50% or higher: fatigue in advanced cancer, COPD, CHF, ESRD, AIDS, and MS; anorexia in advanced cancer, COPD, ESRD, and AIDS; pain in advanced cancer,

COPD, ESRD, AIDS, Parkinson's disease, MND, and MS; dyspnea in COPD, CHF, ESRD, AIDS, and MND; insomnia in COPD, AIDS, and Parkinson's disease; constipation in ESRD and MND; diarrhea in AIDS; depression in COPD and AIDS; anxiety in AIDS and Parkinson's disease; worry in advanced cancer, COPD, ESRD, and AIDS; and lack of well-being in advanced cancer.

Within the group of 143 articles, there were only five studies that directly compared the prevalence of palliative care—related problems with another diagnostic group while using the same measurement tool. ^{73,79,80,152,153} In these five studies, COPD patients were compared with CHF and dementia patients, ^{73,79,80} and Parkinson's disease patients to MND and MS patients. ^{152,153} In these studies, the range of the minimum and maximum prevalence ranges was not wide.

Discussion

This study showed that there are commonalities in problem prevalence across diagnostic groups. Some problems were rarely studied, for example, social and spiritual problems; and for certain diagnostic groups, we found a very limited number of problem prevalence studies, for example, for people living with MND and MS.

There are a balanced number of strengths, weaknesses, or limitations in this systematic review. One of the limitations of our study is that each disease group we selected has its specific pathology process with its specific (other) symptoms occurring throughout the trajectory. For example, muscle cramps, dry skin, muscle soreness, headaches, and restless legs are specific for people living with ESRD, but these are not included in this review as we aimed to summarize the generally prevalent problems across all our studied diagnostic groups. We also recognize publication bias while looking at certain domains of interest; for example, social and spiritual problems may receive lower priority from peer-reviewed publications. Hence, problems for which we could not find (sufficient) evidence does not mean that these problems are not present. Another limitation is that systematic reviews rely on a broad yet accurate search strategy to identify relevant evidence. We tried to limit this by implementing

a search strategy using synonyms for specific palliative care—related problems. Lastly, from the beginning of our study, we were aware that by collecting data on our research topic, we had to deal with presenting a heterogeneous amount of information. This, for example, resulted in our presenting the data in a very descriptive way using minimum and maximum prevalence ranges.

We believe that this review has the following strengths: using the PRISMA 2009 flowchart to execute our search in a stepwise way, giving us a structured overview of which studies were correctly excluded; assessing the quality of the review using the PRISMA 2009 checklist and the AMSTAR measurement tool; reviewing the evidence on palliative care-related problems not only focusing on the physical and psychological problems but also using a total care view and including prevalence studies of spiritual and social palliative care-related problems; using several search or keywords to describe the same single problem; and excluding intervention studies to prevent selection bias. We expanded the variety of palliative care-related problems and non-cancer conditions in comparison with the review of Solano et al, allowing us to provide a broader view on the palliative care—related problems in palliative care patients in accord with the WHO definition of palliative care.8

One of the study's objectives was to review the prevalence studies across all palliative care domains, but only two studies investigated well-being, 70,79 one study the presence of spiritual pain, 61 one study the lack of spiritual well-being, 84 and one study the problems with receiving support. 71 We can understand this situation given the current availability and accessibility of spiritual outcome measurement tools. There are only nine spiritual outcome measurement tools that have been validated in cross-cultural palliative care populations. 157

The variations in the minimum and maximum prevalence ranges were likely caused by the differences in time points of data collection (point vs. period prevalence, or a mix of both), the period prevalence time ranges, the varying definitions for a problem, the sample size proportions, and the various different measurement methods used across studies. In light of the wide variations in the minimum and maximum prevalence

ranges and the methodological heterogeneity across studies, and the insights gained by looking at the five direct comparison studies, we can state that there is as much variation in prevalence between diagnostic groups as there is similarity. Also, understanding the prevalence of palliative care-related problems of non-cancer patients in comparison with cancer patients is complex as a result of the varying trajectories of functional decline and dependency in non-cancer diagnostic groups, 158 and their comorbidities and changing disease stages. Therefore, although there are commonalities in prevalence across diagnostic groups, the methodological heterogeneity across studies and the limited number of non-cancer studies must be taken into consideration when interpreting our results.

Our review identified some gaps that need to be addressed in future research. More research is needed that directly compares palliative care—related problems between cancer and non-cancer patients using the same measurement tools and time points. Our results also call for an intensification of the measurement and reporting of spiritual and social palliative care—related problems.

We also found that pain, fatigue, anorexia, dyspnea, and anxiety were highly prevalent problems across advanced cancer and non-cancer diagnostic groups. Health care providers should be aware of these problems and be more sensitive to their assessment and treatment.

Future prevalence studies should build on the findings of our review in terms of the most often used measurement methods across studies, and our identified evidence gaps, particularly in the under-researched non-cancer patient groups.

In conclusion, our study reveals that there is as much variation within diagnostic groups as between groups in relation to prevalence of palliative care—related problems. Our study can be considered as an epidemiological needs assessment while using the summarized prevalence data of palliative care—related problems among advanced cancer and non-cancer patients indicating that both are in need of palliative care. These summarized prevalence figures across diagnostic groups support the conclusion that palliative care should be made available based on the problems and needs patients exhibit, and not based on the diagnostic group to which a patient belongs.

Disclosures and Acknowledgments

Funding and sponsorship for this study were made available by EURO IMPACT, **Euro**pean Intersectorial and **M**ultidisciplinary **Pa**lliative Care Research Training, funded by the European Union Seventh Framework Programme (FP7/2007-2013, under grant agreement number [264697]). The authors declare no conflicts of interest.

The EURO IMPACT project aims to develop a multidisciplinary, multiprofessional, and intersectorial educational and research training framework for palliative care research in Europe. EURO IMPACT is coordinated by Prof. Luc Deliens and Prof. Lieve Van den Block of the End-of-Life Care Research Group, Ghent University & Vrije Universiteit Brussel, Brussels, Belgium. Other partners are: VU University Medical Center, EMGO Institute for Health Care Research, Amsterdam, and Netherlands; King's College London, Cicely Saunders Institute, London, Cicely Saunders International, London, and International Observatory on End-of-Life Care, Lancaster University, Lancaster, United Kingdom; Norwegian University of Science and Technology, and EAPC Research Network, Trondheim, Norway; Regional Palliative Care Network, IRCCS AOU San Martino-IST, Genoa, and Cancer Research and Prevention Institute, Florence, EUGMS European Union Geriatric Medicine Society, Geneva, Switzerland; and Springer Science and Business Media, Houten, The Netherlands.

In recognition of the collaborative nature of EURO IMPACT, the authors thank the following EURO IMPACT members for their contribution, namely: Sarah Brearley, Augusto Caraceni, Joachim Cohen, Massimo Costantini, Luc Deliens, Anneke L. Francke, Stein Kaasa, Karen Linden, Koen Meeussen, Guido Miccinesi, Bregje Onwuteaka-Philipsen, Koen Pardon, Roeline Pasman, Sophie Pautex, Sheila Payne, and Lieve Van den Block.

References

- 1. Edmonds P, Karlsen S, Khan S, Addington-Hall J. A comparison of the palliative care needs of patients dying from chronic respiratory disease and lung cancer. Palliat Med 2001;15:287—295.
- 2. Higginson IJ, Jarman B, Astin P, Dolan S. Do social factors affect where patients die: an analysis

- of 10 years of cancer deaths in England. J Public Health Med 1999;21:22-28.
- 3. Murray C, Lopez A. Mortality by cause for eight regions of the world: global burden of disease study. Lancet 1997;349:1269—1276.
- 4. Seale C. Changing patterns of death and dying. Soc Sci Med 2000;51:917–930.
- **5.** Potter J, Hami F, Bryan T, Quigley C. Symptoms in 400 patients referred to palliative care service: prevalence and patterns. Palliat Med 2003;17: 310–314.
- 6. Addington-Hall JM, Higginson IJ. Palliative care for non-cancer patients, 1st ed. New York: Oxford University Press, 2001.
- 7. Solano J, Gomes B, Higginson IJ. A comparison of symptom prevalence in far advanced cancer, AIDS, heart disease, chronic obstructive pulmonary disease and renal disease. J Pain Symptom Manage 2006;31:58—69.
- 8. World Health Organization. WHO definition of "palliative care." Available from http://www.who.int/cancer/palliative/definition/en/. Accessed May 14, 2012.
- 9. Elman L, Houghton D, Wu G, et al. Palliative care in amyotrophic lateral sclerosis, Parkinson's disease, and multiple sclerosis. J Palliat Med 2007; 10:433–457.
- **10.** United Nations Population Division. World population prospects. New York: United Nations, 2007.
- 11. Centre for Reviews and Dissemination. Systematic reviews: CRD's guidance for undertaking reviews in health care. York: CRD, University of York, 2009.
- 12. Moher D, Liberati A, Tetzlaff J, Altman D, The PRISMA Group. Preferred reporting items for systematic reviews and meta-analyses: the PRISMA statement. PLoS Med 2009;6:e1000097.
- 13. Shea J, Grimshaw M, Wells A, et al. Development of AMSTAR: a measurement tool to assess the methodological quality of systematic reviews. BMC Med Res Methodol 2007;7:7.
- 14. Simms V, Higginson I, Harding R. What palliative care-related problems do patients experience at HIV diagnosis? A systematic review of the evidence. J Pain Symptom Manage 2011;42:734–753.
- 15. Lipman A, Jackson K, Tyler L, eds. Evidence based symptom control in palliative care. Systematic reviews and validated clinical practice guidelines for 15 common problems in patients with life limiting disease. New York: Pharmaceutical Products Press, 2000.
- 16. Chang V, Hwang S, Feuerman M, Kasimis B. Symptom and quality of life survey of medical oncology patients at a veteran's affairs medical center: a role for symptom assessment. Cancer 2000;88: 1175–1183.

- 17. Chen M, Chang H. Physical symptom profiles of depressed and nondepressed patients with cancer. Palliat Med 2004;18:712–718.
- 18. Dunlop G. A study of the relative frequency and importance of gastrointestinal symptoms, and weakness in patients with far advanced cancer: student paper. Palliat Med 1989;4:31–43.
- 19. Edmonds P, Stuttaford J, Penny J, Lynch A, Chamberlain J. Do hospital palliative care teams improve symptom control? Use of a modified STAS as an evaluation tool. Palliat Med 1998;12: 345–351.
- **20.** Grond S, Zech D, Diefenbach C, Bischoff A. Prevalence and pattern of symptoms in patients with cancer pain: a prospective evaluation of 1635 cancer patients referred to a pain clinic. J Pain Symptom Manage 1994;9:377–382.
- 21. Hoekstra J, de Vos R, van Duijn N, Schade E, Bindels P. Using the symptom monitor in a randomized controlled trial: the effect on symptom prevalence and severity. J Pain Symptom Manage 2006; 31:22–30.
- 22. Kutner J, Kassner C, Nowels D. Symptom burden at the end of life: hospice providers' perceptions. J Pain Symptom Manage 2001;21:473–480.
- 23. Lidstone V, Butters E, Seed P, et al. Symptoms and concerns amongst cancer outpatients: identifying the need for specialist palliative care. Palliat Med 2003;17:588–595.
- 24. Lo R, Ding A, Chung T, Woo J. Prospective study of symptom control in 133 cases of palliative care inpatients in Shatin Hospital. Palliat Med 1999;13:335–340.
- 25. Lundh Hagelin C, Seiger A, Furst C. Quality of life in terminal care—with special reference to age, gender and marital status. Support Care Cancer 2006;14:320–328.
- **26.** Lutz S, Norrell R, Bertucio C, et al. Symptom frequency and severity in patients with metastatic or locally recurrent lung cancer: a prospective study using the Lung Cancer Symptom Scale in a community hospital. J Palliat Med 2001;4:157–165.
- 27. McMillan S, Small B. Symptom distress and quality of life in patients with cancer newly admitted to hospice home care. Oncol Nurs Forum 2002;29: 1421–1428.
- 28. Modonesi C, Scarpi E, Maltoni M, et al. Impact of palliative care unit admission on symptom control evaluated by the Edmonton Symptom Assessment System. J Pain Symptom Manage 2005;30: 367–377.
- 29. Portenoy R, Thaler H, Kornblith A, et al. Symptom prevalence, characteristics and distress in a cancer population. Qual Life Res 1994;3:183–189.
- 30. Schuit K, Sleijfer D, Meijler W, et al. Symptoms and functional status of patients with disseminated cancer visiting outpatient departments. J Pain Symptom Manage 1998;16:290—297.

- **31.** Tamburini M, Brunelli C, Rosso S, Ventafridda V. Prognostic value of quality of life scores in terminal cancer patients. J Pain Symptom Manage 1996;11:32–41.
- 32. Tranmer J, Heyland D, Dudgeon D, et al. Measuring the symptom experience of seriously ill cancer and noncancer hospitalized patients near the end of life with the memorial symptom assessment scale. J Pain Symptom Manage 2003;25: 420–429.
- 33. Bedard J, Dionne A, Dionne L. The experience of La Maison Michel Sarrazin (1985-1990): profile analysis of 952 terminal-phase cancer patient. J Palliat Care 1991;7:42—53.
- 34. Conill C, Verger E, Henriquez I. Symptom prevalence in the last week of life. J Pain Symptom Manage 1997;14:328—331.
- 35. Ellershaw J, Peat S, Boys L. Assessing the effectiveness of a hospital palliative care team. Palliat Med 1995;9:145–152.
- **36.** Grosvenor M, Bulcavage L, Chlebowski R. Symptoms potentially influencing weight loss in a cancer population. Correlations with primary site, nutritional status, and chemotherapy administration. Cancer 1989;63:330–334.
- 37. Krech R, Walsh D. Symptoms of pancreatic cancer. J Pain Symptom Manage 1991;6:360–367.
- 38. Krech R, Davis J, Walsh D, Curtis E. Symptoms of lung cancer. Palliat Med 1992;6:309—315.
- **39.** Maltoni M, Pirovano M, Scarpi E, et al. Prediction of survival of patients terminally ill with cancer. Results of an Italian prospective multicentric study. Cancer 1995;75:2613—2622.
- **40.** Mercadante S, Fulfaro F, Casuccio A. The impact of home palliative care on symptoms in advanced cancer patients. Support Care Cancer 2000;8:307–310.
- 41. Meuser T, Pietruck C, Radbruch L, et al. Symptoms during cancer pain treatment following WHO guidelines: a longitudinal follow-up study of symptom prevalence, severity and etiology. Pain 2001; 93:247–257.
- **42.** Morita T, Tsunoda J, Inoue S, Chihara S. Contributing factors to physical symptoms in terminally ill cancer patients. J Pain Symptom Manage 1999;18:338–346.
- 43. Radbruch L, Sabatowski R, Loick G, et al. Cognitive impairment and its influence on pain and symptom assessment in a palliative care unit: development of a Minimal Documentation System. Palliat Med 2000;14:266–276.
- 44. Reuben D, Mor V, Hiris J. Clinical symptoms and length of survival in patients with terminal cancer. Arch Intern Med 1998;148:1586—1591.
- 45. Sebastian P, Varghese C, Sankaranarayanan R, et al. Evaluation of symptomatology in planning palliative care. Palliat Med 1993;7:27–34.

- **46.** Simpson K. The use of research to facilitate the creation of a hospital palliative care team. Palliat Med 1991;5:122–129.
- 47. Tay W, Shaw R, Goh C. A survey of symptoms in hospice patients in Singapore. Ann Acad Med Singapore 1994;23:191–196.
- 48. Teunissen S, de Graeff A, de Haes J, Voest E. Prognostic significance of symptoms of advanced cancer patients. Eur J Cancer 2006;42:2510–2516.
- 49. Vainio A, Auvinen A. Prevalence of symptoms among patients with advanced cancer: an international collaborative study. Symptom Prevalence Group. J Pain Symptom Manage 1996;12:3—10.
- **50.** Walsh D, Donnelly S, Rybicki L. The symptoms of advanced cancer: relationship to age, gender, and performance status in 1,000 patients. Support Care Cancer 2000;8:175–179.
- **51.** Brescia F, Adler D, Gray G, et al. Hospitalized advanced cancer patients: a profile. J Pain Symptom Manage 1990;5:221–227.
- 52. Coyle N, Adelhardt J, Foley K, Portenoy R. Character of terminal illness in the advanced cancer patient: pain and other symptoms during the last four weeks of life. J Pain Symptom Manage 1990;5: 83–93
- **53.** Dunn A, Carter J, Carter H. Anemia at the end of life: prevalence, significance, and causes in patients receiving palliative care. J Pain Symptom Manage 2003;26:1132–1139.
- **54.** Dunphy K, Amesbury B. A comparison of hospice and home care patients: patterns of referral, patient characteristics and predictors of place of death. Palliat Med 1990;4:105–111.
- 55. Forbes K. Palliative care in patients with cancer of the head and neck. Clin Otolaryngol Allied Sci 1997;22:117–122.
- **56.** Klepstad P, Kaasa S, Cherny N, Hanks G, de Conno F, Research Steering Committee of the EAPC. Pain and pain treatments in European palliative care units. A cross sectional survey from the European Association for Palliative Care Research Network. Palliat Med 2005;19:477–484.
- **57.** Stromgren A, Groenvold M, Pedersen L, Olsen A, Sjogren P. Symptomatology of cancer patients in palliative care: content validation of self-assessment questionnaires against medical records. Eur J Cancer 2002;38:788–794.
- 58. Twycross R, Lack S. Control of alimentary symptoms in advanced cancer. Edinburgh: Churchill Livingstone, 1986:1–11.
- **59.** Oi-Ling K, Man-Wah D, Kam-Hung D. Symptom distress as rated by advanced cancer patients, caregivers and physicians in the last week of life. Palliat Med 2005;19:228–233.
- **60.** Wachtel T, Allen-Masterson S, Reuben D, Goldberg R, Mor V. The end stage cancer patient: terminal common pathway. Hosp J 1988;4:43–80.

- **61.** Delgado-Guay M, Hui D, Parsons H, et al. Spirituality, religiosity, and spiritual pain in advanced cancer patients. J Pain Symptom Manage 2011;41: 986–994.
- **62.** Caraceni A, Nanni O, Maltoni M, et al. Impact of delirium on the short term prognosis of advanced cancer patients. Cancer 2000;89:1145–1149.
- **63.** Gómez-Batiste X, Porta-Sales J, Espinosa-Rojas J, et al. Effectiveness of palliative care services in symptom control of patients with advanced terminal cancer: a Spanish, multicenter, prospective, quasi-experimental, pre-post study. J Pain Symptom Manage 2010;40:652–660.
- **64.** Dumitrescu L, van den Heuvel-Olaroiu M, van den Heuvel W. Changes in symptoms and pain intensity of cancer patients after enrollment in palliative care at home. J Pain Symptom Manage 2007;34: 488–496.
- 65. Ogawa A, Shimizu K, Akizuki N, Uchitomi Y. Involvement of a psychiatric consultation service in a palliative care team at the Japanese cancer center hospital. Jpn J Clin Oncol 2010;40: 1139–1146.
- **66.** Spencer R, Nilsson M, Wright A, Pirl W, Prigerson H. Anxiety disorder in advanced cancer patients. Cancer 2010;116:1810—1819.
- **67.** Tsai J-S, Wu C-H, Chiu T-Y, Chen C-Y. Significance of symptom clustering in palliative care of advanced cancer patients. J Pain Symptom Manage 2010;39:655–662.
- **68.** Delgado-Guay M, Yennurajalingam S, Parsons H, Palmer L, Bruera E. Association between self-reported sleep disturbance and other symptoms in patients with advanced cancer. J Pain Symptom Manage 2011;41: 819–827.
- **69.** Harding R, Selman L, Agupio G, et al. The prevalence and burden of symptoms amongst cancer patients attending palliative care in two African countries. Eur J Cancer 2011;47:51–56.
- **70.** Jiménez A, Madero R, Alonso A, et al. Symptom clusters in advanced cancer. J Pain Symptom Manage 2011;42:24–31.
- 71. Ostgathe C, Alt-Epping B, Golla H, et al. Non-cancer patients in specialized palliative care in Germany: what are the problems? Palliat Med 2010;25: 148–152.
- 72. Claessens M, Lynn J, Zhong Z, et al. Dying with lung cancer or chronic obstructive pulmonary disease: insights from SUPPORT. Study to understand prognoses and preferences for outcomes and risks of treatments. J Am Geriatr Soc 2000;48:146–153.
- 73. Lynn J, Teno J, Phillips R, et al. Perceptions by family members of the dying experience of older and seriously ill patients. SUPPORT Investigators. Study to understand prognoses and preferences for outcomes and risks of treatments. Ann Intern Med 1997;126:97–106.

- **74.** Elkington H, White P, Addington-Hall J, Higgs R, Edmonds P. The healthcare needs of chronic obstructive pulmonary disease patients in the last year of life. Palliat Med 2005;19:485–491.
- 75. Skilbeck J, Mott L, Page H, et al. Palliative care in chronic obstructive airways disease: a needs assessment. Palliat Med 1998;12:245–254.
- 76. Stapleton R, Nielsen E, Engelberg R, Patrick D, Curtis J. Association of depression and life-sustaining treatment preferences in patients with COPD. Chest 2005;127:328–334.
- 77. Bausewein C, Booth S, Gysels M, et al. Understanding breathlessness: cross-sectional comparison of symptom burden and palliative care needs in chronic obstructive pulmonary disease and cancer. J Palliat Med 2010;13:1109—1118.
- 78. Lacasse Y, Rousseau L, Maltais F. Prevalence of depressive symptoms and depression in patients with severe oxygen-dependent chronic obstructive pulmonary disease. J Cardiopulm Rehabil 2001;21: 80–86.
- **79.** Walke L, Gallo W, Tinetti M, Fried T. The burden of symptoms among community-dwelling older persons with advanced chronic disease. Arch Intern Med 2004;164:2321–2324.
- **80.** Janssen D, Spruit M, Uszko-Lencer N, Schols J, Wouters E. Symptoms, comorbidities, and health care in advanced chronic obstructive pulmonary disease or chronic heart failure. J Palliat Med 2011;14: 735–743.
- **81.** McCarthy M, Lay M, Addington-Hall J. Dying from heart disease. J R Coll Physicians Lond 1996; 30:325–328.
- **82.** Zambroski C, Moser D, Roser L, Heo S, Chung M. Patients with heart failure who die in hospice. Am Heart J 2005;149:558–564.
- 83. Nordgren L, Sorensen S. Symptoms experienced in the last six months of life in patients with end-stage heart failure. Eur J Cardiovasc Nurs 2003;2:213–217.
- 84. Anderson H, Ward C, Eardley A, et al. The concerns of patients under palliative care and a heart failure clinic are not being met. Palliat Med 2001; 15:279–286.
- **85.** Formiga F, Olmedo C, Lopez-Soto A, et al. Dying in hospital of terminal heart failure or severe dementia: the circumstances associated with death and the opinions of caregivers. Palliat Med 2007; 21:35–40.
- **86.** Chater S, Davison S, Germain M, Cohen L. Withdrawal from dialysis: a palliative care perspective. Clin Nephrol 2006;66:364–372.
- 87. Weisbord S, Carmody S, Brun F, et al. Symptom burden, quality of life, advance care planning and the potential value of palliative care in severely ill haemodialysis patients. Nephrol Dial Transplant 2003;18:1345—1352.

- 88. Saini T, Murtagh F, Dupont P. Comparative pilot study of symptoms and quality of life in cancer patients and patients with end stage renal disease. Palliat Med 2006;20:631–636.
- 89. Cohen L, Dobscha S, Hails K, Pekow P, Chochinov H. Depression and suicidal ideation in patients who discontinue the life-support treatment of dialysis. Psychosom Med 2002;64:889–896.
- **90.** Parfrey P, Vavasour H, Henry S, Bullock M, Gault M. Clinical features and severity of nonspecific symptoms in dialysis patients. Nephron 1988; 50:121–128.
- **91.** McCann K, Boore J. Fatigue in persons with renal failure who require maintenance haemodialysis. J Adv Nurs 2000;32:1132—1142.
- **92.** Yasuda G, Shibata K, Takizawa T, et al. Prevalence of constipation in continuous ambulatory peritoneal dialysis patients and comparison with hemodialysis patients. Am J Kidney Dis 2002;39: 1292–1299.
- 93. Kimmel P, Emont S, Newmann J, et al. ESRD patient quality of life: symptoms, spiritual beliefs, psychosocial factors, and ethnicity. Am J Kidney Dis 2003;42:713—721.
- **94.** Davison S. Pain in hemodialysis patients: prevalence, cause, severity, and management. Am J Kidney Dis 2003;42:1239—1247.
- 95. Loos C, Briançon S, Frimat L, Hanesse B, Kessler M. Effect of end-stage renal disease on the quality of life of older patients. J Am Geriatr Soc 2003;51:229–233.
- **96.** Martin C, Thompson D. Prediction of quality of life in patients with end-stage renal disease. Br J Health Psychol 2000;5:41–55.
- **97.** Lopes A, Bragg J, Young E, et al. Depression as a predictor of mortality and hospitalization among hemodialysis patients in the United States and Europe. Kidney Int 2002;62:199–207.
- 98. Iliescu E, Coo H, McMurray M, et al. Quality of sleep and health-related quality of life in haemodialysis patients. Nephrol Dial Transplant 2003;18: 126–132.
- 99. Aghanwa H, Morakinyo O. Psychiatric complications of hemodialysis at a kidney center in Nigeria. J Psychosom Res 1997;42:445–452.
- 100. Cohen L, Germain M, Woods A, Mirot A, Burleson J. The family perspective of ESRD death. Am J Kidney Dis 2005;45:154–161.
- **101.** Yong D, Kwok A, Wong D, et al. Symptom burden and quality of life in end-stage renal disease: a study of 179 patients on dialysis and palliative care. Palliat Med 2009;23:111–119.
- 102. Tyrrell J, Paturel L, Cadec B, Capezzali E, Poussin G. Older patients undergoing dialysis treatment: cognitive functioning, depressive mood and health-related quality of life. Aging Ment Health 2005;9:374—379.

- 103. Chan C-K, Wong SS-H, Ho T-LE, et al. Supportive management in patients with end-stage renal disease: local experience in Hong Kong. Hong Kong J Nephrol 2010;12:31—36.
- 104. al Hihi E, Awad A, Hagedorn A. Screening for depression in chronic hemodialysis patients. Mo Med 2003;100:266–268.
- 105. Craven J, Rodin G, Littlefield C. The Beck Depression Inventory as a screening device for major depression in renal dialysis patients. Int J Psychiatry Med 1988;18:365–374.
- 106. Curtin R, Bultman D, Thomas-Hawkins C, et al. Hemodialysis patients' symptom experiences: effects on physical and mental functioning. Nephrol Nurs J 2002;29:567–574.
- 107. Farmer C, Snowden S, Parsons V. The prevalence of psychiatric illness among patients on home haemodialysis. Psychol Med 1979;9:509—514.
- 108. Hong B, Smith M, Robson A, et al. Depressive symptomatology and treatment in patients with end-stage renal disease. Psychol Med 1987;17:185—190.
- 109. Kutner N, Fair P, Kutner M. Assessing depression and anxiety in chronic dialysis patients. J Psychosom Res 1985;29:23–31.
- 110. O'Donnell K, Chung J. The diagnosis of major depression in end-stage renal disease. Psychother Psychosom 1997;66:38–43.
- 111. Rodin G, Voshart K. Depressive symptoms and functional impairment in the medically ill. Gen Hosp Psychiatry 1987;9:251–258.
- 112. Smith M, Hong B, Robson A. Diagnosis of depression in patients with end-stage renal disease. Comparative analysis. Am J Med 1985;79:160–166.
- 113. Walters B, Hays R, Spritzer K, et al. Health-related quality of life, depressive symptoms, anemia, and malnutrition at hemodialysis initiation. Am J Kidney Dis 2002;40:1185–1194.
- 114. Watnick S, Kirwin P, Mahnensmith R, et al. The prevalence and treatment of depression among patients starting dialysis. Am J Kidney Dis 2003;41: 105–110.
- 115. Wuerth D, Finkelstein S, Ciarcia J, et al. Identification and treatment of depression in a cohort of patients maintained on chronic peritoneal dialysis. Am J Kidney Dis 2001;37:1011–1017.
- 116. Wuerth D, Finkelstein S, Finkelstein F. The identification and treatment of depression in patients maintained on dialysis. Semin Dial 2005;18: 142–146.
- 117. Barrett B, Vavasour H, Major A, et al. Clinical and psychological correlates of somatic symptoms in patients on dialysis. Nephron 1990;55:10–15.
- 118. Devins G, Edworthy S, Paul L, et al. Restless sleep, illness intrusiveness, and depressive symptoms in three chronic illness conditions: rheumatoid arthritis, end-stage renal disease, and multiple sclerosis. J Psychosom Res 1993;37:163—170.

- 119. Han S, Yoon J, Jo S, et al. Insomnia in diabetic hemodialysis patients. Prevalence and risk factors by a multicenter study. Nephron 2002;92:127–132.
- 120. Holley J, Nespor S, Rault R. A comparison of reported sleep disorders in patients on chronic hemodialysis and continuous peritoneal dialysis. Am J Kidney Dis 1992;19:156–161.
- 121. Locking-Cusolito H, Huyge L, Strangio D. Sleep pattern disturbance in hemodialysis and peritoneal dialysis patients. Nephrol Nurs J 2001;28: 40–44.
- 122. Mucsi I, Molnar M, Ambrus C, et al. Restless legs syndrome, insomnia and quality of life in patients on maintenance dialysis. Nephrol Dial Transplant 2005;20:571–577.
- 123. Sabbatini M, Minale B, Crispo A, et al. Insomnia in maintenance haemodialysis patients. Nephrol Dial Transplant 2002;17:852–856.
- 124. Stepanski E, Faber M, Zorick F, et al. Sleep disorders in patients on continuous ambulatory peritoneal dialysis. J Am Soc Nephrol 1995;6:192—197.
- 125. Venmans B, van Kralingen K, Chandi D, et al. Sleep complaints and sleep disordered breathing in hemodialysis patients. Neth J Med 1999;54: 207–212.
- 126. Walker S, Fine A, Kryger M. Sleep complaints are common in a dialysis unit. Am J Kidney Dis 1995;26:751–756.
- 127. Winkelman J, Chertow G, Lazarus J. Restless legs syndrome in end-stage renal disease. Am J Kidney Dis 1996;28:372—378.
- 128. Curtis B, Barret B, Jindal K, et al. Canadian survey of clinical status at dialysis initiation 1998—1999: a multicenter prospective survey. Clin Nephrol 2002;58:282—288.
- 129. Frank A, Auslander G, Weissgarten J. Quality of life of patients with end-stage renal disease at various stages of the illness. Soc Work Health Care 2003;38:1–27.
- 130. Hammer J, Oesterreicher C, Hammer K, et al. Chronic gastrointestinal symptoms in hemodialysis patients. Wien Klin Wochenschr 1998;110(10): 287–291.
- 131. Murtagh F, Addington-Hall J, Edmonds P, et al. Symptoms in the month before death for stage 5 chronic kidney disease patients managed without dialysis. J Pain Symptom Manage 2010;40:342—352.
- 132. Black B, Finucane T, Baker A, et al. Health problems and correlates of pain in nursing home residents with advanced dementia. Alzheimer Dis Assoc Disord 2006;20:283—290.
- 133. Mitchell S, Teno J, Kiely D, et al. The clinical course of advanced dementia. N Engl J Med 2009; 361:1529–1538.
- 134. Di Giulio P, Toscani F, Villani D, et al. Dying with advanced dementia in long-term care geriatric

- institutions: a retrospective study. J Palliat Med 2008;11:1023–1028.
- 135. Aminoff B, Adunsky A. Dying dementia patients: too much suffering, too little palliation. Am J Hosp Palliat Care 2005;22:344–348.
- 136. Kverno K, Black B, Blass D, Geiger-Brown J, Rabins P. Neuropsychiatric symptom patterns in hospice-eligible nursing home residents with advanced dementia. J Am Med Dir Assoc 2008;9: 509–515.
- 137. Pautex S, Herrmann F, Le Lous P, et al. Symptom relief in the last week of life: is dementia always a limiting factor? Letters to the Editor. J Am Geriatr Soc 2007;55:1316–1317.
- 138. Vogl D, Rosenfeld B, Breitbart W, et al. Symptom prevalence, characteristics, and distress in AIDS outpatients. J Pain Symptom Manage 1999;18: 253–262.
- 139. Karus D, Raveis V, Alexander C, et al. Patient reports of symptoms and their treatment at three palliative care projects servicing individuals with HIV/AIDS. J Pain Symptom Manage 2005;30: 408–417.
- 140. Rabkin J, Ferrando S, Jacobsberg L, Fishman B. Prevalence of axis I disorders in an AIDS cohort: a cross-sectional, controlled study. Compr Psychiatry 1997;38:146—154.
- 141. Breitbart W, McDonald M, Rosenfeld B, Monkman N, Passik S. Fatigue in ambulatory AIDS patients. J Pain Symptom Manage 1998;15:159–167.
- 142. Kimball L, McCormick W. The pharmacologic management of pain and discomfort in persons with AIDS near the end of life: use of opioid analgesia in the hospice setting. J Pain Symptom Manage 1996;11:88–94.
- 143. Norval D. Symptoms and sites of pain experienced by AIDS patients. S Afr Med J 2004;94: 450–454.
- 144. Schreiner R, McCormick W. Challenges in pain management among persons with AIDS in a long-term-care facility. J Am Med Dir Assoc 2002;3: 51–56.
- **145.** Vincent I, Herouville D, Moulin P, et al. Modalities of palliative care in hospitalized patients with advanced AIDS. AIDS Care 2000;12:211–219.
- 146. Lapane K, Fernandez H, Friedman J, SAGE Study Group. Prevalence, clinical characteristics, and pharmacologic treatment of Parkinson's disease in residents in long-term care facilities. Pharmacotherapy 1999;19:1321–1327.
- 147. Goetz C, Tanner C, Levy M, Wilson R, Garron D. Pain in Parkinson's disease. Mov Disord 1986;1:45–49.
- 148. Quittenbaum B, Grahn B. Quality of life and pain in Parkinson's disease: a controlled cross-sectional study. Parkinsonism Relat Disord 2004; 10:129–136.

- 149. Schulman L, Taback R, Rabinstein A, Weiner W. Non-recognition of depression and other non-motor symptoms in Parkinson's disease. Parkinsonism Relat Disord 2002;6:193—197.
- 150. Starkstein S, Robinson R, Leiguarda R, Preziosi T. Anxiety and depression in Parkinson's disease. Behav Neurol 1993;6:151—154.
- **151.** Menza M, Robertson-Hoffman D, Bonapace A. Parkinson's disease and anxiety: comorbidity with depression. Biol Psychiatry 1993;34:465–470.
- 152. Goy E, Carter J, Ganzini L. Neurologic disease at the end of life: caregiver descriptions of Parkinson disease and amyotrophic lateral sclerosis. J Palliat Med 2008;11:548–554.
- **153.** Kristjanson L, Aoun S, Oldham L. Palliative care and support for people with neurodegenerative conditions and their carers. Int J Palliat Nurs 2005;12:368–377.

- 154. Lee M, Prentice W, Hildreth A, Walker R. Measuring symptom load in idiopathic Parkinson's disease. Parkinsonism Relat Disord 2007;13: 284–289.
- **155.** Oliver D. The quality of care and symptom control—the effects on the terminal phase of ALS/MNS. J Neurol Sci 1996;139:134–136.
- 156. Higginson I, Hart S, Silber E, Burman R, Edmonds P. Symptom prevalence and severity in people severely affected by multiple sclerosis. I Palliat Care 2006;22:158–165.
- **157.** Selman L, Harding R, Gysels M, Speck P, Higginson IJ. The measurement of spirituality in palliative care and the content of tools validated cross-culturally: a systematic review. J Pain Symptom Manage 2011;41:728–753.
- 158. Addington-Hall JM, Fakhoury W, McCarthy M. Specialist palliative care in non-malignant disease. Palliat Med 1998;12:417—427.