

# Curved Yang-Mills gauge theories

based on my preprint [arXiv:2210.02924](https://arxiv.org/abs/2210.02924)

Simon-Raphael Fischer


National Center for Theoretical Sciences

30 December 2022

# Table of contents

- 1 Infinitesimal curved Yang-Mills-Higgs gauge theories
  - Motivation and short introduction
- 2 Integration: Ansatz
  - Principal bundles based on Lie group bundle actions
  - Connections as parallel transport
- 3 Connection
  - Basic notions
  - Definitions
  - Gauge transformation
- 4 Curvature
- 5 Conclusion

# Infinitesimal gauge theory



# Guide: Infinitesimal curved Yang-Mills-Higgs gauge theory

Classical formalism	CYMH GT
Lie algebra $\mathfrak{g}$ as $M \times \mathfrak{g}$	Lie algebroid $E \rightarrow N$
$\mathfrak{g}$ -action $\gamma$	Anchor $\rho$ of $E$ & $E$ -connections
Canonical flat connection $\nabla^0$ on $M \times \mathfrak{g}$	General connection $\nabla$ on $E$

# Guide: Infinitesimal curved Yang-Mills-Higgs gauge theory

Classical formalism	CYMH GT
Lie algebra $\mathfrak{g}$ as $M \times \mathfrak{g}$	Lie algebroid $E \rightarrow N$
$\mathfrak{g}$ -action $\gamma$	Anchor $\rho$ of $E$ & $E$ -connections
Canonical flat connection $\nabla^0$ on $M \times \mathfrak{g}$	General connection $\nabla$ on $E$

## Remarks (Why a "curved theory"?)

Usually, the field strength  $F$  is given by (abelian, for simplicity)

$$F := dA = d^{\nabla^0} A.$$

$\leadsto$  We will use a general connection  $\nabla$  instead of  $\nabla^0$ , and  $\nabla$  may not be flat.

	Classical	Curved
Infinitesimal	Lie algebra $\mathfrak{g}$	LAB $\mathcal{G}$
Integrated	Lie group $G$	LGB <sup>1</sup> $\mathcal{G}$

$$G \longrightarrow \mathcal{G}$$

$$M$$


---

<sup>1</sup>LGB = Lie group bundle

## Definition (LGB actions, simplified)

$$\begin{array}{ccc} & \mathcal{G} & \\ & \downarrow & \\ \mathcal{P} & \xrightarrow{\pi} & M \end{array}$$

$\mathcal{P} \xrightarrow{\pi} M$  a fibre bundle. A **right-action of  $\mathcal{G}$  on  $\mathcal{P}$**  is a smooth map  $\mathcal{P} * \mathcal{G} := \pi^* \mathcal{G} = \mathcal{P} \times_M \mathcal{G} \rightarrow \mathcal{P}$ ,  $(p, g) \mapsto p \cdot g$ , satisfying the following properties:

$$\pi(p \cdot g) = \pi(p), \quad (1)$$

$$(p \cdot g) \cdot h = p \cdot (gh), \quad (2)$$

$$p \cdot e_{\pi(p)} = p \quad (3)$$

for all  $p \in \mathcal{P}$  and  $g, h \in \mathcal{G}_{\pi(p)}$ , where  $e_{\pi(p)}$  is the neutral element of  $\mathcal{G}_{\pi(p)}$ .

# Examples

## Example

$\mathcal{G}$  acts canonically on itself:

$$\begin{aligned}\mathcal{G} * \mathcal{G} &\rightarrow \mathcal{G}, \\ (q, h) &\mapsto qh.\end{aligned}$$

## Example (Recovering Lie group action)

- Either by  $M = \{*\}$ .
- Or by  $\mathcal{G} \cong M \times G$ , then also  $\mathcal{P} * \mathcal{G} \cong \mathcal{P} \times G$ , and we can define

$$\begin{aligned}\mathcal{P} \times G &\rightarrow \mathcal{P}, \\ (p, g) &\mapsto p \cdot g := p \cdot (\pi(p), g),\end{aligned}$$

which is equivalent to  $\mathcal{P} * \mathcal{G} \rightarrow \mathcal{P}$ .



# Examples

## Example

$\mathcal{G}$  acts canonically on itself:

$$\begin{aligned}\mathcal{G} * \mathcal{G} &\rightarrow \mathcal{G}, \\ (q, h) &\mapsto qh.\end{aligned}$$

## Example (Recovering Lie group action)

- Either by  $M = \{*\}$ .
- Or by  $\mathcal{G} \cong M \times G$ , then also  $\mathcal{P} * \mathcal{G} \cong \mathcal{P} \times G$ , and we can define

$$\begin{aligned}\mathcal{P} \times G &\rightarrow \mathcal{P}, \\ (p, g) &\mapsto p \cdot g := p \cdot (\pi(p), g),\end{aligned}$$

which is equivalent to  $\mathcal{P} * \mathcal{G} \rightarrow \mathcal{P}$ .

$\Rightarrow$  Think of the "classical" theory as coming from a trivial LGB

# Examples

## Example

$\mathcal{G}$  acts canonically on itself:

$$\begin{aligned}\mathcal{G} * \mathcal{G} &\rightarrow \mathcal{G}, \\ (q, h) &\mapsto qh.\end{aligned}$$

## Example (Recovering Lie group action)

- Either by  $M = \{*\}$ .
- Or by  $\mathcal{G} \cong M \times G$ , then also  $\mathcal{P} * \mathcal{G} \cong \mathcal{P} \times G$ , and we can define

$$\begin{aligned}\mathcal{P} \times G &\rightarrow \mathcal{P}, \\ (p, g) &\mapsto p \cdot g := p \cdot (\pi(p), g),\end{aligned}$$

which is equivalent to  $\mathcal{P} * \mathcal{G} \rightarrow \mathcal{P}$ .

$\Rightarrow$  Think of the "classical" theory as coming from a trivial LGB

## Definition (Principal bundle)

Still a fibre bundle

$$\begin{array}{ccc} G & \longrightarrow & \mathcal{P} \\ & & \downarrow \pi \\ & & M \end{array}$$


but with  $\mathcal{G}$ -action

$$\begin{array}{ccc} \cancel{\mathcal{P} * G} & \longrightarrow & \mathcal{P} \\ \mathcal{P} * \mathcal{G} & & \end{array}$$

simply transitive on fibres of  $\mathcal{P}$ , and "suitable" atlas.

Connections as parallel transport

# Connection on $\mathcal{P}$ : Idea




But:

$$r_g : \mathcal{P}_x \rightarrow \mathcal{P}_x$$

 $\Rightarrow$ 
 $D_p r_g$  only defined on vertical structure

# Connection on $\mathcal{P}$ : Idea




But:

$$r_g : \mathcal{P}_x \rightarrow \mathcal{P}_x$$

$\Rightarrow$

$D_p r_g$  only defined on vertical structure


# Connection on $\mathcal{P}$ : Idea



Use  $\sigma \in \Gamma(\mathcal{G}) : r_\sigma(p) := p \cdot \sigma_x$


# Connection on $\mathcal{P}$ : Revisiting the classical setup

If  $\mathcal{P}$  a typical principal bundle  
 ( $\mathcal{G}$  trivial,  $\sigma \equiv g$  constant),  
 and  $H$  a connection:



# Connection on $\mathcal{P}$ : Revisiting the classical setup

If  $\mathcal{P}$  a typical principal bundle  
 ( $\mathcal{G}$  trivial,  $\sigma \equiv g$  constant),  
 and  $H$  a connection:



## Remarks (Integrated case)


Parallel transport  $\text{PT}_\alpha^{\mathcal{P}}$  in  $\mathcal{P}$ :

$$\text{PT}_\alpha^{\mathcal{P}}(p \cdot g) = \text{PT}_\alpha^{\mathcal{P}}(p) \cdot g$$

where  $\alpha : I \rightarrow M$  is a base path



# Connection on $\mathcal{P}$ : General case




## Remarks (Integrated case)

Ansatz:

$$\text{PT}_\alpha^{\mathcal{P}}(p \cdot g) = \text{PT}_\alpha^{\mathcal{P}}(p) \cdot \text{PT}_\alpha^{\mathcal{G}}(g)$$

# Connection on $\mathcal{P}$ : General case



## Remarks (Integrated case)

Ansatz:

$$\text{PT}_\alpha^{\mathcal{P}}(p \cdot g) = \text{PT}_\alpha^{\mathcal{P}}(p) \cdot \text{PT}_\alpha^{\mathcal{G}}(g)$$

$\Rightarrow$  Introduce connection on  $\mathcal{G}$

# Classical situation: Differential of Lie group action

## Remarks (Lie group $G$ situation with Lie algebra $\mathfrak{g}$ )

In the case of a right  $G$ -action on  $\mathcal{P}$ ,  $\Phi : \mathcal{P} \times G \rightarrow \mathcal{P}$ , we have

$$D_{(p,g)}\Phi(X, Y) = D_p r_g(X) + \overline{(\mu_G)_g(Y)} \Big|_{p \cdot g}$$

for all  $p \in \mathcal{P}$ ,  $g \in G$ ,  $X \in T_p \mathcal{P}$  and  $Y \in T_g G$ , where

- $\bar{\nu}$  denotes the fundamental vector field on  $\mathcal{P}$  of  $\nu \in \mathfrak{g}$ ,
- $\mu_G$  is the Maurer-Cartan form of  $G$ .

## Definition (Fundamental vector fields)

**Fundamental vector fields** defined by

$$\bar{\nu}_p := \left. \frac{d}{dt} \right|_{t=0} (p \cdot e^{t\nu_x})$$

for all  $\nu \in \Gamma(\mathfrak{g})$  and  $p \in \mathcal{P}_x$ , where  $\mathfrak{g}$  is the LAB<sup>a</sup> of  $\mathcal{G}$ .

---

<sup>a</sup>Lie algebra bundle

## Definition (Darboux derivative)

For  $\sigma \in \Gamma(\mathcal{G})$  we define the **Darboux derivative**  $\Delta\sigma \in \Omega^1(M; \mathfrak{g})$

$$\Delta\sigma = \sigma^! \mu_{\mathcal{G}},$$

where  $\mu_{\mathcal{G}}$  is the connection 1-form on  $\mathcal{G}$ .

## Definition (Darboux derivative)

For  $\sigma \in \Gamma(\mathcal{G})$  we define the **Darboux derivative**  $\Delta\sigma \in \Omega^1(M; \mathfrak{g})$

$$\Delta\sigma = \sigma^! \mu_{\mathcal{G}},$$

where  $\mu_{\mathcal{G}}$  is the connection 1-form on  $\mathcal{G}$ .

## Remarks

If  $\mathcal{G}$  a trivial LGB with canonical flat connection, then  $\mu_{\mathcal{G}}$  is the Maurer-Cartan form.

If Lie group additionally a matrix group, then

$$\Delta\sigma = \sigma^{-1} d\sigma.$$

# Proposition (Differential of LGB action $\Phi$ , [S.-R. F.])

We have

$$D_{(p,g)}\Phi(X, Y) = D_p r_\sigma(X) - \overline{(\pi^! \Delta \sigma)|_p(X)} \Big|_{p \cdot g} + \overline{(\mu_{\mathcal{G}})_g(Y)} \Big|_{p \cdot g}$$

for all  $(p, g) \in \mathcal{P}_x \times \mathcal{G}_x$ ,  $(X, Y) \in T_{(p,g)}(\mathcal{P} * \mathcal{G})$ , where  $\sigma$  is any section of  $\mathcal{G}$  with  $\sigma_x = g$ .

## Proposition (Differential of LGB action $\Phi$ , [S.-R. F.])

We have

$$D_{(p,g)}\Phi(X, Y) = D_p r_\sigma(X) - \overline{(\pi^! \Delta \sigma)|_p(X)} \Big|_{p \cdot g} + \overline{(\mu_{\mathcal{G}})_g(Y)} \Big|_{p \cdot g}$$

for all  $(p, g) \in \mathcal{P}_x \times \mathcal{G}_x$ ,  $(X, Y) \in T_{(p,g)}(\mathcal{P} * \mathcal{G})$ , where  $\sigma$  is any section of  $\mathcal{G}$  with  $\sigma_x = g$ .

## Definition (Modified right-pushforward, [S.-R. F.])

$$\mathcal{r}_{g*}(X) := D_p r_\sigma(X) - \overline{(\pi^! \Delta \sigma)|_p(X)} \Big|_{p \cdot g},$$

$$\mathcal{r}_{\sigma*}(X) := \mathcal{r}_{\sigma_x*}(X).$$



# Proposition (Well-defined isomorphism, [S.-R. F.])

We have that

$$\begin{aligned} T\mathcal{P}|_{\mathcal{P}_x} &\rightarrow T\mathcal{P}|_{\mathcal{P}_x}, \\ X &\mapsto r_{g*}(X), \end{aligned}$$

is a well-defined automorphism over  $r_g$ . Similarly,

$$\begin{aligned} T\mathcal{P} &\rightarrow T\mathcal{P}, \\ X &\mapsto r_{\sigma*}(X), \end{aligned}$$

is an automorphism over  $r_\sigma$ .

## Definition (Ehresmann connection, [S.-R. F.]

$H$  a horizontal distribution of  $T\mathcal{P}$  with

$$\tau_{g*}(H_p) = H_{p \cdot g}$$

### Definition (Ehresmann connection, [S.-R. F.])

$H$  a horizontal distribution of  $T\mathcal{P}$  with

$$\tau_{g*}(H_p) = H_{p \cdot g}$$

### Definition (Connection 1-form, [S.-R. F.])

$A \in \Omega^1(\mathcal{P}; \pi^*\mathcal{G})$  with

$$\tau_{\sigma}^! A = \text{Ad}_{\sigma^{-1}} \circ A,$$

$$A(\bar{\nu}) = \pi^* \nu$$

for all  $\sigma \in \Gamma(\mathcal{G})$  and  $\nu \in \Gamma(\mathcal{G})$ .

### Remarks

$$\left( \tau_{\sigma}^! A \right)_p (X) = A_{p\sigma_x} (\tau_{\sigma*}(X)).$$

## Theorem (Equivalence of both definitions, [S.-R. F.]

*There is the usual 1:1 correspondence between both definitions:*

- *Given  $H$ , define  $A$  by*

$$A_p(\bar{\nu}_p + X) := (\pi^* \nu)_p,$$

*where  $X \in H_p$ .*

- *Given  $A$ , define  $H$  by*

$$H_p := \text{Ker}(A_p).$$

Infinitesimal curved Yang-Mills-Higgs gauge theories  
○○

Integration: Ansatz  
○○○○○○○○

**Connection**  
○○○○○○○●

Curvature  
○

Conclusion  
○○

Gauge transformation



# Summary


	Locally	Globally
Curved Yang-Mills	Pre-classical	$\text{ad}(\mathbb{S}^7 \rightarrow \mathbb{S}^4)$ curved

## Remarks (Integrated point of view)

This is probably linked to that an LGB is locally trivial  
 $\leadsto$  LGB action locally equivalent to a Lie group action

# Hope: Structural Lie groupoids

Gauge theory	Structure
Yang-Mills	Lie group $G$
Curved Yang-Mills	Lie group bundle $\mathcal{G}$
Curved Yang-Mills-Higgs	Lie groupoid $\mathcal{G}$ ?



## Remarks

- Richer set of principal bundles, containing LGBs.
- May result into obstruction statements for curved Yang-Mills-Higgs gauge theories.



**Thank you!**