

Integrating curved Yang-Mills gauge theories

Gauge theories related to principal bundles equipped with Lie group bundle actions

Simon-Raphael FISCHER^{*†}

October 6, 2022: First version

May 2, 2025: More examples and classifying setup with simple Lie group bundles

June 12, 2026: Major rewrite (without adding new statements), in particular for shortening the paper

^{*} 2021–2024: National Center for Theoretical Sciences (國家理論科學研究中心),
Mathematics Division, National Taiwan University
No. 1, Sec. 4, Roosevelt Rd., Taipei City 106, Room 407, Cosmology Building, Taiwan
106319 臺北市羅斯福路四段1號 (國立臺灣大學次震宇宙館407室)

[†] 2024–today: Mathematical Institute, Georg-August University Göttingen
Bunsenstr. 3–5, 37073 Göttingen, Germany

simon.fischer@mathematik.uni-goettingen.de

ORCID: 0000-0002-5859-2825

Abstract^{*}

We construct a gauge theory based on principal bundles $\mathcal{P}$ equipped with a right $\mathcal{G}$ -action, where $\mathcal{G}$ is a Lie group bundle instead of a Lie group. Due to the fact that a $\mathcal{G}$ -action acts fibre by fibre, pushforwards of tangent vectors via a right-translation act now only on the vertical structure of $\mathcal{P}$. Thus, we generalize pushforwards using a connection on $\mathcal{G}$ which will modify the pushforward. A horizontal distribution on $\mathcal{P}$ invariant under such a modified pushforward will provide a proper notion of Ehresmann connection. For achieving gauge invariance we impose conditions on the connection 1-form μ on $\mathcal{G}$: μ has to be a multiplicative form, *i.e.* closed w.r.t. a certain simplicial differential δ on $\mathcal{G}$, and the curvature R_μ of μ has to be δ -exact with primitive ζ ; μ will be the generalization of the Maurer-Cartan form of the classical gauge theory, while the δ -exactness of R_μ will generalize the role of the Maurer-Cartan equation. This introduces the notion of multiplicative Yang-Mills connections, a connection which helped classifying singular foliations and (topological) symmetry breaking. For allowing curved connections on $\mathcal{G}$ in the dynamical theory we will need to generalize the typical definition of the curvature/field strength F on $\mathcal{P}$ by adding ζ to F .

Several examples for a gauge theory with a curved μ will be provided, including the inner group bundle of the Hopf fibration $\mathbb{S}^7 \rightarrow \mathbb{S}^4$, and including a classification for whether these theories admit a classical description.

^{*}Abbreviations used in this paper: **LGB(s)** for Lie group bundle(s), **LAB(s)** for Lie algebra bundle(s).

CONTENTS

1	Introduction and summary	1
1.1	Summary	5
1.2	Conclusion	13
1.3	Basic notations and remarks	13
1.4	Assumed background knowledge	15
2	Lie group bundles (LGBs)	15
2.1	Definition	15
2.2	Associated Lie group bundles	17
2.3	Pullbacks of LGBs	18
3	LGB actions, part I	19
3.1	Definition	19
3.2	Examples of LGB actions	21
4	Lie algebra bundles (LABs)	23
4.1	Definition	23
4.2	From LGBs to LABs	25
4.3	Vertical Maurer-Cartan form of LGBs	27
4.4	Exponential map of LGBs	28
5	LGB actions, part II	29
5.1	LGB and LAB representations	29
5.2	Fundamental vector fields	31
5.3	Differential of smooth LGB actions	32
6	Connections and curvature on principal LGB-bundles	34
6.1	Principal bundles with structural LGB	34
6.2	Generalized distributions and connections	43
6.3	Gauge transformations	65
6.4	Generalized curvature/field strength	74
7	Curved Yang-Mills gauge theory	95
7.1	Definition and gauge invariance	95
7.2	Field redefinitions	97
7.3	Examples	99
7.4	Future prospects	102
	List of References	103
	Appendix A Double tangent bundle and its canonical flip map	103

1. INTRODUCTION AND SUMMARY

This paper's research concerns curved Yang-Mills-Higgs gauge theories, originally introduced by Alexei Kotov and Thomas Strobl in [1], where essentially the structural Lie algebra together with its action on the manifold N of values of the Higgs fields is replaced by a general Lie algebroid $E \rightarrow N$:

Definition: Lie algebroid

e.g. [2, reduced definition of §16.1, page 113]

Let $E \rightarrow N$ be a real vector bundle. Then E is a smooth **Lie algebroid** if there is a bundle map $\rho : E \rightarrow TN$, called the **anchor**, and a Lie algebra structure on $\Gamma(E)$ with Lie bracket $[\cdot, \cdot]_E$ satisfying^a

$$[\mu, f\nu]_E = f[\mu, \nu]_E + \mathcal{L}_{\rho(\mu)}(f) \nu$$

for all $f \in C^\infty(N)$ and $\mu, \nu \in \Gamma(E)$, where $\mathcal{L}_{\rho(\mu)}(f)$ is the action of the vector field $\rho(\mu)$ on the function f by derivation.

^aWith $\Gamma(E)$ I denote the space of sections of a vector bundle E , and with TN the tangent bundle of N .

The idea of replacing Lie algebras with Lie algebroids was proposed by Thomas Strobl in [3], with further understanding of the involved gauge transformations in [4]; eventually, this type of infinitesimal gauge theory got summarised and finalised in [1]. My Ph.D. thesis was devoted to this type of infinitesimal gauge theory, attempting to find new (physical) examples and understanding the geometry of this infinitesimal gauge theory; see [5] and [6].

A short summary of this type of generalized infinitesimal gauge theory follows; we have the following ingredients:

- M a spacetime;
- N a smooth manifold, serves as set for the values of the Higgs field $\Xi : M \rightarrow N$;
- $E \rightarrow N$ a Lie algebroid with anchor ρ , replacing the structural Lie algebra $\mathfrak{g}$ and its action $\gamma : \mathfrak{g} \rightarrow \mathfrak{X}(N)$ of the classical formulation, where $\mathfrak{X}$ denotes the set of vector fields in this work;
- a vector bundle connection ∇ on E ;
- a fibre metric κ on E , as a substitute of the ad-invariant scalar product on $\mathfrak{g}$;
- a Riemannian metric g on N , replacing the scalar product on the vector space in which the Higgs field usually has values in and which is invariant under the action of γ , used for the kinetic term of Ξ which is minimally coupled to the field of gauge bosons $A \in \Omega^1(M; \Xi^*E)$ (1-form on M with values in Ξ^*E);
- a 2-form on N with values in E , $\zeta \in \Omega^2(N; E)$, an additional contribution to the field strength of A .

Infinitesimal gauge invariance of the Yang-Mills type functional leads to two **infinitesimal compatibility conditions** and two metric compatibilities to be satisfied between these structures; we will present those compatibility conditions later in this introduction. If the connection on E is flat, the compatibilities imply that the Lie algebroid is locally an action Lie algebroid:

Definition: Action Lie algebroids

e.g. [2, §16.2, Example 5; page 114]

Let $(\mathfrak{g}, [\cdot, \cdot]_{\mathfrak{g}})$ be a Lie algebra equipped with a Lie algebra action $\gamma : \mathfrak{g} \rightarrow \mathfrak{X}(N)$ on a smooth manifold N . A **transformation Lie algebroid** or **action Lie algebroid** is defined as the bundle $E := N \times \mathfrak{g}$ over N with anchor

$$\rho(p, v) := \gamma(v)|_p$$

for $(p, v) \in E$, and Lie bracket

$$[\mu, \nu]_E|_p := [\mu_p, \nu_p]_{\mathfrak{g}} + (\mathcal{L}_{\gamma(\mu(p))}(\nu^a) - \mathcal{L}_{\gamma(\nu(p))}(\mu^a))|_p e_a$$

for all $p \in N$ and $\mu, \nu \in \Gamma(E)$, where one views a section $\mu \in \Gamma(E)$ as a map $\mu : N \rightarrow \mathfrak{g}$, and $(e_a)_a$ is some arbitrary frame of constant sections.

As it is shown in the mentioned references for this type of infinitesimal gauge theory, one gets back to the standard Yang-Mills-Higgs gauge theory if additionally $\zeta \equiv 0$. Thus, the theory represents a covariantized version of gauge theory equipped with an additional 2-form ζ . If ∇ is flat we say in general that we have a **pre-classical** gauge theory, and if additionally $\zeta \equiv 0$ we have a **classical** gauge theory. In fact, we will observe in this work that the integration of ∇ is a generalisation of the Maurer-Cartan form, while ζ sort of reflects the curvature of ∇ . Thus, the theory presented here will represent a gauge theory covariantised w.r.t. the Maurer-Cartan form; see later.

The 2-form ζ is needed to allow non-flat ∇ , because otherwise only flat ∇ could satisfy the compatibility conditions. But ζ is not just an auxiliary map, in [6] I have shown that there is also a class of **field redefinitions**² for the classical formulation of gauge theory. This field redefinition breaks the gauge invariance; so, in order to keep gauge invariance, one needs to add ζ to the field strength, and at the same time one achieves a richer framework for gauge theories. In fact, these field redefinitions are not just any transformation preserving the physics. As pointed out in [6], curved Yang-Mills gauge theory is a reformulation of classical gauge theory in such a way that gauge theory is form-invariant w.r.t. the field redefinitions, similar to how one reformulates Classical Mechanics so that it is form-invariant w.r.t. Galilei or Lorentz transformations. This reformulation precisely introduces those extra terms, ζ and connection 1-forms of ∇ , that is, every classical gauge theory is equivalent to a possibly curved theory with a non-zero ζ contributing to the field strength. Thus, this gauge theory is not just a “simple” generalisation where one simply adds extra terms, and henceforth it is natural to study whether there is a gauge theory where ζ is non-zero and cannot be transformed to zero by the mentioned field redefinitions; see the Tables [Table 1](#), [Table 2](#), and [Table 3](#).

TABLE 1. The idea of covariantisation as for classical mechanics; the classical and covariantised flat theory are equivalent.

Classical theory	Covariantised flat theory	Curved Theory
Vector space V	Trivial vector bundle $M \times V$	Vector bundle $V \rightarrow M$
$\frac{\partial}{\partial x^i}$	Canonical flat connection ∇^0	Vector bundle connection ∇
Coordinate changes may lead to extra terms	Coordinate expressions form-invariant under coordinate changes	



TABLE 2. Similarly, one can covariantise gauge theory w.r.t. the structural Lie group, leading to the notion of curved Yang-Mills (YM) gauge theory.

YM theory	Covariantised YM theory	Curved YM theory
Lie group G	Trivial Lie group bundle $M \times G$	Lie group bundle $G \rightarrow M$
Maurer-Cartan form	Fibre-wise Maurer-Cartan	Multiplicative YM connection
Field redefinitions lead to extra terms in gauge transformations and field strength	Expressions form-invariant under field redefinitions, but curvature transforms non-trivially	



²So far, there is no proper name for them.

TABLE 3. The same can be done for Yang-Mills-Higgs theories, leading to the notion of curved Yang-Mills-Higgs (YMH) theory; see also [7].

YMH theory	Covariantised YMH theory	Curved YMH theory
Lie group G with right-action on N	Action groupoid $N \times G$	Lie groupoid $G \rightrightarrows N$
Maurer-Cartan form	Fibre-wise Maurer-Cartan	Covariant adjustments
Field redefinitions lead to extra terms in gauge transformations and field strength	Expressions form-invariant under field redefinitions, but curvature transforms non-trivially	

In particular, we will study and introduce the notions for curved Yang-Mills gauge theories (Table 2), and we will provide a full family of examples of new gauge theories which cannot be flattened via field redefinitions. For completeness, gauge theories with a structural groupoid, curved Yang-Mills-Higgs theory, were introduced in [7] after this work. In fact, many of the examples provided here were first introduced in [7], and later added to this work here; these examples are a direct consequence of [8], a first application of the kinematics of curved Yang-Mills gauge theories.

Remark 1.1: References for connections on principal LGB-bundles

It is important to mention beforehand that there is another great paper, [9], which also discusses the notion of connections on principal LGB-bundles. When I wrote this paper, I was not aware of this paper so that there will be some shared results which I found independently of the other authors. This will be obvious due to the fact that our approach is different. Both, this and the other paper, lead to the same sense of Ehresmann connection (based on the same idea of the behaviour of the parallel transport), but are explicitly not directly the same formulation. More like equivalent definitions of the same object, two sides of the same coin, so that it should be worth it to read both research works. Hence, our approach will be different with some shared and some different results. [9] sole purpose was in defining connections on principal bundles with Lie group bundle action (with an application in variational calculus), while we will also introduce and discuss Darboux derivatives, Yang-Mills gauge theory, and related concepts in this context, also pinpointing what happens for Yang-Mills-Higgs theories. Furthermore, our field strength on the principal bundle will be different and more general as already pointed out previously; it will consist of the typical curvature measuring the failure of a distribution to be integrable, plus an adjustment term ζ .

In [6] I first focused on Lie algebroids where the anchor map is an isomorphism and, thus, the Lie algebroid is just the tangent bundle of M . Locally, such examples can be excluded: The curvature can be always transformed to zero by the field redefinitions. However, globally, this is not always true: I have shown that the tangent bundle of the seven-dimensional sphere $\mathbb{S}^7$ is an example of a curved gauge theory which cannot be transformed to a flat pre-classical gauge theory, otherwise it would be a Lie group which is not possible.

However, I also studied the other edge case of having Lie algebra bundles (LABs) instead of Lie algebroids: [6] (also in [5]) points out that locally, *i.e.* over a contractible base manifold N , there is always a field redefinition transforming the initial gauge theory to a pre-classical one. Hence, we arrive at a similar situation as for tangent bundles. But globally there are examples like the adjoint bundle of the Hopf fibration $\mathbb{S}^7 \rightarrow \mathbb{S}^4$, where $\mathbb{S}^n$ ($n \in \mathbb{N}$) denotes the n -dimensional sphere.

This is now the starting point of this paper. We want to understand why the adjoint bundle of $\mathbb{S}^7 \rightarrow \mathbb{S}^4$ as a structural LAB gives rise to a gauge theory which cannot be described classically, and where it differs from the classical formalism of gauge theory. To truly understand this, and also

to possibly get new information about the general case regarding Lie algebroids, we will integrate curved Yang-Mills-Higgs theories in the case of LABs in this paper. Using LABs means that there is no coupling to a Higgs field in the usual sense and therefore we will now just speak instead of a curved Yang-Mills gauge theory. It will turn out, that the Hopf fibration is not just a nice coincidence, but a member of a family of canonical examples: The adjoint bundles of simple non-flat principal bundles as structural Lie group bundles.

Due to the length of this paper, we will start outlining the paper's main results in a summary; we will focus on an easy presentation in the introduction, especially we will not always restate our assumptions, and the notation will be simpler than the actual formulation.

1.1. Summary. The gauge theory presented here will be based on principal bundles $\mathcal{P} \xrightarrow{\pi} M$, M a smooth manifold, related to a right action of a Lie group bundle (LGB) $\mathcal{G} \xrightarrow{\pi_{\mathcal{G}}} M$. The most important distinction to classical gauge theory is that the action Φ is a map

$$\begin{aligned} \mathcal{P} * \mathcal{G} &:= \pi^* \mathcal{G} \rightarrow \mathcal{P}, \\ (p, g) &\mapsto p \cdot g, \end{aligned}$$

where $\pi^* \mathcal{G}$ is the pullback LGB of $\mathcal{G}$ along π . Observe that $\mathcal{G}_x$ only acts on $\mathcal{P}_x$ as a Lie group, thus an element $g \in \mathcal{G}_x$ cannot act on all of $\mathcal{P}$, where notations like $\mathcal{G}_x$ denote the fibre of $\mathcal{G}$ over an $x \in M$. This leads to certain difficulties for defining gauge theory which we will resolve in this paper.

However, this paper aims to be accessible for all people knowing the basics of "classical" gauge theory. Thus, we start with an extensive introduction to LGBs, their actions and their infinitesimal analogues, the LABs, in [Section 2](#) to [Section 5](#). These sections introduce and generalize all the notions known in the "classical" Lie group situations in the context of gauge theory, most of which is straight-forward to generalize. We will especially follow [\[10\]](#), but since this and other references are usually introducing these basic notions on (Lie) groupoids, we decided to reintroduce all the needed notions and proofs so that it should not be required for the reader to follow and learn about groupoids because we just need the much simpler situation regarding LGBs. The experienced reader may start directly with [Section 6](#), but there is one partially new result in these preliminary sections: For gauge theory we will need to understand the differential of the $\mathcal{G}$ -action Φ which we will derive in [Proposition 5.16](#) and which will form the fundament for many calculations. For this we will need notations like TM which will denote a tangent bundle with fibre $T_x M$ at $x \in M$; the fibres of other bundles with similar notation are denoted in the same fashion.

Proposition: Differential of LGB action Φ , simplified situation

We have

$$D_{(p,g)} \Phi(X, Y) = D_p r_{\sigma}(X) + \overline{(\mu_{MC})_g}(Y - D_x \sigma(\omega)) \Big|_{p \cdot g}$$

for all $(p, g) \in \mathcal{P} * \mathcal{G}$ and $(X, Y) \in T_{(p,g)}(\mathcal{P} * \mathcal{G})$, where $x := \pi(p)$, σ is any (local) section of $\mathcal{G}$ with $\sigma_x = g$ and r_{σ} denotes the right-translation of σ in $\mathcal{P}$, μ_{MC} is the fibre-wise defined Maurer-Cartan form of the fibres, $\overline{(\mu_{MC})_g}(\dots)$ denotes its generated fundamental vector field on $\mathcal{P}_x$, and ω is an element of $T_x M$ given by

$$\omega := D_p \pi(X) = D_g \pi_{\mathcal{G}}(Y).$$

In [Section 6](#) we will introduce connections and generalized curvatures on principal LGB-bundles: [Section 6.1](#) starts with introducing principal $\mathcal{G}$ -bundles $\mathcal{P}$. Such principal bundles were already introduced in [\[11, §5.7, page 144f.\]](#), but also here we rephrased it in such a way that these bundles' definition has a more familiar shape for readers not so proficient with the study of groupoids. The main difference to Lie group based principal bundles is the previously-mentioned LGB action Φ . Since these bundles were not studied a lot before we will rephrase [\[11\]](#) and introduce basic notions like morphisms and easy examples, for example LGBs $\mathcal{G}$ themselves are principal $\mathcal{G}$ -bundles, so that

the gauge theory presented in this paper can also be understood as a gauge theory based not only on "classical" principal bundles but also on LGBs which were excluded in general in the classical formalism. Furthermore, as in [11], we generalize certain statements known about classical principal bundles, *e.g.* we observe that a section of $\mathcal{P}$ does not trivialize $\mathcal{P}$ in general, however it introduces an isomorphism to $\mathcal{G}$, as will be pointed out in Lemma 6.7.

In Section 6.2 we finally turn to the notion of connections on $\mathcal{P}$, described as horizontal subbundle $H\mathcal{P}$ of the tangent bundle $T\mathcal{P}$, complementary to the vertical bundle $V\mathcal{P}$, and equivalently we want to describe $H\mathcal{P}$ as a connection 1-form A . As one can already see in the previous proposition about the derivative of the action Φ , due to the fact that right-translations r_g of an element $g \in \mathcal{G}_x$ now only acts on $\mathcal{P}_x$, its derivative Dr_g only acts on the tangent bundle of $\mathcal{P}_x$, the vertical tangent vectors of $\mathcal{P}$ (over x). Hence, in order to define the pushforward of non-vertical vectors we make use of auxiliary sections. However, this leads to the problem that there are many different sections with the same value over a fixed base point $x \in M$, all of whose pushforwards of non-vertical vectors are in general different. To get rid of the ambiguity in the choice of section we are going to modify and generalize the pushforward of tangent vectors by right-translations. To do so, we fix a horizontal distribution $H\mathcal{G}$ on $\mathcal{G}$, without further assumptions than the bare-bones for horizontal distributions. The **(modified) right-pushforward** of $X \in T_p\mathcal{P}$ ($p \in \mathcal{P}_x$) by $g \in \mathcal{G}_x$ is then defined as

$$\boldsymbol{r}_{g*}(X) := D_{(p,g)}\Phi(X, Y),$$

where $Y \in H_g\mathcal{G}$ is the unique horizontal vector at g lifting $D_p\pi(X)$.

Then we start to construct the horizontal distribution $H\mathcal{P}$ initially by its associated parallel transport, shortly denoted now by $PT^{\mathcal{P}}$; similarly we denote the parallel transport associated to $H\mathcal{G}$ by $PT^{\mathcal{G}}$. We define an Ehresmann connection on $\mathcal{P}$ by

$$PT^{\mathcal{P}}(p \cdot g) = PT^{\mathcal{P}}(p) \cdot PT^{\mathcal{G}}(g)$$

for all $(p, g) \in \mathcal{P} * \mathcal{G}$. However, in order to define the connection 1-form it is useful to find a definition of such $H\mathcal{P}$ via a symmetry w.r.t. a certain action on $T\mathcal{P}$: As already high-lighted, this will be a modification of the pushforward via right-translations. Differentiating the condition about parallel transports w.r.t. the curve parameter achieves an alternative version of this modification, summarised in Proposition 6.40; we will explain the second summand of the following proposition afterwards.

Proposition: The modified right-pushforward

For $g \in \mathcal{G}_x$ ($x \in M$) we have

$$T\mathcal{P}|_{\mathcal{P}_x} \rightarrow T\mathcal{P}|_{\mathcal{P}_x},$$

$$X \mapsto \boldsymbol{r}_{g*}(X) = D_p r_{\sigma}(X) - \overline{(\pi^! \Delta \sigma)|_p(X)} \Big|_{p \cdot g},$$

where $p \in \mathcal{P}_x$, $X \in T_p\mathcal{P}$, $\pi^!$ denotes the pull-back of forms with π , and σ is any (local) section of $\mathcal{G}$ with $\sigma_x = g$. $\boldsymbol{r}_{g*}$ is independent of the choice of the local section σ by definition, and it is a vector bundle automorphism over the right-translation r_g .

Furthermore, if we have a horizontal distribution $H\mathcal{P}$ on $\mathcal{P}$, then $H_p\mathcal{P}$ is isomorphic via $\boldsymbol{r}_{g*}$ to a complement of $V_{p \cdot g}\mathcal{P}$ in $T_{p \cdot g}\mathcal{P}$ (this complement is not necessarily $H_{p \cdot g}\mathcal{P}$).

This can be generalized to sections σ of $\mathcal{G}$ in a straight-forward manner, giving rise to an automorphism $\boldsymbol{r}_{\sigma*}$ of $T\mathcal{P}$ over the right-translation r_{σ} . $\Delta\sigma$ is the generalised version of the Darboux derivative, often simply denoted as $\sigma^{-1}d\sigma$ in the classical formalism, but in some works like [10, §5.1, page 182ff.] also already written as $\Delta\sigma$ in the classical formalism. As expected, this derivative plays an important role in gauge theory, which is why we will also discuss and introduce the Darboux derivative; usually it only appears in the gauge transformations, but in our case it will already appear now. To no surprise, one can introduce the connection 1-form $\mu_{\mathcal{G}}$ on $\mathcal{G}$ corresponding to $H\mathcal{G}$

with values in the LAB $\pi_{\mathcal{G}}^* \mathcal{G}$ of $\mathcal{G}$ by defining it as μ_{MC} contracted by the projection to the vertical part. The Darboux derivative is then the form-pullback of $\mu_{\mathcal{G}}$ (Definition 6.24) by a section.

Definition: Darboux derivative

For (local) $\sigma \in \Gamma(\mathcal{G})$ we define the **Darboux derivative** $\Delta\sigma \in \Omega^1(M; \mathcal{G})$ by

$$\Delta\sigma = \sigma^! \mu_{\mathcal{G}}.$$

Using the Darboux derivative, we defined the modified pushforward via right-translations (also called modified right-pushforward) which eventually leads to the definition of an Ehresmann connection on the principal $\mathcal{G}$ -bundle $\mathcal{P}$, as pinpointed in Definition 6.44.

Definition: Ehresmann connection on principal LGB-bundles

We call $H\mathcal{P}$ an **Ehresmann connection** or a **connection on $\mathcal{P}$** if it is **right-invariant (w.r.t. modified right-pushforward)**, i.e.

$$\mathcal{R}_{g*}(\mathcal{H}_p \mathcal{P}) = \mathcal{H}_{p \cdot g} \mathcal{P}$$

for all $p \in \mathcal{P}_x$ and $g \in \mathcal{G}_x$ ($x := \pi(p)$).

$H\mathcal{P}$ will now always be an Ehresmann connection in the following, and one may also say that $H\mathcal{P}$ is an Ehresmann connection modified by $H\mathcal{G}$. The basic idea now is to replace the typical right-pushforward with the modified one in all the involved definitions which is straight-forward. Definition 6.50 introduces connection 1-forms on principal bundles:

Definition: Connection 1-forms on principal LGB-bundles, simplified notation

A **connection 1-form** or **gauge field** on $\mathcal{P}$ is a 1-form $A \in \Omega^1(\mathcal{P}; \pi^* \mathcal{G})$ satisfying:

- (**$\mathcal{G}$ -equivariance, but w.r.t. modified right-pushforward**)

$$\mathcal{R}_{\sigma}^! A = \text{Ad}_{\sigma^{-1}} \circ A$$

for all (local) $\sigma \in \Gamma(\mathcal{G})$, where Ad is the adjoint representation of $\mathcal{G}$ on $\mathcal{G}$.

- (**Identity on $V\mathcal{P}$**)

$$A(\tilde{\nu}) = \pi^* \nu$$

for all (local) $\nu \in \Gamma(\mathcal{G})$, where notations like π^* denote the pull-back of sections.

Of course we will achieve a 1:1 correspondence to Ehresmann connections in Theorem 6.52.

Theorem: 1:1 correspondence of Ehresmann connections and connection 1-forms

There is a 1:1 correspondence between Ehresmann connections and connection 1-forms on $\mathcal{P}$:

- *Let $H\mathcal{P}$ be an Ehresmann connection on $\mathcal{P}$. Then $H\mathcal{P}$ defines a connection 1-form $A \in \Omega^1(\mathcal{P}; \pi^* \mathcal{G})$ by*

$$A_p(\tilde{v}_p + X_p) = (p, v)$$

for all $p \in \mathcal{P}_x$ ($x \in M$), $v \in \mathcal{G}_x$ and $X \in \mathcal{H}_p \mathcal{P}$.

- *Let $A \in \Omega^1(\mathcal{P}; \pi^* \mathcal{G})$ be a connection 1-form on $\mathcal{P}$. Then A defines an Ehresmann connection $H\mathcal{P}$ on $\mathcal{P}$ via its kernel $\text{Ker}(A)$, that is,*

$$\mathcal{H}_p \mathcal{P} = \text{Ker}(A_p)$$

for all $p \in \mathcal{P}$.

Not only these results are straight-forward once one has the definition of the modified right-pushforward, also other results will be simple to guess. For example in Section 6.3 we turn to the

gauge transformations which will have a familiar shape but with the generalized Darboux derivative appearing, see Proposition 6.58.

Proposition: Gauge transformations of connection 1-forms, simplified notation

Let H be a (base-preserving) automorphism of $\mathcal{P}$. We then have that $H^!A$ is a connection 1-form on $\mathcal{P}$ and

$$H^!A = \text{Ad}_{(\sigma^H)^{-1}} \circ A + (\pi^*\Delta)\sigma^H,$$

where $\sigma^H \in \Gamma(\pi^*\mathcal{G})$ is uniquely defined by

$$H(p) = p \cdot \sigma_p^H$$

for all $p \in \mathcal{P}$, and $\pi^*\Delta$ is the Darboux derivative on $\pi^*\mathcal{G}$ naturally inherited by the pullback of the connection on $\mathcal{G}$.

The 1:1 correspondence between H and σ^H arising here leads to similar statements as in the classical gauge theory which we will also shortly point out. Afterwards we will show in Proposition 6.64 that this theorem about gauge transformations implies a pullback version, that is, one looks at the pullback $A_{s_i} := s_i^!A$ of A w.r.t. a section (also called gauge) s_i of $\mathcal{P}$ defined over an open subset $U_i \subset M$, and then the change of A by changing the gauge s_i to another gauge follows by the previous theorem.

Proposition: Gauge transformations as a change of gauge in the local gauge field

Let U_i and U_j be two open subsets of M so that $U_i \cap U_j \neq \emptyset$, two gauges $s_i \in \Gamma(\mathcal{P}|_{U_i})$ and $s_j \in \Gamma(\mathcal{P}|_{U_j})$, and the unique $\sigma_{ji} \in \Gamma(\mathcal{G}|_{U_i \cap U_j})$ with $s_i = s_j \cdot \sigma_{ji}$ on $U_i \cap U_j$. Then we have over $U_i \cap U_j$ that

$$A_{s_i} = \text{Ad}_{\sigma_{ji}^{-1}} \circ A_{s_j} + \Delta\sigma_{ji}.$$

This will finish the discussion about A , but of course we also need its field strength F and its gauge transformations. Up until now $H\mathcal{G}$ was an arbitrary horizontal distribution, but we will need to fix certain conditions on it to assure gauge invariance later. We will discuss this in Section 6.4. On one hand, viewing $\mathcal{G}$ as the principal $\mathcal{G}$ -bundle $\mathcal{P}$ itself whose horizontal distribution $H\mathcal{P}$ aligns with $H\mathcal{G}$, it is natural to guess that $H\mathcal{G}$ should be an Ehresmann connection itself. Ehresmann connections on LGBs modifying themselves were already discussed in works like [12, §4.4, implication of Lemma 4.14]; there is also [13] related to a similar subject. One of the results of these works is that such Ehresmann connections on LGBs are characterized by the fact that the connection 1-form $\mu_{\mathcal{G}}$ on $\mathcal{G}$ has to be a multiplicative form:

Definition: Multiplicative forms, simplified situation

[14, §2.1, special situation of Def. 2.1]

We call an $\omega \in \Omega^1(\mathcal{G}; \pi_{\mathcal{G}}^*\mathcal{G})$ a **multiplicative form** if

$$\omega_{gq}(\text{D}_{(g,q)}\Phi(X, Y)) = \text{Ad}_{q^{-1}}(\omega_g(X)) + \omega_q(Y)$$

for all $(g, q) \in \mathcal{G} * \mathcal{G}$ and $(X, Y) \in \text{T}_{(g,q)}(\mathcal{G} * \mathcal{G})$, where $\Phi: \mathcal{G} * \mathcal{G} \rightarrow \mathcal{G}$ is the multiplication in $\mathcal{G}$.

Assuming that $H\mathcal{G}$ is multiplicative will also allow us to formulate certain technical identities for the Darboux derivative like a Leibniz rule. In order to calculate the gauge transformation for the curvature/field strength F of A we need to understand the curvature of $\mu_{\mathcal{G}}$. Classically, this is described by the Maurer-Cartan equation, however, we generalize this condition, allowing non-flat

connections on $\mathcal{G}$. For this we will study a connection on the LAB $\mathcal{Q}$ of $\mathcal{G}$, naturally induced by $H\mathcal{G}$; we will construct this connection in [Proposition 6.31](#) and [Definition 6.33](#), and our construction aligns with [12, §4.5, Prop. 4.22] even though we argue the following differently since we will show the following statement actually for all horizontal distributions $H\mathcal{G}$ instead of just Ehresmann connections.

Proposition: LGB connection induces LAB connection, simplified notation

The map $\nabla^{\mathcal{G}}: \Gamma(\mathcal{Q}) \rightarrow \Omega^1(M; \mathcal{Q})$, $\nu \mapsto \nabla^{\mathcal{G}}\nu$ denoted as an element of $\Omega^1(M; \mathcal{Q})$ by $X \mapsto \nabla_X^{\mathcal{G}}\nu$, defined by

$$\nabla_X^{\mathcal{G}}\nu|_x := \left. \frac{d}{dt} \right|_{t=0} \left((\Delta e^{t\nu})_x(X) \right)$$

for all $x \in M$, $X \in T_x M$ and $\nu \in \Gamma(\mathcal{Q})$, is a vector bundle connection on $\mathcal{Q}$, where $t \in \mathbb{R}$. We will call $\nabla^{\mathcal{G}}$ the **$\mathcal{G}$ -connection (on its LAB $\mathcal{Q}$)**.

In fact, $\nabla^{\mathcal{G}}$ will play the role of ∇ from the beginning of this introduction. Furthermore, this connection will allow us to define the generalized Maurer-Cartan equation in [Theorem 6.81](#) and [Definition 6.85](#).

Definition: Multiplicative Yang-Mills connection, simplified notation

We say that $H\mathcal{G}$, and thus $\mu_{\mathcal{G}}$, is a **multiplicative Yang-Mills connection (w.r.t. a $\zeta \in \Omega^2(M; \mathcal{Q})$)**, if it satisfies the **compatibility conditions**:

- (1) $\mu_{\mathcal{G}}$ is multiplicative (i.e. $H\mathcal{G}$ is an Ehresmann connection modifying itself),
- (2) $\mu_{\mathcal{G}}$ satisfies the **generalized Maurer-Cartan equation**, that is,

$$\left(d^{\pi_{\mathcal{G}}^* \nabla^{\mathcal{G}}} \mu_{\mathcal{G}} + \frac{1}{2} [\mu_{\mathcal{G}} \hat{\wedge} \mu_{\mathcal{G}}]_{\pi_{\mathcal{G}}^* \mathcal{Q}} \right) \Big|_g = \text{Ad}_{g^{-1}} \circ \pi_{\mathcal{G}}^! \zeta|_g - \pi_{\mathcal{G}}^! \zeta|_g$$

for all $g \in \mathcal{G}_x$ ($x \in M$), where $[\cdot, \cdot]_{\pi_{\mathcal{G}}^* \mathcal{Q}}$ is the Lie bracket of the pullback-LAB $\pi_{\mathcal{G}}^* \mathcal{Q}$, and the $\hat{\wedge}$ denotes the typical graded extension of tensors, that is, the second summand on the left hand side is an element of $\Omega^2(\mathcal{G}; \pi_{\mathcal{G}}^* \mathcal{Q})$ given here by

$$\left(\frac{1}{2} [\mu_{\mathcal{G}} \hat{\wedge} \mu_{\mathcal{G}}]_{\pi_{\mathcal{G}}^* \mathcal{Q}} \right) (X, Y) = [\mu_{\mathcal{G}}(X), \mu_{\mathcal{G}}(Y)]_{\pi_{\mathcal{G}}^* \mathcal{Q}}$$

for all $X, Y \in \mathfrak{X}(\mathcal{G})$.

Making a pull-back of the generalized Maurer-Cartan equation w.r.t. a section $\sigma \in \Gamma(\mathcal{G})$ is straight-forward and will be provided in [Corollary 6.84](#).

Corollary: Pullback of generalized Maurer-Cartan equation

Let $H\mathcal{G}$ be a multiplicative Yang-Mills connection w.r.t. a $\zeta \in \Omega^2(M; \mathcal{Q})$. Then

$$d^{\nabla^{\mathcal{G}}} \Delta \sigma + \frac{1}{2} [\Delta \sigma \hat{\wedge} \Delta \sigma]_{\mathcal{Q}} + \zeta = \text{Ad}_{\sigma^{-1}} \circ \zeta$$

for all $\sigma \in \Gamma(\mathcal{G})$.

We will now argue that those compatibility conditions are the integrals of the infinitesimal compatibility conditions proposed by Alexei Kotov and Thomas Strobl, mentioned earlier (without details). In [12] it was already observed that an Ehresmann connection on an LGB naturally implies that $\nabla^{\mathcal{G}}$ is a Lie bracket derivation.

Lemma: Ehresmann connections induce Lie bracket derivations

[12, §4.5, Prop. 4.21]

If $H\mathcal{G}$ is a multiplicative connection on $\mathcal{G}$, then

$$\nabla^{\mathcal{G}}([\mu, \nu]_{\mathcal{G}}) = [\nabla^{\mathcal{G}}\mu, \nu]_{\mathcal{G}} + [\mu, \nabla^{\mathcal{G}}\nu]_{\mathcal{G}}$$

for all $\mu, \nu \in \Gamma(\mathcal{G})$.

That $\nabla^{\mathcal{G}}$ is a Lie bracket derivation is one of the infinitesimal compatibility conditions. Furthermore, we will show in [Theorem 6.81](#) that the generalized Maurer-Cartan equation is actually equivalent to its infinitesimal version.

Theorem: Generalized Maurer-Cartan equation, changed formulation

Let $H\mathcal{G}$ be a multiplicative connection on $\mathcal{G}$ with connected fibres, and $\zeta \in \Omega^2(M; \mathcal{G})$. Then $H\mathcal{G}$ satisfies the generalized Maurer-Cartan equation w.r.t. ζ if and only if

$$R_{\nabla^{\mathcal{G}}}(X, Y)\mu = [\zeta(X, Y), \mu]_{\mathcal{G}}$$

for all $X, Y \in \mathfrak{X}(M)$ and $\mu \in \Gamma(\mathcal{G})$, where $R_{\nabla^{\mathcal{G}}}$ is the curvature of $\nabla^{\mathcal{G}}$.

The condition about the curvature is in fact the second infinitesimal compatibility condition proposed by Alexei Kotov and Thomas Strobl, so that we can conclude that Yang-Mills connections on LGBs serve as an integral of the infinitesimal compatibility conditions which we will point out in [Definition 6.67](#) and [Remark 6.86](#).

Theorem: The infinitesimal conditions of multiplicative Yang-Mills connections

Every multiplicative Yang-Mills connection (w.r.t. $\zeta \in \Omega^2(M; \mathcal{G})$) $H\mathcal{G}$ is an **infinitesimal multiplicative Yang-Mills connection (on $\mathcal{G}$)**, that is, it satisfies the **infinitesimal compatibility conditions**

$$\nabla^{\mathcal{G}}([\mu, \nu]_{\mathcal{G}}) = [\nabla^{\mathcal{G}}\mu, \nu]_{\mathcal{G}} + [\mu, \nabla^{\mathcal{G}}\nu]_{\mathcal{G}},$$

$$R_{\nabla^{\mathcal{G}}}(X, Y)\mu = [\zeta(X, Y), \mu]_{\mathcal{G}}$$

for all $\mu, \nu \in \Gamma(\mathcal{G})$ and $X, Y \in \mathfrak{X}(M)$.

Additionally we will mention in [Remark 6.87](#) that multiplicativity of $\mu_{\mathcal{G}}$ is in fact a closedness condition w.r.t. a certain simplicial differential δ , introduced in [15, beginning of §1.2]. The generalized Maurer-Cartan equation is then an exactness-condition of the "classical" curvature of $\mu_{\mathcal{G}}$, its δ -primitive given by ζ . This is in alignment with the infinitesimal compatibility conditions which are also equivalent to statements about closedness and exactness w.r.t. the Chevalley-Eilenberg complex.

This will finish the discussion about the connection on $\mathcal{G}$, and it will be important for defining the field strength which will be discussed afterwards, starting with [Definition 6.91](#). For this we will always assume now that $H\mathcal{G}$ is a multiplicative Yang-Mills connection w.r.t. a $\zeta \in \Omega^2(M; \mathcal{G})$ (while $H\mathcal{P}$ is an Ehresmann connection modified by such $H\mathcal{G}$).

Definition: (Generalized) Field strength

We define the **(generalized) curvature** or **(generalized) field strength F (of A)** as an element of $\Omega^2(\mathcal{P}; \pi^*\mathcal{G})$ by

$$F := d\pi^*\nabla^{\mathcal{G}} A \circ (\pi^{H\mathcal{P}}, \pi^{H\mathcal{P}}) + \pi^!\zeta,$$

where $d^{\pi^* \nabla^{\mathcal{G}}}$ is the exterior covariant derivative related to the pullback connection $\pi^* \nabla^{\mathcal{G}}$, and $\pi^{\text{H}\mathcal{P}}: T\mathcal{P} \rightarrow \text{H}\mathcal{P}$ is the canonical projection onto the associated Ehresmann connection $\text{H}\mathcal{P}$ on $\mathcal{P}$; that is,

$$F(X, Y) = d^{\pi^* \nabla^{\mathcal{G}}} A(\pi^{\text{H}\mathcal{P}}(X), \pi^{\text{H}\mathcal{P}}(Y)) + (\pi^* \zeta)(D\pi(X), D\pi(Y))$$

for all $X, Y \in \mathfrak{X}(\mathcal{P})$.

By usual arguments, the first summand measures the failure of $\text{H}\mathcal{P}$ being a regular foliation; that is, the gauge theory we will present here not only modifies Ehresmann connections but the modification's curvature additionally modifies the curvature. To avoid confusion with the ordinary curvature (the first summand) one may now need to distinguish between curvatures and field strengths.

With a lengthy calculation evolving around that $\text{H}\mathcal{G}$ is a multiplicative Yang-Mills connection, especially using the generalized Maurer-Cartan equation, we will show in [Proposition 6.92](#) that F transforms in a suitable way when making a pull-back w.r.t. the modified right-pushforward, which is the last step needed for defining the physical theory and its associated Lagrangian.

Proposition: Properties of the generalized field strength, simplified notation

We have the following properties of the field strength:

- **(Form of type Ad)**

$$\tau_\sigma^! F = \text{Ad}_{\sigma^{-1}} \circ F$$

for all (local) $\sigma \in \Gamma(\mathcal{G})$.

- **(Horizontal form)**

For $X, Y \in T_p \mathcal{P}$ ($p \in \mathcal{P}$) we have

$$F(X, Y) = 0$$

if either of X and Y is vertical.

We will also derive a structure equation in [Proposition 6.93](#).

Proposition: Structure equation of the generalized field strength

We have the **structure equation**

$$F = d^{\pi^* \nabla^{\mathcal{G}}} A + \frac{1}{2} [A \wedge A]_{\pi^* \mathcal{G}} + \pi^! \zeta.$$

We will argue that ζ of course also affects the Bianchi identity, which we have in fact already derived in earlier works.

Lemma: Generalized Bianchi identity

[5, §7, Thm. 7.3] & [6, §5, Thm. 5.1.42]

We have the **(generalized) Bianchi identity**

$$d^{\pi^* \nabla^{\mathcal{G}}} F + [A \wedge F]_{\pi^* \mathcal{G}} = \pi^! d^{\nabla^{\mathcal{G}}} \zeta.$$

Similarly to the discussion about connections, [Section 6.4](#) will be concluded with a discussion about the gauge transformation of F in [Proposition 6.96](#); as for A the gauge transformations of F are given by form-pullbacks with a (base-preserving) principal bundle automorphism H with associated unique $\sigma^H \in \Gamma(\pi^* \mathcal{G})$ given by $H(p) = p \cdot \sigma_p^H$ for all $p \in \mathcal{P}$.

Proposition: Gauge transformation of the field strength, simplified notation

We have that $H^1 F$ is the field strength related to $H^1 A$ and

$$H^1 F = \text{Ad}_{(\sigma^H)^{-1}} \circ F.$$

Again as for A , we are interested about what this formula for the gauge transformation implies for pull-backs $F_s := s^! F$ w.r.t. a gauge s of $\mathcal{P}$ when changing the choice of s . This is straight-forward to calculate and will be stated in [Proposition 6.99](#).

Proposition: Gauge transformations again as a change of gauge

Let U_i and U_j be two open subsets of M so that $U_i \cap U_j \neq \emptyset$, two gauges $s_i \in \Gamma(\mathcal{P}|_{U_i})$ and $s_j \in \Gamma(\mathcal{P}|_{U_j})$, and the unique $\sigma_{ji} \in \Gamma(\mathcal{G}|_{U_i \cap U_j})$ with $s_i = s_j \cdot \sigma_{ji}$ on $U_i \cap U_j$. Then we have for the field strength of A over $U_i \cap U_j$ that

$$F_{s_i} = \text{Ad}_{\sigma_{ji}^{-1}} \circ F_{s_j}.$$

Eventually in [Section 7](#) we will start to introduce the physical theory, that is, first of all defining the Lagrangian of a gauge theory based on principal LGB-bundles in [Section 7.1](#). Due to the fact that we will have pointed out that this theory integrates the infinitesimal curved Yang-Mills-Higgs gauge theory developed by Alexei Kotov and Thomas Strobl (in the case of LABs), we are going to label this theory **curved Yang-Mills gauge theory** in [Definition 7.2](#). To do so, M is now a spacetime, so that we can define the Hodge star operator $*$, and we assume that we have an Ad-invariant fibre metric κ on $\mathcal{G}$.

Definition: Curved Yang-Mills gauge theory

Let $(U_i)_i$ be an open covering of M so that there are subordinate gauges $s_i \in \Gamma(\mathcal{P}|_{U_i})$. Then the top-degree form $\mathfrak{L}_{\text{CYM}}[A] \in \Omega^{\dim(M)}(M; \mathbb{R})$, defined locally by

$$(\mathfrak{L}_{\text{CYM}}[A])|_{U_i} := -\frac{1}{2} \kappa(F_{s_i} \wedge *F_{s_i}),$$

is called the **curved Yang-Mills Lagrangian**.

We will have argued in [Corollary 7.1](#) that this Lagrangian is well-defined; a trivial consequence of the previously-highlighted gauge transformations. Similarly, we will achieve gauge invariance in [Theorem 7.3](#).

Theorem: Gauge invariance of the curved Yang-Mills Lagrangian

We have

$$\mathfrak{L}_{\text{CYM}}[H^1 A] = \mathfrak{L}_{\text{CYM}}[A]$$

for all principal bundle automorphisms H .

Finally we will discuss new examples of gauge theory in [Section 7.3](#), after pointing out that this gauge theory generalizes the classical formalism. We will also discuss this matter in the context of my previously-mentioned Ph.D. thesis, [6], and the field redefinitions. Especially, the paper concludes with the reason why this project started: The inner group bundle of the Hopf fibration $\mathbb{S}^7 \rightarrow \mathbb{S}^4$ is one of many examples of curved Yang-Mills gauge theory, so that $\nabla^{\mathcal{G}}$ is not curved; we will present this in [Example 7.14](#).

Example: How to cook new gauge theories

First an explicit example: Let P be the Hopf bundle

$$\begin{array}{ccc} \mathrm{SU}(2) \cong \mathbb{S}^3 & \longrightarrow & \mathbb{S}^7 \\ & & \downarrow \\ & & \mathbb{S}^4 \end{array}$$

We define the principal bundle $\mathcal{P}$ and LGB $\mathcal{G}$ as the inner group bundle of P

$$\mathcal{P} := \mathcal{G} := c_{\mathrm{SU}(2)}(P) := (P \times G) / G,$$

where $(P \times G)/G$ is the associated Lie group bundle, where the action of G on itself is given by its conjugation, $c_g(q) = gqg^{-1}$ for all $g, q \in G$. This principal $c_{\mathrm{SU}(2)}(P)$ -bundle admits the structure as curved Yang-Mills gauge theory, which is neither classical nor pre-classical. Furthermore, all possible (curved) Yang-Mills gauge theory structures lead to a curved multiplicative Yang-Mills connection and non-zero ζ , which also implies that this gauge theory cannot be described with the typical formalism of gauge theory.

Similarly, using non-flat principal H -bundles P for a Lie group H with no centre leads to curved multiplicative Yang-Mills connections on $c_H(P)$ given by associated connections; those cannot be flattened by field redefinitions. Thus, one quickly achieves curved Yang-Mills gauge theories on other principal $c_H(P)$ -bundles like $c_H(P)$ itself, but also P or the Atiyah groupoid of P , both of which can also be viewed as principal $c_H(P)$ -bundles.

The example section will also provide a neat geometric interpretation using the notion of singular foliations; see Table 5.

1.2. Conclusion. Since a curved Yang-Mills connection which cannot be flattened by redefinitions only exists globally, one might say that this theory may be just a toy model for physics, but thinking of symplectic geometry, the global aspects may be at least interesting in mathematics; as already mentioned, these connections helped classifying singular foliations and topological symmetry breaking ([8, 16]). Alternatively, gauge theory is also a theory about connections, so one could investigate what the sense of connection and curvature given here implies for the plethora of other theories based on connections. More applications will follow, pointing out the importance of such constructions like the generalised Maurer-Cartan equation.

1.3. Basic notations and remarks.

- The appendix serves for providing extra information or background knowledge which did not fit in the flow of this paper's text. We will sometimes refer to the appendix, but the experienced reader may be able to ignore the appendix.
- In the main text we usually repeat and reintroduce needed objects for the statements (like "**Let M be a manifold [...]**" in every statement) (almost) allowing to just read the statements without having read the text introducing it, while the appendix is written as a continuous text which has to be read as a whole in order to understand the essential statements.
- As usual, there will be definitions of certain objects depending on other elements, and for keeping notations simple we will not always explicitly denote all dependencies. It will be clear by context on which it is based on, that is for example, if we define an object A using the notion of Lie algebra actions γ and we write "Let X be an object A ", then it will be clear by context which Lie algebra action is going to be used, for example given in a previous sentence writing "Let γ be a Lie algebra action".
- Throughout this work we always use Einstein's sum convention if suitable.
- If not mentioned explicitly, we always assume finite dimensions and morphisms denote base-preserving ones.

- With f^*F we denote the pullback/pull-back of the fibre bundles $F \rightarrow M$ under a smooth map $f : N \rightarrow M$. Similarly we denote the pullbacks of sections of a fibre bundle.
- For $V \rightarrow M$ a vector bundle over M : Do not confuse the pull-back of sections with the pull-back of forms $\omega \in \Omega^l(M; V)$ ($l \in \mathbb{N}_0$), here denoted by $f^!\omega$, which is an element of $\Gamma\left(\left(\bigwedge_{m=1}^l T^*N\right) \otimes f^*V\right) \cong \Omega^l(N; f^*V)$, and not of $\Gamma\left(\left(\bigotimes_{m=1}^l (f^*TM)^*\right) \otimes f^*V\right)$ like $f^*\omega$.
- Let $F \xrightarrow{\pi_F} M$ and $G \xrightarrow{\pi_G} N$ be two fibre bundles over smooth manifolds M and N , respectively, and let $\phi : N \rightarrow M$ be a smooth map. Furthermore, let us assume we have a morphism $\Phi : G \rightarrow F$ of fibre bundles over ϕ , that is, Φ is a smooth map such that the following diagram commutes

$$\begin{array}{ccc} G & \xrightarrow{\Phi} & F \\ \downarrow \pi_G & & \downarrow \pi_F \\ N & \xrightarrow{\phi} & M \end{array}$$

especially, $\pi_F \circ \Phi = \phi \circ \pi_G$. We make often use of that such morphisms have a 1:1 correspondence to **base-preserving** fibre bundle morphisms $\tilde{\Phi} : G \rightarrow \phi^*F$, i.e. $\tilde{\Phi}$ is a smooth map with $\phi^*\pi_F \circ \tilde{\Phi} = \pi_G$. For $p \in N$ the morphism $\tilde{\Phi}$ has the form

$$\tilde{\Phi}_p = (p, \Phi_p),$$

that is,

$$\tilde{\Phi}_p(g) = (p, \Phi_p(g))$$

for all $g \in G_p$, which is well-defined since $\Phi_p(g) \in F_{\phi(p)}$. The map $\tilde{\Phi} \mapsto \Phi := \text{pr}_2 \circ \tilde{\Phi}$ is then a bijective map between base-preserving morphisms $G \rightarrow \phi^*F$ and morphisms $G \rightarrow F$ over ϕ , where pr_2 is the projection onto the second component.

In total, $\tilde{\Phi}$ is a base-preserving morphism if and only if Φ is a morphism over ϕ . Observe that $\tilde{\Phi}$ is an isomorphism (diffeomorphism) if and only if Φ is a fibre-wise isomorphism (diffeomorphism).

One can extend all of this similarly for more specific types of morphisms like vector bundle-morphisms, and very often we will not mention this 1:1 correspondence explicitly, it should be clear by context. Hence, we will also denote $\tilde{\Phi}$ by Φ .

- We will often have connections on a bundle, that is, a splitting into a horizontal and vertical bundle of its tangent bundle. When we write "Let π be the projection onto the vertical/horizontal bundle", then this will be always w.r.t. to the fixed connection even though we are not explicitly mention it again.
- We also need to extend contractions of tensors to graded extensions:

Definition 1.2: Graded extension of products

Let $l \in \mathbb{N}$ and $E_1, \dots, E_{l+1} \rightarrow M$ be vector bundles over a smooth manifold M , and $F \in \Gamma\left(\left(\bigotimes_{m=1}^l E_m^*\right) \otimes E_{l+1}\right)$. Then we define the **graded extension of F** as

$$\begin{aligned} \Omega^{k_1}(M; E_1) \times \dots \times \Omega^{k_l}(M; E_l) &\rightarrow \Omega^k(M; E_{l+1}), \\ (A_1, \dots, A_l) &\mapsto F(A_1 \wedge \dots \wedge A_l), \end{aligned}$$

where $k := k_1 + \dots + k_l$ and $k_i \in \mathbb{N}_0$ for all $i \in \{1, \dots, l\}$. $F(A_1 \wedge \dots \wedge A_l)$ is defined as an element of $\Omega^k(M; E_{l+1})$ by

$$\begin{aligned} F(A_1 \wedge \dots \wedge A_l)(Y_1, \dots, Y_k) &:= \\ \frac{1}{k_1! \dots k_l!} \sum_{\sigma \in S_k} \text{sgn}(\sigma) &F\left(A_1(Y_{\sigma(1)}, \dots, Y_{\sigma(k_1)}), \dots, A_l(Y_{\sigma(k-k_l+1)}, \dots, Y_{\sigma(k)})\right) \end{aligned}$$

for all $Y_1, \dots, Y_k \in \mathfrak{X}(M)$, where S_k is the group of permutations of $\{1, \dots, k\}$ and $\text{sgn}(\sigma)$ the signature of a given permutation σ .
 $\hat{\wedge}$ may be written just as a comma when a zero-form is involved.

In case of antisymmetric tensors we get of course:

Proposition 1.3: Graded extensions of antisymmetric tensors

Let $E_1, E_2 \rightarrow N$ be real vector bundles of finite rank over a smooth manifold N , $F \in \Omega^2(E_1; E_2)$. Then

$$F(A \hat{\wedge} B) = -(-1)^{km} F(B \hat{\wedge} A)$$

for all $A \in \Omega^k(N; E_1)$ and $B \in \Omega^m(N; E_1)$ ($k, m \in \mathbb{N}_0$). Similarly extended to all $F \in \Omega^l(E_1; E_2)$.

Remarks 1.4.

These are a common definitions and identities; see for example similar relations for the Lie algebra bracket $[\cdot, \cdot]_{\mathfrak{g}}$ of a Lie algebra $\mathfrak{g}$, see [17, §5].

Proof.

Trivial, for example by using local coordinates. ■

- References are not only given in the text, the references of referenced statements and definitions are most of the time given below the title of those statements. It will also mention whether the statement as written is a variation or generalization; if it is a strong generalization, then the reference will be mentioned in a remark after the statement or its proof.

1.4. Assumed background knowledge. It is highly recommended to have basic knowledge about differential geometry and gauge theory as presented in [17, especially Chapter 1 to 5], and we will follow the style and labeling as in [17] when we generalize certain notions; however, sometimes we will still give explicit references to help with more technical details. It can be useful to have knowledge about Lie algebra and Lie group bundles, and even Lie algebroids and Lie groupoids, but we will introduce the basic notions of LGBs and LABs such that it is not necessarily needed to have knowledge about these upfront.

We often give references about Lie group bundles (LGBs), but the given references are most of the time about Lie groupoids. If the reader has no knowledge about Lie groupoids, then it is important to know that LGBs are a special example of Lie groupoids; Lie groupoids carry “two projections,” called **source** and **target**. An LGB is a special example of a Lie groupoid whose source equals the target.³ If you look into such a reference, then the source and target are often denoted by α and β , or by s and t , respectively; simply put both to be the same and identify these with our bundle projection which we often denote by π or $\pi_{\mathcal{G}}$, with $\mathcal{G}$ the corresponding LGB. In that way it should be possible to read the references without the need to know Lie groupoids. However, we try to re-prove the needed statements such that these types of references could be avoided by the reader.

2. LIE GROUP BUNDLES (LGBs)

2.1. Definition.

³But not every Lie groupoid with equal source and target is an LGB, those are in general bundles of Lie groups which is not completely the same; this nuance will not be important here.

Definition 2.1: Lie group bundle

[10, §1.1, Def. 1.1.19; p. 11]

Let $G, \mathcal{G}, M$ be smooth manifolds. A fibre bundle

$$\begin{array}{ccc} G & \longrightarrow & \mathcal{G} \\ & & \downarrow \pi \\ & & M \end{array}$$

is called a **Lie group bundle (LGB)** if:

- (1) G and each fibre $\mathcal{G}_x := \pi^{-1}(\{x\})$, $x \in M$, are Lie groups;
- (2) there exists a bundle atlas $\{(U_i, \phi_i)\}_{i \in I}$ such that the induced maps

$$\phi_{ix} := \text{pr}_2 \circ \phi_i|_{\mathcal{G}_x} : \mathcal{G}_x \rightarrow G$$

are Lie group isomorphisms, where I is an (index) set, U_i are open sets covering M , $\phi_i : \mathcal{G}|_{U_i} \rightarrow U_i \times G$ subordinate trivializations, and pr_2 the projection onto the second factor. This atlas will be called **Lie group bundle atlas** or **LGB atlas**.

We also often say that $\mathcal{G}$ is an **LGB (over M)**, whose structural Lie group is either clear by context or not explicitly needed; and we may also denote LGBs by $G \rightarrow \mathcal{G} \xrightarrow{\pi} M$ or $\pi : \mathcal{G} \rightarrow M$.

The global section e of $\mathcal{G}$ so that e_p is the neutral element of $\mathcal{G}_p$ will be labelled as the **neutral (element)/identity section (of $\mathcal{G}$)**; we may also denote it by $e^{\mathcal{G}}$.

If M is a point we recover the notion of Lie groups, and, as usual, we have the notion of trivial LGBs:

Example 2.2: Trivial LGB

The **trivial LGB** is given as the product manifold $M \times G \rightarrow M$ with canonical multiplication $(x, g) \cdot (x, q) := (x, gq)$.

In a straightforward manner one defines LGB bundle morphisms:

Definition 2.3: LGB morphism

[10, §1.2, special situation of Def. 1.2.1 & 1.2.3, page 12]

Let $\mathcal{H} \xrightarrow{\pi_{\mathcal{H}}} N$ and $\mathcal{G} \xrightarrow{\pi_{\mathcal{G}}} M$ be two LGBs over two smooth manifolds N and M . An **LGB morphism F over f** is a pair of smooth maps $F : \mathcal{H} \rightarrow \mathcal{G}$ and $f : N \rightarrow M$ such that

$$\begin{aligned} \pi_{\mathcal{G}} \circ F &= f \circ \pi_{\mathcal{H}}, \\ F(gq) &= F(g) F(q) \end{aligned}$$

for all $g, q \in \mathcal{H}$ with $\pi_{\mathcal{H}}(g) = \pi_{\mathcal{H}}(q)$. We then also say that F is an **LGB morphism over f** . If $N = M$ and $f = \text{id}_M$, then we omit mentioning f explicitly and just write that F is a **(base-preserving) LGB morphism**.

We speak of an **LGB isomorphism (over f)** if F is a diffeomorphism.

Remark 2.4: Typical basics for LGB morphisms

As usual we have

- It is clear that condition 2 in Definition 2.1 is equivalent to say that $\mathcal{G}$ is locally isomorphic to a trivial LGB; as one may have expected already.

- If F is a diffeomorphism, then also f , and F^{-1} is an LGB isomorphism, too.

Similar to the case of Lie groups, *the* example of an LGB are the automorphisms of a vector bundle.

Example 2.5: Automorphisms of a vector bundle

[10, §1.1, special situation of Ex. 1.1.12, page 8]

Let $V \rightarrow M$ be a vector bundle and $\text{Aut}(V) \rightarrow M$ its bundle of fibre-wise automorphisms (not to be confused with the sections of $\text{Aut}(V)$ which are the base-preserving automorphisms of V). Denote with W the structural vector space of V , then $\text{Aut}(V)$ is an LGB with structural Lie group $\text{Aut}(W)$. It is clear that each fibre of $\text{Aut}(V)$ is a Lie group, and the LGB atlas is directly inherited by a vector bundle atlas $\{(U_i, L_i)\}_{i \in I}$ of V , where we use a similar notation as for LGB atlases, especially we have vector bundle trivializations $L_i : V|_{U_i} \rightarrow U_i \times W$. Then define an LGB atlas over the same open covering $(U_i)_i$ by

$$\begin{aligned} \text{Aut}(V)|_{U_i} &\rightarrow U_i \times \text{Aut}(W), \\ T &\mapsto L_i \circ T \circ L_i^{-1}|_{\{x\} \times W}, \end{aligned}$$

where $T \in \text{Aut}(V)|_x = \text{Aut}(V_x)$, and $U_i \times \text{Aut}(W)$ acts canonically on $U_i \times W$ in a fibre-wise sense. Then it is trivial to check that these give local trivializations such that $\text{Aut}(V)$ carries the structure as an LGB.

2.2. Associated Lie group bundles. For another important example recall that there is the notion of associated fibre bundles and related concepts; following references like [10, §1, Construction 1.3.8, page 20] and [17, §4.7, page 237ff.; see also Rem. 4.7.8, page 242f.]: Let $P \xrightarrow{\pi_P} M$ be a principal bundle with structural Lie group G over a smooth manifold M , N another smooth manifold, and Ψ a smooth left G -action denoted by

$$\begin{aligned} G \times N &\rightarrow N, \\ (g, v) &\mapsto \Psi(g, v) := g \cdot v. \end{aligned}$$

Then we have a right G -action on $P \times N$ given by

$$\begin{aligned} (P \times N) \times G &\rightarrow P \times N, \\ (p, v, g) &\mapsto (p \cdot g, g^{-1} \cdot v), \end{aligned}$$

and one can show that the quotient under this action, $P \times_{\Psi} N := (P \times N)/G$, yields the structure of a fibre bundle

$$\begin{array}{ccc} N & \longrightarrow & P \times_{\Psi} N \\ & & \downarrow \pi_{P \times_{\Psi} N} \\ & & M \end{array}$$

such that the map to the equivalence classes $P \times N \rightarrow P \times_{\Psi} N$, $(p, v) \mapsto [p, v]$, is a smooth surjective submersion, where the projection $\pi_{P \times_{\Psi} N} : P \times_{\Psi} N \rightarrow M$ is given by

$$\pi_{P \times_{\Psi} N}([p, v]) := \pi_P(p)$$

for all $[p, v] \in P \times_{\Psi} N$. For $x \in M$, the fibre $(P \times_{\Psi} N)_x$ is given by $(P_x \times N)/G = P_x \times_{\Psi} N$, and the fibre is diffeomorphic to N by $N \ni v \mapsto [p, v] \in (P \times_{\Psi} N)_x$ for a fixed $p \in P_x$. We will frequently use this diffeomorphism in the following without further notice.

A very important example are of course associated vector bundles, related to N being a vector space, and Ψ a Lie group representation. Similarly one has associated Lie group bundles: If G acts

on another Lie group H by Lie group automorphisms, then $P \times_\psi H$ canonically admits an LGB structure given by

$$[p, h] \cdot [p, q] := [p, hq]$$

for all $h, q \in H$ and $p \in P_x$ (that is, $\pi_P(p) = x$). The structural Lie group is given by H , and one can check that the identity section is point-wise well-defined by

$$e_x = [p, e] ,$$

where $p \in P_x$ is arbitrary and e is the neutral element of H ; the inverse of $[p, h] \in \mathcal{H}_x$ is clearly given by

$$([p, h])^{-1} = [p, h^{-1}] .$$

The special situation of $H = G$ is already an important example:

Example 2.6: Inner group bundle

[10, §1, paragraph after Def. 1.1.19, page 11; comment after Construction 1.3.8, page 20]

The **inner group bundle** or **inner LGB** of a principal bundle $P \rightarrow M$, denoted by $c_G(P)$, is defined by

$$c_G(P) := P \times_{c_G} G,$$

where $c_G : G \times G \rightarrow G$ is the left action of G on itself given by **conjugation**

$$c_G(g, h) := c_g(h) = (L_g \circ R_{g^{-1}})(h) = ghg^{-1}$$

for all $g, h \in G$, where we also denote left- and right-multiplications (with g) by L_g and R_g , respectively.

2.3. Pullbacks of LGBs. For this recall that there is the notion of pullbacks of fibre bundles, see e.g. [17, §4.1.4, page 203ff.; especially Thm. 4.1.17, page 204f.]. That is, if we additionally have a smooth manifold N and a smooth map $f : N \rightarrow M$, then we have the pullback $f^*\mathcal{G}$ of an LGB $\mathcal{G} \rightarrow M$ as a fibre bundle defined as usual by

$$f^*\mathcal{G} := N \times_{f \times \pi} \mathcal{G} := \{(x, g) \in N \times \mathcal{G} \mid f(x) = \pi(g)\} .$$

It is an embedded submanifold of $N \times \mathcal{G}$, and the structural fibre is the same Lie group as for $\mathcal{G}$. That is, the following diagram commutes

$$\begin{array}{ccc} f^*\mathcal{G} & \xrightarrow{\pi_2} & \mathcal{G} \\ \downarrow \pi_1 & & \downarrow \pi \\ N & \xrightarrow{f} & M \end{array}$$

where π_1 and π_2 are the projections onto the first and second factor, respectively, of $N \times \mathcal{G}$. Actually, $f^*\mathcal{G}$ carries a natural structure as an LGB.

Corollary 2.7: Pullbacks of LGBs are LGBs

[10, §2.3, simplified situation of the discussion around Prop. 2.3.1, page 63ff.]

Let M, N be smooth manifolds, $\mathcal{G} \xrightarrow{\pi} M$ an LGB over M and $f : N \rightarrow M$ a smooth map. Then $f^*\mathcal{G}$ has a unique (up to isomorphisms) LGB structure such that the projection $\pi_2 : f^*\mathcal{G} \rightarrow \mathcal{G}$ onto the second factor is an LGB morphism over f with $\pi_2|_x : (f^*\mathcal{G})_x \rightarrow \mathcal{G}_{f(x)}$ being a Lie group isomorphism for all $x \in N$.

Remark 2.8: Other references

The mentioned reference, [10], is rather general, formulated for Lie groupoids. If the reader is only interested into LGBs, then see e.g. [18, §3, Thm. 3.1].

Definition 2.9: Pullback LGB

Let M, N be smooth manifolds, $\mathcal{G} \rightarrow M$ an LGB over M and $f: N \rightarrow M$ a smooth map. Then we call the LGB structure on $f^*\mathcal{G}$ as given in Cor. Corollary 2.7 the **pullback LGB of $\mathcal{G}$ (under f)**. By writing $f^*\mathcal{G}$ we will often refer to this structure without further mention.

3. LGB ACTIONS, PART I

3.1. Definition. As for Lie groups, we are interested into their actions: We have an LGB $\mathcal{G} \rightarrow M$ over a smooth manifold M , and we want to construct an action of $\mathcal{G}$ on another smooth manifold N . Each fibre of $\mathcal{G}$ is a Lie group, and we have a notion of Lie groups acting on a manifold N . Therefore one could define an LGB action as a collection of Lie group actions, that is, only sections of $\mathcal{G}$ act on N ; however, one then expects that the general outcome of a product of $\Gamma(\mathcal{G})$ on N would be smooth maps from M to N . In order to recover a typical structure of action one could instead introduce a "multiplication rule," i.e. each point $p \in N$ can only be multiplied with elements of a specific fibre of $\mathcal{G}$. This "multiplication rule" will be described by a smooth map $f: N \rightarrow M$ in the sense that only the fibre over $f(p)$ will act on p .

Definition 3.1: Lie group bundle actions

[10, §1.6, special case of Def. 1.6.1, page 34]

Let M, N be smooth manifolds, $\mathcal{G} \xrightarrow{\pi} M$ an LGB over M and $f: N \rightarrow M$ a smooth map. Then a **right-action of $\mathcal{G}$ on N** is a smooth map

$$\begin{aligned} f^*\mathcal{G} &\rightarrow N, \\ (p, g) &\mapsto p \cdot g, \end{aligned}$$

satisfying the following properties:

$$\begin{aligned} f(p \cdot g) &= \pi(g), \\ (p \cdot g) \cdot h &= p \cdot (gh), \\ p \cdot e_{f(p)} &= p \end{aligned}$$

for all $p \in N$ and $g, h \in \mathcal{G}_{f(p)}$, where e is the neutral section of $\mathcal{G}$.

We analogously define left-actions, and we often write (left or right) **$\mathcal{G}$ -action on N** . Furthermore, in order to increase readability as long as the dependency on f is not important, we introduce the notation

$$N * \mathcal{G} := f^*\mathcal{G},$$

such that the action's notation has the typical shape $N * \mathcal{G} \rightarrow N$; one may also write $N * \mathcal{G} = N \times_M \mathcal{G}$. For left actions similarly $\mathcal{G} * N \rightarrow N$; even though $\mathcal{G} * N$ is the same pullback LGB as for right-actions; we also change the order of notation in this case, that is, $(g, p) \in \mathcal{G} * N$ reads $g \in \mathcal{G}_{f(p)}$ and $p \in N$.

Remark 3.2: Relation to the structure of the canonical pullback $f^*\mathcal{G}$

Observe that by the definition of $f^*\mathcal{G}$ we can also write

$$f(p \cdot g) = f(p),$$

so, the $\mathcal{G}$ -action is defined in such a way that f is invariant under it. If f is the projection of N as a bundle over M , then this means that an LGB action is fibre-preserving. Moreover, the fibre-wise group structure on $\mathcal{G}$ naturally defines a $\mathcal{G}$ -action on $\mathcal{G}$; in this situation f would be π itself, see also Example 3.8 later. Furthermore, having $M = \{*\}$ recovers the

notion of a Lie group action.

The other conditions are the typical conditions for actions, especially such that we get a $\mathcal{G}$ -action on $f^*\mathcal{G}$ by

$$(p, g) \cdot q := (p \cdot q, q^{-1}g)$$

for all $p \in N$ and $g, q \in \mathcal{G}_{f(p)}$.^a As usual, this gives rise to an equivalence relation, whose set of equivalence classes denoted by $f^*\mathcal{G}/\mathcal{G}$ is isomorphic to N (as a set) by $[p, g] \mapsto p \cdot g$, where we denote equivalence classes of $(p, g) \in f^*\mathcal{G}$ by $[p, g]$.

^aIn alignment to Definition 3.1, this action is a map $(f \circ \pi_1)^*\mathcal{G} \rightarrow f^*\mathcal{G}$, where π_1 is the projection onto the first factor in $f^*\mathcal{G}$.

Remark 3.3: Localizing LGB actions

We can localize the LGB action in the following sense: Fix any open neighbourhood U of M . Then $f^{-1}(U)$ is an open neighbourhood of N , and we can restrict the action to $f^{-1}(U)$, resulting into a map

$$(N * \mathcal{G})|_{f^{-1}(U)} \rightarrow f^{-1}(U),$$

because of $f(p \cdot g) = \pi(g) = f(p)$, that is, if $(p, g) \in (N * \mathcal{G})|_{f^{-1}(U)}$, then $p \in f^{-1}(U)$ and so $p \cdot g \in f^{-1}(U)$. In fact, by the definition of $N * \mathcal{G}$ as $f^*\mathcal{G}$, this describes a $\mathcal{G}|_U$ -action on $f^{-1}(U)$

$$f^{-1}(U) * \mathcal{G}|_U := (f|_{f^{-1}(U)})^*(\mathcal{G}|_U) = (N * \mathcal{G})|_{f^{-1}(U)} \rightarrow f^{-1}(U).$$

If $x \in M$ is a regular value of f , then $f^{-1}(\{x\})$ is an embedded submanifold of N due to the regular value theorem. In that case one could apply the same arguments to restrict the action on $f^{-1}(\{x\})$. Since $\mathcal{G}_x$ is a Lie group this describes a Lie group action on $f^{-1}(\{x\})$.

One can probably see that it is straightforward to extend a lot of the typical notions of Lie group actions to LGB actions; hence, we mainly focus on the definitions and properties which we need in this paper.

Definition 3.4: Left and right translations

[10, §1.4, special situation of Def. 1.4.1 and its discussion, page 22]

Let M, N be smooth manifolds, $\mathcal{G} \xrightarrow{\pi} M$ an LGB over M and $f: N \rightarrow M$ a smooth map. Furthermore assume that we have a right action $N * \mathcal{G} \rightarrow N$. We define the **right translation** over $x \in M$ with $g \in \mathcal{G}_x$ as a map r_g defined by

$$\begin{aligned} f^{-1}(\{x\}) &\rightarrow f^{-1}(\{x\}), \\ p &\mapsto p \cdot g, \end{aligned}$$

and we define the **orbit map** through $p \in f^{-1}(\{x\})$ as a map Φ_p given by

$$\begin{aligned} \mathcal{G}_x &\rightarrow N, \\ g &\mapsto p \cdot g. \end{aligned}$$

For $\sigma \in \Gamma(\mathcal{G})$ we define the **right translation** on N as a map r_σ by

$$\begin{aligned} N &\rightarrow N, \\ p &\mapsto p \cdot \sigma_{f(p)}. \end{aligned}$$

If one has a section of f , *i.e.* a smooth map $\tau : M \rightarrow N$, $x \mapsto \tau_x$, with $f \circ \tau = \text{id}_M$, then we can define the **orbit map** through τ as a map Φ_τ given by

$$\begin{aligned}\mathcal{G} &\rightarrow N, \\ g &\mapsto \tau_{\pi(g)} \cdot g.\end{aligned}$$

Remark 3.5: Left action and translation

Similarly we define left translations for left actions, which we similarly denote by l_g and l_σ . By Remark 3.3 we can define r_σ (and l_σ) also for local sections $\sigma \in \Gamma(\mathcal{G}|_U)$ by restricting N onto $f^{-1}(U)$, where U is some open subset of M ; then $r_\sigma, l_\sigma : f^{-1}(U) \rightarrow f^{-1}(U)$. In the same manner one achieves a restriction for $\Phi_\tau : \mathcal{G}|_U \rightarrow f^{-1}(U)$, if $\tau : U \rightarrow f^{-1}(U)$. In case of N being $\mathcal{G}$ itself we will introduce right (and left) translations later via capital letters.

Remark 3.6: Group action on sections

Assume that N is a fibre bundle over M with f as its projection. Also observe that $\Gamma(\mathcal{G})$ is clearly a group and it may be possible to endow it with an infinite-dimensional Lie group structure (but this is not important here now). In the same fashion as before, we can define a $\Gamma(\mathcal{G})$ -action on $\Gamma(N)$ given by

$$(\tau \cdot \sigma)_x := \tau_x \cdot \sigma_x$$

for all $\tau \in \Gamma(N)$, $\sigma \in \Gamma(\mathcal{G})$, and $x \in M$. It is straightforward to show that this is well-defined.

Remark 3.7: Smoothness of translations and orbit maps

It is a straightforward exercise to show that Φ_p and Φ_τ are smooth, and that r_σ and l_σ are diffeomorphisms with inverse $r_{\sigma^{-1}}$ and $l_{\sigma^{-1}}$, respectively; the same holds for r_g and l_g if x is a regular value of f , or more general, if $f^{-1}(\{x\})$ is an embedded submanifold.

Motivated by the previous remark, it might be useful to require that f is a submersion, or that N is actually some bundle over M and f its projection. In fact, this will be the case later.

3.2. Examples of LGB actions. If M is a point or f a constant map, then we recover the typical notion of a Lie group action acting on N . Additionally, we have the following examples. The notation will be as in Definition 3.4.

Example 3.8: LGB acting on itself

Each LGB $\mathcal{G} \xrightarrow{\pi} M$ acts on itself from the left and right, having $N := \mathcal{G}$ and $f := \pi$,

$$\begin{aligned}\mathcal{G} * \mathcal{G} &:= \pi^* \mathcal{G} \rightarrow \mathcal{G}, \\ (g, h) &\mapsto gh.\end{aligned}$$

That this satisfies all properties for an LGB action is clear by definition of LGBs; and using the localisation of this action and the local trivialisation of $\mathcal{G}$ it is clearly smooth. We will also call the $\mathcal{G}$ -action on itself the **multiplication in $\mathcal{G}$** .

Similarly one can argue that the **inverse map** $\mathcal{G} \rightarrow \mathcal{G}$, $g \mapsto g^{-1}$, is smooth, too.

Example 3.9: Trivial action

[10, §1.6, special situation of Ex. 1.6.3, page 35]

The projection π_1 onto the first factor of $f^*\mathcal{G} \xrightarrow{\pi_1} N$ satisfies the properties of a right $\mathcal{G}$ -action on N , that is, the action is given by

$$\begin{aligned} N * \mathcal{G} &\rightarrow N, \\ (p, g) &\mapsto p \cdot g := p. \end{aligned}$$

That this action satisfies the properties of an action for all f is trivial, hence we call it the **trivial action**.

Example 3.10: Actions of trivial LGBs

Assume that $\mathcal{G}$ is trivial, that is $\mathcal{G} \cong M \times G$ as LGBs, where G is the structural Lie group of $\mathcal{G}$. In that case, for $p \in N$, the product $p \cdot q$ is only defined if $q \in \mathcal{G}$ is of the form $(f(p), g)$, where $g \in G$. We also have

$$f^*\mathcal{G} \cong N \times G,$$

the trivial G -LGB over N . Hence, let us define a map by

$$\begin{aligned} N \times G &\rightarrow N, \\ (p, g) &\mapsto p \cdot g := p \cdot (f(p), g), \end{aligned}$$

which is clearly a smooth map since it is a composition of the $\mathcal{G}$ -action on N and

$$\begin{aligned} N \times G &\rightarrow f^*\mathcal{G}, \\ (p, g) &\mapsto (p, f(p), g). \end{aligned}$$

The latter is smooth because $(p, g) \mapsto (p, f(p), g)$ is a smooth map $N \times G \rightarrow N \times M \times G$ and $f^*\mathcal{G}$ is an embedded submanifold of $N \times M \times G$. Using [Definition 3.1](#), it is trivial to see that $p \cdot e = p$, and

$$\begin{aligned} p \cdot (gh) &= p \cdot \underbrace{(f(p), gh)}_{=(f(p), g) \cdot (f(p), h)} = (p \cdot (f(p), g)) \cdot (f(p), h) = (p \cdot g) \cdot h \end{aligned}$$

for all $g, h \in G$. Hence, we have a G -action on N , and by construction it is equivalent to the $\mathcal{G}$ -action on N .

Observe, that one can therefore recover the notion of Lie group actions not only via $M = \{*\}$, the point manifold, (or equivalently via constant maps f) but also via trivial LGBs. Translations with constant sections of $M \times G$ w.r.t. the action $N * \mathcal{G} \rightarrow N$ are trivially to be seen as translations with an element of G w.r.t. the action $N \times G \rightarrow N$.

Regarding general $\mathcal{G}$, not just trivial ones: Due to the discussion in [Remark 3.3](#) we can thence conclude that every $\mathcal{G}$ -action is locally a typical Lie group action. If f is a submersion, then the action is also a G -action on each fibre of f .

Hence, if one wants something "truly" new regarding LGB actions, then one has to look at global structures of non-trivial LGBs and their actions. In fact, the following type of example can provide such a new structure, which we will understand later once we have introduced the physical theory.

Example 3.11: Inner group bundle acting on associated fibre bundles

[10, §1.6, simplified version of Ex. 1.6.4, page 35]

Let $P \xrightarrow{\pi_P} M$ be a principal G -bundle with structural Lie group G over a smooth manifold M , and recall [Example 2.6](#). Furthermore, let F be another smooth manifold, equipped with a smooth left G -action $\Psi: G \times F \rightarrow F$. In total we have two associated bundles over M :

$$\begin{array}{ccc} G & \longrightarrow & c_G(P) \\ & \downarrow \pi_{c_G(P)} & \\ & M & \end{array} \qquad \begin{array}{ccc} F & \longrightarrow & \mathcal{F} := P \times_{\Psi} F \\ & \downarrow \pi_{\mathcal{F}} & \\ & M & \end{array}$$

the inner group bundle of P and an associated F -bundle, respectively. Then we have a left $c_G(P)$ -action on $\mathcal{F}$ given by

$$\begin{aligned} c_G(P) * \mathcal{F} &:= \pi_{\mathcal{F}}^* c_G(P) \rightarrow \mathcal{F}, \\ ([p, g], [p, v]) &\mapsto [p, \Psi(g, v)] = [p \cdot g, v] \end{aligned}$$

for all $p \in P_x$ ($x \in M$), $g \in G$ and $v \in F$. Recall that the group of sections of $c_G(P)$ is isomorphic to the group of principal bundle automorphisms of P as a group. In other words, this action is coming from the well-known action of gauge transformations on associated bundles; see for example [\[17, §5.3\]](#).

4. LIE ALGEBRA BUNDLES (LABs)

4.1. Definition. Lie algebras are the infinitesimal version of Lie groups, hence, we expect something similar for LGBs, the Lie algebra bundles:

Definition 4.1: Lie algebra bundle (LAB)

[\[10, §3.3, Definition 3.3.8, page 104\]](#)

Let $\mathfrak{g}$ be a Lie algebra, and $\mathcal{g}, M$ be smooth manifolds. A vector bundle

$$\begin{array}{ccc} \mathfrak{g} & \longrightarrow & \mathcal{g} \\ & \downarrow \pi & \\ & M & \end{array}$$

is called a **Lie algebra bundle (LAB)** if:

- (1) $\mathfrak{g}$ and each fibre $\mathcal{g}_x$, $x \in M$, are Lie algebras;
- (2) there exists a bundle atlas $\{(U_i, \phi_i)\}_{i \in I}$ such that the induced maps

$$\phi_{ix} := \text{pr}_2 \circ \phi_i|_{\mathcal{g}_x} : \mathcal{g}_x \rightarrow \mathfrak{g}$$

are Lie algebra isomorphisms, where I is an (index) set, U_i are open sets covering M , $\phi_i : \mathcal{g}|_{U_i} \rightarrow U_i \times \mathfrak{g}$ subordinate trivializations, and pr_2 the projection onto the second factor. This atlas will be called **Lie algebra bundle atlas** or **LAB atlas**.

We often say that $\mathcal{g}$ is an **LAB (over M)**, whose structural Lie algebra is either clear by context or not explicitly needed; and we may also denote LABs by $\mathfrak{g} \rightarrow \mathcal{g} \xrightarrow{\pi} M$.

Of course, we have the typical trivial examples:

Example 4.2: Trivial examples

We recover the notion of a Lie algebra, if M consist of just one point. Moreover, the **trivial LAB** is given as the product manifold $\mathcal{g} := M \times \mathfrak{g} \rightarrow M$. We have obviously a canonical smooth field of Lie brackets on this bundle $[\cdot, \cdot]_{\mathcal{g}} : \Gamma(\mathcal{g}) \times \Gamma(\mathcal{g}) \rightarrow \Gamma(\mathcal{g})$, *i.e.* $[\cdot, \cdot]_{\mathcal{g}} \in \Gamma(\wedge^2 \mathcal{g}^* \otimes \mathcal{g})$ which restricts to the Lie algebra bracket $[\cdot, \cdot]_{\mathfrak{g}}$ of $\mathfrak{g}$ on each fibre. The

bracket is given by

$$[(x, X), (x, Y)]_{\mathfrak{g}} := \left(x, [X, Y]_{\mathfrak{g}} \right)$$

for all $(x, X), (x, Y) \in M \times \mathfrak{g}$. Smoothness is an immediate consequence.

The definition of LAB morphisms is straight-forward:

Definition 4.3: LAB morphism

[10, §. 4.3, simplified version of Def. 4.3.1, page 158]

Let $\mathcal{G} \xrightarrow{\pi_{\mathcal{G}}} M$ and $\mathcal{H} \xrightarrow{\pi_{\mathcal{H}}} N$ be two LGBs over two smooth manifolds M and N . An **LAB morphism** is a pair of smooth maps $F: \mathcal{H} \rightarrow \mathcal{G}$ and $f: N \rightarrow M$ such that

$$\begin{aligned} \pi_{\mathcal{G}} \circ F &= f \circ \pi_{\mathcal{H}}, \\ F &\text{ linear,} \end{aligned}$$

$$F\left([g, q]_{\mathcal{H}_p}\right) = [F(g), F(q)]_{\mathcal{G}_{f(p)}}$$

for all $g, q \in \mathcal{H}_p$ ($p \in N$), where $[\cdot, \cdot]_{\mathcal{H}_p}$ and $[\cdot, \cdot]_{\mathcal{G}_{f(p)}}$ are Lie brackets of $\mathcal{H}_p$ and $\mathcal{G}_{f(p)}$, respectively. We also say that F is an **LAB morphism over f** . If $N = M$ and $f = \text{id}_M$, then we often omit mentioning f explicitly and just write that F is a **(base-preserving) LAB morphism**. We speak of an **LAB isomorphism (over f)** if F is a diffeomorphism.

Remark 4.4: Smooth field of Lie brackets

We have similar remarks as in Remark 2.4. Additionally, we have locally a canonical smooth field of Lie brackets which restricts to a Lie bracket on each fibre because every LAB is locally isomorphic to a trivial LAB as in Example 4.2. Define a field of Lie brackets $[\cdot, \cdot]_{\mathcal{G}}: \Gamma(\mathcal{G}) \times \Gamma(\mathcal{G}) \rightarrow \Gamma(\mathcal{G})$, i.e. $[\cdot, \cdot]_{\mathcal{G}} \in \Gamma\left(\bigwedge^2 \mathcal{G}^* \otimes \mathcal{G}\right)$, by

$$[X, Y]_{\mathcal{G}} := [X, Y]_{\mathcal{G}_x}$$

for all $X, Y \in \mathcal{G}_x$ ($x \in M$). Using a local trivialization, this bracket is locally of the form as in Rem. Remark 2.4 such that smoothness follows. In fact, as also argued in [2, §16.2, Example 2, page 114], every vector bundle equipped with a smooth field of Lie brackets is an LAB.

Endomorphisms of a vector bundle are of course another important example of LABs.

Example 4.5: Endomorphisms of a vector bundle

[10, §3.3, part of Ex. 3.3.4]

Let $V \rightarrow M$ be a vector bundle, and denote with $\text{End}(V) \rightarrow M$ its bundle of fibre-wise endomorphisms (its sections are the base-preserving bundle endomorphisms of V). This is clearly an LAB whose field of Lie brackets is given by the commutator.

As we have seen for LGBs, the pullback of LABs is again an LAB.

Corollary 4.6: Pullbacks of LABs are LABs

[18, §3, Thm. 3.2]

Let M, N be smooth manifolds, $\mathcal{G} \xrightarrow{\pi} M$ an LAB over M and $f: N \rightarrow M$ a smooth map. Then the pullback vector bundle $f^*\mathcal{G}$ has a unique (up to isomorphisms) LAB structure such that the projection $\pi_2: f^*\mathcal{G} \rightarrow \mathcal{G}$ onto the second factor is an LAB morphism over f with $\pi_2|_x: (f^*\mathcal{G})_x \rightarrow \mathcal{G}_{f(x)}$ being a Lie algebra isomorphism for all $x \in N$.

Definition 4.7: Pullback LAB

Let M, N be smooth manifolds, $\mathcal{G} \rightarrow M$ an LGB over M and $f: N \rightarrow M$ a smooth map. Then we call the LAB structure on $f^*\mathcal{G}$ as given in [Corollary 4.6](#) the **pullback LAB of $\mathcal{G}$ (under f)**.

By writing $f^*\mathcal{G}$ we will often refer to this structure without further mention.

4.2. From LGBs to LABs. Let us now quickly discuss how LGBs and LABs are related; it is very similar to the relation of Lie groups and algebras, now somewhat fibre-wise. We will follow the style of [17, §1.5.2, page 40ff.] and [10, §3.5, page 119ff.]; our approach will be using left-invariant vector fields but the mentioned latter reference actually uses right-invariant vector fields.

Let us start with introducing the basic notations; in contrast to the discussion around [Definition 3.4](#) we will denote the following maps with capital letters when referring to $\mathcal{G}$ acting on itself.

Definition 4.8: Left and right translation and conjugation

Let $\mathcal{G} \rightarrow M$ be an LGB over a smooth manifold M . For $g \in \mathcal{G}_x$ ($x \in M$) we define the following maps:

- **Left translation** given by

$$\begin{aligned} L_g: \mathcal{G}_x &\rightarrow \mathcal{G}_x, \\ h &\mapsto gh. \end{aligned}$$

- **Right translation** given by

$$\begin{aligned} R_g: \mathcal{G}_x &\rightarrow \mathcal{G}_x, \\ h &\mapsto hg. \end{aligned}$$

- **Conjugation** given by

$$\begin{aligned} c_g: \mathcal{G}_x &\rightarrow \mathcal{G}_x, \\ h &\mapsto ghg^{-1}. \end{aligned}$$

Remark 4.9: Conjugation by sections

[10, §1.4, Def. 1.4.6 and its discussion afterwards, page 24f.]

By definition of $\mathcal{G}$, all these maps are smooth. Furthermore, they clearly satisfy the typical properties as known for these maps since $\mathcal{G}_x$ is a Lie group for all $x \in M$. In particular, while the conjugation c_g is a Lie group automorphism of $\mathcal{G}_x$, it describes an LGB automorphism of $\mathcal{G}$ if extended to sections. That is for $\sigma \in \Gamma(\mathcal{G})$ we define the conjugation c_σ as a smooth map by

$$\begin{aligned} \mathcal{G} &\rightarrow \mathcal{G}, \\ q &\mapsto c_\sigma(q) := (L_\sigma \circ R_{\sigma^{-1}})(q) = (R_{\sigma^{-1}} \circ L_\sigma)(q) = \sigma_{\pi(q)} \cdot q \cdot \sigma_{\pi(q)}^{-1}. \end{aligned}$$

It is clear that $c_\sigma(gq) = c_\sigma(g) \cdot c_\sigma(q)$ for all $g, q \in \mathcal{G}$ with $\pi(g) = \pi(q)$, and that a smooth inverse is given by $c_{\sigma^{-1}}$; thence, c_σ is an LGB isomorphism of $\mathcal{G}$ on itself, an automorphism,

in sense of [Definition 2.3](#). It is also trivial to check that we have $c_{\sigma \cdot \tau} = c_\sigma \circ c_\tau$, where τ is another section of $\mathcal{G}$. Here we defined R_σ as r_σ as in [Definition 3.4](#). Similarly for left translations.

Since these are diffeomorphism of the fibres, it makes sense to say that a left-invariant vector field of $\mathcal{G}$ has to be a vertical vector field, that is, it is in the kernel of $D\pi$, the total differential/tangent map of the projection of $\mathcal{G} \xrightarrow{\pi} M$. For this recall that there is the notion of a **vertical bundle** for fibre bundles $F \xrightarrow{\varpi} M$, which is defined as a subbundle $VF \rightarrow F$ of the tangent bundle $TF \rightarrow F$ given as the kernel of $D\varpi: TF \rightarrow TM$. The fibres $V_v F$ of VF at $v \in F$ are then given by

$$V_v F = T_v F_x,$$

where $x := \varpi(v) \in M$ and F_x is the fibre of F at x . F_x is an embedded submanifold of F , thence, by definition a section $X \in \Gamma(VF)$ restricts to a vector field on the fibres, that is,

$$X|_{F_x} \in \mathfrak{X}(F_x).$$

In our case $F = \mathcal{G}$, hence $F_x = \mathcal{G}_x$ is a Lie group, so, the vertical bundle just consists of the tangent bundles of Lie groups of all fibres. All of these are generated by their Lie algebra at e_x , the identity element of $\mathcal{G}_x$. Hence, it is natural to guess that the LAB for $\mathcal{G}$ will be $V\mathcal{G}|_{e_M}$, where e_M is the image of M under the identity/neutral section of $\mathcal{G}$, thus, an embedding of M into $\mathcal{G}$. $V\mathcal{G}|_{e_M}$ is a fibre bundle, and clearly a vector bundle. Equivalently, since the identity section e is an embedding, we think of $V\mathcal{G}|_{e_M}$ as the pullback vector bundle $e^*V\mathcal{G}$, which is conveniently a vector bundle over M . That is, the construction of an LAB out of an LGB is precisely the same as in the Lie group setting: Restricting left-invariant vector fields to the identity. For all $\nu_x \in V_{e_x} \mathcal{G}$ ($x \in M$) we define its induced left-invariant vector field X_ν as a tangent vector of $\mathcal{G}_x$ given by

$$X_\nu|_g := D_{e_x} L_g(\nu_x)$$

for all $g \in \mathcal{G}_x$. Similarly for sections of $e^*V\mathcal{G}$, giving rise to a left-invariant vertical vector field on $\mathcal{G}$.

Definition 4.10: The LAB of an LGB

[10, §3.5, special situation of Def. 3.5.1, page 120]

Let $\mathcal{G} \rightarrow M$ be an LGB over a smooth manifold M , and denote with e the identity section of $\mathcal{G}$. Then we define the **LAB** $\mathcal{q}$ of $\mathcal{G}$ as the vector bundle $e^*V\mathcal{G}$ with bracket

$$[\mu_x, \nu_x]_{\mathcal{q}} := [X_\mu, X_\nu]|_{e_x}$$

for all $x \in M$ and $\mu_x, \nu_x \in V_{e_x} \mathcal{G}$.

In the following $\mathcal{q}$ will usually denote the LAB of $\mathcal{G}$.

Example 4.11: Endomorphisms as LAB of fibre-wise automorphisms

E.g. canonically extend the proof for why the Lie algebra of automorphisms are endomorphisms of a vector space as in [17, §1.5.4, page 45ff.]

Recall [Example 2.5](#) and [Example 4.5](#). For a vector bundle $V \rightarrow M$ one can show that the LAB of $\text{Aut}(V)$ is given by $\text{End}(V)$.

Example 4.12: Associated LAB as LAB of associated LGB

Similar to [10, §3.1, Prop. 3.1.4, page 90]

Recall associated LGBs of Section 2.2. So, let G, H be Lie groups, $P \rightarrow M$ a principal G -bundle over a smooth manifold M , and ψ a G -action on H by Lie group isomorphisms. Then we have the associated LGB $\mathcal{H} := P \times_{\psi} H$.

$\psi_* := [g \mapsto D_{e_G} \psi_g]$ is the induced G -representation on $\mathfrak{h}$ by Lie algebra isomorphisms, the Lie algebra of H , where e_G is the neutral element of G . Hence, we also have the associated bundle $\mathcal{H} := P \times_{\psi_*} \mathfrak{h}$ which is naturally the LAB of $\mathcal{H}$. As a special example, the LAB of the inner group bundle $c_G(P)$, Example 2.6, is the **adjoint bundle** $\text{Ad}(P)$ of P , $\text{Ad}(P) := P \times_{\text{Ad}} \mathfrak{g}$, where Ad is the adjoint representation of G .

Example 4.13: LAB of pullback LGB is pullback LAB

[18, §3, Thm. 3.5, page 21]

Let $\mathcal{G} \rightarrow M$ be an LGB over a smooth manifold M , and let $f: N \rightarrow M$ be a smooth map defined on another smooth manifold N . Then the LAB of $f^*\mathcal{G}$ is isomorphic to the pullback LAB $f^*\mathcal{g}$.

Either directly or with the previous example it is straightforward to show the following:

Corollary 4.14: Differentials of LGB morphisms are LAB morphisms

[10, §3.5, section about morphisms, page 124f.]

Let $\mathcal{H} \rightarrow N$ and $\mathcal{G} \rightarrow M$ be two LGBs over two smooth manifolds N and M , and we denote with $\mathcal{h}$ and $\mathcal{g}$ the LABs of $\mathcal{H}$ and $\mathcal{G}$, respectively. Furthermore, assume that we have an LGB morphism $F: \mathcal{H} \rightarrow \mathcal{G}$ over a smooth map $f: N \rightarrow M$. Then

$$DF|_{\mathcal{h}}: \mathcal{h} \rightarrow \mathcal{g}$$

is an LAB morphism over f .

4.3. Vertical Maurer-Cartan form of LGBs. Naturally, we have the canonical Maurer-Cartan form on each fibre; as a 1-form on the total space of $\mathcal{G}$ it is only defined on the vertical bundle $V\mathcal{G}$ and it is a straight-forward exercise to show smoothness of the following:

Definition 4.15: Vertical Maurer-Cartan form of LGBs

Straight-forward generalisation of the Maurer-Cartan form.

Let $\mathcal{G} \xrightarrow{\pi} M$ be an LGB over a smooth manifold M . We define the **vertical Maurer-Cartan form** μ_{MC} as an element of $\Gamma(V^*\mathcal{G} \otimes \pi^*\mathcal{g})$ given by

$$(\mu_{\text{MC}})_g(v) := (D_g L_{g^{-1}})(v)$$

for all $g \in \mathcal{G}$ and $v \in V_g\mathcal{G}$, where $V^*\mathcal{G}$ is the dual bundle of $V\mathcal{G}$.

Recall Section 1.3, we have a 1:1 correspondence of μ_{MC} to the following commuting diagram

$$\begin{array}{ccc} V\mathcal{G} & \xrightarrow{\mu_{\text{MC}}} & \mathcal{g} \\ \downarrow & & \downarrow \\ \mathcal{G} & \xrightarrow{\pi} & M \end{array}$$

which is the same diagram as in [10, §3.5, special situation of Prop. 3.5.3, page 121]. Since μ_{MC} restricts to the Maurer-Cartan form of the fibres, this induces a canonical isomorphism:

Corollary 4.16: Vertical tangent space of $\mathcal{G}$

[10, §3.5, a reformulation of Proposition 3.5.3, page 121]

Let $\mathcal{G} \xrightarrow{\pi} M$ be an LGB over a smooth manifold M . Then we have an isomorphism of vector bundles

$$V\mathcal{G} \cong \pi^* \mathcal{g}.$$

Observe that we know by Corollary 4.6 that $V\mathcal{G}$ admits a unique LAB structure such that $V\mathcal{G} \cong \pi^* \mathcal{g}$ is an isomorphism of LABs. By $e^* V\mathcal{G} = \mathcal{g}$, this implies that $V\mathcal{G}$ is trivial if and only if $\mathcal{g}$ is trivial; as also argued in [10, §3.5, discussion after Cor. 3.5.4, page 121]. Last but not least, sections of $V\mathcal{G}$ are therefore generated by sections of $\mathcal{g}$, the left-invariant vector fields.

4.4. Exponential map of LGBs. Similarly there is a natural exponential map, extending the one defined in each fibre.

Definition 4.17: Exponential map

[10, §3.6, second part of Example 3.6.2, page 133f.]

Let $\mathcal{G} \rightarrow M$ be an LGB over a smooth manifold M , and $\mathcal{g} \rightarrow M$ its LAB. Then we define the **exponential map** $\exp: \mathcal{g} \rightarrow \mathcal{G}$ by

$$\exp(X) := e^X := \exp_{\mathcal{G}_x}(X)$$

for all $x \in M$ and $X \in \mathcal{g}_x$, where $\exp_{\mathcal{G}_x}$ is the exponential map of the Lie group $\mathcal{G}_x$. Its extension to sections, also denoted by $\exp: \Gamma(\mathcal{g}) \rightarrow \Gamma(\mathcal{G})$, is canonically given by

$$\exp(\nu)|_x := \exp(\nu_x)$$

for all $x \in M$ and $\nu \in \Gamma(\mathcal{g})$. Usually, the LGB is given by context, otherwise we will denote the exponential map by $\exp_{\mathcal{G}}$ instead.

The exponential map is smooth because it describes the flow of left-invariant vector fields as it also happens for Lie groups. Recall that we denote left-invariant vector fields by X_ν where $\nu \in \Gamma(\mathcal{g})$.

Corollary 4.18: The exponential map as flow

[10, discussion at the beginning of §3.6, Prop. 3.6.1 and its discussion afterwards; page 132f.]

Let $\mathcal{G} \xrightarrow{\pi} M$ be an LGB over a smooth manifold M , and $\mathcal{g} \rightarrow M$ its LAB. Then the flow of a left-invariant vector field $X_\nu \in L(\mathcal{G})$ ($\nu \in \Gamma(\mathcal{g})$) is a complete flow $\phi: \mathbb{R} \times \mathcal{G} \rightarrow \mathcal{G}$ given by

$$\phi(t, g) = g \cdot e^{t\nu_{\pi(g)}}$$

for all $t \in \mathbb{R}$ and $g \in \mathcal{G}$. Especially, the map

$$\begin{aligned} \mathbb{R} \times M &\rightarrow \mathcal{G}, \\ (t, x) &\mapsto e^{t\nu_x} \end{aligned}$$

is smooth.

Remark 4.19: Simplifying notation related to the exponential map

Due to Definition 4.17 we recover a lot of the typical properties of the exponential map, if we understand these properties point-wise. As in [10, §3.6, discussion after Prop. 3.6.1, page 133], we say that $\mathbb{R} \ni t \mapsto e^{t\nu}$ is smooth in the sense of the second statement of Corollary 4.18,

and by construction we have

$$\nu_x = \left. \frac{d}{dt} \right|_{t=0} e^{t\nu_x}$$

for all $x \in M$, so that we also write

$$\nu = \left. \frac{d}{dt} \right|_{t=0} e^{t\nu}.$$

5. LGB ACTIONS, PART II

Finally, we come to the last part of the *basics* for LGBs and their notions which we require.

5.1. LGB and LAB representations. As usual, representations are a special type of group action, with an infinitesimal analogue.

Definition 5.1: LGB representations

[10, §1.7, special situation of the remark before Def. 1.7.1, page 43]

Let $\mathcal{G} \xrightarrow{\pi} M$ be an LGB over a smooth manifold M , $V \xrightarrow{p} M$ be a vector bundle, and assume that we have a left $\mathcal{G}$ -action on V denoted by $\Psi: \mathcal{G} * V := p^*\mathcal{G} \rightarrow V$. Then we say that Ψ is a **$\mathcal{G}$ -representation on V** if it is linear, that is,

$$\begin{aligned} \Psi(g, \alpha v) &= \alpha \Psi(g, v), \\ \Psi(g, v + w) &= \Psi(g, v) + \Psi(g, w) \end{aligned}$$

for all $\alpha \in \mathbb{R}$, and $(g, v), (g, w) \in \mathcal{G} * V$. In alignment with previous notations we may also write $\Psi(g, v) = \Psi_g(v)$, or $\Psi(g, v) = g \cdot v$.

By fixing a base point $x \in M$ we clearly have the typical notion of a $\mathcal{G}_x$ -representation on V_x . Equivalently, we could therefore obviously define LGB representations as a base-preserving LGB morphism $\Psi: \mathcal{G} \rightarrow \text{Aut}(V)$ as also in [10, §1.7, Def. 1.7.1, page 43].

Example 5.2: Representations of trivial LGBs

Similarly as to [Example 3.10](#), if $\mathcal{G} = M \times G$ is a trivial LGB over M , G its structural Lie group, and $V = M \times W$ a trivial vector bundle with structural vector space W , then Ψ is equivalent to a G -representation on W . Vice versa, every Lie group representation gives a representation of trivial LGBs on trivial vector bundles.

Also recall [Ex. Example 2.5](#): We can similarly construct another LGB needed for the adjoint representation.

Example 5.3: Automorphisms of LABs as LGB

[10, §1.7, special situation of Ex. 1.7.12, page 46]

Let $\mathcal{g}$ be an LAB over the smooth manifold M . We denote with $\text{Aut}(\mathcal{g}) \rightarrow M$ the bundle of fibre-wise Lie algebra automorphisms of $\mathcal{g}$, where the sections of $\text{Aut}(\mathcal{g})$ are the base-preserving LAB automorphisms of $\mathcal{g}$. As in [Example 2.5](#) one can show that $\text{Aut}(\mathcal{g})$ is an LGB by using LAB trivializations of $\mathcal{g}$ instead of vector bundle trivializations.

The notation of $\text{Aut}(\mathcal{g})$ may be confusing with the notation for vector bundle automorphisms. We usually refer to LAB automorphism when speaking of LABs, and state it explicitly if we just mean the vector bundle version.

Recall the notation introduced in [Definition 4.8](#):

Example 5.4: Adjoint LGB representation

[10, §3.5, special situation of Prop. 3.5.20, page 131]

For $g \in \mathcal{G}_x$ ($x \in M$) we have the conjugation c_g which is a $\mathcal{G}_x$ -automorphism. In the usual way we define the **adjoint representation of $\mathcal{G}$** as a base-preserving LGB morphism $\mathcal{G} \rightarrow \text{Aut}(\mathcal{G})$ given by

$$\text{Ad}_g := D_{e_x} c_g$$

for all $g \in \mathcal{G}_x$ ($x \in M$), which restrict fibre-wise to the typical adjoint representation ([Example 5.2](#)).

Infinitesimally, we have LAB actions and representations; recall Cor. [Corollary 4.6](#).

Definition 5.5: LAB actions

[10, §4.1, reformulated version for LABs of Def. 4.1.1, page 149]

Let M, N be smooth manifolds, $\mathcal{G} \xrightarrow{\pi} M$ an LAB, and $f: N \rightarrow M$ a smooth map. Then a **$\mathcal{G}$ -action on N** is a base-preserving vector bundle morphism

$$\begin{aligned} f^* \mathcal{G} &\rightarrow \text{TN} , \\ \nu &\mapsto \rho(\nu) , \end{aligned}$$

satisfying

$$Df \circ \rho = 0$$

and such that the induced map

$$\begin{aligned} \Gamma(\mathcal{G}) &\rightarrow \Gamma(f^* \mathcal{G}) \rightarrow \mathfrak{X}(N) , \\ \mu &\mapsto f^* \mu \mapsto \rho(f^* \mu) \end{aligned}$$

is a homomorphism of Lie algebras, that is,

$$\rho\left(f^*\left([\mu, \nu]_{\mathcal{G}}\right)\right) = [\rho(f^* \mu), \rho(f^* \nu)]$$

for all $\mu, \nu \in \Gamma(\mathcal{G})$.

As for Lie group and algebra actions, an LGB action induces an LAB action.

Lemma 5.6: LGB actions induce LAB actions

[10, §4.1, special situation of Thm. 4.1.6, page 152]

Let M, N be smooth manifolds, $\mathcal{G} \xrightarrow{\pi} M$ an LGB over M and $f: N \rightarrow M$ a smooth map. Then any right $\mathcal{G}$ -action $\Phi: N * \mathcal{G} = f^* \mathcal{G} \rightarrow N$ on N induces a $\mathcal{G}$ -action ρ on N by

$$\rho := D\Phi|_{f^* \mathcal{G}} ,$$

i.e.

$$\rho(\eta) = D_{(p, e_{f(p)})} \Phi(\eta)$$

for all $\eta \in (f^* \mathcal{G})_p$ ($p \in N$).

Remark 5.7: Induced LAB action by left LGB actions

Similarly to the discussion as in [17, §3.4, page 141ff.], one can show the same for left actions, but one has to use the multiplication with the inverse, that is,

$$\rho := D\Phi'|_{f^*\mathcal{G}} ,$$

where

$$\begin{aligned} \Phi' : \mathcal{G} * N &\rightarrow N , \\ (g, p) &\mapsto g^{-1} \cdot p . \end{aligned}$$

Remark 5.8: LAB actions as fibrewise actions

If f is a submersion, then one can think of the LAB action as a “collection” of Lie algebra actions on the fibres $f^{-1}(\{x\})$ ($x \in M$).

Of course, by restricting to linear vector fields one achieves representations.

Definition 5.9: LAB representations

[10, §3.3, Def. 3.3.13, page 107]

Let $\mathcal{G} \rightarrow M$ be an LAB over a smooth manifold M , and $V \rightarrow M$ be a vector bundle. A **$\mathcal{G}$ -representation on V** is an LAB morphism $\psi : \mathcal{G} \rightarrow \text{End}(V)$.

Corollary 5.10: LAB representations are specific LAB actions

[10, §4.1, special consequence of Prop. 4.1.7, page 153]

Let $\mathcal{G} \rightarrow M$ be an LAB over a smooth manifold M , $V \xrightarrow{p} M$ be a vector bundle, and $\psi : \mathcal{G} \rightarrow \text{End}(V)$ a $\mathcal{G}$ -representation on V . Then ψ defines an LAB action $\bar{\psi} : p^*\mathcal{G} \rightarrow \text{TV}$ by

$$\bar{\psi}(v, \nu) := -\psi(\nu)(v)$$

for all $(v, \nu) \in p^*\mathcal{G}$, where one makes use of the identification $T_v V_x = V_v V \cong V_x$ ($x := p(v)$).

This is a known relationship in the case of Lie algebra representations on vector spaces and their associated actions; usually this is proven by assuming a Lie group representation, however, one can show it without assuming integrability of the Lie algebra action. See for example [6, §2.1, proof of Prop. 2.1.16, page 22] and extend this to LABs.

Of course, the adjoint representation is an important example inherited by an LGB representation.

Example 5.11: Adjoint LAB representations

[10, §3.3, special situation of Ex. 3.3.15, page 108]

Let us assume the same situation as in Example 5.4 for the adjoint representation of the LGB $\mathcal{G} \rightarrow M$, $\text{Ad} : \mathcal{G} \rightarrow \text{Aut}(\mathcal{G})$. As discussed earlier, its infinitesimal version is a $\mathcal{G}$ -representation on itself, the **adjoint representation** **ad of $\mathcal{G}$** . By construction,

$$\text{ad}_\nu(\mu) := \text{ad}(\nu)(\mu) = [\nu, \mu]_{\mathcal{G}_x}$$

for all $\nu, \mu \in \mathcal{G}_x$ ($x \in M$).

5.2. Fundamental vector fields. The LAB actions inherited by LGB actions are also called fundamental vector fields; we are following [17, §3.4, generalization of Def. 3.4.1, page 143] for the labelling:

Definition 5.12: Fundamental vector fields

Let M and N be two smooth manifolds, $\mathcal{G} \rightarrow M$ an LGB over M , $f: N \rightarrow M$ a smooth map, and assume we have a right $\mathcal{G}$ -action on N , denoted as $\Phi: N * \mathcal{G} \rightarrow N$. For $\nu \in \Gamma(\mathcal{G})$ we define its induced **fundamental vector field** $\bar{\nu}$ as an element of $\mathfrak{X}(N)$ by

$$\bar{\nu}_p := D_{e_{f(p)}} \Phi_p(\nu_{f(p)})$$

for all $p \in N$, where Φ_p is the orbit map defined in [Definition 3.4](#). For left actions we define fundamental vector fields similarly by

$$\bar{\nu}_p := D_{e_{f(p)}} \Phi'_p(\nu_{f(p)})$$

for all $p \in N$, where Φ'_p is a slightly adjusted orbit map given by

$$\begin{aligned} \mathcal{G}_{f(p)} &\rightarrow N, \\ g &\mapsto g^{-1} \cdot p. \end{aligned}$$

Remark 5.13: Fundamental vectors pointwise

Again, point-wise over $x := f(p)$ this is just the typical definition of a fundamental vector field with respect to $\nu_x \in \mathcal{G}_x$ (except that $f^{-1}(\{x\})$ may not be a manifold). Hence, one has also a point-wise definition which we will also denote similarly by $\bar{\nu}_x$. If $f^{-1}(\{x\})$ is not a manifold, then $\bar{\nu}_x|_p$ is just a formal notation, and it only defines an element of $T_p N$ which may not be related to a vector field tangent to $f^{-1}(\{x\})$, not even to a vector field on N if ν_x does not formally come from a fixed section of $\mathcal{G}$.

However, if f is *e.g.* the projection of a bundle, then we have a $\mathcal{G}$ -action on the fibre $f^{-1}(\{x\})$ as manifold, and therefore $\bar{\nu}|_{f^{-1}(\{x\})}$ and $\bar{\nu}_x$ give rise to a vector field on $f^{-1}(\{x\})$. In other words, fundamental vector fields are vertical vector fields as expected and are fibre-wise fundamental vector fields coming from a Lie group action.

Remark 5.14: Map to fundamental vector fields an LAB action

As anticipated, this is related to the LAB action coming from the right $\mathcal{G}$ -action $\Phi: N * \mathcal{G} \rightarrow N$. The following can also be shown for left actions in a similar manner by recalling [Remark 5.7](#). By [Lemma 5.6](#) we have an LAB action $\rho: f^* \mathcal{G} \rightarrow TN$ given by $\rho := D\Phi|_{f^* \mathcal{G}}$. We have

$$\rho(p, \nu) = \left. \frac{d}{dt} \right|_{t=0} \Phi(p, e^{t\nu}) = \left. \frac{d}{dt} \right|_{t=0} (p \cdot e^{t\nu}) = D_{e_{f(p)}} \Phi_p(\nu) = \bar{\nu}_p$$

for all $(p, \nu) \in f^* \mathcal{G}$, where $t \in \mathbb{R}$, and e is the exponential map of $\mathcal{G}$. Thus, the map to fundamental vector fields

$$\begin{aligned} \Gamma(\mathcal{G}) &\mapsto \mathfrak{X}(N), \\ \nu &\mapsto \bar{\nu}, \end{aligned}$$

is equivalent to the map

$$\begin{aligned} \Gamma(\mathcal{G}) &\rightarrow \mathfrak{X}(N), \\ \nu &\mapsto \rho(f^* \nu) \end{aligned}$$

induced by ρ as in [Definition 5.5](#), which also implies that the map to the fundamental vector fields is a homomorphism of Lie algebras. Last but not least, by [Definition 5.5](#) it follows that fundamental vector fields are in the kernel of Df .

5.3. Differential of smooth LGB actions. We will often need to differentiate LGB actions, and thus we have to understand the tangent space of pullback manifolds:

Lemma 5.15: Tangent bundle of pullback fibre bundles

Let M and N be two smooth manifolds, $F \xrightarrow{\pi} M$ a fibre bundle over M , and $f: N \rightarrow M$ a smooth map. Then we have for its tangent spaces

$$T_{(p,v)}(f^*F) = \{(X, Y) \mid X \in T_p N, Y \in T_v F \text{ with } D_p f(X) = D_v \pi(Y)\} = TN \times_{Df \times D\pi} TF$$

for all $(p, v) \in f^*F$.

This is a straight-forward exercise to show by making use of the fact that f and π are transverse to each other. With this we can finally show the following proposition; also recall the notations introduced in [Definition 3.4](#), [Definition 4.8](#), [Definition 4.15](#) and [Definition 5.12](#).

Proposition 5.16: Differential of smooth LGB actions

Let M and N be smooth manifolds, $\pi_{\mathcal{G}}: \mathcal{G} \rightarrow M$ an LGB over M , $\Phi: N \rightarrow M$ a smooth map, and assume we have a right $\mathcal{G}$ -action on N , $\Psi: N * \mathcal{G} \rightarrow N$. Then we have

$$D_{(p,g)}\Psi(X, Y) = D_p r_{\sigma}(X) + \overline{(\mu_{MC})_g(Y - D_x \sigma(\omega))} \Big|_{p,g}$$

for all $(p, g) \in N \times_{\Phi \times \pi_{\mathcal{G}}} \mathcal{G}$ and $(X, Y) \in TN \times_{T\Phi \times T\pi_{\mathcal{G}}} T\mathcal{G}$, where $x := \Phi(p) = \pi_{\mathcal{G}}(g)$, σ is any (local) section of $\mathcal{G}$ with $\sigma_x = g$, and ω is an element of $T_x M$ given by

$$\omega := D_p \Phi(X) = D_g \pi_{\mathcal{G}}(Y) .$$

Proof. Due to $N * \mathcal{G} = f^* \mathcal{G}$ we are going to use [Lemma 5.15](#). That is, fix $(p, g) \in N * \mathcal{G}$ and $X \in T_p N$, $Y \in T_g \mathcal{G}$ with

$$D_p f(X) = D_g \pi(Y) =: \omega \in T_{f(p)} M.$$

Recall that we can localize LGB actions in sense of [Remark 3.3](#); so, let $x := f(p) = \pi(g)$, and fix a trivialization of $\mathcal{G}$ around x . Then it is clear that there is a (local)⁴ section $\sigma \in \Gamma(\mathcal{G})$ with $\sigma_x = g$. Observe that we can write

$$Y = Y - D_x \sigma(\omega) + D_x \sigma(\omega).$$

The two first summands result into a vertical tangent vector due to

$$D_g \pi(Y - D_x \sigma(\omega)) = D_g \pi(Y) - D_x \underbrace{(\pi \circ \sigma)}_{= \text{id}_M}(\omega) = D_g \pi(Y) - \omega = 0.$$

So, we write

$$Y^v := Y - D_x \sigma(\omega)$$

We have proven that $Y^v \in V_g \mathcal{G}$, and thus $(0_p, Y^v) \in T_{(p,g)}(f^* \mathcal{G})$ by [Lemma 5.15](#), where 0_p is the zero tangent vector of $T_p N$. In the same fashion we also have

$$(X, D_x \sigma(\omega)) \in T_{(p,g)}(f^* \mathcal{G})$$

because of

$$D_g \pi(D_x \sigma(\omega)) = \omega = D_g \pi(Y) = D_p f(X),$$

thence we can write in $T_{(p,g)}(f^* \mathcal{G})$

$$(X, Y) = (X, D_x \sigma(\omega) + Y^v) = (X, D_x \sigma(\omega)) + (0_p, Y^v).$$

Hence also

$$D_{(p,g)}\Phi(X, Y) = D_{(p,g)}\Phi(X, D_x \sigma(\omega)) + D_{(p,g)}\Phi(0_p, Y^v) \quad (1)$$

⁴For simplicity of notation we omit the notation of restricting on some open subset of M .

The second summand is quickly calculated as

$$D_{(p,g)}\Phi(0_p, Y^v) = \left. \frac{d}{dt} \right|_{t=0} \Phi(p, \gamma) = \left. \frac{d}{dt} \right|_{t=0} \underbrace{(p \cdot \gamma)}_{\Phi_p(\gamma)} = D_g \Phi_p(Y^v)$$

where $\gamma : I \rightarrow \mathcal{G}_x$ (I open interval of $\mathbb{R}$ containing 0) is a curve with $\gamma(0) = g$ and $d/dt|_{t=0} \gamma = Y^v$; this is due to the verticality of Y^v , and so $D_g \Phi_p(Y^v)$ is also well-defined since Φ_p is a map $\mathcal{G}_x \rightarrow N$. Now recall [Definition 4.15](#) and [Definition 5.12](#), and then observe that we can write

$$D_g \Phi_p(Y^v) = (D_g \Phi_p \circ D_{e_x} L_g \circ D_g L_{g^{-1}})(Y^v) = D_{e_x} \underbrace{(\Phi_p \circ L_g)}_{\mathcal{G} \ni q \mapsto p \cdot g q = \Phi_{p \cdot g}(q)}((\mu_{MC})_g(Y^v)) = \overline{(\mu_{MC})_g(Y^v)} \Big|_{p \cdot g} \quad (2)$$

making use of the verticality of Y^v such that operators like DL_g can act on Y^v . For the first summand in Eq. (1) we use

$$\begin{aligned} D_{(p,g)}\Phi(X, D_x \sigma(\omega)) &= D_{(p,g)}\Phi(X, D_x \sigma(D_p f(X))) \\ &= D_p \underbrace{(\Phi \circ (\text{id}_N, \sigma_f))}_{p \mapsto p \cdot \sigma_f(p) = r_\sigma(p)}(X) \\ &= D_p r_\sigma(X) . \end{aligned}$$

So, we get in total

$$D_{(p,g)}\Phi(X, Y) = D_p r_\sigma(X) + \overline{(\mu_{MC})_g(Y^v)} \Big|_{p \cdot g} .$$

■

Remark 5.17: Comparison to “classical” formula

[Proposition 5.16](#) is very similar to the “classical” formula used in gauge theory, see e.g. [17, §3.5, Prop. 3.5.4, page 149]: In the case of a Lie group action on N we have

$$D_{(p,g)}\Phi(X, Y) = D_p r_g(X) + \overline{(\mu_G)_g(Y)} \Big|_{p \cdot g}$$

for all $p \in N$, $g \in G$, $X \in T_p N$ and $Y \in T_g G$, where μ_G is the Maurer-Cartan form of G . However, in our general case the vector Y is deformed by ω , due to the fact that the action Φ has no “constant” Lie group factor anymore. This will be important later when we are going to derive the gauge transformations. Furthermore, already the first summand is now different than the classical formula, because we need to use LGB sections in order to define the push-forward of tangent vectors which are not vertical, that is, X may not be a tangent vector of $f^{-1}(\{x\})$ (which is in the general case not even an embedded submanifold) such that $D_p r_g(X)$ is in general not well-defined anymore.

By Ex. [Example 3.10](#) we expect still a typical Leibniz rule once a trivialization of $\mathcal{G}$ around g is fixed. Indeed, as we will understand and see also later, this is the case; fix such a trivialization, equip it with a canonical flat connection, and take σ to be a parallel section, that is, g as a constant section. Then one can calculate that the typical Leibniz rule is recovered. Hence, one can view this Leibniz rule as the “covariantized” version of the “classical” Leibniz rule, independent of a choice of “coordinate”/section on $\mathcal{G}$.

6. CONNECTIONS AND CURVATURE ON PRINCIPAL LGB-BUNDLES

6.1. Principal bundles with structural LGB.

6.1.1. Definition. The principal bundle $\mathcal{P}$ we are interested into is still a fibre bundle related to the same Lie group as the one behind the LGB $\mathcal{G}$, especially also $\dim(\mathcal{P}) = \dim(\mathcal{G})$, but it is equipped with an LGB action.

Definition 6.1: Principal bundles with structural LGB, [11, simplification of the beginning of §5.7, page 144f.]

Let G be a Lie group, M a smooth manifold, and an LGB $\mathcal{G} \rightarrow M$ which acts on the right on another G -fibre bundle $\mathcal{P} \rightarrow M$

$$\begin{array}{ccccc} G & \longrightarrow & \mathcal{P} & \xleftarrow{\quad} & \mathcal{G} & \longleftarrow & G \\ & & \downarrow \pi_{\mathcal{P}} & & \downarrow \pi_{\mathcal{G}} & & \\ & & M & & M & & \end{array}$$

where the right-action is defined on $\mathcal{P} * \mathcal{G}$ given by $\pi_{\mathcal{P}}^* \mathcal{G}$. Then we call $\mathcal{P}$ a **principal $\mathcal{G}$ -bundle** if

- (1) The right $\mathcal{G}$ -action on $\mathcal{P}$ is **simply transitive on the fibres**, that is, the restriction of $\mathcal{P} * \mathcal{G} \rightarrow \mathcal{P}$ on $x \in M$

$$\mathcal{P}_x \times \mathcal{G}_x \rightarrow \mathcal{P}_x$$

induces bijective orbit maps

$$\begin{aligned} \mathcal{G}_x &\rightarrow \mathcal{P}_x, \\ g &\mapsto p \cdot g \end{aligned}$$

for $p \in \mathcal{P}_x$.

- (2) There exist base-preserving **$\mathcal{G}$ -equivariant diffeomorphisms** $\varphi_i : \mathcal{P}|_{U_i} \rightarrow \mathcal{G}|_{U_i}$ subordinate to an open covering $(U_i)_i$ of M , that is,

$$\begin{aligned} \pi_{\mathcal{G}} \circ \varphi_i &= \pi_{\mathcal{P}}, \\ \varphi_i(p \cdot g) &= \varphi_i(p) \cdot g, \end{aligned}$$

where the multiplication on the right hand side is inherited by $\mathcal{G} * \mathcal{G} \rightarrow \mathcal{G}$, the multiplication on $\mathcal{G}$.

The LGB $\mathcal{G}$ is the **structural LGB** of the principal bundle $\mathcal{P}$.

Remark 6.2: Discussion about the definition of $\mathcal{G}$ -principal bundles

There are several things in need to be discussed:

- (1) The mentioned reference, [11, simplification of the beginning of §5.7, page 144f.], introduces these principal bundles in a different manner; we will come back later to this in Remark [Theorem 6.13](#).
- (2) By Remark [Remark 3.2](#) the $\mathcal{G}$ -action on $\mathcal{P}$ is fibre-preserving, thus, the restriction of $\mathcal{P} * \mathcal{G} \rightarrow \mathcal{P}$ is indeed a map $\mathcal{P}_x \times \mathcal{G}_x \rightarrow \mathcal{P}_x$.
- (3) The $\mathcal{G}$ -equivariant diffeomorphisms φ_i give rise to a **bundle atlas of $\mathcal{G}$ -equivariant bundle charts** $\xi_i : \mathcal{P}|_{U_i} \rightarrow U_i \times G$. Assume w.l.o.g. that U_i is small enough so that there is an LGB chart $\phi_i : \mathcal{G}|_{U_i} \rightarrow U_i \times G$ of $\mathcal{G}$; recall Def. [Definition 2.1](#). Then $\xi_i := \phi_i \circ \varphi_i$, which clearly satisfies

$$\xi_i(p \cdot g) = \xi_i(p) \cdot \phi_i(g),$$

where the multiplication on the right hand side is the canonical one for $U_i \times G$ as a trivial LGB (recall Ex. [Example 2.2](#)), and this can be viewed as $\mathcal{G}$ -equivariance (under the trivialization induced by ϕ_i). We usually then speak of **$\mathcal{G}$ -equivariance w.r.t. the LGB morphism ϕ_i** .

We will call such an atlas and its charts a **principal bundle atlas** and **principal bundle charts** for $\mathcal{P}$, respectively. For simplicity we may also refer to φ_i as principal bundle chart giving rise to the principal bundle atlas.

- (4) Due to the existence of φ_i one does not need to claim upfront that $\mathcal{P}$ is a G -fibre bundle. However, for readability, we decided to structure the definition like this. We

kept a similar style as for "typical" principal bundles as provided in [17, §4.2, Def. 4.2.1, page 207f.].

- (5) Since $\pi_{\mathcal{P}}$ is a surjective submersion we know by Remark [Remark 3.7](#) that right-translations r_g ($g \in \mathcal{G}_x$) are diffeomorphisms on $\mathcal{P}_x$. Furthermore, following [17, §4.2, discussion after Def. 4.2.1, page 208f.], by definition we have a simply transitive $\mathcal{G}_x$ -action (as a Lie group) on $\mathcal{P}_x$, and the isotropy group for each $p \in \mathcal{P}_x$ is trivial; the isotropy group consists in general of $g \in \mathcal{G}_x$ with $p \cdot g = p$, see *e.g.* [17, §3.2, third part of Def. 3.2.4, page 132]. Therefore, and by [17, §3.8, Thm. 3.8.8, page 165], the orbit map Φ_p gives rise to a $\mathcal{G}_x$ -equivariant diffeomorphism

$$\mathcal{G}_x \rightarrow \mathcal{P}_x,$$

where the action on $\mathcal{G}_x$ is the right-action on itself regarding the $\mathcal{G}_x$ -equivariance.

- (6) Similarly by definition, for each $x \in U_i$ we know that $(\varphi_i)|_x : \mathcal{P}_x \rightarrow \mathcal{G}_x$ is a $\mathcal{G}_x$ -equivariant diffeomorphism. In fact, together with $\pi_{\mathcal{G}} \circ \varphi_i = \pi_{\mathcal{P}}$ this clearly gives an equivalent definition of φ_i which we may make use in the following without further mention.

6.1.2. *Examples.* Let us provide examples of such principal bundles.

Example 6.3: The "classical" principal bundle

We recover principal G -bundles as principal $\mathcal{G} := M \times G$ -bundles. Recall Ex. [Example 3.10](#), the LGB action of a trivial LGB $\mathcal{G} \cong M \times G$ is equivalent to a G -action, the right-translation with an element $g \in G$ is then the right-translation of the corresponding constant section in $\mathcal{G}$; this action is clearly simply transitive. The principal G -bundle atlas is then naturally inherited by the existing principal $\mathcal{G}$ -bundle atlas. In fact, reverting this argument proves that principal G -bundles are equivalent to principal $\mathcal{G} := M \times G$ -bundles.

We often refer to principal bundles related to trivial LGBs as **typical** or **classical principal bundles**, or, as usual, principal G -bundle.

Example 6.4: The "trivial" principal $\mathcal{G}$ -bundle

$\mathcal{G}$ itself is a principal $\mathcal{G}$ -bundle, equipped with its canonical right-action inherited by its multiplication $\mathcal{G} * \mathcal{G} \rightarrow \mathcal{G}$. The principal bundle atlas then just consists of the identity map $\text{id}_{\mathcal{G}}$.

By Ex. [Example 6.3](#), it may be natural to call this the **trivial $\mathcal{G}$ -principal bundle** even if $\mathcal{G}$ itself might not be trivial. It will be clearer later why one may choose to do so.

The remarkable property is that this example shows that we are going to define a gauge theory for which LGBs themselves are allowed as principal bundles. One might have wondered why classical gauge theory uses classical principal bundles instead of LGBs (especially including non-trivial LGBs), because LGBs could be viewed as a more natural choice due to the fact that they are an analogue to how vector bundles are the "bundle-construction" of vector spaces. The problems described in Subsection [Section 6.2.1](#) show why it was easier to choose classical principal bundles, since these avoid the difficulties regarding the definition of a connection; however, we are going to solve these problems in such a way that one can either use LGBs or classical bundles or even something more general.

Our main example will be the inner bundle of a classical principal bundle, recall Ex. [Example 2.6](#). To explain the "triviality" of Ex. [Example 6.4](#) we need to introduce morphisms of principal bundles.

6.1.3. *Morphism of principal LGB-bundles.* Let us define morphisms of LGB-principal bundles

Definition 6.5: Morphism of principal bundles with structural LGB

Let M and N be smooth manifolds, $\mathcal{H} \xrightarrow{\pi_{\mathcal{H}}} N$ and $\mathcal{G} \xrightarrow{\pi_{\mathcal{G}}} M$ LGBs, and $\mathcal{P}' \xrightarrow{\pi'} N$ and $\mathcal{P} \xrightarrow{\pi} M$ principal $\mathcal{H}$ - and $\mathcal{G}$ -bundles, respectively. A **principal bundle morphism** between $\mathcal{P}'$ and $\mathcal{P}$ is a triple of smooth maps $F : \mathcal{H} \rightarrow \mathcal{G}$, $f : N \rightarrow M$ and $H : \mathcal{P}' \rightarrow \mathcal{P}$ such that the pair (F, f) is an LGB morphism as in Def. Definition 2.3 and

$$\pi \circ H = f \circ \pi', \quad (3)$$

$$H(p \cdot h) = H(p) \cdot F(h) \quad (4)$$

for all $(p, h) \in \mathcal{P}' * \mathcal{H} = (\pi')^* \mathcal{H}$. We also speak of a **principal bundle morphism over f w.r.t. the LGB morphism F** .

We speak of a **principal bundle isomorphism (over f , w.r.t. F)** if H is a diffeomorphism. If H is a base-preserving isomorphism $\mathcal{P} \rightarrow \mathcal{P}$ w.r.t. $F = \mathcal{K}_{\mathcal{G}}$, then we say that H is a **(global) gauge transformation** or **principal bundle automorphism**, and the set of all such automorphisms is denoted by $\text{Aut}(\mathcal{P})$. If H is defined on an open subset of M , then we may also speak of a **local gauge transformation**.

Remarks 6.6.

- Observe that the right hand side of Eq. (4) is well-defined because of Eq. (3), $(p, h) \in \mathcal{P}' * \mathcal{H}$ and Def. Definition 2.3, that is,

$$(\pi_{\mathcal{G}} \circ F)(h) \stackrel{\text{Definition 2.3}}{=} (f \circ \pi_{\mathcal{H}})(h) \stackrel{\pi'(p) = \pi_{\mathcal{H}}(h)}{=} (f \circ \pi')(p) \stackrel{(3)}{=} (\pi \circ H)(p),$$

thus,

$$(H(p), F(h)) \in \mathcal{P} * \mathcal{G} = \pi^* \mathcal{G}.$$

Furthermore, by additionally using Remark Remark 6.2 we know that LGB actions preserve the fibres of the principal bundles, therefore both, $H(p \cdot h)$ and $H(p) \cdot F(h)$, are over the same base point $(f \circ \pi')(p)$, so that Eq. (4) as a whole is well-defined.

- Also observe that one can conclude that (F, f) has to be an LGB morphism in order to have a satisfied and well-defined Eq. (4), assuming that H is a map over f . To well-define the right hand side, F has to be a map over f , since H is defined over f . Assuming Eq. (4), for $h' \in \mathcal{H}_{\pi'(p)}$ we have

$$H(p \cdot h'h) = H(p) \cdot F(h'h),$$

but we also get by associativity

$$H(p \cdot h'h) = H(p \cdot h') \cdot F(h) = H(p) \cdot F(h') \cdot F(h).$$

Using that $\mathcal{G}$ acts simply transitive on $\mathcal{P}$, we derive

$$F(h'h) = F(h') \cdot F(h).$$

- Assume H is a diffeomorphism, then F is an LGB isomorphism. For this we only have to show that F is a diffeomorphism by Def. Definition 2.3; by Remark Remark 2.4, also f is then a diffeomorphism. Thence, let us show that F is a diffeomorphism. This follows by the fact that the LGB actions are simply transitive on the fibres of $\mathcal{P}'$ and $\mathcal{P}$, *i.e.* orbit maps are diffeomorphisms, also recall Remark Remark 6.2. Denoting the orbit maps inherited by the action on $\mathcal{P}'$ and $\mathcal{P}$ by $\Phi^{\mathcal{P}'}$ and $\Phi^{\mathcal{P}}$, respectively, we can write

$$F(h) = \left(\left(\Phi_{H(p)}^{\mathcal{P}} \right)^{-1} \circ H \circ \Phi_p^{\mathcal{P}'} \right)(h)$$

for arbitrary p by rewriting Eq. (4). Hence, F is a diffeomorphism as the composition of diffeomorphisms.

Last but not least, it follows that H^{-1} is then $\mathcal{G}$ -equivariant in the sense of

$$H^{-1}(q \cdot g) = H^{-1}(q) \cdot F^{-1}(g)$$

for all $(q, g) \in \mathcal{P} * \mathcal{G}$, which is well-defined by similar arguments as before, especially because the inverses are maps over f^{-1} . To prove this, observe that there is a unique $(p, h) \in \mathcal{P}' * \mathcal{H}$ such that $H(p) = q$ and $F(h) = g$ due to the fact that both are bijective maps over f .⁵ Then

$$H^{-1}(q \cdot g) = H^{-1}\left(\underbrace{H(p) \cdot F(h)}_{=H(p \cdot h)}\right) = p \cdot h = H^{-1}(q) \cdot F^{-1}(g).$$

Thence, Def. Definition 6.5 is a valid definition for a principal bundle morphism, because it is easy to check that the principal bundle atlas on $\mathcal{P}$ (consisting of φ_i related to an open covering $(U_i)_i$ of M) is related to the principal bundle atlas on $\mathcal{P}'$ by

$$(F^{-1} \circ \varphi_i \circ H)|_{f^{-1}(U_i)}.$$

• Finally, observe that we have locally an isomorphism of every principal bundle $\mathcal{P}$ to $\mathcal{G}$: Fix a principal bundle chart $\varphi : \mathcal{P}|_U \rightarrow \mathcal{G}|_U$ (U some open subset of M). Then take $H := \varphi$, $F := \mathbb{K}_{\mathcal{G}|_U}$ and $f := \mathbb{K}_U$; by the definition of a principal bundle chart this gives a principal bundle isomorphism. Therefore one could say that every principal bundle is locally "trivial" in the sense of being an LGB; recall Ex. Example 6.4.

Additionally using Remark Remark 6.2 we have locally an isomorphism of $\mathcal{P}$ to a trivial LGB; keeping the same notation as in Remark Remark 6.2, choose $H := \xi_i$, $F := \phi_i$ and $f := \mathbb{K}_{U_i}$. Hence, locally every principal bundle is also classical in the sense of Ex. Example 6.3.

As expected, there is a natural isomorphism induced by local sections of $\mathcal{P}$, which are, however, no trivializations in general. In other words, by Remark Remark 6.2 we know that orbit maps through the fibres of $\mathcal{P}$ are equivariant diffeomorphisms, we want to show the same for the orbit map through a section of $\mathcal{P}$.

Lemma 6.7: Local sections of principal bundles induce isomorphisms to the structural LGB

Let $\mathcal{G} \xrightarrow{\pi_{\mathcal{G}}} M$ be an LGB over a smooth manifold M , and $\mathcal{P} \rightarrow M$ a principal $\mathcal{G}$ -bundle. Let $s : U \rightarrow \mathcal{P}$ be a smooth local section of $\mathcal{P}$ over an open subset U of M . Then the orbit map Φ_s through s , given as in Def. Definition 3.4 by

$$\begin{aligned} \mathcal{G}|_U &\rightarrow \mathcal{P}|_U, \\ g &\mapsto s_{\pi_{\mathcal{G}}(g)} \cdot g, \end{aligned}$$

is a base-preserving principal bundle isomorphism w.r.t. $\mathbb{K}_{\mathcal{G}|_U}$.

Remarks 6.8.

As in the typical formulation of gauge theory, we have an isomorphism induced by sections, but it is not necessarily a trivialization as fibre bundle, so that we do not necessarily also have a "classical" bundle; it is a trivialization in the sense of Ex. Example 6.4. Due to the similarity with the "classical" statement, we therefore were speaking of the "trivial" principal bundle in Ex. Example 6.4. Of course, since every LGB is locally trivial, we can find a typical trivialization by taking a "local-enough" section.

The nomenclature about calling LGBs "trivial" principal bundles is also introduced and discussed in [11, §5.7, third and fourth part of Remark 5.34, page 145] for groupoid-based principal bundles.

Proof of Lemma Lemma 6.7.

The proof is similar to the "classical" statement as e.g. given in [17, §4.2, proof of Lemma 4.2.7, page 210ff.], but generalized since we have to treat a possible non-triviality of $\mathcal{G}|_U$. As already discussed, the orbit map Φ_s is well-defined because of $\pi_{\mathcal{P}} \circ s = \mathbb{K}_U$ such that $(s_{\pi_{\mathcal{G}}(g)}, g) \in \mathcal{P} * \mathcal{G}$. Via Rem. Remark 3.7 we also know that Φ_s is also smooth, and it is point-wise the orbit map Φ_{s_x} through s_x which is a $\mathcal{G}_x$ -equivariant diffeomorphism by Remark Remark 6.2. So, Φ_s is fibre-wise bijective and therefore bijective as a whole since it is base-preserving, and it is $\mathcal{G}$ -equivariant w.r.t. $\mathbb{K}_{\mathcal{G}}$.

⁵In fact, it is easy to prove that $(H, F) : \mathcal{P}' * \mathcal{H} \rightarrow \mathcal{P} * \mathcal{G}$ is an LGB isomorphism (over H); also recall Cor. Corollary 2.7.

Now we want to use the inverse function theorem to show that its inverse is also smooth. Once we know that the tangent map/total derivative $D_g\Phi_s : T_g\mathcal{G} \rightarrow T_{s_x \cdot g}\mathcal{P}$ is an isomorphism of vector spaces for all $g \in \mathcal{G}|_U$, then we know by the inverse function theorem that Φ_s^{-1} is smooth. Hence, we will now show that $D_g\Phi_s$ is injective, then it has to be bijective by dimensional reasons ($\dim(\mathcal{G}) = \dim(\mathcal{P})$) so that we are done. Let us denote with $\Phi : \mathcal{P} * \mathcal{G} \rightarrow \mathcal{G}$ the right $\mathcal{G}$ -action $(p, g) \mapsto p \cdot g$ on $\mathcal{P}$, then

$$\Phi_s(g) = \Phi(s_{\pi_{\mathcal{G}}(g)}, g) = (\Phi \circ (\pi_{\mathcal{G}}^* s, \mathbb{K}_{\mathcal{G}}))(g, g).$$

Then $D_g\Phi_s$ is given by Thm. [Proposition 5.16](#),

$$\begin{aligned} D_g\Phi_s(Y) &= D_{s_x} r_{\sigma}(D_x s(\omega)) + \overline{(\mu_{MC})_g(Y - D_x \sigma(\omega))} \Big|_{s_x \cdot g} \\ &= D_x(r_{\sigma} \circ s)(\omega) + \overline{(\mu_{MC})_g(Y - D_x \sigma(\omega))} \Big|_{s_x \cdot g} \end{aligned} \quad (5)$$

for all $Y \in T_g\mathcal{G}$, where $x := \pi_{\mathcal{G}}(g)$, $\sigma \in \Gamma(\mathcal{G}|_U)^6$ with $\sigma_x = g$ and $\omega := D_g\pi_{\mathcal{G}}(Y) \in T_x M$. We want to decompose Φ_s now. Observe that $Y - D_x \sigma(\omega) \in V_g\mathcal{G}$, *i.e.* it is vertical in $\mathcal{G}$ due to the fact that

$$D_g\pi_{\mathcal{G}}(Y - D_x \sigma(\omega)) = \underbrace{D_g\pi_{\mathcal{G}}(Y)}_{=\omega} - \underbrace{D_g\pi_{\mathcal{G}}(D_x \sigma(\omega))}_{=\omega} = 0 \quad (6)$$

because of

$$\pi_{\mathcal{G}} \circ \sigma = \mathbb{K}_U$$

such that

$$D\pi_{\mathcal{G}} \circ D\sigma = \mathbb{K}_{TM|_U}. \quad (7)$$

Usually, we just make use of that without further mention; we repeat this trivial fact in order to emphasize that $D\sigma$ is injective because $\mathbb{K}_{TM|_U}$ is bijective, and to emphasize that we have

$$\text{Im}(D_x \sigma) \cap \text{Ker}(D_g\pi_{\mathcal{G}}) = \{0\}$$

(the image of $D_x \sigma$ intersects trivially with the kernel of $D_g\pi_{\mathcal{G}}$). The injectivity of $D\sigma$ implies that the dimension of its image satisfies

$$\dim(\text{Im}(D_x \sigma)) = \dim(M),$$

so that, in total, we know by dimensional reasons

$$T_g\mathcal{G} \cong \text{Im}(D_x \sigma) \oplus \text{Ker}(D_g\pi_{\mathcal{G}}) = \text{Im}(D_x \sigma) \oplus V_g\mathcal{G}.$$

We can decompose Y accordingly by Eq. (6),

$$Y = \underbrace{D_x \sigma(\omega)}_{=: Y^h} + \underbrace{Y - D_x \sigma(\omega)}_{=: Y^v} = Y^h + Y^v,$$

v stands for the vertical part and h for its complementary part ("horizontal"). In fact, as it is well-known, a section σ of a bundle induces a splitting of the short exact sequence of vector bundles

$$\sigma^*V\mathcal{G}|_U \hookrightarrow \sigma^*T\mathcal{G}|_U \xrightarrow{D\pi_{\mathcal{G}}} TM|_U,$$

and $D\sigma$ is a splitting/section of this sequence so that we have $\sigma^*T\mathcal{G} \cong \text{Im}(D\sigma) \oplus \sigma^*V\mathcal{G}$.⁷ Furthermore, again due to Eq. (7), we have $\text{Im}(D_x \sigma) \cong T_x M$ as vector spaces by $Y^h \mapsto D_g\pi_{\mathcal{G}}(Y^h) = D_g\pi_{\mathcal{G}}(Y) = \omega$ since Y^v is vertical. Hence, we can rewrite Eq. (5) as a map

$$\begin{aligned} T_x M \oplus V_g\mathcal{G} &\rightarrow T_{s_x \cdot g}\mathcal{P}, \\ (\omega, Y^v) &\mapsto D_x(r_{\sigma} \circ s)(\omega) + \overline{(\mu_{MC})_g(Y^v)} \Big|_{s_x \cdot g}. \end{aligned}$$

⁶W.l.o.g. we assume that σ is defined on U , otherwise "make U smaller around x ".

⁷ σ is actually an embedding, and thus $\text{Im}(D\sigma)$ is a well-defined subbundle isomorphic to TM .

These arguments apply to any section of a bundle, hence, we repeat this now. Observe that $r_\sigma \circ s$ is also a section of $\mathcal{P}|_U$ due to Def. [Definition 3.1](#), that is,

$$\pi_{\mathcal{P}}(s_y \cdot \sigma_y) = \pi_{\mathcal{P}}(s_y) = y$$

for all $y \in M$. As before, we split

$$T_{s_x \cdot g} \mathcal{P}|_U \cong \text{Im}(D_x(r_\sigma \circ s)) \oplus V_{s_x \cdot g} \mathcal{P},$$

and we identify $T_x M \cong \text{Im}(D_x(r_\sigma \circ s))$ as vector spaces, now via $\omega \mapsto D_x(r_\sigma \circ s)(\omega)$ due to Eq. (7) but with $r_\sigma \circ s$ playing the role as section. Additionally by Def. [Definition 5.12](#) and Remark [Remark 5.13](#) we know that fundamental vector fields are vertical, and so we can further rewrite Eq. (5) as a map

$$\begin{aligned} T_x M \oplus V_g \mathcal{G} &\rightarrow T_x M \oplus V_{s_x \cdot g} \mathcal{P}, \\ (\omega, Y^v) &\mapsto \left(\omega, \overline{(\mu_{\text{MC}})_g(Y^v)} \Big|_{s_x \cdot g} \right). \end{aligned}$$

Thus, $D_g \Phi_s$ is the pair of two linear independent maps $T_x M \rightarrow T_x M$ and $V_g \mathcal{G} \rightarrow V_{s_x \cdot g} \mathcal{P}$. The former is clearly an isomorphism, and the latter is by definition of fundamental vector fields and the Maurer-Cartan form of the shape as in Eq. (2), *i.e.*

$$Y^v \mapsto D_g \Phi_{s_x}(Y^v).$$

By Remark [Remark 6.2](#), Φ_{s_x} is a $\mathcal{G}_x$ -equivariant diffeomorphism $\mathcal{G}_x \rightarrow \mathcal{P}_x$, thence, $D_g \Phi_{s_x} : T_g \mathcal{G}_x = V_g \mathcal{G} \rightarrow T_{s_x \cdot g} \mathcal{P}_x = V_{s_x \cdot g} \mathcal{P}_x$ is an isomorphism of vector spaces. Finally we can conclude that $D_g \Phi_s$ is a linear isomorphism as the sum of two linear independent isomorphisms. As argued earlier, the inverse function theorem finishes now the proof. $\blacksquare$

Usually, one likes to think about the choice of sections as a choice of a coordinate transformation as in [17, §4.2, Remark 4.2.21, page 220]. This is due to that the gauge theory usually corresponds to a formulation via a trivial LGB, which we will understand later, but already have an idea of by *e.g.* Ex. [Example 3.10](#) and [Example 6.3](#). Then we have a local trivialization of $\mathcal{P}$ such that one usually thinks of the choice of a section as a choice of coordinate system.

However, we now learned that on a more general scale this is not completely what is happening. The idea of LGBs and principal bundles are very similar; both are fibre bundles related to a Lie group and they carry an action which also restricts on each fibre. But the fibres of an LGB are Lie groups themselves, while the fibres of a principal bundle are "just" diffeomorphic to a Lie group in an equivariant way as outlined in Remark [Remark 6.2](#) and as given in their definition Def. [Definition 6.1](#). One could view the fibres of $\mathcal{P}$ as having an "almost" Lie group structure, a Lie group structure without a designated neutral element.

Lemma [Lemma 6.7](#) shows that the choice of a section of $\mathcal{P}$ is actually the choice of a designated neutral element, naturally inducing a Lie group structure in each fibre and thus an LGB structure, which may not be trivial. Aligning the more general definition of principal bundles with the definition of LGBs. For example set $\mathcal{P} = \mathcal{G} = c_G(P)$ as in Ex. [Example 2.6](#), where P is a non-trivial classical principal bundle and the underlying Lie group G is non-abelian; such an LGB is likely non-trivial but always carries a global section, for examples the neutral section e . We will see such an example later.

We want to use this now in order to study a certain pullback of principal bundles. For this we need to introduce general pullbacks of principal bundles; also recall again Cor. [Corollary 2.7](#).

Corollary 6.9: Pullbacks of principal LGB-bundles are principal LGB-bundles

Let M, N be smooth manifolds, $f : N \rightarrow M$ a smooth map, $\mathcal{G} \rightarrow M$ an LGB, and $\mathcal{P} \rightarrow M$ a principal $\mathcal{G}$ -bundle. Then there is a unique (up to isomorphisms) principal $f^ \mathcal{G}$ -bundle structure on $f^* \mathcal{P}$, such that the projection $\pi_2 : f^* \mathcal{P} \rightarrow \mathcal{P}$ onto the second factor is a principal bundle morphism (over f) w.r.t. the projection $\pi_2^{\mathcal{G}} : f^* \mathcal{G} \rightarrow \mathcal{G}$ onto the second factor as LGB*

morphism, and such that $\pi_2|_x : (f^*\mathcal{P})_x \rightarrow \mathcal{P}_{f(x)}$ is a $\mathcal{G}_x$ -equivariant diffeomorphism w.r.t. the LGB isomorphism $\pi_2^\mathcal{G}|_x$.

Remarks 6.10.

This was also stated in [11, §5.7, second argument in Remark 5.34, page 145] for an even more general type of principal bundle, but without proof.

Proof.

As in the previous statements about pullback structures, this is a rather trivial and canonical construction. We have a right $f^*\mathcal{G}$ -action on $f^*\mathcal{P}$ defined by

$$(x, p) \cdot (x, g) := (x, p \cdot g)$$

for all $(x, p) \in f^*\mathcal{P}$ and $(x, g) \in f^*\mathcal{G}$, that is, $x \in N$, $p \in \mathcal{P}_{f(x)}$ and $g \in \mathcal{G}_{f(x)}$. This is clearly an action $f^*\mathcal{P} * f^*\mathcal{G} := \pi_1^* f^*\mathcal{G} \rightarrow f^*\mathcal{P}$, where π_1 is the canonical projection of $f^*\mathcal{P} \rightarrow N$ as fibre bundle. This action's restriction onto the fibres is simply transitive: A fibre of $f^*\mathcal{P}$ at x is $\{x\} \times \mathcal{P}_{f(x)} \cong \mathcal{P}_{f(x)}$, similarly $(f^*\mathcal{G})_x \cong \mathcal{G}_{f(x)}$. Hence, the restriction of that action at $x \in N$ is the right $\mathcal{G}_{f(x)}$ -action on $\mathcal{P}_{f(x)}$ such that the action is simply transitive.

A principal bundle atlas can be constructed by a pullback of principal bundle charts of $\mathcal{P}$, that is, let $(U_i)_i$ be an open covering of M over which we have $\mathcal{G}$ -equivariant diffeomorphisms $\varphi_i : \mathcal{P}|_{U_i} \rightarrow \mathcal{G}|_{U_i}$. Then define

$$\begin{aligned} f^*\mathcal{P}|_{f^{-1}(U_i)} &\rightarrow f^*\mathcal{G}|_{f^{-1}(U_i)}, \\ (x, p) &\mapsto (f^*\varphi_i)(x, p) := \left(x, (\varphi_i)|_{f(x)}(p)\right), \end{aligned}$$

which is well-defined and by construction a base-preserving $f^*\mathcal{G}$ -equivariant smooth map, and this map is equivalent to $(\mathcal{K}_{f^{-1}(U_i)}, \varphi_i) : N \times \mathcal{P} \rightarrow N \times \mathcal{G}$ restricted onto $f^*\mathcal{P}$ as an embedded submanifold of $N \times \mathcal{P}$. Therefore it is clearly a diffeomorphism, so that we can conclude that $f^*\mathcal{P}$ admits the structure as a principal $f^*\mathcal{G}$ -bundle.

The last part of the proof about the uniqueness of the structure is precisely as in the proof of Cor. Corollary 2.7, just replace the property of being an LGB morphism with being a principal bundle morphism w.r.t. the LGB morphism $\pi_2^\mathcal{G}$ (essentially, replace homomorphism with equivariance). Keeping the same notation as in the proof of Cor. Corollary 2.7, we get analogously

$$\varphi_i \circ \pi_2 \circ \psi_i^{-1} = \pi_2^\mathcal{G} \circ f^*\varphi_i \circ \psi_i^{-1}.$$

Then start by making use of the point-wise behaviour of π_2 and $\pi_2^\mathcal{G}$ and proceed similarly as in the proof of Cor. Corollary 2.7 to conclude the proof. ■

Definition 6.11: Pullback principal bundle

Let M, N be smooth manifolds, $f : N \rightarrow M$ a smooth map, $\mathcal{G} \rightarrow M$ an LGB, and $\mathcal{P} \rightarrow M$ a principal $\mathcal{G}$ -bundle. Then we call the principal $f^*\mathcal{G}$ -bundle structure on $f^*\mathcal{P}$ as given in Cor. Corollary 6.9 the **pullback principal bundle of $\mathcal{P}$ (under f)**. We will refer to this structure often without further mention.

We can actually show that $\mathcal{P} * \mathcal{G}$ is not only $\pi^*\mathcal{G}$ by definition but it is also isomorphic to $\pi^*\mathcal{P}$. For this recall Ex. Example 6.4, i.e. LGBs are "trivially" also principal bundles, hence we know that $\mathcal{P} * \mathcal{G}$ is a principal $\pi^*\mathcal{G}$ -bundle.

Corollary 6.12: $\mathcal{P} * \mathcal{G}$ is the pullback of $\mathcal{P}$ along its projection

Let $\mathcal{G} \rightarrow M$ be an LGB over a smooth manifold M and $\mathcal{P} \xrightarrow{\pi} M$ a principal $\mathcal{G}$ -bundle. Then we have a base-preserving principal bundle isomorphism

$$\mathcal{P} * \mathcal{G} \cong \pi^* \mathcal{P}$$

w.r.t. the LGB isomorphism given as the identity map on $\mathcal{P} * \mathcal{G} = \pi^* \mathcal{G}$.

Proof.

We have a global section of the pullback principal bundle $\pi^* \mathcal{P} \rightarrow \mathcal{P}$ given by

$$\begin{aligned} \mathcal{P} &\rightarrow \pi^* \mathcal{P}, \\ p &\mapsto (p, p). \end{aligned}$$

This is clearly a well-defined global section of $\pi^* \mathcal{P}$, so that by Lemma [Lemma 6.7](#) we achieve the desired isomorphism $\pi^* \mathcal{P} \cong \pi^* \mathcal{G} = \mathcal{P} * \mathcal{G}$. ■

Remarks 6.13.

By Lemma [Lemma 6.7](#) this isomorphism is explicitly given by

$$\begin{aligned} \mathcal{P} * \mathcal{G} &\rightarrow \pi^* \mathcal{P}, \\ (p, g) &\mapsto (p, p) \cdot (p, g) = (p, p \cdot g). \end{aligned}$$

In fact, the cited reference of Def. [Definition 6.1](#), [11, simplification of the beginning of §5.7, page 144f.], takes the existence of such a diffeomorphism as the essential sole part of defining principal bundles. Such an approach essentially avoids the second part of Def. [Definition 6.1](#).

Remark 6.14: Why "principal"?

The last result and remark also outline why we are speaking of a *principal* bundle; this is similar to [17, §4.2.2, page 212ff.]. As in [17, §3.7, Def. 3.7.24, page 159] one could say that a **principal** $\mathcal{G}$ -action on a manifold N (recall Def. [Definition 3.1](#)) is a **free** action, *i.e.* orbit maps are injective, such that

$$\begin{aligned} N * \mathcal{G} &\rightarrow N \times N, \\ (p, g) &\mapsto (p, p \cdot g) \end{aligned}$$

is a closed map. In our case, $N = \mathcal{P}$, we clearly have a free action, and we just have shown that that map is closed, because we have the composition of maps

$$\begin{aligned} \mathcal{P} * \mathcal{G} &\rightarrow \pi^* \mathcal{P} \rightarrow \mathcal{P} \times \mathcal{P}, \\ (p, g) &\mapsto (p, p \cdot g) \mapsto (p, p \cdot g). \end{aligned}$$

By Remark [Theorem 6.13](#) the first arrow is a diffeomorphism and thence a closed map. The second arrow is the inclusion, an embedding because $\pi^* \mathcal{P}$ is a closed embedded submanifold of $\mathcal{P} \times \mathcal{P}$ as the restriction of the fibre bundle $\mathcal{P} \times \mathcal{P} \xrightarrow{\mathcal{K}_{\mathcal{P}} \times \pi} \mathcal{P} \times M$ along the graph of π ; see *e.g.* [17, §4.1, proof of Thm. 4.1.17, page 204ff.]. Additionally, by the continuity of π , the graph Γ of π is closed and thus $\pi^* \mathcal{P} = (\mathcal{K}_{\mathcal{P}} \times \pi)^{-1}(\Gamma)$, too. Using this, it is a quick exercise to show that a set closed in $\pi^* \mathcal{P}$ is also closed in $\mathcal{P} \times \mathcal{P}$, thus the second arrow as inclusion is also closed. Hence, the whole composition is closed.

This knowledge should allow us to carry over a lot of similar results related to principal actions, as in [17, §3.7.5, page 159ff.] and [17, §4.2.2, page 212ff.]. Essentially Remark [Theorem 6.13](#) allows us to look at the quotient $\mathcal{P}/\mathcal{G}$ by using Godement's Theorem as given

in [17, §3.7, Thm. 3.7.10, page 155], leading to a manifold structure on $\mathcal{P}/\mathcal{G}$ and maybe similarly leading to a principal bundle structure on $\mathcal{P} \rightarrow \mathcal{P}/\mathcal{G}$.

Let us conclude our discussion about principal bundles by introducing a typical label.

Definition 6.15: Gauges of a principal bundle

Let $\mathcal{G} \rightarrow M$ be an LGB over a smooth manifold M , and $\mathcal{P} \rightarrow M$ a principal $\mathcal{G}$ -bundle. A **gauge** of $\mathcal{P}$ is a section of $\mathcal{P}$. If the section is globally defined on M , then we speak of a **global gauge**, otherwise we may just say **local gauge** or just **gauge**.

By Lemma Lemma 6.7, a gauge corresponds to a " $\mathcal{G}$ -ization" of $\mathcal{P}$, not necessarily a trivialization.

6.2. Generalized distributions and connections. Finally, let us now define the gauge theory, starting with horizontal distributions. We expect a basic understanding of horizontal distributions and their relationship to what we call connections (on principal and vector bundles). The bare-bones start with horizontal distributions.

Definition 6.16: Horizontal distribution,[17, §5.1.2, Def. 5.1.6, page 260; without the symmetry along right-translations here]

Let $F \rightarrow M$ be a fibre bundle over a smooth manifold M . Then a **horizontal distribution/bundle of F** is a smooth subbundle HF of TF with

$$TF = HF \oplus VF.$$

For $p \in M$ the fibre is denoted by $H_p F$, which we may call a **horizontal tangent space**.

As usual for gauge theory we will understand connections as horizontal distributions with a certain symmetry along the fibres in order to assure a certain behaviour of the gauge transformation of what physicists call minimal coupling; in mathematical words, in order to assure to be able to define a connection on associated vector bundles, and to assure a certain transformation of the associated curvature. To do so it is useful if this symmetry is similar to the symmetry carried in the vertical structure; recall Def. Definition 5.12 and its remark Remark 5.13. The following is a straightforward generalization of what one knows in the typical formulation of gauge theory, see *e.g.* [17, §5.1, part 2 to 4 of Prop. 5.1.3, page 258f].

Corollary 6.17: The natural invariance of the vertical bundle of $\mathcal{P}$

Let $\mathcal{G} \rightarrow M$ be an LGB over a smooth manifold M , $\mathcal{G}$ its LAB, and $\mathcal{P} \xrightarrow{\pi} M$ a principal $\mathcal{G}$ -bundle. Then

$$D_p r_g(V_p \mathcal{P}) = V_{p \cdot g} \mathcal{P} \quad (8)$$

for all $(p, g) \in \mathcal{P} * \mathcal{G}$, and we have an isomorphism of vector bundles

$$V\mathcal{P} \cong \pi^* \mathcal{G} \quad (9)$$

given by

$$\begin{aligned} \pi^* \mathcal{G} &\rightarrow V\mathcal{P}, \\ (p, \nu) &\mapsto \bar{\nu}_p. \end{aligned} \quad (10)$$

Remark 6.18: Extending the notation of fundamental vector fields

For $\mu := (p, \nu)$ we may also write

$$\bar{\mu}_p := \bar{\nu}_p,$$

which simplifies the notation in certain circumstances.

Proof of Cor. Corollary 6.17.

Recall Rem. Remark 6.2 for this proof. By definition it is clear that each fibre $\mathcal{P}_x$ ($x \in M$) is a principal $\mathcal{G}_x$ -bundle over $\{x\}$ whose $\mathcal{G}_x$ -action is the $\mathcal{G}$ -action restricted to x , and thus we know

$$D_p r_g(V_p \mathcal{P}_x) = V_{p \cdot g} \mathcal{P}_x$$

for all $p \in \mathcal{P}_x$ and $g \in \mathcal{G}_x$; see e.g. [17, §5.1, fourth part of Prop. 5.1.3, page 258f.] for such statements about principal Lie group bundles. Due to that $\mathcal{P}_x$ is a bundle over a point, we have $V \mathcal{P}_x = T \mathcal{P}_x = V \mathcal{P}|_{\mathcal{P}_x}$. Thus,

$$D_p r_g(V_p \mathcal{P}) = V_{p \cdot g} \mathcal{P}.$$


Due to the fact that the $\mathcal{G}$ -action is simply transitive we can derive that its induced $\mathcal{G}$ -action ρ is a vector bundle isomorphism: By Rem. Remark 5.14 this LAB action ρ is precisely (10) and ρ has values in $V \mathcal{P}$ by Eq. (??). We know that the orbit maps $\Phi_p : \mathcal{G}_x \rightarrow \mathcal{P}_x$ through $p \in \mathcal{P}_x$ are $\mathcal{G}_x$ -equivariant diffeomorphisms, so that $D_{e_x} \Phi_p : \mathcal{G}_x \rightarrow V_p \mathcal{P}$ is a vector space isomorphism. But we also have

$$\rho(p, \nu) = D_{e_x} \Phi_p(\nu)$$

for all $\nu \in \mathcal{G}_x$, and therefore ρ is fibre-wise an isomorphism of vector spaces such that it is an isomorphism of vector bundles. ■


By this result, we would like to have Eq. (8) also for a chosen horizontal distribution. However, its formulation leads to certain problems which we now want to discuss.

6.2.1. Idea and motivation.



For $\pi_{\mathcal{G}} : \mathcal{G} \rightarrow M$ an LGB over a smooth manifold M and $\mathcal{P} \xrightarrow{\pi} M$ a principal $\mathcal{G}$ -bundle we fix a point $p \in \mathcal{P}_x$ ($x \in M$) and can multiply that with an element $g \in \mathcal{G}_x$. Infinitesimally, we are interested into how this multiplication by g affects tangent vectors, especially non-vertical ones.

However, as we have seen in Def. Definition 3.4, Rem. ?? and Thm. Proposition 5.16 (and its proof) the push-forward of horizontal vectors is not well-defined anymore on non-vertical vectors if one uses a fixed element of an LGB; r_g is just a map $\mathcal{P}_x \rightarrow \mathcal{P}_x$. In order to study push-forwards of non-vertical vectors, we need information of the $\mathcal{G}$ -action in an open neighbourhood U around the fibres over x . Hence we want to use a section $\sigma \in \Gamma(\mathcal{G}|_U)$ with $\sigma_x = g$ instead.



But there are in general a plethora of sections with $\sigma_x = g$, thenceforth one expects that a definition of connections based on that may depend on the choice of sections and thus leading to conflicts once one looks at push-forwards with all possible $g \in \mathcal{G}$. Especially the tangential behaviour of the section's image (as an embedding of the base) may contribute to the push-forward; given a horizontal distribution, a horizontal vector may be still horizontal after a push-forward with one LGB section but not with respect to another LGB section. A similar problem may arise if we would work with local trivializations instead in order to use Ex. [Example 3.10](#). Thus, we need to adjust the typical definition of connections on principal bundles.

In order to understand what has to be changed, let us revisit the "typical" situation, that is, let $\mathcal{P}$ be a classical principal bundle as in Ex. [Example 6.3](#), *i.e.* $\mathcal{G} = M \times G$ is trivial with Lie group G . The $\mathcal{G}$ -action is equivalent to a G -action on $\mathcal{P}$ by Ex. [Example 3.10](#), a push-forward with $g \in G$ w.r.t. the latter action is equivalent to the push-forward with a constant section in $\mathcal{G}$ for which we may still simply write g .

Equip $\mathcal{P}$ with a "typical" (Ehresmann) connection as in [17, §5.1, Def. 5.1.6, page 260], that is, a horizontal distribution $H\mathcal{P}$ of $\mathcal{P}$ with

$$D_p r_g(H\mathcal{P}_p) = H\mathcal{P}_{p \cdot g} \quad (11)$$

for all $p \in \mathcal{P}$ and $g \in G$ as constant section. Recall that a connection has a 1:1 correspondence to a parallel transport, as presented in [17, §5.8, page 286ff.]. This means corresponding to a piece-wise smooth base curve $\alpha : [0, t] \rightarrow M$ ($t > 0$) with $\alpha(0) = x$ we have a **parallel transport along α** as a map $PT_\alpha^\mathcal{P} : \mathcal{P}_x \rightarrow \Gamma(\alpha^*\mathcal{P})$ satisfying

$$PT_{\alpha * \alpha'}^\mathcal{P} \Big|_t = PT_{\alpha'}^\mathcal{P} \Big|_t \circ PT_\alpha^\mathcal{P} \Big|_t, \quad (12)$$

$$PT_{\alpha^-}^\mathcal{P} \Big|_t = \left(PT_\alpha^\mathcal{P} \right)^{-1} \Big|_t, \quad (13)$$

especially parallel transport is a diffeomorphism between the fibres, where α' is just another similarly defined base curve with $\alpha'(0) = \alpha(t)$, $\alpha * \alpha'$ their concatenation (α coming first and with a suitable parametrization such that $\alpha * \alpha'$ is a map $[0, t] \rightarrow M$), and α^- denotes α traversed backwards. Moreover, Eq. (11) integrates and is equivalent to

$$PT_\alpha^\mathcal{P}(p \cdot g) \Big|_t = PT_\alpha^\mathcal{P}(p) \Big|_t \cdot g. \quad (14)$$

Thinking of the associated $\mathcal{G}$ -action, g is an element of $\mathcal{G}_x$ such that the right hand side is in general not well-defined⁸ anymore due to the fact that $PT_\alpha^\mathcal{P}(p) \Big|_t \in \mathcal{P}_{\alpha(t)}$; it is well-defined if interpreting g as a constant section of $\mathcal{G} = M \times G$, denoted now by $\bar{g} \in \Gamma(\mathcal{G})$ for bookkeeping reasons. The left

⁸Also here one could work with trivializations, especially since $\alpha^*\mathcal{P}$ is trivial as a fibre bundle due to the fact that the image of α is contractible. As before, this would just lead to other problems on a global scale, and we aim to provide a definition of connections on $\mathcal{P}$ without making use of trivializations.

hand side uses $\bar{g}_{\alpha(0)} = \bar{g}_x = (x, g)$, the right hand side $\bar{g}_{\alpha(t)} = (\alpha(t), g)$, and by Ex. [Example 3.10](#) we rewrite the G -action to a $\mathcal{G}$ -action:

$$p \cdot g = p \cdot (x, g) = p \cdot \bar{g}_x, \quad \text{PT}_\alpha^\mathcal{P}(p) \cdot g = \text{PT}_\alpha^\mathcal{P}(p) \cdot (\alpha, g) = \text{PT}_\alpha^\mathcal{P}(p) \cdot \alpha^* \bar{g},$$

where we recall that $\alpha^* \mathcal{P}$ is a principal $\alpha^* \mathcal{G}$ -bundle in sense of Def. [Definition 6.11](#).

Now we equip $\mathcal{G} \xrightarrow{\pi_\mathcal{G}} M$ with its **canonical flat connection** $\text{H}\mathcal{G} := \pi_\mathcal{G}^* \text{TM}$ which induces a parallel transport $\text{PT}_\alpha^\mathcal{G} : \mathcal{G}_x \rightarrow \Gamma(\alpha^* \mathcal{G})$ with $\text{PT}_\alpha^\mathcal{G}|_t(x, g) = (\alpha(t), g)$, especially $\text{PT}_\alpha^\mathcal{G}(\bar{g}_x) = \alpha^* \bar{g}$. In total, we can rewrite Eq. (14) to

$$\text{PT}_\alpha^\mathcal{P}(p \cdot \bar{g}_x) = \text{PT}_\alpha^\mathcal{P}(p) \cdot \alpha^* \bar{g} = \text{PT}_\alpha^\mathcal{P}(p) \cdot \text{PT}_\alpha^\mathcal{G}(\bar{g}_x).$$

This opens a gateway to define Eq. (14) on general principal $\mathcal{G}$ -bundles. That is, now let $\mathcal{P}$ be again a general principal $\mathcal{G}$ -bundle. Fix any horizontal distribution $\text{H}\mathcal{G}$ on $\mathcal{G}$ inducing a parallel transport $\text{PT}_\alpha^\mathcal{G}$, then a connection on $\mathcal{P}$ should be equivalent to a parallel transport $\text{PT}_\alpha^\mathcal{P}$ on $\mathcal{P}$ satisfying Eq. (12), (13) and

$$\text{PT}_\alpha^\mathcal{P}(p \cdot g) = \text{PT}_\alpha^\mathcal{P}(p) \cdot \text{PT}_\alpha^\mathcal{G}(g) \quad (15)$$

for all $p \in \mathcal{P}_x$ and $g \in \mathcal{G}_x$. The right hand side is now well-defined since both, $\text{PT}_\alpha^\mathcal{P}(p)$ and $\text{PT}_\alpha^\mathcal{G}(g)$, are elements of $\Gamma(\alpha^* \mathcal{P})$ and $\Gamma(\alpha^* \mathcal{G})$, respectively. We now want to derive its infinitesimal analogue similar to Eq. (11). Recall that we can view the parallel transports like $\text{PT}_\alpha^\mathcal{P}(p)$ also as a map $[0, t] \rightarrow \mathcal{P}$ with $\pi \circ \text{PT}_\alpha^\mathcal{P}(p) = \alpha$ (recall Subsection [Section 1.3](#); alternatively use Lemma [Lemma 5.15](#) and project onto the second component for the following derivatives). Then by construction

$$Y := \left. \frac{d}{dt} \right|_{t=0} \text{PT}_\alpha^\mathcal{G}(g) \in \text{H}_g \mathcal{G}$$

for all $g \in \mathcal{G}$, that is, it is horizontal in $\mathcal{G}$; similarly for the parallel transport on $\mathcal{P}$,

$$X := \left. \frac{d}{dt} \right|_{t=0} \text{PT}_\alpha^\mathcal{P}(p) \in \text{H}_p \mathcal{P}.$$

We differentiate both sides of Eq. (15) with $d/dt|_{t=0}$; the left hand side implies that the right hand side is an element of $\text{H}_{p \cdot g} \mathcal{P}$, while the right hand side gives

$$\left. \frac{d}{dt} \right|_{t=0} \left(\text{PT}_\alpha^\mathcal{P}(p) \cdot \text{PT}_\alpha^\mathcal{G}(g) \right) = \text{D}_{(p, g)} \Phi(X, Y).$$

We can now already derive an infinitesimal definition of $\text{H}\mathcal{P}$, that is, an Ehresmann connection $\text{H}\mathcal{P}$, given $\text{H}\mathcal{G}$, is precisely such a horizontal distribution such that $\text{D}\Phi$ restricts to the horizontal distributions:

$$\text{D}\Phi|_{\text{H}\mathcal{P}_{\text{D}\pi \times \text{D}\pi_\mathcal{G}} \text{H}\mathcal{G}} : \text{H}\mathcal{P}_{\text{D}\pi \times \text{D}\pi_\mathcal{G}} \text{H}\mathcal{G} \rightarrow \text{H}\mathcal{P},$$

where $\Phi : \mathcal{P} * \mathcal{G} \rightarrow \mathcal{P}$ denotes the $\mathcal{G}$ -action on $\mathcal{P}$; one could say that “horizontal vectors are a closed algebra”.

We want to use Thm. [Proposition 5.16](#) now in order to understand what Eq. (15) implies about the corresponding horizontal distribution of $\mathcal{P}$ in sense of right-translations:

$$\text{D}_{(p, g)} \Phi(X, Y) = \text{D}_p r_\sigma(X) + \overline{(\mu_{\text{MC}})_g \left(Y - \text{D}_x \sigma(\dot{\alpha}(0)) \right)} \Big|_{p \cdot g}$$

where σ is any (local) section of $\mathcal{G}$ with $\sigma_x = g$ and

$$\dot{\alpha}(0) = \left. \frac{d}{dt} \right|_{t=0} \alpha = \text{D}_p \pi(X)$$

because of $\pi \circ \text{PT}_\alpha^\mathcal{P}(p) = \alpha$ (similarly w.r.t. $\text{PT}_\alpha^\mathcal{G}$ so that it follows that (X, Y) is a tangent vector of $\mathcal{P} * \mathcal{G}$). As we already have shown several times, for example recall the beginning of the proof of Thm. [Proposition 5.16](#), we have

$$Y - \text{D}_x \sigma(\dot{\alpha}(0)) \in \text{V}_g \mathcal{G},$$




FIGURE 1. Push-forward of a horizontal tangent vector X with constant section (left) and general section (right), where $\mathcal{P}$ is a classical principal bundle as in Ex. [Example 6.3](#) equipped with a "typical" connection $H\mathcal{P}$ of principal G -bundles (G the structural Lie group).

so that the canonical projection $\pi^{\text{vert}, \mathcal{G}} : T\mathcal{G} \rightarrow V\mathcal{G}$ onto the vertical structure of $\mathcal{G}$ acts as identity on it. By making use of the horizontality of Y , we can write

$$\begin{aligned} (\mu_{\text{MC}})_g(Y - D_x \sigma(\dot{\alpha}(0))) &= (\mu_{\text{MC}} \circ \pi^{\text{vert}, \mathcal{G}})_g(Y - D_p(\sigma \circ \pi)(X)) \\ &= -(\mu_{\text{MC}} \circ \pi^{\text{vert}, \mathcal{G}})_g(D_p(\sigma \circ \pi)(X)). \end{aligned}$$

It may not surprise that $\mu_{\text{MC}} \circ \pi^{\text{vert}, \mathcal{G}}$ is by construction actually *the* connection 1-form on $\mathcal{G}$ corresponding to $H\mathcal{G}$, therefore we will denote $\mu_{\text{MC}} \circ \pi^{\text{vert}, \mathcal{G}}$ as the **total Maurer-Cartan form** $\mu_{\mathcal{G}}$ of $\mathcal{G}$. Hence we get in total (see also Figure [Figure 1](#))

$$D_p r_\sigma(X) - \overline{(\mu_{\mathcal{G}})_g(D_p(\sigma \circ \pi)(X))} \Big|_{p \cdot g} \in H_{p \cdot g} \mathcal{P} \quad (16)$$

for all $X \in H_p \mathcal{P}$, and we would like to take this as the starting point of defining a connection on $\mathcal{P}$, denoting this by $\mathcal{r}_{g*}(X)$, the **modified right-pushforward**.

By construction, $\mathcal{r}_{g*}(X)$ is independent of the chosen section σ because we have seen that it is equivalent to $D_{(p,g)} \Phi(X, Y)$, where Y is the unique element of $H_g \mathcal{G}$ lifting $D_p \pi(X)$; and thence this gives a well-defined fundamental formula for the following parts of this paper. Last but not least, our starting point was Eq. (8); there is no contradiction between the approaches of Eq. (8) and (16): If X is a vertical vector, then

$$D_p \pi(X) = 0,$$

and

$$D_p r_\sigma(X) = D_p r_g(X)$$

by Remark ???. Thence, the symmetry behind Eq. (16) will be compatible with the one of Eq. (8) on the vertical bundle. Let us first study the second summand in Eq. (16) via the *Darboux derivative*.

6.2.2. Darboux derivative on LGBs. By Cor. [Corollary 6.17](#) we know that $V\mathcal{P}$ is the pullback of $\mathcal{g}$. Since connections are projections onto the vertical bundle we are interested into the pullback situation, and therefore we will put some remarks after some definitions in the following in order to discuss the situation of pullback LGBs. Hence let us start with a general remark about the pullback situation to which we will later refer in the other remarks regarding this situation.

Remark 6.19: Pullback LGBs and their connections

Let $\mathcal{G} \xrightarrow{\pi_{\mathcal{G}}} M$ be an LGB over a smooth manifold M , and $H\mathcal{G}$ be a horizontal distribution of $\mathcal{G}$, where we denote with $\pi^{\text{vert}} : T\mathcal{G} \rightarrow V\mathcal{G}$ the corresponding (base-preserving) projection onto its vertical bundle; we will view π^{vert} as an element of $\Omega^1(\mathcal{G}; V\mathcal{G})$. Furthermore, let $f : N \rightarrow M$ be a smooth map defined on another smooth manifold N . By Lemma 5.15 we know that $T(f^*\mathcal{G})$ consists of pairs of tangent vectors $(X, Y) \in T_{(p,g)}(f^*\mathcal{G})$ ($(p, g) \in f^*\mathcal{G}$) with $X \in T_p N$, $Y \in T_g \mathcal{G}$ and

$$D_p f(X) = D_g \pi_{\mathcal{G}}(Y).$$

In the following we denote with pr_i ($i \in \{1, 2\}$) the projections onto the i -th component of $f^*\mathcal{G}$.

The projection of $f^*\mathcal{G} \rightarrow N$ is pr_1 , thus $D\text{pr}_1(X, Y) = X$. The vertical bundle $V(f^*\mathcal{G})$ as the kernel of $D\text{pr}_1$ then consists of pairs $(X = 0, Y)$, and therefore $D_g \pi_{\mathcal{G}}(Y) = 0$ which implies $Y \in V_g \mathcal{G}$. Thence,

$$V(f^*\mathcal{G}) \cong \text{pr}_2^*(V\mathcal{G}).$$

It is then trivial to check that

$$\text{pr}_2^! \pi^{\text{vert}} \in \Omega^1(f^*\mathcal{G}; \text{pr}_2^*(V\mathcal{G})) \cong \Omega^1(f^*\mathcal{G}; V(f^*\mathcal{G}))$$

is a projection onto $V(f^*\mathcal{G})$, and gives therefore rise to a horizontal distribution on $f^*\mathcal{G}$, the **pullback connection on $f^*\mathcal{G}$** . Especially we have

$$(\text{pr}_2^! \pi^{\text{vert}})_{(p,g)}(X, Y) = (0, \pi_g^{\text{vert}}(Y)),$$

and thus

$$((D\text{pr}_1, D\text{pr}_2) \circ \text{pr}_2^! \pi^{\text{vert}})|_{(p,g)}(X, Y) = (0, \pi_g^{\text{vert}}(Y)) = (\text{pr}_2^! \pi^{\text{vert}})_{(p,g)}(X, Y).$$

As we have seen, we need a slight adjustment of the vertical Maurer-Cartan as presented in Def. Definition 4.15.

Definition 6.20: Total Maurer-Cartan form

Let $\mathcal{G} \xrightarrow{\pi_{\mathcal{G}}} M$ be an LGB over a smooth manifold M , and $H\mathcal{G}$ be a horizontal distribution of $\mathcal{G}$, where we denote with $\pi^{\text{vert}} : T\mathcal{G} \rightarrow V\mathcal{G}$ the corresponding projection onto its vertical bundle. Then we define the **total Maurer-Cartan form** $\mu_{\mathcal{G}} \in \Omega^1(\mathcal{G}; \pi_{\mathcal{G}}^* \mathcal{G})$ of $\mathcal{G}$ as the connection 1-form corresponding to $H\mathcal{G}$, i.e.

$$(\mu_{\mathcal{G}})_g(Y) := (\mu_{\text{MC}} \circ \pi^{\text{vert}})|_g(Y) = (D_g L_{g^{-1}})(\pi^{\text{vert}}(Y))$$

for all $g \in \mathcal{G}$ and $Y \in T_g \mathcal{G}$.

This is clearly well-defined by construction; also recall our discussion of the vertical Maurer-Cartan form, especially Cor. ??.

Remark 6.21: Pullback situation: Part I

Given the situation as in Remark 6.19, then we have

$$L_{(p,g)}((p, q)) = (p, L_g(q))$$

for all $(p, g), (p, q) \in f^*\mathcal{G}$, where the left-translation on the left and right hand side are the ones of $f^*\mathcal{G}$ and $\mathcal{G}$, respectively. Thence,

$$L_{(p,g)} = (\text{pr}_1, L_{\text{pr}_2(p,g)} \circ \text{pr}_2),$$

and hence,

$$\begin{aligned} D_{(p,g)} L_{(p,g)}^{-1} &= (D_{(p,g)} \text{pr}_1, D_{\text{pr}_2(p,g)} L_{\text{pr}_2(p,g)} \circ D_{(p,g)} \text{pr}_2) \big|_{V_{(p,g)}(f^* \mathcal{G})} \\ &= \left(0, (\mu_{\text{MC}})_{\text{pr}_2(p,g)} \circ D_{(p,g)} \text{pr}_2 \right) \big|_{V_{(p,g)}(f^* \mathcal{G})}, \end{aligned}$$

making use of that $L_{(p,g)} : (f^* \mathcal{G})_p \rightarrow (f^* \mathcal{G})_p$ so that $D_{(p,g)} L_{(p,g)}^{-1} : V_{(p,g)}(f^* \mathcal{G}) \rightarrow V_{(p,e_x)}(f^* \mathcal{G}) = (f^* \mathcal{G})_p$, where $x := f(p)$ and e_x is the neutral element of $\mathcal{G}_x$. Altogether we get

$$\begin{aligned} (\mu_{f^* \mathcal{G}})_{(p,g)} &= D_{(p,g)} L_{(p,g)}^{-1} \circ \text{pr}_2^! \pi^{\text{vert}} \big|_{(p,g)} \\ &= \left(0, (\mu_{\text{MC}})_{\text{pr}_2(p,g)} \circ D_{(p,g)} \text{pr}_2 \right) \circ \text{pr}_2^! \pi^{\text{vert}} \big|_{(p,g)} \\ &= \left(0, (\mu_{\text{MC}} \circ \pi^{\text{vert}}) \big|_{\text{pr}_2(p,g)} \circ D_{(p,g)} \text{pr}_2 \right) \\ &= \left(0, (\mu_{\mathcal{G}})_{\text{pr}_2(p,g)} \circ D_{(p,g)} \text{pr}_2 \right) \\ &= (\text{pr}_2^! \mu_{\mathcal{G}}) \big|_{(p,g)}. \end{aligned}$$

Remark 6.22: Total Maurer-Cartan form just typical form on trivial LGBs

Let G be a Lie group with Lie algebra $\mathfrak{g}$, and $\mathcal{G} := M \times G \xrightarrow{\text{pr}_1} M$ be the trivial LGB equipped with its canonical flat connection $\text{H}\mathcal{G} := \text{pr}_1^* TM$, where pr_1 is the projection onto the first component in $M \times G$. Its LAB $\mathcal{G}$ is also a trivial bundle, $M \times \mathfrak{g}$, and we have several identities (recall Cor. Corollary 4.16)

$$T\mathcal{G} \cong \text{pr}_1^* TM \oplus \text{pr}_2^* TG \cong \text{pr}_1^* TM \oplus \mathfrak{g} \cong \text{pr}_1^* TM \oplus \text{pr}_1^* \mathcal{G} = \text{H}\mathcal{G} \oplus V\mathcal{G},$$

where $\text{pr}_2 : M \times G \rightarrow G$ is the projection onto the second component, and the projection onto the vertical bundle is then equivalent to $D\text{pr}_2 \in \Omega^1(\mathcal{G}; \text{pr}_2^* TG)$.

Now let us view G as an LGB over a point $\{*\}$, then $\mathcal{G} = f^* G$, where $f : M \rightarrow \{*\}$. Making use of the uniqueness of $\pi^{\text{vert}} = \mathbb{K}_{TG}$ for $G \rightarrow \{*\}$, we have for the pullback connection $\text{pr}_2^! \pi^{\text{vert}} = D\text{pr}_2$ which is precisely the projection for $\mathcal{G}$, and therefore we can use Remark Remark 6.21 to derive

$$\mu_{\mathcal{G}} = \text{pr}_2^! \mu_G,$$

where μ_G is the Maurer-Cartan form of G .

The Maurer-Cartan form is important for gauge transformations because it induces a derivative, which we also need now.

Remark 6.23: Maurer-Cartan form inducing a natural derivative: Part I

If G is a Lie group and $\mathfrak{g}$ its Lie algebra, then there is actually some stance that the typical Maurer-Cartan form $\mu_G \in \Omega^1(G; \mathfrak{g})$ describes *the* generalization of the total derivative of smooth maps $M \rightarrow \mathbb{R}^n$ ($n \in \mathbb{N}$), given by the **Darboux derivative** Δ as given in [10, §5.1, page 182ff.]. For a smooth section $\sigma : M \rightarrow G$ of the trivial LGB over M , this is $\Delta\sigma \in \Omega^1(M; \mathfrak{g})$ given by

$$\Delta\sigma := \sigma^! \mu_G,$$

that is,

$$(\Delta\sigma)_p(X) = D_{\sigma_x} L_{\sigma_x^{-1}}(D_x \sigma(X))$$

for all $p \in M$ and $X \in T_p M$. For $G = \mathbb{R}^n$ one usually shows that $D\sigma$ is the actual total derivative (Jacobian) by making use of $T\mathbb{R}^n \cong \mathbb{R}^n \times \mathbb{R}^n$; however, if one views the triviality of $T\mathbb{R}^n$ as the trivialization of general TG as given by $G \times \mathfrak{g}$, then it is more natural to think of the total derivative as $D\sigma$ followed by $D_{\sigma_x} L_{\sigma_x^{-1}}$ in order to receive information about the essential Lie algebra element. Then the classical total derivative of $\mathbb{R}^n$ -valued maps is actually more naturally given by the Darboux derivative.

We now have a similar behaviour in our case but related to arbitrary connections on $\mathbb{R}^n$ (as a trivial bundle over M).

Definition 6.24: Generalised Darboux derivative

Let $\mathcal{G} \xrightarrow{\pi_{\mathcal{G}}} M$ be an LGB over a smooth manifold M , and $H\mathcal{G}$ be a horizontal distribution of $\mathcal{G}$. For $\sigma \in \Gamma(\mathcal{G})$ we define the **Darboux derivative** $\Delta\sigma \in \Omega^1(M; \mathfrak{g})$

$$\Delta\sigma = \sigma^! \mu_{\mathcal{G}} = (\sigma^* \mu_{\mathcal{G}}) \circ D\sigma.$$

We may also write $\Delta^{\mathcal{G}}$ instead of Δ in order to accentuate the LGB.

Remarks 6.25.

The notation with the pullback σ^* is only needed if one wants to view $\Delta\sigma$ as a $C^\infty(M)$ -linear map $\mathfrak{X}(M) \rightarrow \Gamma(\mathfrak{g})$; in this case it is a composition of base-preserving vector bundle morphisms $D\sigma : TM \rightarrow \sigma^* T\mathcal{G}$ and $\sigma^* \mu_{\mathcal{G}} : \sigma^* T\mathcal{G} \rightarrow \sigma^* \pi_{\mathcal{G}}^* \mathfrak{g} \cong \mathfrak{g}$ which can be extended to sections, where π is the projection of $\mathcal{G}$.

Of course one can view $\Delta\sigma$ as a map $TM \rightarrow \mathfrak{g}$ which one can also write as the composition of $D\sigma : TM \rightarrow T\mathcal{G}$ and $\mu_{\mathcal{G}} : T\mathcal{G} \rightarrow \mathfrak{g}$, that is,

$$\Delta\sigma = \mu_{\mathcal{G}} \circ D\sigma.$$

If one wants to emphasize base points of the involved pullback bundles, that is, one views $\mu_{\mathcal{G}}$ as a map $T\mathcal{G} \rightarrow \pi_{\mathcal{G}}^* \mathfrak{g}$, then one can write point-wise

$$(\mu_{\mathcal{G}})_{\sigma_x} \circ D_x \sigma = (\sigma_x, (\Delta\sigma)|_x)$$

for all $x \in M$.

As a derivative we expect a Leibniz rule as for the classical Darboux derivative (see [10, §5.1, Eq. 2, page 182]). However, we cannot make a general statement about that yet, as long as we do not fix a more specific type of connection on $\mathcal{G}$. We will come back to this later; at this point just be sure of that there is of course a certain Leibniz rule, simply due to that it generalizes the "classical" Darboux derivative, as we are going to see.

Let us now discuss the pullback situation for the Darboux derivative.

Remark 6.26: Maurer-Cartan form inducing a natural derivative: Part II

We have now something similar to Remark 6.23. Let $\mathcal{G} = M \times \mathbb{R}^n$ ($n \in \mathbb{N}$) be the trivial abelian LGB, and ∇ a vector bundle connection on $M \times \mathbb{R}^n$, for which we have the associated projection onto the vertical bundle $\pi^{\text{vert}} : T\mathcal{G} \rightarrow V\mathcal{G}$. For $\sigma \in \Gamma(\mathcal{G})$ we then have

$$\nabla_{X_x} \sigma = (\sigma^* \pi^{\text{vert}})(D_x \sigma(X_x)) \quad (17)$$

for all $x \in M$ and $X_x \in T_x M$, by making use of $\sigma^* V\mathcal{G} \cong M \times \mathbb{R}^n$ as vector bundles such that the right hand side has again values in $\mathcal{G}$ due to "enough triviality" and equals the left hand side.

Alternatively, one could use the isomorphism as given in Cor. 4.16. This is also more natural in the sense of that one wants that $\nabla_{X_x} \sigma$ is a section over M ; M can be viewed as the image of the neutral section e (the zero vector here), so that $\nabla_{X_x} \sigma$ should have values

in $e^*V\mathcal{G} = \mathcal{G}$. Henceforth one could say it is more natural to "pull $(\sigma^*\pi^{\text{vert}})(D_x\sigma(X_x))$ back" to $\mathcal{G}$ by left-translation in order to define $\nabla_{X_x}\sigma$, *i.e.*

$$\begin{aligned} D_{\sigma_x} L_{\sigma_x^{-1}} \left((\sigma^*\pi^{\text{vert}})(D_x\sigma(X_x)) \right) &= (\sigma^*\mu_{\text{MC}} \circ \sigma^*\pi^{\text{vert}})|_x (D_x\sigma(X_x)) \\ &= (\Delta\sigma)|_x(X_x). \end{aligned}$$

Thus, as in Remark [Remark 6.23](#) one may say that $\Delta\sigma$ is *the* generalization of vector bundle connections to general LGBs $\mathcal{G}$. Furthermore, this argument is not based on a given (local) trivialization to handle the arising pullback in Eq. (17).

Remark 6.27: Pullback situation: Part II

Following Remark [Remark 6.21](#) and its notation and results (see Remark [Remark 6.19](#) for the initial setup), we can calculate $\Delta^{f^*\mathcal{G}}(f^*\sigma) \in \Omega^1(N; f^*\mathcal{G})$ for $\sigma \in \mathcal{G}$, first we write

$$f^*\sigma|_p = (p, \sigma_{f(p)})$$

for all $p \in N$, and thus

$$\begin{aligned} \Delta^{f^*\mathcal{G}}(f^*\sigma) &= (f^*\sigma)^! \mu_{f^*\mathcal{G}} \\ &= (f^*\sigma)^! \text{pr}_2^! \mu_{\mathcal{G}} \\ &= (\text{pr}_2 \circ f^*\sigma)^! \mu_{\mathcal{G}} \\ &= (\sigma \circ f)^! \mu_{\mathcal{G}} \\ &= ((\sigma \circ f)^* \mu_{\mathcal{G}}) \circ D(\sigma \circ f) \\ &= \underbrace{(f^*\sigma^* \mu_{\mathcal{G}}) \circ f^*(D\sigma)}_{= f^*((\sigma^* \mu_{\mathcal{G}}) \circ D\sigma)} \circ Df \\ &= f^!((\sigma^* \mu_{\mathcal{G}}) \circ D\sigma) \\ &= f^!(\Delta^{\mathcal{G}}\sigma), \end{aligned} \tag{18}$$

where we rewrote the chain rule

$$D(\sigma \circ f) = f^*(D\sigma) \circ Df,$$

but as in Remark [Theorem 6.25](#) one can decide to omit this notation. Such Darboux derivatives related to the pullback connection on $f^*\mathcal{G}$ we call the **pullback Darboux derivative**, and similar to the notation of pullback vector bundle connections we write

$$f^*\Delta^{\mathcal{G}} := \Delta^{f^*\mathcal{G}}.$$

Remark 6.28: Canonical flat Darboux derivative

Given the situation as in Remark [Remark 6.22](#) we can use Remark [Remark 6.27](#) to derive

$$\Delta^{\mathcal{G}} = f^*\Delta^0,$$

where Δ^0 on the right hand side is the Darboux derivative of the Lie group G as a bundle over point. Trivially, $\Delta^0 g \equiv 0$ for all $g \in G$, and thus

$$\Delta^{\mathcal{G}}(f^*g) = f^!(\Delta^0 g) = 0$$

for all $g \in G$ viewed as a section of $G \rightarrow \{0\}$, so f^*g are the constant sections of $\mathcal{G}$. In sense of Remark [Remark 6.26](#) it makes sense to say that σ is **parallel w.r.t. Δ** . Since constant sections generate all sections of $\mathcal{G}$, we then speak of the **canonical flat Darboux derivative**, and constant sections are its parallel sections.

However, just because Δ^0 is a zero map, does not mean that also $\Delta^{\mathcal{G}}$ is zero. On one hand because of a Leibniz rule which we will discuss later in more detail, and on the other hand, as mentioned in Subsection [Section 1.3](#), we can view general sections $\sigma \in \Gamma(\mathcal{G})$ (not necessarily constant) equivalently as a smooth map $\text{pr}_2 \circ \sigma : M \rightarrow G$, denoted by $\bar{\sigma}$ for bookkeeping reasons. By [Remark 6.22](#) we then get

$$\Delta^{\mathcal{G}} \sigma = \sigma^! \text{pr}_2^! \mu_G = (\text{pr}_2 \circ \sigma)^! \mu_G = \Delta^G \bar{\sigma}$$

for all $\sigma \in \Gamma(\mathcal{G})$, where Δ^G is the "classical" Darboux derivative as in [Remark 6.23](#) related to μ_G . This emphasizes why we can speak of a canonical flat derivative due to the fact that μ_G is flat (Maurer-Cartan equation), and we may simply write $\Delta^{\mathcal{G}} = \Delta^G$.

We can actually rewrite some important equations now; recall the notation introduced in [Remark 6.18](#).

Remark 6.29: Darboux derivative in the infinitesimal LGB action

Recall [Thm. Proposition 5.16](#); keeping the same notation as in this theorem but denoting the projection of $\mathcal{P}$ by π , we checked several times that $Y - D_x \sigma(\omega)$ is vertical, and thus we can now write

$$\begin{aligned} D_{(p,g)} \Phi(X, Y) &= D_p r_\sigma(X) + \overline{(\mu_{MC})_g(Y - D_x \sigma(\omega))} \Big|_{p \cdot g} \\ &= D_p r_\sigma(X) + \overline{(\mu_{MC} \circ \pi^{\text{vert}})_g(Y - D_x \sigma(\omega))} \Big|_{p \cdot g} \\ &= D_p r_\sigma(X) + \overline{(\mu_{MC} \circ \pi^{\text{vert}})_g(Y)} \Big|_{p \cdot g} - \overline{(\mu_{MC} \circ \pi^{\text{vert}})_{\sigma_x}(D_x \sigma(\omega))} \Big|_{p \cdot g} \\ &= D_p r_\sigma(X) + \overline{(\mu_{\mathcal{G}})_g(Y)} \Big|_{p \cdot g} - \overline{(\Delta \sigma)_x(\omega)} \Big|_{p \cdot g} \\ &= D_p r_\sigma(X) - \overline{(\pi^! \Delta \sigma)_p(X)} \Big|_{p \cdot g} + \overline{(\mu_{\mathcal{G}})_g(Y)} \Big|_{p \cdot g} \end{aligned}$$

If $\mathcal{G}$ is a trivial LGB, then this emphasizes again that we recover the typical Leibniz rule by choosing a constant section σ , using [Remarks 6.22](#) and [Remark 6.28](#).

Remark 6.30: Idea behind the notion of connection on principal bundles using the Darboux derivative

Recall the notation and discussion around the terms in [Eq. \(16\)](#) which will be important for our definition of a connection on principal LGB-bundles. The terms in [Eq. \(16\)](#) can be similarly rewritten as in

$$\begin{aligned} D_p r_\sigma(X) - \overline{(\mu_{\mathcal{G}})_g(D_p(\sigma \circ \pi)(X))} \Big|_{p \cdot g} &= D_p r_\sigma(X) - \overline{((\Delta \sigma)_x \circ D_p \pi)(X)} \Big|_{p \cdot g} \\ &= D_p r_\sigma(X) - \overline{(\pi^! \Delta \sigma)_p(X)} \Big|_{p \cdot g}. \end{aligned}$$

We will use this form for the definition of the principal bundle connection. As in [Remark 6.29](#), if $\mathcal{G}$ is trivial and σ a constant section corresponding to a Lie group element g , then these terms are just

$$D_p r_g(X),$$

making use of [Ex. Example 3.10](#).

In fact, the Darboux derivative naturally induces a connection on $\mathcal{G}$ as LAB of $\mathcal{G}$. One may have expected that since $H\mathcal{G}$ should infinitesimally induce a horizontal distribution on $\mathcal{G}$; recall the exponential map introduced in Subsection [Section 4.4](#). Also recall that by definition of vertical bundles we know for the vertical bundle of the LAB $\mathcal{G} \xrightarrow{\pi_{\mathcal{G}}} M$ that $V\mathcal{G} \cong \pi_{\mathcal{G}}^* \mathcal{G}$, making use of

that LABs are vector bundles; in the following we will use the natural projection onto the second component of $\pi_{\mathcal{G}}^* \mathcal{G}$ but now defined on $V\mathcal{G}$, denoted by $\text{pr}_2 : V\mathcal{G} \rightarrow \mathcal{G}$.

Proposition 6.31: LGB connection induces LAB connection

Let $\mathcal{G} \xrightarrow{\pi_{\mathcal{G}}} M$ be an LGB over a smooth manifold M , and $H\mathcal{G}$ be a horizontal distribution of $\mathcal{G}$. Then the map $\nabla^{\mathcal{G}} : \Gamma(\mathcal{G}) \rightarrow \Omega^1(M; \mathcal{G})$, $\nu \mapsto \nabla^{\mathcal{G}} \nu$ denoted as an element of $\Omega^1(M; \mathcal{G})$ by $X \mapsto \nabla_X^{\mathcal{G}} \nu$, defined by

$$\nabla_X^{\mathcal{G}} \nu|_x := \text{pr}_2 \left(\frac{d}{dt} \Big|_{t=0} \left((\Delta e^{t\nu})_x(X) \right) \right)$$

for all $x \in M$, $X \in T_x M$ and $\nu \in \Gamma(\mathcal{G})$, is a well-defined vector bundle connection on $\mathcal{G}$, where $t \in \mathbb{R}$, and $\text{pr}_2 : V\mathcal{G} \rightarrow \mathcal{G}$ is the projection onto the second component of $V\mathcal{G}$ naturally viewed as pullback bundle.

Remarks 6.32.

The notation of pr_2 is usually omitted in such constructions due to the fact that

$$t \mapsto (\Delta e^{t\nu})_x(X)$$

is a curve with values in $\mathcal{G}_x$, a vector space, so that one canonically uses the identification of tangent spaces of vector spaces with itself to show that $\nabla_X^{\mathcal{G}} \nu|_x \in \mathcal{G}_x$. We will keep pr_2 for the proof for the sake of rigorousness, but we will drop it after this proposition for simplicity without further mention.

In order to prove this we need to apply Schwarz's Theorem in order to switch Δ with d/dt . To do this rigorously we need to introduce the canonical involution/flip on double tangent bundles; since this does not completely fit into this paper's subject and may be already known by the reader, you can learn about the double tangent bundle and its flip map in Appendix [Section A](#) if needed.

Proof of Prop. Proposition 6.31.

Let us begin with well-definedness. Similar to Diagram (A.1) we have the double vector bundle

$$\begin{array}{ccc} TT\mathcal{G} & \xrightarrow{D\pi_{T\mathcal{G}}} & T\mathcal{G} \\ \downarrow \pi_{TT\mathcal{G}} & & \downarrow \pi_{T\mathcal{G}} \\ T\mathcal{G} & \xrightarrow{\pi_{T\mathcal{G}}} & \mathcal{G} \end{array} \quad (19)$$

We also have

$$\begin{aligned} \frac{d}{dt} \Big|_{t=0} \left((\Delta e^{t\nu})_x(X) \right) &= \frac{d}{dt} \Big|_{t=0} \left((\mu_{\mathcal{G}} \circ D_x e^{t\nu})(X) \right) \\ &= D_{D_x e(X)} \mu_{\mathcal{G}} \left(\frac{d}{dt} \Big|_{t=0} \left(D_x e^{t\nu}(X) \right) \right) \\ &= D_{D_x e(X)} \mu_{\mathcal{G}} \left(S_{\mathcal{G}}(D_x \nu(X)) \right) \\ &= D_{D_x e(X)} \mu_{\mathcal{G}} (\nu_T(X)) \\ &\in T_{\mu_{\mathcal{G}}(D_x e(X))} \mathcal{G} \end{aligned} \quad (20)$$

for all $x \in M$ and $X \in T_x M$, where e is the neutral section of $\mathcal{G}$, and we viewed $\mu_{\mathcal{G}}$ as a map $T\mathcal{G} \rightarrow \mathcal{G}$ when applying the chain rule; recall Remark [Theorem 6.25](#). $S_{\mathcal{G}}$ is the linear canonical flip map on $TT\mathcal{G}$, especially see the last part of Remark [Remark A.2](#). We also introduced the notation ν_T similar to Remark [Remark A.5](#) for simplicity; in fact ν_T is a vector field on $T\mathcal{G}$ only defined over TM which is canonically embedded into $T\mathcal{G}$ by embedding M into $\mathcal{G}$ via e . We also naturally embed $T\mathcal{G}$ into $TT\mathcal{G}$. In total, we will work with these embeddings now so that everything is embedded into $TT\mathcal{G}$ as the "total space"; hence, you will also see e and its total derivative De acting as an embedding several times.

We get by Eq. (19)

$$\begin{aligned} D\pi_{T\mathcal{G}}(\nabla_X^\mathcal{G}\nu|_x) &= D_{D_x e(X)}(\underbrace{\pi_{T\mathcal{G}} \circ \mu_\mathcal{G}}_{=e \circ \pi_\mathcal{G} \circ \pi_{T\mathcal{G}}})(\nu_T(X)) \\ &= (D_x e \circ D_{e_x} \pi_\mathcal{G})(\nu_x) \\ &= 0 \in T_{e_x} \mathcal{G} \end{aligned}$$

viewing $S_\mathcal{G}$ as a base-preserving isomorphism from $\pi_{TT\mathcal{G}} : TT\mathcal{G} \rightarrow T\mathcal{G}$ to $D\pi_{T\mathcal{G}} : TT\mathcal{G} \rightarrow T\mathcal{G}$, and using that $\mathcal{G} = e^*V\mathcal{G}$. The projection of $\mathcal{G} \rightarrow M$ is canonically the restriction of $\pi_{T\mathcal{G}}$, hence we can conclude that

$$\left. \frac{d}{dt} \right|_{t=0} \left((\Delta e^{t\nu})_x(X) \right) \in V_{\mu_\mathcal{G}(D_x e(X))} \mathcal{G},$$

and so we can derive that $\text{pr}_2 : V\mathcal{G} \rightarrow \mathcal{G}$ is defined on this; in total $\nabla^\mathcal{G}\nu$ is well-defined. By Eq. (20) we also trivially know that

$$\pi_{TT\mathcal{G}} \left(\left. \frac{d}{dt} \right|_{t=0} \left((\Delta e^{t\nu})_x(X) \right) \right) = \mu_\mathcal{G}(D_x e(X)) \in \mathcal{G}_x.$$

Viewing $V\mathcal{G}$ naturally as the pullback of $\mathcal{G}$ along its projection we can therefore write

$$D_{D_x e(X)} \mu_\mathcal{G}(\nu_T(X)) \stackrel{\text{Eq. (20)}}{=} \left. \frac{d}{dt} \right|_{t=0} \left((\Delta e^{t\nu})_x(X) \right) = (\mu_\mathcal{G}(D_x e(X)), \nabla_X^\mathcal{G}\nu|_x),$$

so that smoothness of $\nabla^\mathcal{G}\nu$ follows. In order to understand whether $\nabla^\mathcal{G}$ is a vector bundle connection we are hence interested into the restriction of Diagram (19) onto

$$\begin{array}{ccc} V\mathcal{G} & \xrightarrow{D\pi_{T\mathcal{G}}} & \overline{M} \\ \downarrow \pi_{TT\mathcal{G}} & & \downarrow \cong \\ \mathcal{G} & \xrightarrow{\pi_{T\mathcal{G}}} & M \end{array}$$

where M is canonically embedded into $\mathcal{G}$ by e , and $\overline{M}$ is the further canonical embedding of this (the image of e) into $T\mathcal{G}$ by the zero section. As in Appendix Section A, the addition of vectors and the scalar multiplication of the left vertical arrow is denoted as usual, while the one of the upper horizontal arrow will be denoted by $\blackplus$ and $\cdot$, respectively. For $Y \in T_x M$ we now have

$$\begin{aligned} \nabla_{\lambda X + \kappa Y}^\mathcal{G}\nu|_x &= \text{pr}_2 \left(D_{D_x e(\lambda X + \kappa Y)} \mu_\mathcal{G}(\nu_T(\lambda X + \kappa Y)) \right) \\ &= \text{pr}_2 \left(D_{D_x e(\lambda X + \kappa Y)} \mu_\mathcal{G}(\lambda \cdot \nu_T(X) \blackplus \kappa \cdot \nu_T(Y)) \right) \\ &= \text{pr}_2 \left(\lambda \cdot D_{D_x e(X)} \mu_\mathcal{G}(\nu_T(X)) \blackplus \kappa \cdot D_{D_x e(Y)} \mu_\mathcal{G}(\nu_T(Y)) \right) \\ &= \lambda \text{pr}_2 \left(D_{D_x e(X)} \mu_\mathcal{G}(\nu_T(X)) \right) + \kappa \text{pr}_2 \left(D_{D_x e(Y)} \mu_\mathcal{G}(\nu_T(Y)) \right) \\ &= \lambda \nabla_X^\mathcal{G}\nu|_x + \kappa \nabla_Y^\mathcal{G}\nu|_x \end{aligned}$$

for all $\lambda, \kappa \in \mathbb{R}$, using Remark Remark A.3, Remark A.4 and Remark A.5. By these remarks we also derive for another section $\mu \in \Gamma(\mathcal{G})$

$$\begin{aligned} \nabla_X^\mathcal{G}(\lambda\nu + \kappa\mu)|_x &= \text{pr}_2 \left(D_{D_x e(X)} \mu_\mathcal{G}(\underbrace{(\lambda\nu + \kappa\mu)_T(X)}_{=\lambda\nu_T + \kappa\mu_T}) \right) \\ &= \lambda \text{pr}_2 \left(D_{D_x e(X)} \mu_\mathcal{G}(\nu_T(X)) \right) + \kappa \text{pr}_2 \left(D_{D_x e(X)} \mu_\mathcal{G}(\mu_T(X)) \right) \\ &= \lambda \nabla_X^\mathcal{G}\nu|_x + \kappa \nabla_X^\mathcal{G}\mu|_x, \end{aligned}$$

and

$$\nabla_X^\mathcal{G}(f\nu)|_x = \text{pr}_2 \left(D_{D_x e(X)} \mu_\mathcal{G}((f\nu)_T(X)) \right)$$

$$\begin{aligned}
&= \text{pr}_2 \left(D_{D_x e(X)} \mu_{\mathcal{G}} \left(f(x) \nu_T(X) + X(f) \nu^a \frac{\partial}{\partial \xi^a} \Big|_{D_x e(X)} \right) \right) \\
&= f(x) \nabla_X^{\mathcal{G}} \nu|_x + X(f) \text{pr}_2 \left(D_{D_x e(X)} \mu_{\mathcal{G}} \left(\nu^a \frac{\partial}{\partial \xi^a} \Big|_{D_x e(X)} \right) \right)
\end{aligned}$$

for all $f \in C^\infty(M)$, where $(\xi^a)_a$ are fibre coordinates of $\mathcal{G}$. It was well-defined to use the linearity of pr_2 , since we know by what we have shown earlier that both, $D_{D_x e(X)} \mu_{\mathcal{G}}((f\nu)_T(X))$ and $D_{D_x e(X)} \mu_{\mathcal{G}}(f\nu_T(X)) = f D_{D_x e(X)} \mu_{\mathcal{G}}(\nu_T(X))$, are elements of $V\mathcal{G}$, so that

$$D_{D_x e(X)} \mu_{\mathcal{G}} \left(\nu^a \frac{\partial}{\partial \xi^a} \Big|_{D_x e(X)} \right)$$

is vertical, too.

Making use of the aforementioned isomorphism $V\mathcal{G} \cong \pi_{T\mathcal{G}}^* \mathcal{G}$, we can write similar as in Remark A.4

$$\frac{\partial}{\partial \xi^a} \Big|_{D_x e(X)} \cong (D_x e(X), e_a),$$

where $(e_a)_a$ is a local frame of $\mathcal{G}$ dual to ξ^a . Thus,

$$\text{pr}_2 \left(\frac{\partial}{\partial \xi^a} \Big|_{D_x e(X)} \right) = e_a.$$

By definition, $\mu_{\mathcal{G}}$ acts as identity on $\mathcal{G} = e^* V\mathcal{G}$, and so $D\mu_{\mathcal{G}}$ is the identity on $T\mathcal{G}$. Thus, we derive

$$\text{pr}_2 \left(D_{D_x e(X)} \mu_{\mathcal{G}} \left(\nu^a \frac{\partial}{\partial \xi^a} \Big|_{D_x e(X)} \right) \right) = \text{pr}_2 \left(\nu^a \frac{\partial}{\partial \xi^a} \Big|_{D_x e(X)} \right) = \nu^a e_a = \nu,$$

thus, finally,

$$\nabla_X^{\mathcal{G}}(f\nu)|_x = f(x) \nabla_X^{\mathcal{G}}(f\nu)|_x + X(f) \nu.$$

This finishes the proof. ■

Definition 6.33: LGB connection on its LAB

Let $\mathcal{G} \xrightarrow{\pi_{\mathcal{G}}} M$ be an LGB over a smooth manifold M , and $H\mathcal{G}$ be a horizontal distribution of $\mathcal{G}$. Then we call the vector bundle connection $\nabla^{\mathcal{G}}$ on $\mathcal{G}$ of Prop. Proposition 6.31, shortly denoted by

$$\nabla^{\mathcal{G}} \nu = \frac{d}{dt} \Big|_{t=0} \Delta e^{t\nu}$$

for all $\nu \in \Gamma(\mathcal{G})$, the **$\mathcal{G}$ -connection (on the LAB $\mathcal{G}$)**.

Remarks 6.34.

Such a construction of vector bundle connections on LABs also arises in [12, §4.5, Prop. 4.22], but w.r.t. a more specific $H\mathcal{G}$, see Subsubsection Section 6.4.1 later; in this context that reference also shows that the parallel transport associated to $\nabla^{\mathcal{G}}$ is the infinitesimal version of the parallel transport associated with $H\mathcal{G}$.

Example 6.35: Canonical flat $\mathcal{G}$ -connection

If we again focus on trivial LGBs with their canonical flat connection as in Remark Remark 6.22 and Remark 6.28, we can quickly derive that $\nabla^{\mathcal{G}}$ is then the canonical flat connection on the trivial LAB $\mathcal{G} = M \times \mathfrak{g}$ of $\mathcal{G} = M \times G$, $\mathfrak{g}$ the Lie algebra of the Lie group G . Let ν be a constant section of $\mathcal{G}$, then $e^{t\nu}$ is a constant section of $\mathcal{G}$ for all $t \in \mathbb{R}$, so that

by Remark Remark 6.28

$$\Delta^{\mathcal{G}} e^{t\nu} = 0,$$

and thus

$$\nabla^{\mathcal{G}} \nu = 0$$

for all constant sections $\nu \in \Gamma(\mathcal{G})$. By the uniqueness of the canonical flat connection (w.r.t. a trivialisation) we conclude that $\nabla^{\mathcal{G}}$ is the canonical flat connection on $\mathcal{G} = M \times \mathfrak{g}$.

Remark 6.36: Pullback situation: Part III

Following Remark Remark 6.27 (see Remark Remark 6.19 for the initial setup) and all the involved notation we can again derive something related to the pullback LGB $f^*\mathcal{G}$. First of all, let us denote the exponential of $f^*\mathcal{G}$ and $\mathcal{G}$ by $e_{f^*\mathcal{G}}$ and $e_{\mathcal{G}}$, respectively, then we clearly have

$$e_{f^*\mathcal{G}}^{(p, \nu_x)} = \exp_{(f^*\mathcal{G})_p}(p, \nu_x) = (p, \exp_{\mathcal{G}_x}(\nu_x)) = (p, e_{\mathcal{G}}^{\nu_x})$$

for all $(p, \nu_x) \in (f^*\mathcal{G})_p$, where $x := f(p)$ and so $\nu_x \in \mathcal{G}_x$, and where we made use of that $(f^*\mathcal{G})_p \cong \mathcal{G}_x$ as Lie groups via the projection pr_2 onto the second component (recall Cor. Corollary 2.7), that is, we have used the well-known relation

$$\text{pr}_2\left(\exp_{(f^*\mathcal{G})_p}(p, \nu_x)\right) = (\exp_{\mathcal{G}_x} \circ D_{(p, e_x)} \text{pr}_2)(p, \nu_x) = \exp_{\mathcal{G}_x}(\nu_x),$$

see e.g. [17, §1.7, Thm. 1.7.16, page 59] for the general formula; then use that $\text{pr}_2|_{(f^*\mathcal{G})_p}$ is bijective. Hence, we get

$$e_{f^*\mathcal{G}}^{f^*\nu} = f^*(e_{\mathcal{G}}^{\nu})$$

for all $\nu \in \Gamma(\mathcal{G})$. Due to clash of notation we relabel the projection $V\mathcal{G} \rightarrow \mathcal{G}$ in Prop. Proposition 6.31 to π_2 , and then observe, by using Eq. (18),

$$\begin{aligned} \nabla_Y^{f^*\mathcal{G}}(f^*\nu)\Big|_p &= \pi_2\left(\frac{d}{dt}\Big|_{t=0}\left(\left(\Delta^{f^*\mathcal{G}} e_{f^*\mathcal{G}}^{t \cdot f^*\nu}\right)_p(Y)\right)\right) \\ &= \pi_2\left(\frac{d}{dt}\Big|_{t=0}\left(\left(\Delta^{f^*\mathcal{G}}(f^*(e_{\mathcal{G}}^{t\nu}))\right)_p(Y)\right)\right) \\ &= \pi_2\left(\frac{d}{dt}\Big|_{t=0}\left(\left(f^!(\Delta^{\mathcal{G}} e_{\mathcal{G}}^{t\nu})\right)_p(Y)\right)\right) \\ &= \pi_2\left(\frac{d}{dt}\Big|_{t=0}\left(p, (\Delta^{\mathcal{G}} e_{\mathcal{G}}^{t\nu})_{f(p)}(D_p f(Y))\right)\right) \\ &= \pi_2\left(0, \frac{d}{dt}\Big|_{t=0}\left((\Delta^{\mathcal{G}} e_{\mathcal{G}}^{t\nu})_{f(p)}(D_p f(Y))\right)\right) \\ &= \pi_2\left(\frac{d}{dt}\Big|_{t=0}\left((\Delta^{\mathcal{G}} e_{\mathcal{G}}^{t\nu})_{f(p)}(D_p f(Y))\right)\right) \\ &= \nabla_{D_p f(Y)}^{\mathcal{G}} \nu\Big|_{f(p)} \end{aligned}$$

for all $\nu \in \Gamma(\mathcal{G})$, $p \in N$ and $Y \in T_p N$. By the uniqueness of pullback vector bundle connections we therefore get

$$\nabla^{f^*\mathcal{G}} = f^*\nabla^{\mathcal{G}}.$$

6.2.3. First step towards associated bundles. In order to provide a concise definition of connection 1-forms on principal bundles, we will need to introduce some canonical form of action which will be obviously related to the action needed for an analogue to the notion of associated

bundles related to typical principal bundles. However, we will neither discuss nor introduce a more general notion of associated bundles in this paper; so, there will be just the "first step". The following action can be seen as another canonical action on the pullback of a principal bundle, but coming from $\mathcal{G}$ itself instead of its pullback, if $\mathcal{G}$ acts on the manifold one makes a pullback to.

Proposition 6.37: Canonical LGB action on pullback bundles over principal LGB-bundles

Let $\mathcal{G} \xrightarrow{\pi_{\mathcal{G}}} M$ be an LGB over a smooth manifold M and $\mathcal{P} \xrightarrow{\pi} M$ a principal $\mathcal{G}$ -bundle. Furthermore let N be another smooth manifold and $f : N \rightarrow M$ a smooth map on which $\mathcal{G}$ acts on the left as in Def. Definition 3.1. Then on the pullback manifold $\mathcal{P} \times_M N := f^*\mathcal{P}$, whose pairs of points are now reordered as in

$$\mathcal{P} \times_M N := \{(p, x) \in \mathcal{P} \times N \mid \pi(p) = f(x)\},$$

we have a right $\mathcal{G}$ -action given by

$$(\mathcal{P} \times_M N) * \mathcal{G} \rightarrow \mathcal{P} \times_M N,$$

$$(p, x, g) \mapsto (p, x) \cdot g := (p \cdot g, g^{-1} \cdot x)$$

where $(\mathcal{P} \times_M N) * \mathcal{G}$ is given as pullback of $\mathcal{G}$ w.r.t. the map $\bar{\pi} : \mathcal{P} \times_M N \rightarrow M$, $(p, x) \mapsto \pi(p)$.

Proof.

By Def. Definition 3.1 it is clear that this action is well-defined, due to

$$\pi(p \cdot g) = \pi_{\mathcal{G}}(g) = f(g^{-1} \cdot x)$$

for all $(p, x, g) \in (\mathcal{P} \times_M N) * \mathcal{G}$, and this also implies

$$\bar{\pi}((p, x) \cdot g) = \pi(p \cdot g) = \pi_{\mathcal{G}}(g),$$

so that Eq. (??) is satisfied. By construction we also have a smooth action, since it is the composition of maps

$$(\mathcal{P} \times_M N) * \mathcal{G} \rightarrow (\mathcal{P} * \mathcal{G}) \times (\mathcal{G} * N) \rightarrow \mathcal{P} \times_M N,$$

$$(p, x, g) \mapsto ((p, g), (g, x)) \mapsto (p \cdot g, g^{-1} \cdot x).$$

The first arrow is clearly an embedding, and the second arrow is just the restriction of the smooth diagonal action,

$$(\mathcal{P} * \mathcal{G}) \times (\mathcal{G} * N) \rightarrow \mathcal{P} \times N,$$

$$((p, g), (q, p)) \mapsto (p \cdot g, q^{-1} \cdot p),$$

onto an embedded submanifold, and for its image make use of that $\mathcal{P} \times_M N$ is an embedded submanifold of $\mathcal{P} \times N$, as we already did several times for pullback bundles.

Associativity follows simply by

$$((p, x) \cdot g) \cdot q = (p \cdot g, g^{-1} \cdot x) \cdot q = (p \cdot gq, (gq)^{-1} \cdot x) = (p, x) \cdot (gq)$$

for all $g, q \in \mathcal{G}_y$ ($y := \pi(p) = f(x)$); it is trivial to check that $(p, x) \cdot e_y = (p, x)$. ■

As mentioned in Remark Remark 6.14, constructions of quotients related to the LGB action may be possible here. Thus, using the last proposition, one should be able to construct associated bundles in this more general setting. We mainly need this proposition for the following examples.

Example 6.38: Adjoint action on the vertical bundle of $\mathcal{P}$

Recall the adjoint representation of $\mathcal{G}$ on its LAB $\mathcal{Q} \xrightarrow{\pi_{\mathcal{Q}}} M$, Ex. Example 5.4. Using the notation of Prop. Proposition 6.37 we have a right $\mathcal{G}$ -action on $\mathcal{P} \times_M \mathcal{Q} = \{(p, v) \in$

$\mathcal{P} \times \mathcal{G} \mid \pi(p) = \pi_{\mathcal{G}}(v)\}$ given by

$$(p, v) \cdot g := (p \cdot g, \text{Ad}_{g^{-1}}(v))$$

for all $p \in \mathcal{P}_x$ ($x \in M$), $v \in \mathcal{G}_x$ and $g \in \mathcal{G}_x$. Observe that $\mathcal{P} \times_M \mathcal{G} = \pi^* \mathcal{G}$ which is isomorphic to $V\mathcal{P}$ by Cor. [Corollary 6.17](#). We will denote this action shortly by

$$\text{Ad}_{g^{-1}}(p, v) := (p, v) \cdot g,$$

the **adjoint representation of $\mathcal{G}$ on $V\mathcal{P}$** ; not to be confused with the adjoint representation of $\pi^* \mathcal{G}$ on $\pi^* \mathcal{G}$. In fact, it is trivial to check that $\text{Ad} : \mathcal{G} \rightarrow \text{Aut}(\pi^* \mathcal{G})$, $g \mapsto \text{Ad}_g$, is a $\mathcal{G}$ -representation on $\mathcal{P} \times_M \mathcal{G} = \pi^* \mathcal{G} \cong V\mathcal{P}$ in the sense of Cor. ??.

The quotient of $\mathcal{P} \times_M \mathcal{G}$ w.r.t. this group action should lead to a structure which is the generalization of the adjoint bundle in typical formulations of gauge theory.

Similarly, we can define the conjugation action on its integrated bundle; recall the conjugation defined in Def. [Definition 4.8](#). Especially observe that the conjugation defines a left $\mathcal{G}$ -action on itself by

$$\begin{aligned} \mathcal{G} * \mathcal{G} &\rightarrow \mathcal{G}, \\ (g, q) &\mapsto c_g(q) = gqg^{-1}. \end{aligned}$$

Example 6.39: Conjugation action over $\mathcal{P}$

Let us look at $\mathcal{P} \times_M \mathcal{G} \cong \pi^* \mathcal{G}$. Then we have a right $\mathcal{G}$ -action on $\mathcal{P} \times_M \mathcal{G}$ given by

$$(p, q) \cdot g := (p \cdot g, c_{g^{-1}}(q)) = (p \cdot g, g^{-1}qg)$$

for all $p \in \mathcal{P}_x$ ($x \in M$), and $q, g \in \mathcal{G}_x$. We will denote this also by

$$c_{g^{-1}}(p, q) := (p, q) \cdot g,$$

the **conjugation of $\mathcal{G}$ on the integral of $V\mathcal{P}$** .

Taking a quotient of $\mathcal{P} \times_M \mathcal{G}$ over this action should lead to a generalization of inner group bundles as introduced in Ex. [Example 2.6](#).

Let us finally define connections on principal bundles.

6.2.4. Generalized connection 1-forms on principal bundles. As also stated in [[17](#), §5.1, Prop. 5.1.5, page 260], for a given horizontal distribution $H\mathcal{P}$ of a principal $\mathcal{G}$ -bundle $\mathcal{P} \xrightarrow{\pi} M$ over a smooth manifold M we have by construction that

$$D_p \pi|_{H_p \mathcal{P}} : H_p \mathcal{P} \rightarrow T_x M$$

is a vector space isomorphism for all $x \in M$ and $p \in \mathcal{P}_x$; similarly for $\mathcal{G}$ itself. Using that, one has some sort of identification between the horizontal tangent spaces; we want to provide another identification in the sense of Eq. (16), also recall Rem. [Remark 6.30](#); especially recall the Darboux derivative introduced in Subsubsection [Section 6.2.2](#). The following proposition and definition will be needed.

Proposition 6.40: The right-pushforward modified by the Darboux derivative is a well-defined isomorphism

Let $\mathcal{G} \rightarrow M$ be an LGB over a smooth manifold M and $\mathcal{P} \xrightarrow{\pi} M$ a principal $\mathcal{G}$ -bundle, and we have horizontal a distribution $H\mathcal{G}$ on $\mathcal{G}$. Furthermore, for $g \in \mathcal{G}_x$ ($x \in M$) define the map

$$\begin{aligned} T\mathcal{P}|_{\mathcal{P}_x} &\rightarrow T\mathcal{P}|_{\mathcal{P}_x}, \\ X &\mapsto r_{g*}^*(X) := D_p r_{\sigma}(X) - \overline{(\pi^! \Delta \sigma)|_p(X)} \Big|_{p \cdot g}, \end{aligned}$$

where $p \in \mathcal{P}_x$, $X \in T_p\mathcal{P}$, and σ is any (local) section of $\mathcal{G}$ with $\sigma_x = g$. Then $\mathfrak{r}_{g*}$ is independent of the choice of the local section σ , and it is a vector bundle automorphism over the right-translation r_g .

Furthermore, if we also have a horizontal distribution $H\mathcal{P}$ on $\mathcal{P}$, then $H_p\mathcal{P}$ is isomorphic via $\mathfrak{r}_{g*}$ to a complement of $V_{p,g}\mathcal{P}$ in $T_{p,g}\mathcal{P}$ (this complement is not necessarily $H_{p,g}\mathcal{P}$).

Proof.

In the following we have $(p, g) \in \mathcal{P} * \mathcal{G}$, $X \in T_p\mathcal{P}$, and σ is any (local) section of $\mathcal{G}$ with $\sigma_x = g$ ($x := \pi(p)$).

• Let $\pi_{\mathcal{G}}$ be the projection of $\mathcal{G}$, then as mentioned before Prop. [Proposition 6.40](#) there is a unique $Y \in H_g\mathcal{G}$ with

$$D_g\pi_{\mathcal{G}}(Y) = D_p\pi(X) \in T_xM.$$

By Remark [Remark 6.29](#) we can derive

$$D_{(p,g)}\Phi(X, Y) = \mathfrak{r}_{g*}(X),$$

where we made use of that Y is horizontal in $\mathcal{G}$ so that $(\mu_{\mathcal{G}})_g(Y) = 0$, and where Φ denotes the right $\mathcal{G}$ -action on $\mathcal{P}$ as a map $\mathcal{P} * \mathcal{G} \rightarrow \mathcal{P}$. Therefore the independence of $\mathfrak{r}_{g*}$ w.r.t. the choice of σ follows.

• $\mathfrak{r}_{g*}$ is clearly linear by construction. Let us now show fibre-wise that $\mathfrak{r}_{g*}$ is injective, then it is also bijective by dimensional reasons. By Remark [Remark 3.7](#) we know that r_{σ} is a diffeomorphism, and thus $D_pr_{\sigma} : T_p\mathcal{P} \rightarrow T_{p,g}\mathcal{P}$ is a vector space isomorphism. Restricted onto the vertical subspace $V_p\mathcal{P}$ we have by Remark ??

$$D_pr_{\sigma}|_{V_p\mathcal{P}} = D_pr_g : V_p\mathcal{P} \rightarrow V_{p,g}\mathcal{P},$$

which is a vector space isomorphism by Cor. [Corollary 6.17](#) and dimensional reasons (which implies that D_pr_g is surjective and thus bijective). We rewrite D_pr_{σ} as

$$\begin{aligned} H_p\mathcal{P} \oplus V_p\mathcal{P} &\rightarrow T_{p,g}\mathcal{P}, \\ (X^H, X^V) &\mapsto D_pr_{\sigma}|_{H_p\mathcal{P}}(X^H) + D_pr_g(X^V), \end{aligned}$$

where we fix just any horizontal distribution $H\mathcal{P}$. By dimensional reasons and due to the bijectivity of D_pr_{σ} and D_pr_g the image $\text{Im}(D_pr_{\sigma}|_{H_p\mathcal{P}})$ of $D_pr_{\sigma}|_{H_p\mathcal{P}}$ has to be a complement subspace of $V_{p,g}\mathcal{P}$ in $T_{p,g}\mathcal{P}$; $\text{Im}(D_pr_{\sigma}|_{H_p\mathcal{P}})$ is not necessarily equal to $H_{p,g}\mathcal{P}$ in general but of the same dimension, especially of the same dimension as $H_p\mathcal{P}$. Thus, $D_pr_{\sigma}|_{H_p\mathcal{P}}$ is a vector space isomorphism onto its image which is complementary to $V_{p,g}\mathcal{P}$. Furthermore, observe

$$(\pi^!\Delta\sigma)|_p(X) = (\Delta\sigma)|_x(D_p\pi(X^H + X^V)) = (\Delta\sigma)|_x(D_p\pi(X^H)) = (\pi^!\Delta\sigma)|_p(X^H),$$

using that the vertical subbundle is given by the kernel of $D\pi$, and where we split again $X = X^H + X^V$ for all $X \in T_p\mathcal{P} = H_p\mathcal{P} \oplus V_p\mathcal{P}$. Hence, we get in total that $\mathfrak{r}_{g*}$ at p is equivalent to

$$\begin{aligned} H_p\mathcal{P} \oplus V_p\mathcal{P} &\rightarrow \text{Im}(D_pr_{\sigma}|_{H_p\mathcal{P}}) \oplus V_{p,g}\mathcal{P}, \\ (X^H, X^V) &\mapsto \left(D_pr_{\sigma}|_{H_p\mathcal{P}}(X^H), D_pr_g(X^V) - \overline{(\pi^!\Delta\sigma)|_p(X^H)} \right). \end{aligned}$$

Let (X^H, X^V) be now in the kernel of $\mathfrak{r}_{g*}$, then $X^H = 0$ by the bijectivity of $D_pr_{\sigma}|_{H_p\mathcal{P}}$ onto its image. The second component of $\mathfrak{r}_{g*}$ is then just $D_pr_g(X^V)$; again by the bijectivity of $D_pr_g : V_p\mathcal{P} \rightarrow V_{p,g}\mathcal{P}$ we also get $X^V = 0$. Thus, $\mathfrak{r}_{g*}$ is injective and therefore defines vector space isomorphisms $T_p\mathcal{P} \rightarrow T_{p,g}\mathcal{P}$. It follows that $\mathfrak{r}_{g*}$ is an automorphism of $T\mathcal{P}|_{\mathcal{P}_x}$ over r_g , independent of the choice of $H\mathcal{P}$.

- It is then clear that $\mathcal{r}_{g*}|_{H_p\mathcal{P}}$, given by

$$\begin{aligned} H_p\mathcal{P} &\rightarrow \text{Im}\left(D_p r_\sigma|_{H_p\mathcal{P}}\right) \oplus V_{p,g}\mathcal{P}, \\ X^H &\mapsto \left(D_p r_\sigma|_{H_p\mathcal{P}}(X^H), -\overline{(\pi^! \Delta \sigma)|_p(X^H)}\Big|_{p,g}\right), \end{aligned}$$

is also an isomorphism onto its image; its image is complementary to $V_{p,g}\mathcal{P}$ because its intersection with $V_{p,g}\mathcal{P}$ would require that

$$D_p r_\sigma|_{H_p\mathcal{P}}(X^H) = 0,$$

which implies that $X^H = 0$ due to the fact that $D_p r_\sigma|_{H_p\mathcal{P}}$ is a vector space isomorphism onto its image, as we discussed before. But then

$$\mathcal{r}_{g*}|_{H_p\mathcal{P}}(0) = 0,$$

which implies that the image of $\mathcal{r}_{g*}|_{H_p\mathcal{P}}$ is a complement of $V_{p,g}\mathcal{P}$ in $T_{p,g}\mathcal{P}$. This finishes the proof. $\blacksquare$

Hence, we formally define:

Definition 6.41: Modified pushforward via right-translation

Let $\mathcal{G} \rightarrow M$ be an LGB over a smooth manifold M and $\mathcal{P} \xrightarrow{\pi} M$ a principal $\mathcal{G}$ -bundle, and we have a horizontal distribution $H\mathcal{G}$ on $\mathcal{G}$. Then we define the **modified right-pushforward**^a $\mathcal{r}_{g*}$ (**with** $g \in \mathcal{G}_x$, $x \in M$) as the vector bundle isomorphism $T\mathcal{P}|_{\mathcal{P}_x} \rightarrow T\mathcal{P}|_{\mathcal{P}_x}$ over r_g as given in Prop. [Proposition 6.40](#) by

$$\mathcal{r}_{g*}(X) := D_p r_\sigma(X) - \overline{(\pi^! \Delta \sigma)|_p(X)}\Big|_{p,g}$$

for all $p \in \mathcal{P}_x$ and $X \in T_p\mathcal{P}$, where σ is any (local) section of $\mathcal{G}$ with $\sigma_x = g$.

Similarly, for a (local) section σ of $\mathcal{G}$ we define the **modified right-pushforward** $\mathcal{r}_{\sigma*}$ **with** σ as a (local) vector bundle isomorphism $T\mathcal{P} \rightarrow T\mathcal{P}$ by

$$\mathcal{r}_{\sigma*}(X) := \mathcal{r}_{\sigma_x*}(X) = D_p r_\sigma(X) - \overline{(\pi^! \Delta \sigma)|_p(X)}\Big|_{p,\sigma_x}$$

for all $X \in T_p\mathcal{P}$ ($p \in \mathcal{P}_x$).

^aThe font of $\mathcal{r}$ is a calligraphic r.

Remark 6.42: Restriction onto vertical bundle gives typical right-pushforward

It is trivial to check that we have

$$\mathcal{r}_{g*}(X) = D_p r_g(X) =: r_{g*}(X)$$

for all $X \in V_p\mathcal{P}$; we actually have proven this directly after Eq. (16). As also pinpointed in Remark [Remark 6.30](#), if $\mathcal{G}$ is a trivial LGB equipped with its canonical flat connection and σ a constant section, then it is easy to see that $\mathcal{r}_{\sigma*} = \mathcal{r}_{g*} = r_{g*}$, where r_g has to be understood as the right-translation of the canonical Lie group action inherited by $\mathcal{G}$ in the sense of Ex. [Example 3.10](#).

Remarks 6.43.

Prop. [Proposition 6.40](#) trivially extends to $\mathcal{r}_{\sigma*}$, that is, $\mathcal{r}_{\sigma*}$ is a (local) automorphism of $T\mathcal{P}$ over r_σ . Thus, we can view $\mathcal{r}_{\sigma*}$ as an element of $\Omega^1(\mathcal{P}; r_\sigma^* T\mathcal{P})$.

Hence, the following definition makes sense, especially if thinking about what we discussed for Eq. (16), also recall Rem. [Remark 6.30](#).

Definition 6.44: Ehresmann connection on principal LGB-bundles

Let $\mathcal{G} \rightarrow M$ be an LGB over a smooth manifold M and $\mathcal{P} \xrightarrow{\pi} M$ a principal $\mathcal{G}$ -bundle, and we have horizontal distributions $H\mathcal{G}$ and $H\mathcal{P}$ on $\mathcal{G}$ and $\mathcal{P}$, respectively. We call $H\mathcal{P}$ an **Ehresmann connection** or a **connection on $\mathcal{P}$** if it is **right-invariant (w.r.t. modified right-pushforward)**, *i.e.*

$$r_{g*}(H_p\mathcal{P}) = H_{p \cdot g}\mathcal{P}$$

for all $p \in \mathcal{P}_x$ and $g \in \mathcal{G}_x$ ($x := \pi(p)$).

Remarks 6.45.

There is also a definition of connections on such and more general principal bundles in [11, §5.7, paragraph before Prop. 5.38, page 148]. However, this reference provides a different type of definition; translated to our situation, it is based on assuming that $\mathcal{G}$ is defined over another base manifold N . In order to define the LGB action on $\mathcal{P}$ this reference introduces a moment map $\mu : \mathcal{P} \rightarrow N$ so that the action of an element $\mathcal{G}_y$ ($y \in N$) is defined on $\mu^{-1}(\{y\})$, especially the infinitesimal action r_g of a fixed LGB element g acts on tangent vectors of $\mu^{-1}(\{y\})$, not necessarily on the vertical structure of $\mathcal{P}$. Hence, in order to circumvent the problem we discussed in Subsubsection Section 6.2.1, the reference's definition of a connection is then based on assuming that the "fibres" $\mu^{-1}(\{y\})$ are complementary to the vertical structure of $\mathcal{P}$, so these fibres' tangent spaces define a horizontal distribution while the infinitesimal action r_g now acts well-defined on the horizontal structure. Henceforth, the reference does not need to look at using sections σ and their actions.

However, this is not a suitable definition for us, because our moment map is the projection of $\mathcal{P}$ itself such that r_g acts on the vertical structure. The reference's definition is rather restrictive, while our definition works for all principal $\mathcal{G}$ -bundles by fixing a connection on $\mathcal{G}$.

Let us discuss several examples.

Example 6.46: Recovering of the classical definition

As we already discussed several times, especially recall Remark Remark 6.42 and Ex. Example 6.3, but ultimately by the discussion for Eq. (16) which also applies to classical principal bundles and their notion of connection and parallel transport: We recover the typical definition of a connection on a principal bundle if $\mathcal{P}$ is a classical principal bundle and if the trivial LGB $\mathcal{G}$ is equipped with its canonical flat connection $H\mathcal{G}$. We call such a connection a **classical connection**.

Example 6.47: Associated LGBs

Recall Def. ??, and recall that LGBs themselves are principal bundles as in Ex. Example 6.4. That is, let G, H be Lie groups, $P \rightarrow M$ a principal G -bundle over a smooth manifold M , and ψ a G -representation on H . Then we have the LGB associated to the principal bundle P and the representation ψ on H

$$\begin{array}{ccc} H & \longrightarrow & \mathcal{H} := P \times_{\psi} H \\ & & \downarrow \\ & & M \end{array}$$

Fix a typical classical connection on P ; as also introduced in Subsubsection Section 6.2.1 we have an associated parallel transport $\text{PT}_{\alpha}^P : P_x \rightarrow \Gamma(\alpha^*P)$ along a curve $\alpha : [0, t] \rightarrow M$ ($t > 0$), where $\alpha(0) =: x$. As proven in [17, §5.9, Thm. 5.9.1, page 289f.], we have a canonical well-defined parallel transport $\text{PT}_{\alpha}^{\mathcal{H}} : \mathcal{H}_x \rightarrow \Gamma(\alpha^*\mathcal{H})$ given by

$$\text{PT}_{\alpha}^{\mathcal{H}}([p, h]) = [\text{PT}_{\alpha}^P(p), h]$$

for all $[p, h] \in \mathcal{H}_x$. This has a 1:1 correspondence to a horizontal distribution $H\mathcal{H}$ on $\mathcal{H}$. Observe that we have

$$\begin{aligned} \text{PT}_\alpha^\mathcal{H}([p, h] \cdot [p, h']) &= \text{PT}_\alpha^\mathcal{H}([p, hh']) \\ &= [\text{PT}_\alpha^P(p), hh'] \\ &= [\text{PT}_\alpha^P(p), h] \cdot [\text{PT}_\alpha^P(p), h'] \\ &= \text{PT}_\alpha^\mathcal{H}([p, h]) \cdot \text{PT}_\alpha^\mathcal{H}([p, h']) \end{aligned}$$

for all $[p, h], [p, h'] \in \mathcal{H}_x$. Now recall that the whole motivation behind Def. [Definition 6.44](#) comes from Eq. (16) (also recall Remark [Remark 6.30](#)) which itself stems from Eq. (15). We see that that Eq. (15) is satisfied here, and thus Eq. (16) follows, *i.e.*

$$\mathcal{H}_{[p, h']_*}(X) \in H_{[p, h] \cdot [p, h']} \mathcal{H}$$

for all $X \in H_{[p, h]} \mathcal{H}$. By Prop. [Proposition 6.40](#) and dimensional reasons (horizontal subspaces are of the same dimension) it follows immediately that $H\mathcal{H}$ is an (Ehresmann) connection on $\mathcal{H}$ in sense of Def. [Definition 6.44](#).

We call such connections on LGBs associated to a classical principal bundle an **associated connection**.

As expected, we have a corresponding connection 1-form. For this we need to formally define the pullback of 1-forms with respect to the modified right-pushforward.

Definition 6.48: The pullback of forms via modified right-pushforward

Let $\mathcal{G} \rightarrow M$ be an LGB over a smooth manifold M and $\mathcal{P} \xrightarrow{\pi} M$ a principal $\mathcal{G}$ -bundle, and we have a fixed horizontal distribution $H\mathcal{G}$ on $\mathcal{G}$. For $\omega \in \Omega^k(\mathcal{P}; \pi^* \mathcal{G})$ ($k \in \mathbb{N}_0$) we define the **pullback via the modified right-pushforward** $\mathcal{H}_g^!(\omega|_{\mathcal{P}_x})$ (with $g \in \mathcal{G}_x$, over $\mathcal{P}_x$, $x \in M$) as an element of $\Gamma(\Lambda^k T^* \mathcal{P}|_{\mathcal{P}_x} \otimes \mathcal{G}_x)$ by

$$\left(\mathcal{H}_g^!(\omega|_{\mathcal{P}_x}) \right) \Big|_p (Y_1, \dots, Y_k) := \omega_{p \cdot g}(\mathcal{H}_{g*}(Y_1), \dots, \mathcal{H}_{g*}(Y_k))$$

for all $p \in \mathcal{P}_x$ and $Y_1, \dots, Y_k \in T_p \mathcal{P}$. Similarly we define the **pullback via the modified right-pushforward** $\mathcal{H}_\sigma^! \omega$ with a (local) section $\sigma \in \Gamma(\mathcal{G})$ as an element of $\Omega^k(\mathcal{P}; \pi^* \mathcal{G})$ by

$$(\mathcal{H}_\sigma^! \omega) \Big|_p (Y_1, \dots, Y_k) := \left(\mathcal{H}_{\sigma_x}^!(\omega|_{\mathcal{P}_x}) \right) \Big|_p (Y_1, \dots, Y_k) = (r_\sigma^* \omega) \Big|_p (\mathcal{H}_{\sigma*}(Y_1), \dots, \mathcal{H}_{\sigma*}(Y_k))$$

for all $p \in \mathcal{P}_x$ and $Y_1, \dots, Y_k \in T_p \mathcal{P}$.

Remarks 6.49.

By Prop. [Proposition 6.40](#) (and Remark [Theorem 6.43](#)) these definitions are well-defined. A short note about the notation on the very right hand side of the second definition: This notation allows us to extend it to vector fields, that is,

$$(\mathcal{H}_\sigma^! \omega)(Y_1, \dots, Y_k) = (r_\sigma^* \omega)(\mathcal{H}_{\sigma*}(Y_1), \dots, \mathcal{H}_{\sigma*}(Y_k))$$

for all $p \in \mathcal{P}_x$ and $Y_1, \dots, Y_k \in \mathfrak{X}(\mathcal{P})$. Observe that we have $\mathcal{H}_{\sigma*}(Y_l) \in \Gamma(r_\sigma^* T \mathcal{P})$ ($l \in \{1, \dots, k\}$) and $r_\sigma^* \omega \in \Gamma(\Lambda^k r_\sigma^* T \mathcal{P} \otimes \pi^* \mathcal{G})$, using $r_\sigma^* \pi^* \mathcal{G} \cong (\pi \circ r_\sigma)^* \mathcal{G} = \pi^* \mathcal{G}$, such that the right hand side is well-defined.

For the following definition recall the adjoint $\mathcal{G}$ -representation on $V\mathcal{P}$, Ex. [Example 6.38](#), and the isomorphism for $V\mathcal{P}$ in Cor. [Corollary 6.17](#).

Definition 6.50: Connection 1-forms on principal LGB-bundles

Let $\mathcal{G} \rightarrow M$ be an LGB over a smooth manifold M and $\mathcal{P} \xrightarrow{\pi} M$ a principal $\mathcal{G}$ -bundle, and we have a fixed horizontal distribution $\mathcal{H}\mathcal{G}$ on $\mathcal{G}$. A **connection 1-form** or **gauge field** on $\mathcal{P}$ is a 1-form $A \in \Omega^1(\mathcal{P}; \pi^*\mathfrak{g})$ satisfying:

- ($\mathcal{G}$ -equivariance, but w.r.t. modified right-pushforward)

$$\mathcal{r}_\sigma^! A = \mathcal{A}_{\sigma^{-1}} \circ A$$

for all (local) $\sigma \in \Gamma(\mathcal{G})$.

- (Identity on $V\mathcal{P}$)

$$A(\bar{\nu}) = \pi^* \nu$$

for all (local) $\nu \in \Gamma(\mathfrak{g})$.

Remarks 6.51.

- Due to the fact that we formulated the $\mathcal{G}$ -equivariance using Ex. [Example 6.38](#), one may already be able to show an analogue of the 1:1 correspondence of typical (classical) connection 1-forms to connection reforms and splittings of the Atiyah sequence; that is, a generalization of [10, §3.2, page 90 ff.].

- The $\mathcal{G}$ -equivariance reads point-wise for $g := \sigma_x$ ($x \in M$)

$$(\mathcal{r}_g^!(A|_{\mathcal{P}_x}))|_p(X) = (p \cdot g, \text{Ad}_{g^{-1}}(\hat{A}_p(X)))$$

for all $p \in \mathcal{P}_x$ and $X \in T_p\mathcal{P}$, where we wrote $A_p = (p, \hat{A}_p)$ with $\hat{A}_p \in \Gamma(T_p^*\mathcal{P} \otimes \mathfrak{g}_x)$, and Ad is the adjoint representation of $\mathcal{G}$. In the typical formulation of gauge theory the base point component is usually omitted due to that $\mathfrak{g}$ is then trivial and so also $V\mathcal{P}(\cong \pi^*\mathfrak{g})$.

The identity on $V\mathcal{P}$ reads pointwise

$$A_p(\bar{v}_p) = (p, v_p)$$

for all $v_p \in \mathfrak{g}_x$. For readability we may also omit the basepoint information and just write

$$A_p(\bar{v}_p) \equiv v_p.$$

Furthermore, recall the notation introduced in [Remark 6.18](#), then for $\mu := (p, v_p)$ we can therefore write

$$A_p(\bar{\mu}_p) = \mu.$$

Finally, we identify (Ehresmann) connections and connection 1-forms on principal LGB-bundles in the typical way; for this recall [Cor. Corollary 6.17](#).

Theorem 6.52: 1:1 correspondence of Ehresmann connections and connection 1-forms

Let $\mathcal{G} \rightarrow M$ be an LGB over a smooth manifold M and $\mathcal{P} \xrightarrow{\pi} M$ a principal $\mathcal{G}$ -bundle, and let $\mathcal{H}\mathcal{G}$ be a horizontal distribution on $\mathcal{G}$. Then there is a 1:1 correspondence between Ehresmann connections and connection 1-forms on $\mathcal{P}$:

- Let $\mathcal{H}\mathcal{P}$ be an Ehresmann connection on $\mathcal{P}$. Then $\mathcal{H}\mathcal{P}$ defines a connection 1-form $A \in \Omega^1(\mathcal{P}; \pi^*\mathfrak{g})$ by

$$A_p(\bar{v}_p + X_p) = (p, v)$$

for all $p \in \mathcal{P}_x$ ($x \in M$), $v \in \mathfrak{g}_x$ and $X \in \mathcal{H}_p\mathcal{P}$.

- Let $A \in \Omega^1(\mathcal{P}; \pi^*\mathfrak{g})$ be a connection 1-form on $\mathcal{P}$. Then A defines an Ehresmann connection $\mathcal{H}\mathcal{P}$ on $\mathcal{P}$ via its kernel $\text{Ker}(A)$, that is,

$$\mathcal{H}_p\mathcal{P} = \text{Ker}(A_p)$$

for all $p \in \mathcal{P}$.

Proof.

The proof is very similar to the proof of "typical connections", as *e.g.* provided in [17, §5.2, Thm. 5.2.2, page 262].

• For the first bullet point we need to show that Def. Definition 6.50 is satisfied, and the identity behaviour on $V\mathcal{P}$ quickly follows by definition: We have

$$A(\bar{\nu})|_p = A_p(\bar{\nu}_p) = (p, \nu_x) = (\pi^* \nu)|_p$$

for all $\nu \in \Gamma(\mathcal{G})$ and $p \in \mathcal{P}_x$ ($x \in M$). Therefore it is only left to show the $\mathcal{G}$ -equivariance. As mentioned in Remark Remark 5.13, for $v \in \mathcal{G}_x$ the vector field $\bar{v}$ on $\mathcal{P}_x$ is a fundamental vector field coming from the $\mathcal{G}_x$ -action on $\mathcal{P}_x$. Hence, we know by [17, §3.4, Prop. 3.4.6, page 145f.] that

$$r_{g*}(\bar{v}) = r_{g*}(\bar{v}) = \overline{\text{Ad}_{g^{-1}}(v)}$$

for all $g \in \mathcal{G}_x$, also using Remark Remark 6.42.⁹ Thus,

$$\begin{aligned} (r_{\sigma}^! A)|_p(\bar{v}_p + X_p) &= A_{p \cdot \sigma_x} \left(\underbrace{r_{\sigma_x*}(\bar{v}_p)}_{= r_{\sigma_x*}(\bar{v})|_{p \cdot \sigma_x}} + \underbrace{r_{\sigma_x*}(X_p)}_{\in H_{p \cdot \sigma_x} \mathcal{P}} \right) \\ &= (p \cdot \sigma_x, \text{Ad}_{\sigma_x^{-1}}(v)) \\ &= \text{Ad}_{\sigma_x^{-1}}(p, v) \\ &= \text{Ad}_{\sigma_x^{-1}}(A_p(\bar{v}_p + X_p)) \\ &= (\text{Ad}_{\sigma^{-1}} \circ A)|_p(\bar{v}_p + X_p) \end{aligned}$$

for all $p \in \mathcal{P}_x$, $\sigma \in \Gamma(\mathcal{G})$, $v \in \mathcal{G}_x$ and $X_p \in H_p \mathcal{P}$. This finishes the proof for the first bullet point.

• For the second bullet point we make use of Cor. Corollary 6.17 and

$$A(\bar{\nu}) = \pi^* \nu$$

for all (local) sections $\nu \in \Gamma(\mathcal{G})$. This implies that A has not only values in $V\mathcal{P}$, but acts also as identity on $V\mathcal{P}$. Thus, $H\mathcal{P} = \text{Ker}(A_p)$ is a complementary subspace of $V_p \mathcal{P}$ in $T_p \mathcal{P}$ for all $p \in \mathcal{P}$. In order to show that $H\mathcal{P}$ is a horizontal distribution of $\mathcal{P}$, we fix a local frame $(e_a)_a$ of $\mathcal{G}$ over U (an open subset of M), so that $(\pi^* e_a)_a$ is a frame of $\pi^* \mathcal{G}|_{\pi^{-1}(U)}$. Therefore we write

$$A = A^a \otimes \pi^* e_a,$$

where $A^a \in \Omega^1(\mathcal{P}|_U)$. Due to $A(\bar{\nu}) = \pi^* \nu$ we get

$$A^a(\bar{e}_b) = \delta_b^a,$$

where δ_b^a is the Kronecker delta. Hence, $(A^a)_a$ is the dual frame of $(\bar{e}_a)_a$, which is a frame of $V\mathcal{P}$ by Cor. Corollary 6.17, and so the A^a are linear independent to each other. Fix an auxiliary fibre metric $\langle \cdot, \cdot \rangle$ on $T\mathcal{P}$, and denote with $(V^a)_a$ the $\langle \cdot, \cdot \rangle$ -dual frame to $(A^a)_a$, *i.e.*

$$A^a = \langle V^a, \cdot \rangle.$$

Due to the smoothness of A and due to that $\langle \cdot, \cdot \rangle$ is a fibre metric, $(V^a)_a$ are smooth (local) vector fields on $\mathcal{P}$, linear independent to each other. Observe that any orthogonal frame $(W^\alpha)_\alpha$ to $(V^a)_a$ satisfies

$$A^a(W^\alpha) = \langle V^a, W^\alpha \rangle = 0.$$

Thus, we derived that $\text{Ker}(A)|_U$ is spanned by such frames $(W^\alpha)_\alpha$, in total, $\text{Ker}(A)$ is spanned by a locally free sheaf of modules of constant rank. By the 1:1 correspondence of vector bundle and locally free sheaf of modules of constant rank, we can conclude that $H\mathcal{P} = \text{Ker}(A)$ is a subbundle of $T\mathcal{P}$; complementary to $V\mathcal{P}$ due to what we have shown earlier.

⁹This is very straightforward to prove.

It is only left to show the right-invariance of $H\mathcal{P}$. So let $X \in H_p\mathcal{P}$ for $p \in \mathcal{P}$. Then

$$A_{p \cdot \sigma_x}(\nu_{\sigma*}(X)) = (\nu_{\sigma}^!(A))_p(X) = \text{Ad}_{\sigma_x^{-1}}(A_p(X)) = 0$$

for all (local) $\sigma \in \Gamma(\mathcal{G})$, where $x := \pi(p)$. Thence, $\nu_{\sigma*}(X) = \nu_{\sigma_x*}(X) \in H_{p \cdot \sigma_x}\mathcal{P}$, and therefore we derive by Prop. [Proposition 6.40](#) and dimensional reasons that $\nu_{\sigma_x*}(H_p\mathcal{P}) = H_{p \cdot \sigma_x}\mathcal{P}$. We eventually conclude that $H\mathcal{P}$ is an Ehresmann connection. $\blacksquare$

We finish this subsection with the following technical and useful corollary.

Corollary 6.53: Commutation of modified push-forward and projections

Let $\mathcal{G} \rightarrow M$ be an LGB over a smooth manifold M and $\mathcal{P} \rightarrow M$ a principal $\mathcal{G}$ -bundle, also let $H\mathcal{G}$ be a horizontal distribution on $\mathcal{G}$ and $H\mathcal{P}$ an Ehresmann connection on $\mathcal{P}$; denote with π_h and π_v the associated projections on the horizontal and vertical bundle of $\mathcal{P}$, respectively. Then we have

$$\nu_{g*} \circ \pi_h = \pi_h \circ \nu_{g*},$$

$$\nu_{g*} \circ \pi_v = \pi_v \circ \nu_{g*}$$

for all $g \in \mathcal{G}$.

Remarks 6.54.

This extends of course to (local) sections $\sigma \in \Gamma(\mathcal{G})$, and as discussed in Remark [Theorem 6.25](#) we can extend these equations to vector fields on $\mathcal{P}$ via pullbacks, that is,

$$\nu_{\sigma*} \circ \pi_h = r_{\sigma}^* \pi_h \circ \nu_{\sigma*},$$

$$\nu_{\sigma*} \circ \pi_v = r_{\sigma}^* \pi_v \circ \nu_{\sigma*},$$

making use of that $\nu_{\sigma*}$ is an isomorphism over $r_{\sigma*}$, recall Prop. [Proposition 6.40](#).

Proof.

By Cor. [Corollary 6.17](#) (also recall Remark [Remark 6.42](#)) and Def. [Definition 6.44](#) we have

$$\nu_{g*}(V_p\mathcal{P}) = V_{p \cdot g}\mathcal{P},$$

$$\nu_{g*}(H_p\mathcal{P}) = H_{p \cdot g}\mathcal{P},$$

thus, ν_{g*} preserves the splitting $T\mathcal{P} = H\mathcal{P} \oplus V\mathcal{P}$. This concludes the proof. $\blacksquare$

6.3. Gauge transformations. Let us now look at how gauge transformations of A look like in this setting; for this recall the definition of gauge transformations in Def. [Definition 6.5](#). As in Remark ?? we expect a relationship between gauge transformations and certain LGB valued maps. For the following also recall Ex. [Example 6.39](#).

Definition 6.55: LGB-valued conjugation maps

Let $\mathcal{G} \rightarrow M$ be an LGB over a smooth manifold M , and $\mathcal{P} \xrightarrow{\pi} M$ a principal $\mathcal{G}$ -bundle. Then we define the group $C^\infty(\mathcal{P}; \mathcal{G})^\mathcal{G}$ of $\mathcal{G}$ -valued conjugation maps as a set by

$$C^\infty(\mathcal{P}; \mathcal{G})^\mathcal{G} = \{\sigma \in \Gamma(\pi^*\mathcal{G}) \mid \sigma_{p \cdot g} = c_{g^{-1}}(\sigma_p) \text{ for all } (p, g) \in \mathcal{P} * \mathcal{G}\}.$$

Its group structure is inherited by the point-wise group structure of $\Gamma(\pi^*\mathcal{G})$; recall Cor. [Corollary 2.7](#).

Remark 6.56: Group structure on $C^\infty(\mathcal{P}; \mathcal{G})^\mathcal{G}$

It is trivial to check that $C^\infty(\mathcal{P}; \mathcal{G})^\mathcal{G}$ is indeed a subgroup of $\Gamma(\pi^*\mathcal{G})$.

$C^\infty(\mathcal{P}; \mathcal{G})^\mathcal{G}$ canonically acts on $\mathcal{P}$ on the right via the given $\mathcal{G}$ -action: Let $\sigma \in C^\infty(\mathcal{P}; \mathcal{G})^\mathcal{G}$, then we define

$$p \cdot \sigma_p := p \cdot \text{pr}_2(\sigma_p) \quad (21)$$

for all $p \in \mathcal{P}$, where $\text{pr}_2 : \pi^* \mathcal{G} \rightarrow \mathcal{G}$ is the projection onto the second component. In essence, we drop the notation of pr_2 , as if we view σ as a map $\mathcal{P} \rightarrow \mathcal{G}$ of fibre bundles over π ; recall Subsection Section 1.3. If we denote the right $\mathcal{G}$ -action on $\mathcal{P}$ by Φ , then observe by $\sigma_p = (p, \text{pr}_2(\sigma_p))$ that we could also write

$$p \cdot \sigma_p = \Phi(p, \text{pr}_2(\sigma_p)) = \Phi(\sigma_p).$$

As in the case of typical/classical principal bundles, we have the following statement.

Proposition 6.57: Gauge transformations as $\mathcal{G}$ -valued conjugation maps

Let $\mathcal{G} \rightarrow M$ be an LGB over a smooth manifold M , and $\mathcal{P} \xrightarrow{\pi} M$ a principal $\mathcal{G}$ -bundle. Then there is a well-defined group isomorphism of gauge transformations and $\mathcal{G}$ -valued conjugation maps given by

$$\begin{aligned} \text{Aut}(\mathcal{P}) &\rightarrow C^\infty(\mathcal{P}; \mathcal{G})^\mathcal{G}, \\ H &\mapsto \sigma^H, \end{aligned}$$

where $\sigma^H \in C^\infty(\mathcal{P}; \mathcal{G})^\mathcal{G}$ is defined by

$$H(p) = p \cdot \sigma_p^H$$

for all $p \in \mathcal{P}$.

Proof.

This result can be proven similarly as for Lie group based principal bundles; however smoothness needs to be discussed a bit. In the following we make use of that pullback manifolds like $\mathcal{P} * \mathcal{G}$ are embedded submanifolds of product manifolds like $\mathcal{P} \times \mathcal{G}$, we will not further mention it.

• First of all, due to the fact that H is base-preserving we know that $H(p)$ is in the same fibre as p ($p \in \mathcal{P}$), and due to that the $\mathcal{G}$ -action on $\mathcal{P}$ is simply transitive there is a unique element of $\pi^* \mathcal{G}|_p$, denoted by σ_p^H , such that

$$H(p) = p \cdot \sigma_p^H = p \cdot \text{pr}_2(\sigma_p^H),$$

where $\text{pr}_2 : \mathcal{P} * \mathcal{G} = \pi^* \mathcal{G} \rightarrow \mathcal{G}$ is the smooth projection onto the second component. Let us first show smoothness of $p \mapsto \sigma_p^H$: We can write

$$H(p) = \Phi_p(\text{pr}_2(\sigma_p^H)),$$

where $\Phi_p : \mathcal{G}_x \rightarrow \mathcal{P}_x$ is the orbit map through $p \in \mathcal{P}_x$ ($x \in M$). By Remark Remark 6.2, Φ_p is a $\mathcal{G}_x$ -equivariant diffeomorphism, especially invertible, so that

$$\text{pr}_2(\sigma_p^H) = (\Phi_p)^{-1}(H(p)).$$

Let us define the map

$$\begin{aligned} \pi^* \mathcal{P} &\rightarrow \mathcal{G}, \\ (p, p') &\mapsto \Phi^{-1}(p, p') := (\Phi_p)^{-1}(p') \end{aligned}$$

which is a map over π by construction. If we can show smoothness of Φ^{-1} , then smoothness of σ^H follows additionally due to smoothness of H and

$$\sigma_p^H = (p, \text{pr}_2(\sigma_p^H)) = \left(p, (\Phi_p)^{-1}(H(p)) \right) = \left(p, \Phi^{-1}(p, H(p)) \right).$$

Observe that we have

$$\text{pr}_2(p, g) = g = (\Phi_p)^{-1}(\Phi_p(g)) = \Phi^{-1}(p, \Phi(p, g))$$

for all $(p, g) \in \mathcal{P} * \mathcal{G}$, where Φ denotes the right $\mathcal{G}$ -action on $\mathcal{P}$. The map

$$\begin{aligned} L : \mathcal{P} * \mathcal{G} &\rightarrow \pi^* \mathcal{P}, \\ (p, g) &\mapsto (p, \Phi(p, g)), \end{aligned}$$

is a base-preserving principal bundle isomorphism (w.r.t. the LGB isomorphism given as the identity map on $\mathcal{P} * \mathcal{G}$), recall Cor. [Corollary 6.12](#) and Remark [Theorem 6.13](#). Thus, we have in total

$$\Phi^{-1} \circ L = \text{pr}_2,$$

and thus

$$\Phi^{-1} = \text{pr}_2 \circ L^{-1},$$

so Φ^{-1} is smooth, and therefore σ^H is smooth, too.

- That σ^H is an element of $C^\infty(\mathcal{P}; \mathcal{G})^\mathcal{G}$ follows as usual:

$$(p \cdot g) \cdot \text{pr}_2(\sigma_{p \cdot g}^H) = H(p \cdot g) = H(p) \cdot g = p \cdot (\text{pr}_2(\sigma_p^H) \cdot g)$$

for all $(p, g) \in \mathcal{P} * \mathcal{G}$, so that, by the simply transitivity of the action,

$$g \cdot \text{pr}_2(\sigma_{p \cdot g}^H) = \text{pr}_2(\sigma_p^H) \cdot g,$$

and thus

$$\sigma_{p \cdot g}^H = (p \cdot g, \text{pr}_2(\sigma_{p \cdot g}^H)) = (p \cdot g, g^{-1} \cdot \text{pr}_2(\sigma_p^H) \cdot g) = c_{g^{-1}}(\sigma_p^H).$$

Thus, $\sigma^H \in C^\infty(\mathcal{P}; \mathcal{G})^\mathcal{G}$.

- The inverse of $H \mapsto \sigma^H$ is clearly given by

$$\begin{aligned} C^\infty(\mathcal{P}; \mathcal{G})^\mathcal{G} &\rightarrow \mathcal{A}ut(\mathcal{P}), \\ \sigma &\mapsto H^\sigma, \end{aligned}$$

where

$$H^\sigma(p) := p \cdot \sigma_p$$

for all $p \in \mathcal{P}$. Smoothness here is now obvious, it is also clearly base-preserving, and we have

$$\begin{aligned} H^\sigma(p \cdot g) &= p \cdot g \cdot \underbrace{\sigma_{p \cdot g}}_{=c_{g^{-1}}(\sigma_p)} = p \cdot g \cdot \underbrace{\text{pr}_2(c_{g^{-1}}(\sigma_p))}_{=c_{g^{-1}}(\text{pr}_2(\sigma_p))} = p \cdot \text{pr}_2(\sigma_p) \cdot g = p \cdot \sigma_p \cdot g = H^\sigma(p) \cdot g \end{aligned}$$

for all $(p, g) \in \mathcal{P} * \mathcal{G}$. Thus, $H^\sigma \in \mathcal{A}ut(\mathcal{P})$, and so this finishes the proof. $\blacksquare$

Using this, we can finally formulate the gauge transformations of connection 1-forms; for this recall Cor. [Corollary 6.9](#) and the Darboux derivative, Def. [Definition 6.24](#). Also recall the pullback Darboux derivative, Remark [Remark 6.27](#).

Proposition 6.58: Gauge transformations of connection 1-forms

Let $\mathcal{G} \rightarrow M$ be an LGB over a smooth manifold M and $\mathcal{P} \xrightarrow{\pi} M$ a principal $\mathcal{G}$ -bundle, also let $H\mathcal{G}$ be a horizontal distribution on $\mathcal{G}$ and $A \in \Omega^1(\mathcal{P}; \pi^* \mathcal{G})$ be a connection 1-form on $\mathcal{P}$. Furthermore, let $H \in \mathcal{A}ut(\mathcal{P})$. We then have that $H^!A$ is a connection 1-form on $\mathcal{P}$ and

$$H^!A = \mathcal{A}d_{\text{pr}_2 \circ (\sigma^H)^{-1}} \circ A + (\pi^* \Delta) \sigma^H,$$

where $\sigma^H \in C^\infty(\mathcal{P}; \mathcal{G})^\mathcal{G}$ is defined as in Prop. [Proposition 6.57](#) and $\text{pr}_2 : \pi^* \mathcal{G} \rightarrow \mathcal{G}$ is the projection onto the second component.

Similar to Def. (21) we may shortly just write

$$H^!A = \mathcal{A}d_{(\sigma^H)^{-1}} \circ A + (\pi^* \Delta) \sigma^H.$$

Proof.

First of all observe that $H^!A \in \Omega^1(\mathcal{P}; H^*\pi^*\mathcal{G})$, and trivially $H^*\pi^*\mathcal{G} \cong (\pi \circ H)^*\mathcal{G} = \pi^*\mathcal{G}$. We will make use of this and similar isomorphisms in the following without further mentioning it. We will often not bookkeep the basepoint component in the following pairs and triples; so, everything has to be read in such a way that we have again values in the correct space. It would just be cumbersome to keep track of all these natural isomorphisms, and we decided for readability to avoid this bookkeeping most of the time.

• We then have

$$\begin{aligned} (\mathcal{r}_\sigma^! H^!A)_p(X) &= (H^!A)_{p \cdot \sigma_x}(\mathcal{r}_{\sigma^*}(X)) \\ &= (H^!A)_{p \cdot \sigma_x} \left(D_p r_\sigma(X) - \overline{(\pi^! \Delta \sigma)|_p(X)} \Big|_{p \cdot \sigma_x} \right) \\ &= A_{H(p \cdot \sigma_x)} \left(D_{p \cdot \sigma_x} H \left(D_p r_\sigma(X) - \overline{(\pi^! \Delta \sigma)|_p(X)} \Big|_{p \cdot \sigma_x} \right) \right) \end{aligned}$$

for all (local) $\sigma \in \Gamma(\mathcal{G})$, $p \in \mathcal{P}_x$ ($x \in M$) and $X \in T_p \mathcal{P}$. We also get by definition of $\mathcal{A}d(\mathcal{P})$

$$H(p \cdot \sigma_x) = H(p) \cdot \sigma_x,$$

that is,

$$H \circ r_\sigma = r_\sigma \circ H$$

and thus

$$D_{p \cdot \sigma_x} H \circ D_p r_\sigma = D_p(H \circ r_\sigma) = D_p(r_\sigma \circ H) = D_{H(p)} r_\sigma \circ D_p H.$$

Now also observe that

$$D_p H(\bar{\nu}_p) = (D_p H \circ D_{e_x} \Phi_p)(\nu) = D_{e_x} \underbrace{(H \circ \Phi_p)}_{\mathcal{G}_x \ni g \mapsto H(p \cdot g) = H(p) \cdot g}(\nu) = D_{e_x} \Phi_{H(p)}(\nu) = \bar{\nu}_{H(p)} \quad (22)$$

for all $\nu \in \mathcal{G}_x$, where Φ_p is the orbit map through p . Due to that H is base-preserving we derive

$$(\pi^! \Delta \sigma)|_{H(p)}(D_p H(X)) = (H^! \pi^! \Delta \sigma)_p(X) = ((\pi \circ H)^! \Delta \sigma)_p(X) = (\pi^! \Delta \sigma)_p(X).$$

In total we get

$$\begin{aligned} (\mathcal{r}_\sigma^! H^!A)_p(X) &= A_{H(p) \cdot \sigma_x} \left(D_{H(p)} r_\sigma(D_p H(X)) - \overline{(\pi^! \Delta \sigma)|_{H(p)}(D_p H(X))} \Big|_{H(p) \cdot \sigma_x} \right) \\ &= A_{H(p) \cdot \sigma_x}(\mathcal{r}_{\sigma^*}(D_p H(X))) \\ &= (\mathcal{r}_\sigma^! A)_{H(p)}(D_p H(X)) \\ &= (\mathcal{A}d_{\sigma^{-1}} \circ A)_{H(p)}(D_p H(X)) \\ &= \left(H(p) \cdot \sigma_x, \text{Ad}_{\sigma_x^{-1}}(\hat{A}_{H(p)}(D_p H(X))) \right) \\ &= \mathcal{A}d_{\sigma^{-1}} \left(H(p), \hat{A}_{H(p)}(D_p H(X)) \right) \\ &= \mathcal{A}d_{\sigma^{-1}} \left((H^!A)_p(X) \right), \end{aligned}$$

where we wrote $A = (p, \hat{A}_p)$ with $\hat{A} := \pi_2 \circ A$ ($\pi_2 : \pi^*\mathcal{G} \rightarrow \mathcal{G}$ the projection onto the second component), and thus

$$\mathcal{r}_\sigma^! H^!A = \mathcal{A}d_{\sigma^{-1}} \circ H^!A.$$

By Eq. (22) we also get $DH(\bar{\nu}) = H^*\bar{\nu}$ for all (local) $\nu \in \Gamma(\mathcal{G})$ and thus

$$(H^!A)(\bar{\nu}) = (H^*A)(DH(\bar{\nu})) = (H^*A)(H^*\bar{\nu}) = H^*(A(\bar{\nu})) = H^*\pi^*\nu = (\pi \circ H)^*\nu = \pi^*\nu.$$

Hence, $H^!A$ is a connection 1-form on $\mathcal{P}$.

• For the last part recall Prop. [Proposition 6.57](#), especially we have a unique $\sigma^H \in C^\infty(\mathcal{P}; \mathcal{G})^{\mathcal{G}}$ such that

$$H(p) = p \cdot \sigma_p^H = p \cdot \text{pr}_2(\sigma_p^H) = \Phi(p, \text{pr}_2(\sigma_p^H)) = \left(\Phi \circ (\kappa_{\mathcal{P}}, \text{pr}_2 \circ \sigma^H) \right)(p)$$

for all $p \in \mathcal{P}_x$ ($x \in M$), where Φ is the right $\mathcal{G}$ -action on $\mathcal{P}$, also recall Def. [Equation \(21\)](#), that is, $\text{pr}_2 : \pi^*\mathcal{G} \rightarrow \mathcal{G}$ is the projection onto the second component. Define $\bar{\sigma} := \text{pr}_2 \circ \sigma^H$, $\mathcal{P} \ni p \mapsto \bar{\sigma}_p \in \mathcal{G}$; by Remark [Remark 6.21](#), [Remark 6.27](#) and Thm. [Proposition 5.16](#), especially Remark [Remark 6.29](#), we can calculate

$$\begin{aligned} D_p H(X) &= D_{(p, \bar{\sigma}_p)} \Phi(X, D_p \bar{\sigma}(X)) \\ &= D_p r_\sigma(X) - \overline{(\pi^! \Delta \sigma)|_p(X)} \Big|_{p \cdot \bar{\sigma}_p} + \overline{(\mu_{\mathcal{G}})_{\bar{\sigma}_p} (D_p \bar{\sigma}(X))} \Big|_{p \cdot \bar{\sigma}_p} \\ &= r_{\bar{\sigma}_p^*}^*(X) + \overline{(\mu_{\mathcal{G}})_{\text{pr}_2(\sigma_p^H)} \left(\left(D_{\sigma_p^H} \text{pr}_2 \circ D_p \sigma^H \right)(X) \right)} \Big|_{H(p)} \\ &= r_{\bar{\sigma}_p^*}^*(X) + \overline{(\text{pr}_2^! \mu_{\mathcal{G}})_{\sigma_p^H} (D_p \sigma^H(X))} \Big|_{H(p)} \\ &= r_{\bar{\sigma}_p^*}^*(X) + \overline{(\mu_{\pi^*\mathcal{G}})_{\sigma_p^H} (D_p \sigma^H(X))} \Big|_{H(p)} \\ &= r_{\bar{\sigma}_p^*}^*(X) + \overline{(\Delta^{\pi^*\mathcal{G}} \sigma^H)_p(X)} \Big|_{H(p)} \\ &= r_{\bar{\sigma}_p^*}^*(X) + \overline{((\pi^* \Delta) \sigma^H)_p(X)} \Big|_{H(p)} \end{aligned} \quad (23)$$

for all $X \in T_p \mathcal{P}$, where σ is a (local) section of $\mathcal{G}$ so that $\sigma_x = \bar{\sigma}_p = \text{pr}_2(\sigma_p^H)$.

Then

$$\begin{aligned} (H^! A)_p(X) &= A_{H(p)}(D_p H(X)) \\ &= A_{H(p)} \left(r_{\bar{\sigma}_p^*}^*(X) + \overline{((\pi^* \Delta) \sigma^H)_p(X)} \Big|_{H(p)} \right) \\ &= A_{p \cdot \bar{\sigma}_p} \left(r_{\bar{\sigma}_p^*}^*(X) \right) + ((\pi^* \Delta) \sigma^H)_p(X) \\ &= \left(r_{\bar{\sigma}_p^*}^! A \right)_p(X) + ((\pi^* \Delta) \sigma^H)_p(X) \\ &= \mathcal{A}d_{\bar{\sigma}_p^{-1}}(A_p(X)) + ((\pi^* \Delta) \sigma^H)_p(X) \\ &= \mathcal{A}d_{\text{pr}_2((\sigma_p^H)^{-1})}(A_p(X)) + ((\pi^* \Delta) \sigma^H)_p(X), \end{aligned}$$

which finishes the proof, where we used the trivial relation (alternatively recall Cor. [Corollary 2.7](#))

$$(\text{pr}_2(\sigma_p^H))^{-1} = \text{pr}_2((\sigma_p^H)^{-1}).$$

■

Of course there is also the sense of gauge transformation with respect to gauges as in typical gauge theory (see *e.g.* [\[17, §5.4, page 270ff.\]](#)), recall Def. [Definition 6.15](#).

Definition 6.59: Local gauge field

Let $\mathcal{G} \rightarrow M$ be an LGB over a smooth manifold M and $\mathcal{P} \xrightarrow{\pi} M$ a principal $\mathcal{G}$ -bundle, also let $H\mathcal{G}$ be a horizontal distribution on $\mathcal{G}$ and $A \in \Omega^1(\mathcal{P}; \pi^*\mathcal{G})$ be a connection 1-form on $\mathcal{P}$. Furthermore, let $s \in \Gamma(\mathcal{P}|_U)$ be a (local) gauge over an open subset $U \subset M$. Then we define the **local connection 1-form of local gauge field** $A_s \in \Omega^1(U; \mathcal{G}|_U)$ (**w.r.t.** s) by

$$A_s := s^! A.$$

Remarks 6.60.

A has values in $\pi^*\mathcal{G}$, and thus A_s has values in $s^*\pi^*\mathcal{G} \cong (\pi \circ s)^*\mathcal{G} = \mathcal{K}_U^*\mathcal{G} \cong \mathcal{G}|_U$.

Gauge transformations now naturally arise in a change of the gauge s . So, let U_i and U_j be two open subsets of M so that $U_i \cap U_j \neq \emptyset$. For two gauges $s_i \in \Gamma(\mathcal{P}|_{U_i})$ and $s_j \in \Gamma(\mathcal{P}|_{U_j})$ there is then a unique $\sigma_{ji} \in \Gamma(\mathcal{G}|_{U_i \cap U_j})$ such that

$$s_i = s_j \cdot \sigma_{ji} = \Phi_{s_j} \circ \sigma_{ji}$$

on $U_i \cap U_j$, where Φ_{s_j} is the orbit through s_j , especially also recall Lemma 6.7. The unique existence is clear by the definition of principal bundles, while the smoothness follows by Lemma 6.7 so that we can write σ_{ji} as the composition of smooth maps

$$\sigma_{ji} = \Phi_{s_j}^{-1} \circ s_i.$$

In the following we also introduce the notation

$$\left(\Phi_{s_j}^{-1}\right)^{-1}(p) := \left(\Phi_{s_j}^{-1}(p)\right)^{-1}$$

for all $p \in \mathcal{P}_{U_j}$, that is, $\Phi_{s_j}^{-1}$ is the inverse of the orbit map Φ_{s_j} , but $(\cdot)^{-1}$ is the inverse of $\Phi_{s_j}^{-1}(p)$ as an element of $\mathcal{G}_{U_j}$.

Instead of calculating directly how A_{s_i} and A_{s_j} are related we want to use Thm. Proposition 6.58. For this it will be useful to understand that gauge transformations $\mathcal{Aut}(\mathcal{P})$ are locally isomorphic to the group of sections $\Gamma(\mathcal{G})$ via a gauge, analogously to the typical formulation of gauge theory as *e.g.* illustrated in [17, §5.3.2, page 268f.]. While the isomorphism of $\mathcal{Aut}(\mathcal{P})$ to $C^\infty(\mathcal{P}; \mathcal{G})^\mathcal{G} \subset \Gamma(\pi^*\mathcal{G})$ is global, the following isomorphism is in general only local.

Proposition 6.61: Gauge transformations and sections of the structural LGB

Let $\mathcal{G} \xrightarrow{\pi_{\mathcal{G}}} M$ be an LGB over a smooth manifold M , and $\mathcal{P} \xrightarrow{\pi} M$ a principal $\mathcal{G}$ -bundle. Also let $s \in \Gamma(\mathcal{P}|_U)$ be a gauge defined over some open subset U of M . Then we have a group isomorphism given by

$$\begin{aligned} C^\infty(\mathcal{P}|_U; \mathcal{G}|_U)^{\mathcal{G}|_U} &\rightarrow \Gamma(\mathcal{G}|_U), \\ \sigma &\mapsto \text{pr}_2 \circ \sigma \circ s, \end{aligned}$$

where $\text{pr}_2 : \pi^*\mathcal{G} \rightarrow \mathcal{G}$ is the projection onto the second component, with inverse

$$\begin{aligned} \Gamma(\mathcal{G}|_U) &\rightarrow C^\infty(\mathcal{P}|_U; \mathcal{G}|_U)^{\mathcal{G}|_U}, \\ \tau &\mapsto \sigma^\tau, \end{aligned}$$

where

$$\sigma_p^\tau := \left(p, c_{(\Phi_s^{-1}(p))^{-1}}(\tau_{\pi(p)})\right) = \left(p, (\Phi_s^{-1}(p))^{-1} \tau_{\pi(p)} \Phi_s^{-1}(p)\right)$$

for all $p \in \mathcal{P}_U$, with Φ_s being the orbit map through s .

Remark 6.62: Other notation

Due to the fact that $s^*\pi^*\mathcal{G} \cong (\pi \circ s)^*\mathcal{G} = \mathcal{K}_U^*\mathcal{G} \cong \mathcal{G}|_U$, and due to what we discussed in Subsection 1.3 about pullback sections, we could rewrite the first map to

$$\begin{aligned} C^\infty(\mathcal{P}|_U; \mathcal{G}|_U)^{\mathcal{G}|_U} &\rightarrow \Gamma(\mathcal{G}|_U), \\ \sigma &\mapsto s^*\sigma. \end{aligned}$$

Proof of Prop. Proposition 6.61.

For $\sigma \in C^\infty(\mathcal{P}|_U; \mathcal{G}|_U)^{\mathcal{G}|_U} \subset \Gamma(\pi^*\mathcal{G}|_U)$ observe that $\text{pr}_2 \circ \sigma \circ s$ is by construction a section of $\mathcal{G}|_U$

due to

$$\pi_{\mathcal{G}} \circ \text{pr}_2 \circ \sigma \circ s = \pi \circ \text{pr}_1 \circ \sigma \circ s = \pi \circ s = \mathbb{K}_U,$$

where $\text{pr}_1 : \pi^* \mathcal{G} \rightarrow \mathcal{P}$ is the projection onto the first component. For the supposed inverse we have similarly that σ^τ is a section of $\Gamma(\pi^* \mathcal{G}|_U)$ by construction; then observe, using Lemma 6.7,

$$\begin{aligned} \sigma_{p \cdot g}^\tau &= \left(p \cdot g, c_{(\Phi_s^{-1}(p \cdot g))}^{-1}(\tau_{\pi(p \cdot g)}) \right) \\ &= \left(p \cdot g, c_{g^{-1}(\Phi_s^{-1}(p))}^{-1}(\tau_{\pi(p)}) \right) \\ &= \left(p \cdot g, (c_{g^{-1}} \circ c_{(\Phi_s^{-1}(p))}^{-1})(\tau_{\pi(p)}) \right) \\ &= c_{g^{-1}} \left(p, c_{(\Phi_s^{-1}(p))}^{-1}(\tau_{\pi(p)}) \right) \\ &= c_{g^{-1}}(\sigma_p^\tau) \end{aligned}$$

for all $(p, g) \in \mathcal{P}|_U * \mathcal{G}|_U$, thus, $\sigma^\tau \in C^\infty(\mathcal{P}|_U; \mathcal{G}|_U)^{\mathcal{G}|_U}$. Recall

$$p = s_{\pi(p)} \cdot \Phi_s^{-1}(p)$$

for all $p \in \mathcal{P}|_U$, so that

$$\Phi_s^{-1} \circ s = e|_U,$$

where e is the neutral section of $\mathcal{G}$. On one hand

$$(\text{pr}_2 \circ \sigma^\tau)(s_x) = \text{pr}_2(\sigma_{s_x}^\tau) = c_{(\Phi_s^{-1}(s_x))}^{-1}(\tau_{\pi(s_x)}) = c_{e_x}(\tau_x) = \tau_x$$

for all $x \in M$ and $\tau \in \Gamma(\mathcal{G}|_U)$, and by Def. Definition 6.55 and Ex. Example 6.39 we have on the other hand

$$\begin{aligned} \sigma_p^{\text{pr}_2 \circ \sigma \circ s} &= \left(p, (c_{(\Phi_s^{-1}(p))}^{-1} \circ \text{pr}_2)(\sigma_{s_{\pi(p)}}) \right) \\ &= \left(p, (c_{(\Phi_s^{-1}(p))}^{-1} \circ \text{pr}_2)(\sigma_{p \cdot (\Phi_s^{-1}(p))}^{-1}) \right) \\ &= \left(p, (c_{(\Phi_s^{-1}(p))}^{-1} \circ \text{pr}_2 \circ c_{\Phi_s^{-1}(p)})(\sigma_p) \right) \\ &= \left(p, (c_{(\Phi_s^{-1}(p))}^{-1} \circ c_{\Phi_s^{-1}(p)} \circ \text{pr}_2)(\sigma_p) \right) \\ &= (p, \text{pr}_2(\sigma_p)) \\ &= \sigma_p \end{aligned}$$

for all $p \in \mathcal{P}|_U$ and $\sigma \in C^\infty(\mathcal{P}|_U; \mathcal{G}|_U)^{\mathcal{G}|_U}$. Thus, $\sigma \mapsto \text{pr}_2 \circ \sigma \circ s$ is bijective with inverse $\tau \mapsto \sigma^\tau$, and so it is only left to show that $\sigma \mapsto \text{pr}_2 \circ \sigma \circ s$ is a group homomorphism. For $\sigma' \in C^\infty(\mathcal{P}|_U; \mathcal{G}|_U)^{\mathcal{G}|_U}$ we have

$$\sigma \sigma'|_p = (p, \text{pr}_2(\sigma_p) \text{pr}_2(\sigma'_p)),$$

thus,

$$(\text{pr}_2 \circ (\sigma \sigma') \circ s)_x = \text{pr}_2(\sigma_{s_x}) \text{pr}_2(\sigma'_{s_x}) = ((\text{pr}_2 \circ \sigma \circ s) \cdot (\text{pr}_2 \circ \sigma' \circ s))_x$$

for all $x \in M$. This finishes the proof. ■

Proposition 6.63: Change of gauge as a local bundle automorphism

Let $\mathcal{G} \rightarrow M$ be an LGB over a smooth manifold M , and $\mathcal{P} \xrightarrow{\pi} M$ a principal $\mathcal{G}$ -bundle. Also let U_i and U_j be two open subsets of M so that $U_i \cap U_j \neq \emptyset$, two gauges $s_i \in \Gamma(\mathcal{P}|_{U_i})$ and $s_j \in \Gamma(\mathcal{P}|_{U_j})$, and the unique $\sigma_{ji} \in \Gamma(\mathcal{G}|_{U_i \cap U_j})$ with $s_i = s_j \cdot \sigma_{ji}$ on $U_i \cap U_j$. Then the map H_{ji} defined by

$$\begin{aligned} \mathcal{P}|_{U_i \cap U_j} &\rightarrow \mathcal{P}|_{U_i \cap U_j}, \\ p &\mapsto s_i|_{\pi(p)} \cdot \Phi_{s_j}^{-1}(p), \end{aligned}$$

is an element of $\mathcal{Aut}(\mathcal{P}|_{U_i \cap U_j})$, where Φ_{s_j} is the orbit map through s_j .

Using the notation of Prop. Proposition 6.57, we also get that $\sigma^{H_{ji}}$ is related to σ_{ji} via s_j in sense of Prop. Proposition 6.61, that is,

$$\sigma_p^{H_{ji}} = \left(p, c_{(\Phi_{s_j}^{-1}(p))}^{-1}(\sigma_{ji}|_{\pi(p)}) \right)$$

for all $p \in \mathcal{P}|_{U_i \cap U_j}$, and

$$\sigma_{ji} = \text{pr}_2 \circ \sigma^{H_{ji}} \circ s_j,$$

where $\text{pr}_2 : \pi^*\mathcal{G} \rightarrow \mathcal{G}$ is the projection onto the second component.

Proof.

Recall Lemma Lemma 6.7 for the following proof.

We have $s_i|_{\pi(p)} \in \mathcal{P}_{\pi(p)}$ and Φ_{s_j} is base-preserving, and thus $\Phi_{s_j}^{-1}(p) \in \mathcal{G}_{\pi(p)}$, so that H_{ji} is well-defined and also base-preserving because the right $\mathcal{G}$ -action on $\mathcal{P}$ preserves the fibres. H_{ji} is clearly smooth by construction. Finally,

$$H_{ji}(p \cdot g) = s_i|_{\pi(p \cdot g)} \cdot \Phi_{s_j}^{-1}(p \cdot g) = s_i|_{\pi(p)} \cdot \Phi_{s_j}^{-1}(p) \cdot g = H_{ji}(p) \cdot g$$

for all $(p, g) \in (\mathcal{P} * \mathcal{G})|_{U_i \cap U_j}$, thus, H_{ji} is a principal bundle morphism. The inverse H_{ji}^{-1} is given by

$$H_{ji}^{-1} = H_{ij}$$

due to

$$(H_{ij} \circ H_{ji})(p) = s_j|_x \cdot \Phi_{s_i}^{-1}(s_i|_x \cdot \Phi_{s_j}^{-1}(p)) = s_j|_x \cdot \underbrace{\Phi_{s_i}^{-1}(s_i|_x)}_{=e_x} \cdot \Phi_{s_j}^{-1}(p) = s_j|_x \cdot \Phi_{s_j}^{-1}(p) = p$$

for all $p \in \mathcal{P}|_{U_i \cap U_j}$, where $x := \pi(p)$; by symmetry of course similarly when swapping j and i . Thus, H_{ji} is bijective and its inverse smooth, so that $H_{ji} \in \mathcal{Aut}(\mathcal{P}|_{U_i \cap U_j})$.

Now write

$$\begin{aligned} H_{ji}(p) &= s_i|_x \cdot \Phi_{s_j}^{-1}(p) \\ &= s_j|_x \cdot \sigma_{ji}|_x \cdot \Phi_{s_j}^{-1}(p) \\ &= s_j|_x \cdot \Phi_{s_j}^{-1}(p) \cdot \left(\Phi_{s_j}^{-1}(p) \right)^{-1} \cdot \sigma_{ji}|_x \cdot \Phi_{s_j}^{-1}(p) \\ &= p \cdot c_{(\Phi_{s_j}^{-1}(p))}^{-1}(\sigma_{ji}|_x), \end{aligned}$$

and due to $H_{ji}(p) = p \cdot \text{pr}_2(\sigma_p^{H_{ji}})$ we can immediately derive

$$\sigma_p^{H_{ji}} = \left(p, c_{(\Phi_{s_j}^{-1}(p))}^{-1}(\sigma_{ji}|_{\pi(p)}) \right),$$

thus, $\sigma^{H_{ji}}$ is related to σ_{ji} w.r.t. s_j in sense of Prop. [Proposition 6.61](#) and so the remaining part of the proposition's statement follows by Prop. [Proposition 6.61](#). ■

Using this gauge transformation we can now relate A_{s_i} and A_{s_j} .

Proposition 6.64: Gauge transformations as a change of gauge

Let $\mathcal{G} \rightarrow M$ be an LGB over a smooth manifold M and $\mathcal{P} \xrightarrow{\pi} M$ a principal $\mathcal{G}$ -bundle, also let $H\mathcal{G}$ be a horizontal distribution on $\mathcal{G}$ and $A \in \Omega^1(\mathcal{P}; \pi^*\mathfrak{g})$ be a connection 1-form on $\mathcal{P}$. Also let U_i and U_j be two open subsets of M so that $U_i \cap U_j \neq \emptyset$, two gauges $s_i \in \Gamma(\mathcal{P}|_{U_i})$ and $s_j \in \Gamma(\mathcal{P}|_{U_j})$, and the unique $\sigma_{ji} \in \Gamma(\mathcal{G}|_{U_i \cap U_j})$ with $s_i = s_j \cdot \sigma_{ji}$ on $U_i \cap U_j$. Then we have over $U_i \cap U_j$ that

$$A_{s_i} = \text{Ad}_{\sigma_{ji}^{-1}} \circ A_{s_j} + \Delta\sigma_{ji}.$$

Proof.

Define $H_{ji} \in \mathcal{Aut}(\mathcal{P}|_{U_i \cap U_j})$ as in Prop. [Proposition 6.63](#) (also following its notation), then observe

$$(H_{ji} \circ s_j)_x = s_i|_x \cdot \underbrace{\Phi_{s_j}^{-1}(s_j|_x)}_{=e_x} = s_i|_x$$

for all $x \in U_i \cap U_j$, and thus

$$s_j^! H_{ji}^! A = (H_{ji} \circ s_j)^! A = A_{s_i}.$$

By Thm. [Proposition 6.58](#) we also have

$$H_{ji}^! A = \mathcal{Ad}_{\text{pr}_2 \circ (\sigma^{H_{ji}})^{-1}} \circ A + (\pi^* \Delta) \sigma^{H_{ji}}.$$

We now want to apply $s_j^!$ on both hand sides. Recall Remark [Theorem 6.60](#) and Remark [6.62](#), we will now use several natural isomorphisms without further mention, especially the base-point component will be treated in a "sloppy" way; the advantage will be a cleaner notation, and this is the reason why $\mathcal{Ad}$ becomes Ad by applying $s_j^!$ on the right hand side. Using Prop. [Proposition 6.63](#) and Remark [Remark 6.27](#) we can derive

$$\Delta\sigma_{ji} = ((\pi \circ s_j)^* \Delta) (s_j^* \sigma^{H_{ji}}) = (s_j^* \pi^* \Delta) (s_j^* \sigma^{H_{ji}}) = s_j^! ((\pi^* \Delta) (\sigma^{H_{ji}})),$$

and

$$s_j^! \left(\mathcal{Ad}_{\text{pr}_2 \circ (\sigma^{H_{ji}})^{-1}} \circ A \right) = \text{Ad}_{\text{pr}_2 \circ (\sigma^{H_{ji}})^{-1} \circ s_j} \circ s_j^! A = \text{Ad}_{(\text{pr}_2 \circ \sigma^{H_{ji}} \circ s_j)^{-1}} \circ s_j^! A = \text{Ad}_{\sigma_{ji}^{-1}} \circ A_{s_j},$$

where we used again the trivial relation (alternatively recall Cor. [Corollary 2.7](#))

$$(\text{pr}_2 \circ \sigma^{H_{ji}})^{-1} = \text{pr}_2 \circ (\sigma^{H_{ji}})^{-1}.$$

Altogether, concluding the proof with

$$A_{s_i} = s_j^! H_{ji}^! A = \text{Ad}_{\sigma_{ji}^{-1}} \circ A_{s_j} + \Delta\sigma_{ji}.$$

■

This allows us to finally show the first relation to the infinitesimal gauge theory developed by Alexei Kotov and Thomas Strobl.

Remark 6.65: Integrating curved Yang-Mills gauge theories, part I

Let the situation be as in Thm. [Proposition 6.64](#), then set

$$\sigma_{ji} := e^{t\varepsilon}$$

for a $\varepsilon \in \Gamma(\mathcal{G}|_{U_i \cap U_j})$ and $t \in \mathbb{R}$. Then we define the **infinitesimal gauge transformation** $\delta_\varepsilon A_{s_j}$ of A along ε (w.r.t. s_j) as an element of $\Omega^1(U_i \cap U_j; \mathcal{G}|_{U_i \cap U_j})$ by

$$\delta_\varepsilon A_{s_j} := \left. \frac{d}{dt} \right|_{t=0} A_{s_i}.$$

The corresponding derivative related to the adjoint representation can be calculated as usual, and for the corresponding term related to the Darboux derivative we use the vector bundle connection introduced in Def. Definition 6.33. Thus, we get^a

$$\delta_\varepsilon A_{s_j} = \nabla^\mathcal{G} \varepsilon - [\varepsilon, A_{s_j}]_{\mathcal{G}}.$$

This is *precisely* the infinitesimal gauge transformation developed by Alexei Kotov and Thomas Strobl, see [1] for a concise summary or [6] for an extended introduction. These references are for the very general situation using Lie algebroids, hence see alternatively [5] for this type of gauge theory restricted to Lie algebra bundles.

Thus, we successfully integrated this definition of infinitesimal gauge transformations.

^aRigorously, one has to define $\delta_\varepsilon A_{s_j}$ via the natural projection $V\mathcal{G} \rightarrow \mathcal{G}$ as we did in Prop. Proposition 6.31.

Remark 6.66: Closure of gauge transformations

The gauge transformations close simply by construction, that is,

$$H'^! H^! A = (H \circ H')^! A$$

for all $H', H \in \text{Aut}(\mathcal{P})$. The reader might wonder why, given that existing literature for the infinitesimal version (like [6, Thm. 4.3.37 and Thm. 4.3.43], [19, Prop. 8, Thm. 1] and [20, Eq. 9, 10 and 11]) prove that one needs a **multiplicative connection** on $\mathcal{G}$, that is, the parallel transport on $\mathcal{G}$ has to be a homomorphism of the $\mathcal{G}$ -multiplication, while we did not assume this in this Subsection. However, the culprit is that A as connection 1-form on $\mathcal{P}$ exists if and only if the connection on $\mathcal{G}$ is a multiplicative connection. It is not in the scope of this paper to prove this but the proof of existence of A is similar to ordinary existence arguments because multiplicative connections are indeed Ehresmann connections, and for the other direction one only has to show that A implies a parallel transport as in (15) which is also straightforward to show. Then

$$\text{PT}_\alpha^\mathcal{P}(p \cdot g \cdot q) = \text{PT}_\alpha^\mathcal{P}(p) \cdot \text{PT}_\alpha^\mathcal{G}(gq)$$

for all $(p, g, q) \in \mathcal{P} * \mathcal{G} * \mathcal{G}$, but also

$$\text{PT}_\alpha^\mathcal{P}(p \cdot g \cdot q) = \text{PT}_\alpha^\mathcal{P}(p) \cdot \text{PT}_\alpha^\mathcal{G}(g) \cdot \text{PT}_\alpha^\mathcal{G}(q).$$

Due to the fact that we have a simply transitive $\mathcal{G}$ -action on $\mathcal{P}$, it follows that the connection on $\mathcal{G}$ is multiplicative. Multiplicative connections will be introduced later in more detail when we discuss the field strength. We conclude that the closure of gauge transformations is equivalent to the existence of A , which is equivalent to the multiplicativity of the connection on $\mathcal{G}$.

Alexei's and Thomas's theory also generalizes the curvature/field strength related to gauge theory. Hence, let us now turn to this notion.

6.4. Generalized curvature/field strength.

6.4.1. *multiplicative Yang-Mills connections on LGBs.* In the following we will introduce the curvature; in order to achieve gauge invariance later in the physical gauge theory, we will now need to assume what we will call **compatibility conditions** on the horizontal distribution of the structural

LGB $\mathcal{G}$. Here in this work we will just present these compatibility conditions, and you will be able to see in the proofs why this leads to gauge invariance later. However, if you want to understand the following conditions coming from a point of view not knowing the solution beforehand, then either see [1] or [6]; the latter also provides a physical motivation using what I called the **field redefinitions** which naturally motivate the gauge theory presented here by transforming a classical gauge theory in such a way that the physics is preserved; now recall Def. Definition 6.33 for the following.

We start with the infinitesimal version of the compatibility conditions, already known and pointed out in the previously-mentioned references.

Definition 6.67: Infinitesimal multiplicative Yang-Mills connection

Let $\mathcal{G} \rightarrow M$ be an LGB over a smooth manifold M , and $H\mathcal{G}$ be a horizontal distribution of $\mathcal{G}$. Then we say that $H\mathcal{G}$ is an **infinitesimal multiplicative Yang-Mills connection (on $\mathcal{G}$)** if there is a $\zeta \in \Omega^2(M; \mathfrak{g})$ such that the $\mathcal{G}$ -connection $\nabla^{\mathcal{G}}$ on the associated LAB $\mathfrak{g}$ with bracket $[\cdot, \cdot]_{\mathfrak{g}}$ satisfies the **infinitesimal compatibility conditions**

$$\nabla^{\mathcal{G}}([\mu, \nu]_{\mathfrak{g}}) = [\nabla^{\mathcal{G}}\mu, \nu]_{\mathfrak{g}} + [\mu, \nabla^{\mathcal{G}}\nu]_{\mathfrak{g}}, \quad (24)$$

$$R_{\nabla^{\mathcal{G}}}(X, Y)\mu = [\zeta(X, Y), \mu]_{\mathfrak{g}} \quad (25)$$

for all $\mu, \nu \in \Gamma(\mathfrak{g})$ and $X, Y \in \mathfrak{X}(M)$, where $R_{\nabla^{\mathcal{G}}}$ is the curvature of $\nabla^{\mathcal{G}}$. We then also call $\nabla^{\mathcal{G}}$ an **(infinitesimal) multiplicative Yang-Mills connection** and denote it by $\nabla^{\mathcal{G}}$.

In the following, ζ is always such a 2-form as given here, and even though it is not uniquely given we will treat it as if it is fixed for a given $\nabla^{\mathcal{G}}$.

Remarks 6.68.

As illustrated in [6, §5.1] and [5] such connections also arise in [10, §7.2, page 271ff.]: There these connections are called **Lie derivation laws covering a coupling** due to the fact that these are important in the construction of "coupling" an LAB to a Lie algebroid via a Whitney sum in such a way that the outcome is again a Lie algebroid. There is a strong relationship of this study with the gauge theory we are going to define, however, we will not need this so that we will not repeat what has been discussed in [6, §5.1] and [5].

One may want to write that $(\nabla^{\mathcal{G}}, \zeta)$ is a multiplicative Yang-Mills connection, giving an emphasis on the non-uniqueness of ζ . However, we are not going to do so, and the mentioned references and relations to Mackenzie's work actually show that there is just one object behind all of that describing this multiplicative Yang-Mills connection and its relation to the following gauge theory. It would take too much time introducing all of this again; see the references.

We want to integrate these infinitesimal compatibility conditions to conditions on $H\mathcal{G}$ directly. Let us initially focus on the first infinitesimal compatibility condition (24). In the following we will also view $\mathcal{G}$ as a principal $\mathcal{G}$ -bundle which is equipped with the same horizontal distribution as $\mathcal{G}$ as LGB; related notions and notations carry over. One can show that Ehresmann connections satisfy (24), and in fact Ehresmann connections on LGBs (and Lie groupoids) were already discussed in [12]; and there is also a rather recent preprint related to a similar subject, see [13]. **Special thanks to Camille Laurent-Gengoux for pointing out these references and discussing related subjects, this helped me tremendously.**

Lemma 6.69: Ehresmann connections induce Lie bracket derivations, [12, §4.5, Prop. 4.21]

Let $\mathcal{G} \rightarrow M$ be an LGB over a smooth manifold M , and $H\mathcal{G}$ a horizontal distribution on $\mathcal{G}$. If $H\mathcal{G}$ is an (Ehresmann) connection on $\mathcal{G}$ as principal bundle, then the infinitesimal compatibility condition (24) is satisfied.

Proof.

This can be quickly answered by recalling how the parallel transport behaves; recall that Def. Definition 6.44 comes from Eq. (16) which itself stems from Eq. (15); that is, the associated parallel transport to an Ehresmann connection $H\mathcal{G}$ satisfies Eq. (15). This means that the corresponding parallel transport $PT^{\mathcal{G}}$ of $\mathcal{G}$ (omitting the notation of the involved curve) is a homomorphism, *i.e.*

$$PT^{\mathcal{G}}(gq) = PT^{\mathcal{G}}(g) PT^{\mathcal{G}}(q)$$

for all $(g, q) \in \mathcal{G} * \mathcal{G}$. Thus, infinitesimally the associated parallel transport on $\mathcal{G}$ is a homomorphism w.r.t. the Lie bracket, and thus $\nabla^{\mathcal{G}}$ is a Lie bracket derivation. ■

Remark 6.70: Base manifold a horizontal leave

Observe that the proof's argument implies that a parallel transported neutral element is again the neutral element of the corresponding final fibre, if $H\mathcal{G}$ is an Ehresmann connection. Thus, M naturally embedded via the neutral section e can then be viewed as a horizontal leave of $H\mathcal{G}$.

Due to this, given compatibility condition (24), the following is clear by definition.

Corollary 6.71: The total Maurer-Cartan form as connection 1-form

Let $\mathcal{G} \rightarrow M$ be an LGB over a smooth manifold M , and $H\mathcal{G}$ an Ehresmann connection on $\mathcal{G}$. Then the total Maurer-Cartan form $\mu_{\mathcal{G}}$ is the connection 1-form corresponding to $H\mathcal{G}$.

Proof.

This simply follows by Def. Definition 6.20 and Thm. Theorem 6.52. ■

Remark 6.72: Darboux derivative as local gauge field

If $H\mathcal{G}$ is an Ehresmann connection on $\mathcal{G}$, we know by Cor. Corollary 6.71 that the Darboux derivative $\Delta\sigma$ of a (local) section σ of $\mathcal{G}$ (Def. Definition 6.24) is a local gauge field (w.r.t. σ) corresponding to $\mu_{\mathcal{G}}$, recall Def. Definition 6.59. That is,

$$\Delta\sigma = (\mu_{\mathcal{G}})_{\sigma}.$$

Furthermore, following [12], we can derive that $\mu_{\mathcal{G}}$ is a multiplicative form if and only if $H\mathcal{G}$ is an Ehresmann connection; let us introduce multiplicative 1-forms. Recall Lemma Lemma 5.15.

Definition 6.73: Multiplicative forms, [14, §2.1, special situation of Def. 2.1]

Let $\mathcal{G} \xrightarrow{\pi_{\mathcal{G}}} M$ be an LGB over a smooth manifold M . Then we call an $\omega \in \Omega^1(\mathcal{G}; \pi_{\mathcal{G}}^* \mathcal{G})$ a **multiplicative form** if

$$\omega_{gq}(D_{(g,q)}\Phi(X, Y)) = \mathcal{A}d_{q^{-1}}(\omega_g(X)) + (gq, \widehat{\omega}_q(Y))$$

for all $(g, q) \in \mathcal{G} * \mathcal{G}$ and $(X, Y) \in T_{(g,q)}(\mathcal{G} * \mathcal{G})$, where $\Phi : \mathcal{G} * \mathcal{G} \rightarrow \mathcal{G}$ is the multiplication in $\mathcal{G}$, and $\widehat{\omega} := \text{pr}_2 \circ \omega$, where $\text{pr}_2 : \pi_{\mathcal{G}}^* \mathcal{G} \rightarrow \mathcal{G}$ is the natural projection onto the second component.

Remarks 6.74.

The shape of the second summand is again just because of technical reasons due to the definition of pullback bundles and their base points. If we do not care so much about the involved base points since these are clear by context, then we can also shortly write

$$(\Phi^! \omega)_{(g,q)}(X, Y) = \mathcal{A}d_{q^{-1}}(\omega_g(X)) + \omega_q(Y).$$

Also recall what we mentioned in Subsection Section 1.3 about pullback sections.

Theorem 6.75: Ehresmann connection 1-forms on LGBs are multiplicative, [12, §4.4, implication of Lemma 4.14]

Let $\mathcal{G} \xrightarrow{\pi_{\mathcal{G}}} M$ be an LGB over a smooth manifold M , and $H\mathcal{G}$ a horizontal distribution on $\mathcal{G}$ with corresponding total Maurer-Cartan form $\mu_{\mathcal{G}}$. Then $H\mathcal{G}$ is an (Ehresmann) connection on $\mathcal{G}$ as principal bundle if and only if $\mu_{\mathcal{G}}$ is multiplicative.

Proof.

The general approach in the mentioned reference is different to ours; hence, let us provide a proof suitable to our constructions. We will view $\mathcal{G}$ as a principal $\mathcal{G}$ -bundle whose horizontal distribution is also $H\mathcal{G}$. We will also use the same notation as in Def. Definition 6.73.

To answer whether or not $H\mathcal{G}$ is an Ehresmann connection we just need to discuss whether $\mu_{\mathcal{G}}$ is a connection 1-form on $\mathcal{G}$ due to the fact that $H\mathcal{G}$ is the kernel of $\mu_{\mathcal{G}}$ by definition (see also Cor. Corollary 6.71), also using Thm. Theorem 6.52. Also by definition, $\mu_{\mathcal{G}}$ already satisfies

$$\mu_{\mathcal{G}}(\bar{\nu}) = \pi_{\mathcal{G}}^* \nu.$$

By additionally using Thm. Proposition 5.16 (especially the associated Rem. Remark 6.29) we show

$$\begin{aligned} (\mu_{\mathcal{G}})_{gq}(\mathcal{D}_{(g,q)}\Phi(X, Y)) &= (\mu_{\mathcal{G}})_{gq} \left(\mathcal{D}_g R_{\sigma}(X) - \overline{(\pi_{\mathcal{G}}^! \Delta \sigma)|_g(X)} \Big|_{gq} + \overline{(\mu_{\mathcal{G}})_q(Y)} \Big|_{gq} \right) \\ &= (\mu_{\mathcal{G}})_{gq}(\mathcal{r}_{q*}(X)) + (gq, (\widehat{\mu_{\mathcal{G}}})_q(Y)) \\ &= (\mathcal{r}_q^! \mu_{\mathcal{G}})_g(X) + (gq, (\widehat{\mu_{\mathcal{G}}})_q(Y)) \end{aligned}$$

for all $(g, q) \in \mathcal{G} * \mathcal{G}$ and $(X, Y) \in T_{(g,q)}(\mathcal{G} * \mathcal{G})$, where σ is a (local) section of $\mathcal{G}$ such that $\sigma_x = q$. Thus, $\mu_{\mathcal{G}}$ is multiplicative if and only if

$$\mathcal{A}d_{q^{-1}} \left((\mu_{\mathcal{G}})_g(X) \right) + (gq, (\widehat{\mu_{\mathcal{G}}})_q(Y)) = (\mathcal{r}_q^! \mu_{\mathcal{G}})_g(X) + (gq, (\widehat{\mu_{\mathcal{G}}})_q(Y))$$

if and only if

$$\mathcal{A}d_{q^{-1}} \left((\mu_{\mathcal{G}})_g(X) \right) = (\mathcal{r}_q^! \mu_{\mathcal{G}})_g(X)$$

if and only if $\mu_{\mathcal{G}}$ is a connection 1-form. This finishes the proof. ■

This Theorem also implies the Leibniz rule of the Darboux derivative.

Proposition 6.76: Leibniz rule of the Darboux derivative

Let $\mathcal{G} \xrightarrow{\pi_{\mathcal{G}}} M$ be an LGB over a smooth manifold M , and $H\mathcal{G}$ an Ehresmann connection on $\mathcal{G}$. Then we have

$$\Delta(\sigma\tau) = \text{Ad}_{\tau^{-1}} \circ \Delta\sigma + \Delta\tau$$

and

$$\Delta(\sigma^{-1}) = -\text{Ad}_{\sigma} \circ \Delta\sigma$$

for all $\sigma, \tau \in \Gamma(\mathcal{G})$.

Remarks 6.77.

These are precisely the formulas one expects, see *e.g.* [10, §5.1, Eq. (2) and (3), page 182] for the "classical" case; beware that the reference uses right- instead of left-invariant vector fields to describe the Lie algebra, so that the referenced formulas look explicitly a bit different. Nevertheless, the similarity of the structure should be obvious.

The mentioned reference for Thm. Theorem 6.75, [12, §4.4, Lemma 4.14], actually uses the Leibniz rule of the Darboux derivative as an equivalent description of Ehresmann connections.

Proof of Prop. Proposition 6.76.

For this calculation we will take care about base points of pullback bundles w.r.t. the Darboux derivative, thus recall the last part of Remark Theorem 6.25. Using the same notation as in Def. Definition 6.73, this proposition follows simply by Thm. Theorem 6.75, that is, set $Y := D_x \sigma(X)$ and $Z := D_x \tau(X)$ for $X \in T_x M$ ($x \in M$), then

$$\begin{aligned} (\sigma_x \tau_x, \Delta(\sigma\tau)_x(X)) &= (\mu_{\mathcal{G}})_{\sigma_x \tau_x} (D_x(\sigma\tau)(X)) \\ &= (\mu_{\mathcal{G}})_{\sigma_x \tau_x} (D_{(\sigma_x, \tau_x)} \Phi(Y, Z)) \\ &= \mathcal{A}d_{\tau_x^{-1}} \left((\mu_{\mathcal{G}})_{\sigma_x} (Y) \right) + \left(\sigma_x \tau_x, (\widehat{\mu}_{\mathcal{G}})_{\tau_x} (Z) \right) \\ &= \left(\sigma_x \tau_x, \text{Ad}_{\tau_x^{-1}} \left((\widehat{\mu}_{\mathcal{G}})_{\sigma_x} (Y) \right) + (\widehat{\mu}_{\mathcal{G}})_{\tau_x} (Z) \right) \\ &= \left(\sigma_x \tau_x, \text{Ad}_{\tau_x^{-1}} ((\Delta\sigma)|_x(X)) + (\Delta\tau)|_x(X) \right). \end{aligned}$$

Hence,

$$\Delta(\sigma\tau) = \text{Ad}_{\tau^{-1}} \circ \Delta\sigma + \Delta\tau.$$

Now recall Remark Remark 6.70; this implies for the neutral element section e that

$$\Delta e \equiv 0,$$

and therefore, using the previous result just derived,

$$0 = \Delta e = \Delta(\sigma\sigma^{-1}) = \text{Ad}_{\sigma} \circ \Delta\sigma + \Delta(\sigma^{-1}),$$

thus, concluding the proof with

$$\Delta(\sigma^{-1}) = -\text{Ad}_{\sigma} \circ \Delta\sigma.$$

■

The Leibniz rule of the Darboux derivative extends to $\nabla^{\mathcal{G}}$.

Lemma 6.78: The conjugation of the canonical LAB connection is a multiplicative Yang-Mills connection

Let $\mathcal{G} \xrightarrow{\pi_{\mathcal{G}}} M$ be an LGB over a smooth manifold M , and $\text{H}\mathcal{G}$ an Ehresmann connection on $\mathcal{G}$. Then we have

$$\text{Ad}_{\sigma^{-1}} \circ \nabla^{\mathcal{G}} \circ \text{Ad}_{\sigma} = \nabla^{\mathcal{G}} + \text{ad}_{\Delta\sigma} \quad (26)$$

for all $\sigma \in \Gamma(\mathcal{G})$, that is,

$$(\text{Ad}_{\sigma^{-1}} \circ \nabla_X^{\mathcal{G}} \circ \text{Ad}_{\sigma})\nu = \nabla_X^{\mathcal{G}}\nu + [\Delta\sigma(X), \nu]_{\mathcal{G}}$$

for all $\nu \in \Gamma(\mathcal{G})$ and $X \in \mathfrak{X}(M)$. If $\text{H}\mathcal{G}$ is an infinitesimal multiplicative Yang-Mills connection, $\nabla^{\mathcal{G}} = \nabla^{\mathcal{G}}$, especially $R_{\nabla^{\mathcal{G}}} = \text{ad}_{\zeta}$ for $\zeta \in \Omega^2(M; \mathcal{G})$, then $\overline{\nabla}^{\sigma} := \text{Ad}_{\sigma^{-1}} \circ \nabla^{\mathcal{G}} \circ \text{Ad}_{\sigma}$ is also an infinitesimal multiplicative Yang-Mills connection with

$$R_{\overline{\nabla}^{\sigma}} = \text{ad}_{\text{Ad}_{\sigma^{-1}}(\zeta)},$$

in particular we get

$$\left[d^{\nabla^{\mathcal{G}}} \Delta\sigma + \frac{1}{2} [\Delta\sigma \wedge \Delta\sigma]_{\mathcal{G}} + \zeta - \text{Ad}_{\sigma^{-1}} \circ \zeta, \nu \right]_{\mathcal{G}} = 0 \quad (27)$$

for all $\nu \in \Gamma(\mathcal{G})$.

Proof.

By Prop. Proposition 6.76 we get

$$\begin{aligned} \Delta(e^{\text{Ad}_{\sigma}(\nu)}) &= \Delta(\sigma e^{\nu} \sigma^{-1}) \\ &= \text{Ad}_{\sigma} \circ \Delta(\sigma e^{\nu}) + \Delta(\sigma^{-1}) \end{aligned}$$

$$\begin{aligned}
&= \text{Ad}_\sigma \circ (\text{Ad}_{e^{-\nu}} \circ \Delta\sigma + \Delta e^\nu) - \text{Ad}_\sigma \circ \Delta\sigma \\
&= \text{Ad}_\sigma \circ (\text{Ad}_{e^{-\nu}} \circ \Delta\sigma - \Delta\sigma + \Delta e^\nu)
\end{aligned}$$

for all $\sigma \in \Gamma(\mathcal{G})$ and $\nu \in \Gamma(\mathcal{Q})$, where the first equality is a well-known fact, see *e.g.* [17, §1.7, Thm. 1.7.16, page 59]. Then

$$\begin{aligned}
\nabla^\mathcal{G}(\text{Ad}_\sigma(\nu)) &= \left. \frac{d}{dt} \right|_{t=0} \left(\Delta(e^{\text{Ad}_\sigma(t\nu)}) \right) \\
&= \left. \frac{d}{dt} \right|_{t=0} \left(\text{Ad}_\sigma \circ (\text{Ad}_{e^{-t\nu}} \circ \Delta\sigma - \Delta\sigma + \Delta e^{t\nu}) \right) \\
&= \text{Ad}_\sigma \circ (\text{ad}_{-\nu} \circ \Delta\sigma + \nabla^\mathcal{G}\nu) \\
&= \text{Ad}_\sigma \circ (\text{ad}_{\Delta\sigma}(\nu) + \nabla^\mathcal{G}\nu)
\end{aligned}$$

making use of that this calculation is pointwise at $x \in M$ just a standard derivative in the vector space $\mathcal{Q}_x$. Thus,

$$\text{Ad}_{\sigma^{-1}} \circ \nabla^\mathcal{G} \circ \text{Ad}_\sigma = \nabla^\mathcal{G} + \text{ad}_{\Delta\sigma}.$$

In order to show $R_{\overline{\nabla}^\sigma} = \text{ad}_{\text{Ad}_{\sigma^{-1}}(\zeta)}$ one uses the definition $\overline{\nabla}^\sigma := \text{Ad}_{\sigma^{-1}} \circ \nabla^\mathcal{G} \circ \text{Ad}_\sigma$ to calculate the curvature,

$$\begin{aligned}
R_{\overline{\nabla}^\sigma}(X, Y)\nu &= \overline{\nabla}_X^\sigma \overline{\nabla}_Y^\sigma \nu - \overline{\nabla}_Y^\sigma \overline{\nabla}_X^\sigma \nu - \overline{\nabla}_{[X, Y]}^\sigma \nu \\
&= \left(\text{Ad}_{\sigma^{-1}} \circ \left(\nabla_X^\mathcal{G} \circ \nabla_Y^\mathcal{G} - \nabla_Y^\mathcal{G} \circ \nabla_X^\mathcal{G} - \nabla_{[X, Y]}^\mathcal{G} \right) \circ \text{Ad}_\sigma \right)(\nu) \\
&= (\text{Ad}_{\sigma^{-1}} \circ R_{\nabla^\mathcal{G}}(X, Y) \circ \text{Ad}_\sigma)(\nu) \\
&= \text{Ad}_{\sigma^{-1}} \left([\zeta(X, Y), \text{Ad}_\sigma(\nu)]_{\mathcal{Q}} \right) \\
&= [\text{Ad}_{\sigma^{-1}}(\zeta(X, Y)), \nu]_{\mathcal{Q}}
\end{aligned}$$

for all $X, Y \in \mathfrak{X}(M)$ and $\nu \in \Gamma(\mathcal{Q})$. The remaining parts of the statement are trivial and straightforward calculations, making use of Def. Definition 6.67; in fact, we have proven all the remaining statements already in [5, §3, part of Thm. 3.6] and in a more general manner in [6, §4.6, Thm. 4.6.9]: That $\overline{\nabla}^\sigma$ is a Lie bracket derivation is trivial to show, and one uses identity (26) to show another identity of its curvature,

$$R_{\overline{\nabla}^\sigma} = \left[d^{\nabla^\mathcal{G}} \Delta\sigma + \frac{1}{2} [\Delta\sigma \wedge \Delta\sigma]_{\mathcal{Q}} + \zeta, \cdot \right]_{\mathcal{Q}}.$$

Combining both identities for $R_{\overline{\nabla}^\sigma}$ we conclude with a proof of Eq. (27). ■

We also get a generalization of statements like [17, §5.5, Lemma 5.5.5, page 276]:

Lemma 6.79: Lie bracket of horizontal and vertical vector

Let $\mathcal{G} \rightarrow M$ be an LGB over a smooth manifold M and $\mathcal{P} \xrightarrow{\pi} M$ a principal $\mathcal{G}$ -bundle, also let $\text{H}\mathcal{G}$ be a horizontal distribution on $\mathcal{G}$ and $A \in \Omega^1(\mathcal{P}; \pi^*\mathcal{Q})$ be a connection 1-form on $\mathcal{P}$. Then we have

$$A([X, \bar{\nu}]) = \left. \frac{d}{dt} \right|_{t=0} \left((\pi^! \Delta e^{t\nu})(X) \circ r_{e^{-t\nu}} \right)$$

for all $\nu \in \Gamma(\mathcal{Q})$ and horizontal vector fields $X \in \mathfrak{X}(\mathcal{P})$. If there is an $\omega \in \mathfrak{X}(M)$ such that $D\pi(X) = \pi^*\omega$, then

$$A([X, \bar{\nu}]) = (\pi^* \nabla^\mathcal{G})_X (\pi^* \nu) = \pi^* (\nabla_\omega^\mathcal{G} \nu).$$

If $H\mathcal{G}$ is an Ehresmann connection, then we can conclude for all horizontal vector fields X that

$$A([X, \bar{\nu}]) = (\pi^* \nabla^{\mathcal{G}})_X (\pi^* \nu).$$

Remarks 6.80.

The last statement got also stated in [13, Eq. (2.8)] in the case of the LGB as principal bundle; we were unaware of this proof, which is why our proof is a bit different.

Proof of Lemma Lemma 6.79.

First of all recall the very basic fact (see e.g. [17, §A.1, Thm. A.1.46, page 615]) that for two vector fields $X, Y \in \mathfrak{X}(\mathcal{P})$ we have¹⁰

$$[X, Y]_p = - \frac{d}{dt} \Big|_{t=0} D_{\phi_t(p)} \phi_{-t} (X_{\phi_t(p)})$$

for all $p \in \mathcal{P}$, where ϕ_t is the local flow of Y (with parameter t in an open interval of $\mathbb{R}$ containing 0). Now let X be a horizontal vector field and $Y = \bar{\nu}$. For such a Y we know by Def. Definition 5.12 that

$$\bar{\nu}_p = D_{e_x} \Phi_p(\nu_x) = \frac{d}{dt} \Big|_{t=0} (p \cdot e^{t\nu_x})$$

for all $p \in \mathcal{P}$, where Φ_p is the orbit map through p , $x := \pi(p)$, and e_x the neutral element of $\mathcal{G}_x$. Thus,

$$\phi_t(p) = p \cdot e^{t\nu_x} = r_{e^{t\nu}}(p),$$

and by rewriting in sense of Def. Definition 6.41 we achieve

$$\begin{aligned} [X, \bar{\nu}]_p &= - \frac{d}{dt} \Big|_{t=0} D_{p \cdot e^{t\nu_x}} r_{e^{-t\nu}} (X_{p \cdot e^{t\nu_x}}) \\ &= - \frac{d}{dt} \Big|_{t=0} \left(D_{p \cdot e^{t\nu_x}} r_{e^{-t\nu}} (X_{p \cdot e^{t\nu_x}}) - \overline{(\pi^! \Delta e^{-t\nu})|_{p \cdot e^{t\nu_x}} (X_{p \cdot e^{t\nu_x}})} \Big|_p \right) \\ &\quad - \frac{d}{dt} \Big|_{t=0} \left(\overline{(\pi^! \Delta e^{-t\nu})|_{p \cdot e^{t\nu_x}} (X_{p \cdot e^{t\nu_x}})} \Big|_p \right) \\ &= - \frac{d}{dt} \Big|_{t=0} \left(r_{e^{-t\nu_x}}^* (X_{p \cdot e^{t\nu_x}}) \right) \\ &\quad - \frac{d}{dt} \Big|_{t=0} \left(\overline{(\pi^! \Delta e^{-t\nu})|_{p \cdot e^{t\nu_x}} (X_{p \cdot e^{t\nu_x}})} \Big|_p \right). \end{aligned}$$

The first summand, $t \mapsto r_{e^{-t\nu_x}}^* (X_{p \cdot e^{t\nu_x}})$, is a curve with values in $H_p \mathcal{P}$ due to the fact that X is a horizontal vector field. Thus, the velocity of this curve at $t = 0$ will also be an element of $H_p \mathcal{P}$, and the first summand is therefore in the kernel of A . Similarly,

$$t \mapsto \eta(t) := (\pi^! \Delta e^{-t\nu})|_{p \cdot e^{t\nu_x}} (X_{p \cdot e^{t\nu_x}}) = (\Delta e^{-t\nu})|_x (D_{p \cdot e^{t\nu_x}} \pi(X_{p \cdot e^{t\nu_x}}))$$

is a curve with values in $\mathcal{G}_x$, and therefore its fundamental vector field at p is a curve with values in $V_p \mathcal{P}$. Hence, we can calculate in the sense of vector spaces to derive

$$- \frac{d}{dt} \Big|_{t=0} \overline{\eta(t)}_p = \frac{d}{dt} \Big|_{t=0} \left(D_{e_x} \Phi_p(\eta(-t)) \right) = D_{e_x} \Phi_p \left(\frac{d}{dt} \Big|_{t=0} \eta(-t) \right) = \overline{\frac{d}{dt} \Big|_{t=0} \eta(-t)} \Big|_p$$

¹⁰Rigorously, similarly to Prop. Proposition 6.31 one has a projection involved, but as explained in Remark Theorem 6.32 we will omit notating this projection; in the context of this proof we can allow ourselves to take the typical "less rigorous" approach for simplicity.

where we substituted $t \leftrightarrow -t$ after the first equality sign. In total we achieve

$$A_p([X, \bar{\nu}]_p) = A_p\left(\overline{\frac{d}{dt}\bigg|_{t=0} \eta(-t)}\bigg|_p\right) = \left(p, \frac{d}{dt}\bigg|_{t=0} \left((\Delta e^{t\nu})|_x \left(D_{p \cdot e^{-t\nu_x}} \pi(X_{p \cdot e^{-t\nu_x}})\right)\right)\right),$$

and thus

$$A([X, \bar{\nu}]) = \frac{d}{dt}\bigg|_{t=0} \left((\pi^! \Delta e^{t\nu})(X) \circ r_{e^{-t\nu}}\right).$$

If there is an $\omega \in \mathfrak{X}(M)$ such that $D\pi(X) = \pi^*\omega$, then

$$A_p([X, \bar{\nu}]_p) = \left(p, \frac{d}{dt}\bigg|_{t=0} \left((\Delta e^{t\nu})|_x (\omega_x)\right)\right),$$

and therefore by Def. [Definition 6.33](#)

$$A_p([X, \bar{\nu}]_p) = (p, \nabla_\omega^\mathcal{G} \nu|_x) = \pi^*(\nabla_\omega^\mathcal{G} \nu)|_p.$$

By the definition of pullback connections we could also write

$$A([X, \bar{\nu}]) = (\pi^* \nabla^\mathcal{G})_X (\pi^* \nu).$$

If $H\mathcal{G}$ is an Ehresmann connection, then recall [Remark 6.70](#); this implies for the neutral element section e that

$$\Delta e \equiv 0,$$

and, again, $t \mapsto (\Delta e^{t\nu})_x$ and $t \mapsto D_{p \cdot e^{-t\nu_x}} \pi(X_{p \cdot e^{-t\nu_x}})$ are curves with values in the vector spaces $T_x^* M \otimes \mathfrak{g}_x$ and $T_x M$, respectively. Thus, we can just apply typical Leibniz rules to calculate the derivative of the contraction

$$\begin{aligned} & \frac{d}{dt}\bigg|_{t=0} \left((\Delta e^{t\nu})|_x \left(D_{p \cdot e^{-t\nu_x}} \pi(X_{p \cdot e^{-t\nu_x}})\right)\right) \\ &= \left(\frac{d}{dt}\bigg|_{t=0} \Delta e^{t\nu}\right)\bigg|_x (D_p \pi(X_p)) + \underbrace{(\Delta e)|_x}_{=0} \left(\frac{d}{dt}\bigg|_{t=0} D_{p \cdot e^{-t\nu_x}} \pi(X_{p \cdot e^{-t\nu_x}})\right) \\ &= \nabla_{D_p \pi(X_p)}^\mathcal{G} \nu, \end{aligned}$$

and as before we conclude

$$A([X, \bar{\nu}]) = (\pi^* \nabla^\mathcal{G})_X (\pi^* \nu).$$

■

This concludes our discussion about the infinitesimal compatibility condition (24) and its integrated version, "H $\mathcal{G}$ = Ehresmann connection". Via Thm. [Theorem 6.75](#) and Lemma [Lemma 6.69](#) we therefore concluded that compatibility condition (24) holds if the total Maurer-Cartan form is multiplicative. Let us therefore now turn to compatibility condition (25), which we can actually integrate to an equivalent statement.

Theorem 6.81: Generalized Maurer-Cartan equation

Let $\mathcal{G} \xrightarrow{\pi_\mathcal{G}} M$ be an LGB over a smooth manifold M with connected fibres, $H\mathcal{G}$ an Ehresmann connection on $\mathcal{G}$, and $\zeta \in \Omega^2(M; \mathfrak{g})$. Then $H\mathcal{G}$ satisfies the infinitesimal compatibility condition (25) w.r.t. ζ if and only if the total Maurer-Cartan form $\mu_\mathcal{G}$ satisfies

$$\left(d\pi_\mathcal{G}^* \nabla^\mathcal{G} \mu_\mathcal{G} + \frac{1}{2}[\mu_\mathcal{G} \wedge \mu_\mathcal{G}]_{\pi_\mathcal{G}^* \mathfrak{g}} + \pi_\mathcal{G}^! \zeta\right)\bigg|_g = (g, \text{Ad}_{g^{-1}} \circ \zeta_x \circ (D_g \pi_\mathcal{G}, D_g \pi_\mathcal{G})) \quad (28)$$

for all $g \in \mathcal{G}_x$ ($x \in M$). We then say that $\mu_{\mathcal{G}}$ solves the **generalized Maurer-Cartan equation** (w.r.t. ζ).

Remarks 6.82.

The structure on the right hand side is again due to denoting the base point in pullback bundles; one may decide to drop the base point for simplicity in the notation. In that case, since the base point is clear by context, one may also just write

$$\mathcal{A}d_{g^{-1}} \circ \pi_{\mathcal{G}}^! \zeta|_g \quad \text{or} \quad \text{Ad}_{g^{-1}} \circ \pi_{\mathcal{G}}^! \zeta|_g$$

on the right hand side.

Remark 6.83: Classical Maurer-Cartan equation recovered

Recall Rem. [Remark 6.28](#). If $\mathcal{G}$ is a trivial LGB, $H\mathcal{G}$ its canonical flat connection and $\zeta \equiv 0$, then $\nabla^{\mathcal{G}}$ is the canonical flat connection on the trivial LAB $\mathcal{G}$ (recall Ex. [Example 6.35](#)) and it clearly satisfies the compatibility conditions in Def. [Definition 6.67](#) w.r.t. $\zeta \equiv 0$. Furthermore, the generalized Maurer-Cartan equation then reduces to the classical Maurer-Cartan equation.

Proof of Thm. Theorem 6.81.

The idea of the proof is similar to the proof of a statement about when two multiplicative 2-forms¹¹ are equal, see e.g. [21, §3, Cor. 3.4]. That is, we want to discuss under what conditions

$$\mathcal{G} \ni g \mapsto \omega_g := \left(d^{\pi_{\mathcal{G}}^* \nabla^{\mathcal{G}}} \mu_{\mathcal{G}} + \frac{1}{2} [\mu_{\mathcal{G}} \wedge \mu_{\mathcal{G}}]_{\pi_{\mathcal{G}}^* \mathcal{G}} + \pi_{\mathcal{G}}^! \zeta \right) \Big|_g - (g, \text{Ad}_{g^{-1}} \circ \zeta_x \circ (D_g \pi_{\mathcal{G}}, D_g \pi_{\mathcal{G}}))$$

is constant along the fibres of $\mathcal{G}$, that is, constant along the flows of vertical tangent vectors, the left-invariant vector fields (recall Cor. [Corollary 4.16](#)).

• First of all observe that ω vanishes if at least one tangent vector is vertical. Let $Y \in V_g \mathcal{G}$ ($g \in \mathcal{G}_x$, $x \in M$), and write $Y = \overline{\nu}_x|_g$ for a $\nu_x \in \mathcal{G}_x$. Due to $D\pi_{\mathcal{G}}(Y) = 0$ it is clear that the contraction of $\pi_{\mathcal{G}}^! \zeta$ with Y is 0. Let $Z \in T_g \mathcal{G}$ for which we write

$$Z := \overline{\mu}_x|_g + X_g$$

for a $\mu_x \in \mathcal{G}_x$ and $X_g \in H_g \mathcal{G}$; then by Cor. [Corollary 6.71](#)

$$\frac{1}{2} [\mu_{\mathcal{G}} \wedge \mu_{\mathcal{G}}]_{\pi_{\mathcal{G}}^* \mathcal{G}} \Big|_g (Y, Z) = \left[(\mu_{\mathcal{G}})_g(Y), (\mu_{\mathcal{G}})_g(Z) \right]_{\pi_{\mathcal{G}}^* \mathcal{G}} = \left(g, [\nu_x, \mu_x]_{\mathcal{G}_x} \right).$$

Extend Y and Z naturally to vector fields on $\mathcal{G}$ (for simplicity also denoted by Y and Z , respectively) by viewing ν_x and μ_x as values of sections ν and μ , respectively, of $\mathcal{G}$ at x , and extending X_g (locally) to a horizontal vector field X ; thence,

$$\begin{aligned} \left(d^{\pi_{\mathcal{G}}^* \nabla^{\mathcal{G}}} \mu_{\mathcal{G}} \right)_g (Y, Z) &= (\pi_{\mathcal{G}}^* \nabla^{\mathcal{G}})_{Y_g} (\mu_{\mathcal{G}}(Z)) - (\pi_{\mathcal{G}}^* \nabla^{\mathcal{G}})_{Z_g} (\mu_{\mathcal{G}}(Y)) - (\mu_{\mathcal{G}})_g([Y, Z]) \\ &= \underbrace{(\pi_{\mathcal{G}}^* \nabla^{\mathcal{G}})_{\overline{\nu}_g} (\pi_{\mathcal{G}}^* \mu)}_{=0} - (\pi_{\mathcal{G}}^* \nabla^{\mathcal{G}})_{\overline{\mu}_g + X_g} (\pi_{\mathcal{G}}^* \nu) \\ &= \pi_{\mathcal{G}}^* \left(\nabla_{D\pi_{\mathcal{G}}(\overline{\nu})} \mu \right) \Big|_g = 0 \\ &\quad - (\mu_{\mathcal{G}})_g([\overline{\nu}, \overline{\mu}]) - (\mu_{\mathcal{G}})_g([\overline{\nu}, X]) \\ &= -(\pi_{\mathcal{G}}^* \nabla^{\mathcal{G}})_{X_g} (\pi_{\mathcal{G}}^* \nu) - \left(g, [\nu_x, \mu_x]_{\mathcal{G}_x} \right) + (\pi_{\mathcal{G}}^* \nabla^{\mathcal{G}})_{X_g} (\pi_{\mathcal{G}}^* \nu) \\ &= - \left(g, [\nu_x, \mu_x]_{\mathcal{G}_x} \right) \end{aligned}$$

¹¹See Rem. [Remark 6.87](#) later; multiplicativity is related to closedness w.r.t. a differential. However, we will not need this general notion.

using Rem. [Remark 5.14](#) (bracket of fundamental/left-invariant vector fields is fundamental/left-invariant) and Lemma [Lemma 6.79](#). Thus,

$$\left(d^{\pi_{\mathcal{G}}^* \nabla^{\mathcal{G}}} \mu_{\mathcal{G}} + \frac{1}{2} [\mu_{\mathcal{G}} \wedge \mu_{\mathcal{G}}]_{\pi_{\mathcal{G}}^* \mathcal{G}} \right) \Big|_g (Y, Z) = 0.$$

It follows that ω vanishes if contracted with a vertical tangent vector.

• Therefore we can study ω just w.r.t. tangent vectors complementary to the vertical structure, these do not necessarily need to be elements of $\mathcal{H}\mathcal{G}$. For a given $g \in \mathcal{G}_x$ ($x \in M$) define

$$\begin{aligned} \mathbb{R} &\rightarrow \Gamma(\mathcal{G}), \\ t &\mapsto \gamma(t) := \sigma e^{t\nu}, \end{aligned}$$

where ν is any section of $\mathcal{G}$ and $\sigma \in \Gamma(\mathcal{G})$ (local) with $\sigma_x = g$; pointwise we denote $\gamma(t)$ as a map $M \ni x \mapsto \gamma_x(t)$. By Cor. [Corollary 4.16](#) the flows of elements of $V_g \mathcal{G}$ can be precisely described by curves like γ , so that the derivative w.r.t. t of pullbacks of tensors with γ characterizes any derivative along a fibre of $\mathcal{G}$. Furthermore, $\gamma(t)$ is by definition a section of $\mathcal{G}$ for all t and thence describes an embedding of M into $\mathcal{G}$. Thus, the t -derivatives of terms like $\gamma(t)^! \omega$ describe the derivatives of ω along fibres of $\mathcal{G}$ contracted with tangent vectors complementary to the vertical structure. Therefore let us calculate $\gamma(t)^! \omega$; we have

$$\gamma(t)^! \pi_{\mathcal{G}}^! \zeta = \underbrace{(\pi_{\mathcal{G}} \circ \gamma(t))^!}_{\equiv \mathcal{K}_M^!} \zeta = \zeta$$

and under the identification $\gamma^* \pi_{\mathcal{G}}^* \mathcal{G} \cong (\pi_{\mathcal{G}} \circ \gamma)^* \mathcal{G} = \mathcal{K}_M^* \mathcal{G} \cong \mathcal{G}$ we get similarly

$$(\gamma(t)^! (g, \text{Ad}_{g^{-1}} \circ \zeta_{\pi_{\mathcal{G}}(g)} \circ D_g \pi_{\mathcal{G}}))_x \cong \text{Ad}_{\gamma_x^{-1}(t)} \circ \zeta_{\pi_{\mathcal{G}}(\gamma_x(t))} \circ D_x (\pi_{\mathcal{G}} \circ \gamma(t)) = \text{Ad}_{\gamma_x^{-1}(t)} \circ \zeta_x$$

for all $x \in M$, where $\gamma^{-1}(t)$ is the $\mathcal{G}$ -inverse of $\gamma(t)$. It is also a straight-forward calculation to show that

$$\gamma(t)^! d^{\pi_{\mathcal{G}}^* \nabla^{\mathcal{G}}} \mu_{\mathcal{G}} = d^{\gamma(t)^* \pi_{\mathcal{G}}^* \nabla^{\mathcal{G}}} (\gamma(t)^! \mu_{\mathcal{G}}) = d^{(\pi_{\mathcal{G}} \circ \gamma(t))^* \nabla^{\mathcal{G}}} (\gamma(t)^! \mu_{\mathcal{G}}) = d^{\nabla^{\mathcal{G}}} (\gamma(t)^! \mu_{\mathcal{G}}),$$

using the definition of pullback vector bundle connections; alternatively see the explicit calculations in [5, Appendix, Prop. A.1, Eq. (A.1)] or [6, Appendix, Prop. A.1.1, Eq. (A.2)], and we have

$$\begin{aligned} &\left(\gamma(t)^! \left(\frac{1}{2} [\mu_{\mathcal{G}} \wedge \mu_{\mathcal{G}}]_{\pi_{\mathcal{G}}^* \mathcal{G}} \right) \right)_x (X, Y) \\ &= \left[(\mu_{\mathcal{G}})_{\gamma_x(t)} (D_x (\gamma(t)) (X)), (\mu_{\mathcal{G}})_{\gamma_x(t)} (D_x (\gamma(t)) (Y)) \right]_{\mathcal{G}_x} \\ &= [(\gamma(t)^! \mu_{\mathcal{G}})_x (X), (\gamma(t)^! \mu_{\mathcal{G}})_x (Y)]_{\mathcal{G}_x} \\ &= \left(\frac{1}{2} [\gamma(t)^! \mu_{\mathcal{G}} \wedge \gamma(t)^! \mu_{\mathcal{G}}]_{\mathcal{G}} \right)_x (X, Y) \end{aligned}$$

for all $x \in M$ and $X, Y \in T_x M$. Thus, we have so far

$$\begin{aligned} \gamma(t)^! \omega &= d^{\nabla^{\mathcal{G}}} (\gamma(t)^! \mu_{\mathcal{G}}) + \frac{1}{2} [\gamma(t)^! \mu_{\mathcal{G}} \wedge \gamma(t)^! \mu_{\mathcal{G}}]_{\mathcal{G}} + \zeta - \text{Ad}_{\gamma^{-1}(t)} \circ \zeta \\ &= d^{\nabla^{\mathcal{G}}} (\Delta \gamma(t)) + \frac{1}{2} [\Delta \gamma(t) \wedge \Delta \gamma(t)]_{\mathcal{G}} + \zeta - \text{Ad}_{\gamma^{-1}(t)} \circ \zeta. \end{aligned} \tag{29}$$

By Prop. [Proposition 6.76](#) we get

$$\Delta \gamma(t) = \text{Ad}_{e^{-t\nu}} \circ \Delta \sigma + \Delta e^{t\nu},$$

and via Lemma [Lemma 6.78](#) we can derive

$$\begin{aligned} &\left(d^{\nabla^{\mathcal{G}}} (\text{Ad}_{e^{-t\nu}} \circ \Delta \sigma) \right) (X, Y) \\ &= \nabla_X^{\mathcal{G}} (\text{Ad}_{e^{-t\nu}} (\Delta \sigma(Y))) - \nabla_Y^{\mathcal{G}} (\text{Ad}_{e^{-t\nu}} (\Delta \sigma(X))) - \text{Ad}_{e^{-t\nu}} (\Delta \sigma([X, Y])) \\ &= \text{Ad}_{e^{-t\nu}} \left(\nabla_X^{\mathcal{G}} (\Delta \sigma(Y)) + [(\Delta e^{-t\nu})(X), \Delta \sigma(Y)]_{\mathcal{G}} \right) \end{aligned}$$

$$\begin{aligned}
& -\nabla_Y^{\mathcal{G}}(\Delta\sigma(X)) - [(\Delta e^{-t\nu})(Y), \Delta\sigma(X)]_{\mathcal{G}} \\
& - \Delta\sigma([X, Y]) \\
& = \text{Ad}_{e^{-t\nu}} \left(\left(d^{\nabla^{\mathcal{G}}} \Delta\sigma \right)(X, Y) + [(\Delta e^{-t\nu})(X), \Delta\sigma(Y)]_{\mathcal{G}} - [(\Delta e^{-t\nu})(Y), \Delta\sigma(X)]_{\mathcal{G}} \right) \\
& = \left(\text{Ad}_{e^{-t\nu}} \circ \left(d^{\nabla^{\mathcal{G}}} \Delta\sigma + [\Delta e^{-t\nu} \frown \Delta\sigma]_{\mathcal{G}} \right) \right)(X, Y)
\end{aligned}$$

for all $X, Y \in \mathfrak{X}(M)$, hence,

$$\begin{aligned}
\frac{d}{dt} \Big|_{t=0} \left(d^{\nabla^{\mathcal{G}}} \Delta\gamma(t) \right) &= [d^{\nabla^{\mathcal{G}}} \Delta\sigma, \nu]_{\mathcal{G}} - [\Delta\sigma \frown \nabla^{\mathcal{G}} \nu]_{\mathcal{G}} + d^{\nabla^{\mathcal{G}}} \nabla^{\mathcal{G}} \nu \\
&= [d^{\nabla^{\mathcal{G}}} \Delta\sigma, \nu]_{\mathcal{G}} - [\Delta\sigma \frown \nabla^{\mathcal{G}} \nu]_{\mathcal{G}} + R_{\nabla^{\mathcal{G}}}(\cdot, \cdot) \nu
\end{aligned}$$

making use of that this calculation is pointwise at $x \in M$ just a standard derivative in the vector space $\mathcal{G}_x$ and that $\Delta e \equiv 0$ by Remark 6.70; we also used Prop. Proposition 1.3. In a similar fashion,

$$\frac{d}{dt} \Big|_{t=0} [\Delta\gamma(t) \frown \Delta\gamma(t)]_{\mathcal{G}} = 2 \left[[\Delta\sigma, \nu]_{\mathcal{G}} \frown \Delta\sigma \right]_{\mathcal{G}} + 2 [\Delta\sigma \frown \nabla^{\mathcal{G}} \nu]_{\mathcal{G}},$$

which we can rewrite by the Jacobi identity

$$\begin{aligned}
\left[[\Delta\sigma, \nu]_{\mathcal{G}} \frown \Delta\sigma \right]_{\mathcal{G}}(X, Y) &= \left[[\Delta\sigma(X), \nu]_{\mathcal{G}}, \Delta\sigma(Y) \right]_{\mathcal{G}} - \left[[\Delta\sigma(Y), \nu]_{\mathcal{G}}, \Delta\sigma(X) \right]_{\mathcal{G}} \\
&= \left[[\Delta\sigma(X), \Delta\sigma(Y)]_{\mathcal{G}}, \nu \right]_{\mathcal{G}} \\
&= \left[\frac{1}{2} [\Delta\sigma \frown \Delta\sigma]_{\mathcal{G}}, \nu \right]_{\mathcal{G}}(X, Y)
\end{aligned}$$

for all $X, Y \in \mathfrak{X}(M)$, and we also trivially have

$$\frac{d}{dt} \Big|_{t=0} \text{Ad}_{\gamma^{-1}(t)} \circ \zeta = \frac{d}{dt} \Big|_{t=0} \text{Ad}_{e^{-t\nu}} \circ \text{Ad}_{\sigma^{-1}} \circ \zeta = [\text{Ad}_{\sigma^{-1}} \circ \zeta, \nu]_{\mathcal{G}}.$$

Collecting everything we derive

$$\begin{aligned}
\frac{d}{dt} \Big|_{t=0} \gamma(t)! \omega &= [d^{\nabla^{\mathcal{G}}} \Delta\sigma, \nu]_{\mathcal{G}} - [\Delta\sigma \frown \nabla^{\mathcal{G}} \nu]_{\mathcal{G}} + R_{\nabla^{\mathcal{G}}}(\cdot, \cdot) \nu \\
&\quad + \left[\frac{1}{2} [\Delta\sigma \frown \Delta\sigma]_{\mathcal{G}}, \nu \right]_{\mathcal{G}} + [\Delta\sigma \frown \nabla^{\mathcal{G}} \nu]_{\mathcal{G}} - [\text{Ad}_{\sigma^{-1}} \circ \zeta, \nu]_{\mathcal{G}} \\
&= R_{\nabla^{\mathcal{G}}}(\cdot, \cdot) \nu + \left[d^{\nabla^{\mathcal{G}}} \Delta\sigma + \frac{1}{2} [\Delta\sigma \frown \Delta\sigma]_{\mathcal{G}} - \text{Ad}_{\sigma^{-1}} \circ \zeta, \nu \right]_{\mathcal{G}} \tag{30}
\end{aligned}$$

for all $\nu \in \Gamma(\mathcal{G})$ and $\sigma \in \Gamma(\mathcal{G})$.

• On the one hand, if $\mu_{\mathcal{G}}$ satisfies Eq. (28), then $\omega \equiv 0$. As previously mentioned, we have $\Delta e \equiv 0$, so that we then get in total for Eq. (30) w.r.t. $\sigma := e$ that

$$R_{\nabla^{\mathcal{G}}}(\cdot, \cdot) \nu = [\zeta, \nu]_{\mathcal{G}}$$

for all $\nu \in \Gamma(\mathcal{G})$, which is the infinitesimal compatibility condition (25).

• On the other hand, if $\text{H}\mathcal{G}$ satisfies the infinitesimal compatibility condition (25), $R_{\nabla^{\mathcal{G}}}(\cdot, \cdot) \nu = [\zeta, \cdot]_{\mathcal{G}}$, then $\text{H}\mathcal{G}$ is an infinitesimal multiplicative Yang-Mills connection by Lemma 6.69, $\nabla^{\mathcal{G}} = \nabla^{\mathcal{G}}$, and we know by Eq. (27) (Lemma 6.78) that

$$\left[d^{\nabla^{\mathcal{G}}} \Delta\sigma + \frac{1}{2} [\Delta\sigma \frown \Delta\sigma]_{\mathcal{G}} - \text{Ad}_{\sigma^{-1}} \circ \zeta, \nu \right]_{\mathcal{G}} = -[\zeta, \nu]_{\mathcal{G}},$$

so that Eq. (30) then has the form

$$\left. \frac{d}{dt} \right|_{t=0} \gamma(t)^! \omega = R_{\nabla^{\mathcal{G}}}(\cdot, \cdot) \nu - [\zeta, \nu]_{\mathcal{G}} \stackrel{(25)}{=} 0$$

for all $\nu \in \Gamma(\mathcal{G})$ and $\sigma \in \Gamma(\mathcal{G})$. As aforementioned in the discussion around the definition of γ , also recall that ω vanishes if contracted with a vertical vector, we can conclude that ω is constant along the fibres of $\mathcal{G}$ which are connected, so that this implies

$$\omega = \pi_{\mathcal{G}}^! \xi$$

for a $\xi \in \Omega^2(M; \mathcal{G})$, especially

$$e^! \omega = (\pi_{\mathcal{G}} \circ e)^! \xi = \xi.$$

But we already have calculated this in Eq. (29), set $t = 0$ and $\sigma := e$, then also again by $\Delta e = 0$ we have

$$\xi = e^! \omega \stackrel{(29)}{=} \zeta - \zeta = 0.$$

Finally we get

$$\omega = \pi_{\mathcal{G}}^! \xi = 0,$$

which finishes the proof. ■

Trivially, this theorem is an extension of Eq. (27) (Lemma Lemma 6.78).

Corollary 6.84: Pullback of generalized Maurer-Cartan equation

Let $\mathcal{G} \xrightarrow{\pi_{\mathcal{G}}} M$ be an LGB over a smooth manifold M , $H\mathcal{G}$ an Ehresmann connection on $\mathcal{G}$ so that the total Maurer-Cartan form $\mu_{\mathcal{G}}$ satisfies the generalized Maurer-Cartan equation (28) w.r.t. a $\zeta \in \Omega^2(M; \mathcal{G})$. Then

$$d^{\nabla^{\mathcal{G}}} \Delta \sigma + \frac{1}{2} [\Delta \sigma \frown \Delta \sigma]_{\mathcal{G}} + \zeta = \text{Ad}_{\sigma^{-1}} \circ \zeta$$

for all $\sigma \in \Gamma(\mathcal{G})$.

Proof.

Keeping the same notation as in the proof of Thm. Theorem 6.81, we have derived Eq. (29), and due to Thm. Theorem 6.81 we get $\omega = 0$; altogether Eq. (29) at $t = 0$ gives then

$$0 = d^{\nabla^{\mathcal{G}}} \Delta \sigma + \frac{1}{2} [\Delta \sigma \frown \Delta \sigma]_{\mathcal{G}} + \zeta - \text{Ad}_{\sigma^{-1}} \circ \zeta.$$

Thus, we can finally define the connection we need on the LGB. ■

Definition 6.85: multiplicative Yang-Mills connection

Let $\mathcal{G} \rightarrow M$ be an LGB over a smooth manifold M , and $H\mathcal{G}$ be a horizontal distribution of $\mathcal{G}$. Then we say that $H\mathcal{G}$ is a **multiplicative Yang-Mills connection (w.r.t. a $\zeta \in \Omega^2(M; \mathcal{G})$)**, if it satisfies the **compatibility conditions**:

- (1) $H\mathcal{G}$ is an Ehresmann connection,
- (2) the total Maurer-Cartan form $\mu_{\mathcal{G}}$ satisfies the generalized Maurer-Cartan equation (w.r.t. a $\zeta \in \Omega^2(M; \mathcal{G})$).

By Thm. Theorem 6.75 we can equivalently define $H\mathcal{G}$ as a multiplicative Yang-Mills connection if

- (1) $\mu_{\mathcal{G}}$ is multiplicative,
- (2) $\mu_{\mathcal{G}}$ satisfies the generalized Maurer-Cartan equation (w.r.t. a $\zeta \in \Omega^2(M; \mathcal{G})$).

We then label $\mu_{\mathcal{G}}$ as a multiplicative Yang-Mills connection, too.

Remark 6.86: Multiplicative Yang-Mills connections integrate infinitesimal multiplicative Yang-Mills connections

By Lemma [Lemma 6.69](#) and Thm. [Theorem 6.81](#), every multiplicative Yang-Mills connection is also an infinitesimal multiplicative Yang-Mills connections, *i.e.* the infinitesimal compatibility conditions are also satisfied.

Remark 6.87: Relation to simplicial differential

In fact, Def. [Definition 6.73](#) comes from a simplicial differential (w.r.t. points on $\mathcal{G}$) defined in [\[15, beginning of §1.2\]](#). We do not have time to introduce it completely, but we give a short sketch about its definitions, also following the style of [\[13, appendix\]](#). On degree one the cited differential is a map

$$\begin{aligned}\Omega^\bullet(\mathcal{G}; \pi_{\mathcal{G}}^* \mathcal{G}) &\rightarrow \Omega^\bullet(\mathcal{G} * \mathcal{G}; \pi_{\mathcal{G}}^* \mathcal{G}), \\ \omega &\mapsto \delta\omega,\end{aligned}$$

with

$$(\delta\omega)_{(g,q)} := \mathcal{A}d_{q^{-1}} \circ (\text{pr}_1^! \omega)_{(g,q)} + (\text{pr}_2^! \omega)_{(g,q)} - (\Phi^! \omega)_{(g,q)}$$

for all $(g, q) \in \mathcal{G} * \mathcal{G}$, where $\Phi : \mathcal{G} * \mathcal{G} \rightarrow \mathcal{G}$ is the $\mathcal{G}$ -multiplication, and $\text{pr}_i : \mathcal{G} * \mathcal{G} \rightarrow \mathcal{G}$ are the canonical projections onto the i -th component ($i \in \{1, 2\}$); $\pi_{\mathcal{G}}$ is canonically extended to $\mathcal{G} \times \mathcal{G}$. Recall Def. [Definition 6.73](#) and its Rem. [Theorem 6.74](#), ω can be defined to be multiplicative if $\delta\omega = 0$, *i.e.* ω is closed w.r.t. δ .

On degree 0 δ is a map $\Omega^\bullet(M; \mathcal{G}) \rightarrow \Omega^\bullet(\mathcal{G}; \pi_{\mathcal{G}}^* \mathcal{G})$ given for $\xi \in \Omega^\bullet(M; \mathcal{G})$ by

$$(\delta\xi)_g := (g, \text{Ad}_{g^{-1}} \circ \xi_x \circ \underbrace{(D_g \pi_{\mathcal{G}}, \dots, D_g \pi_{\mathcal{G}})}_{\text{times the degree of } \xi}) - (\pi_{\mathcal{G}}^! \xi)|_g$$

for all $g \in \mathcal{G}$. Then observe that we can rewrite the generalized Maurer-Cartan equation to

$$F_{\mathcal{G}} = \delta\zeta,$$

where

$$F_{\mathcal{G}} = d^{\pi_{\mathcal{G}}^* \nabla^{\mathcal{G}}} \mu_{\mathcal{G}} + \frac{1}{2} [\mu_{\mathcal{G}} \wedge \mu_{\mathcal{G}}]_{\pi_{\mathcal{G}}^* \mathcal{G}} \in \Omega^2(\mathcal{G}; \pi_{\mathcal{G}}^* \mathcal{G}).$$

It is straight-forward to check that $\delta F_{\mathcal{G}} = 0$ (see *e.g.* [\[12, §4.6, Thm. 4.27, Eq. \(53\)\]](#) or [\[13, §2.5, Prop. 2.22\]](#)) by making use of that $H\mathcal{G}$ is an Ehresmann connection. In total we concluded that compatibility condition (24) is implied, if $\mu_{\mathcal{G}}$ is multiplicative, which also implies that $F_{\mathcal{G}}$ is multiplicative and thus δ -closed; additionally with compatibility condition (25) we achieve δ -exactness of $F_{\mathcal{G}}$. This is no wonder, the compatibility conditions already tell us by definition that the connection-1-form of ∇ is closed w.r.t. the Chevalley-Eilenberg complex, which also implies the same for the curvature, however the curvature has to be exact by (25). We depict that as $\delta^{\text{CE}} \nabla^{\mathcal{G}} = 0$ and $R_{\nabla^{\mathcal{G}}} = \delta^{\text{CE}} \zeta$; a summary is shown in Table [Table 4](#).

TABLE 4. Infinitesimal compatibility conditions on the LAB $\mathcal{G}$

and integrated compatibility conditions on the overlying LGB $\mathcal{G}$

	$\mathcal{G}$	$\mathcal{G}$
Closedness	$\delta^{\text{CE}} \nabla^{\mathcal{G}} = 0 \ (\Rightarrow \delta^{\text{CE}} R_{\nabla^{\mathcal{G}}} = 0)$	$\delta\mu_{\mathcal{G}} = 0 \ (\Rightarrow \delta F_{\mathcal{G}} = 0)$
Exactness	$R_{\nabla^{\mathcal{G}}} = \delta^{\text{CE}} \zeta$	$F_{\mathcal{G}} = \delta\zeta$

To conclude this subsection, let us provide canonical examples of multiplicative Yang-Mills connections; we will make use of Thm. [Theorem 6.81](#) in order to prove that the generalized Maurer-Cartan equation is satisfied.

Example 6.88: (Pre-)Classical multiplicative Yang-Mills connection

Let us look at a classical principal bundle P with trivial LGB $\mathcal{G} = M \times G$, G a Lie group; recall Ex. [Example 6.3](#) and [Example 6.46](#). The canonical flat connection $H\mathcal{G}$ on $\mathcal{G}$ induces the canonical flat connection $\nabla^{\mathcal{G}}$ on its LAB $\mathcal{Q} = M \times \mathfrak{g}$, recall Ex. [Example 6.35](#). Thus,

$$R_{\nabla^{\mathcal{G}}} = 0$$

and so we can set $\zeta \equiv 0$. By Ex. [Example 6.46](#) (viewing $\mathcal{G}$ also as a principal bundle equipped with the same horizontal distribution) we know that $H\mathcal{G}$ is an Ehresmann connection, so that its total Maurer-Cartan form is multiplicative.

We therefore call flat $\nabla^{\text{cYM}} := \nabla^{\mathcal{G}}$ a **classical multiplicative Yang-Mills connection**, regardless of whether or not $\mathcal{G}$ is trivial.

However, observe that ζ can be centre-valued and non-zero while $\nabla^{\mathcal{G}}$ is still flat. In such an occasion we speak of a **pre-classical multiplicative Yang-Mills connection**, denoted by ∇^{pYM} .

Example 6.89: Associated connections of inner group bundles are multiplicative Yang-Mills connections

Recall Ex. [Example 6.47](#), its setup and notation. We have discussed a canonical connection on associated LGBs $\mathcal{H} = P \times_{\psi} H$, called associated connection which is an Ehresmann connection.

As it is well-known, the curvature measures the holonomy, that is, the parallel transport over closed (contractible) curves. Together with Eq. (25), the parallel transport over a closed (contractible) curve α should act as conjugation with some group element. Let us see whether this holds here; there is a unique $g \in G$ for $p \in P$ such that

$$\text{PT}_{\alpha}^P(p) = p \cdot g,$$

and thus for $[p, h] \in \mathcal{H}$

$$\text{PT}_{\alpha}^{\mathcal{H}}([p, h]) = [p \cdot g, h] = [p, \psi_g(h)].$$

Thus, if $\mathcal{H} = c_G(P)$, the inner group bundle of Ex. [Example 2.6](#), we get

$$\text{PT}_{\alpha}^{\mathcal{H}}([p, h]) = [p, ghg^{-1}] = [p, g] \cdot [p, h] \cdot [p, g^{-1}],$$

the conjugation of $[p, h]$ with $[p, g]$ in $c_G(P)$. It follows that there is a $\zeta \in \Omega^2(M; \mathfrak{g})$ such that Eq. (25) is satisfied. This also implies that the associated connection of an inner group bundle is a multiplicative Yang-Mills connection. We will not prove this approach, instead the next approach provides an alternative proof with an explicit expression for ζ .

Example 6.90: multiplicative Yang-Mills connections on inner group bundles: Alternative point of view and explanation

Alternatively one can prove that the total Maurer-Cartan form solves the generalized Maurer-Cartan equation in Ex. [Example 6.89](#) by using the aforementioned relationship of the compatibility conditions with Mackenzie's study about extending Lie algebroids via LABs; recall Remark [Theorem 6.68](#). The following is an information for the experienced reader: The LAB of $c_G(P)$ is given by the adjoint bundle $\text{Ad}(P)$; recall Ex. [Example 4.12](#). $\text{Ad}(P)$ is the kernel of the Atiyah sequence (see *e.g.* [10, §3.2, page 90ff.]), a short exact sequence of Lie algebroids over M

$$0 \longrightarrow \text{Ad}(P) \longrightarrow \text{At}(P) \longrightarrow TM \longrightarrow 0,$$

where $\text{At}(P)$ is the Atiyah bundle of P . A connection on P in the "classical" sense (that is, related to a canonical flat connection on $M \times G$ in our sense) has a 1:1 correspondence to a splitting $\chi : TM \rightarrow \text{At}(P)$ to that sequence, *i.e.* a section of the right non-trivial arrow. We can then construct a vector bundle connection $\nabla^{\mathcal{G}}$ on the adjoint bundle by using the Lie algebroid bracket $[\cdot, \cdot]_{\text{At}(P)}$ on $\text{At}(P)$, that is,

$$\nabla_X^{\mathcal{G}} \nu = [\chi(X), \nu]_{\text{At}(P)}$$

for all $X \in \mathfrak{X}(M)$ and $\nu \in \Gamma(\text{Ad}(P))$. It follows by straight-forward calculations as in [10, §7.3, Prop. 7.3.2 and Lemma 7.3.3, page 278] that $\nabla^{\mathcal{G}}$ is indeed a multiplicative Yang-Mills connection with $\zeta \in \Omega^2(M; \text{Ad}(P))$ for example given by

$$\zeta(Y, Z) := [\chi(Y), \chi(Z)]_{\text{At}(P)} - \chi([Y, Z])$$

for all $Y, Z \in \mathfrak{X}(M)$. In other words, $\nabla^{\mathcal{G}}$ is the **adjoint connection** induced by a connection on P (see for example [10, §5.3, especially Prop. 5.3.13, page 199]). Thence, by the discussion in [17, §5.9, page 289ff.] $\nabla^{\mathcal{G}}$ corresponds to the natural parallel transport $\text{PT}^{\text{Ad}(P)}$ on $\text{Ad}(P)$ (omitting the notation of the corresponding curve) given by

$$\text{PT}^{\text{Ad}(P)}([p, v]) := [\text{PT}^P(p), v]$$

for all $[p, v] \in \text{Ad}(P)$. This is clearly just the infinitesimal parallel transport inherited by $\text{PT}^{c_G(P)}$, so that we can conclude that $\nabla^{\mathcal{G}}$ corresponds to the multiplicative Yang-Mills connection on $c_G(P)$ in sense of Def. Definition 6.33 (also recall Rem. Theorem 6.34).

This proof gives a certain interpretation of $\nabla^{\mathcal{G}}$ and ζ : In the case of $\mathcal{G}$ being the inner group bundle of a typical gauge theory related to a principal G -bundle P , one can view $\nabla^{\mathcal{G}}$ as the adjoint connection on the adjoint bundle, induced by χ a classical connection on P , this implies by Eq. (25) that ζ is the field strength related to χ (as a field strength with values in the adjoint bundle). Thus one may say in this case that the generalised sense of connection as in Def. Definition 6.50 is modified by a classical connection from a classical principal bundle, thence also the label as multiplicative Yang-Mills connection to reflect its relation to the classical theory.

6.4.2. *Field strength related to multiplicative Yang-Mills connections.* Let us now introduce and discuss the field strength.

Definition 6.91: (Generalized) Field strength

Let $\mathcal{G} \rightarrow M$ be an LGB over a smooth manifold M and $\mathcal{P} \xrightarrow{\pi} M$ a principal $\mathcal{G}$ -bundle, also let $\text{H}\mathcal{G}$ be a multiplicative Yang-Mills connection on $\mathcal{G}$ (w.r.t. a $\zeta \in \Omega^2(M; \mathfrak{g})$) and $A \in \Omega^1(\mathcal{P}; \pi^*\mathfrak{g})$ be a connection 1-form on $\mathcal{P}$. Then we define the **(generalized) curvature** or **(generalized) field strength F (of A)** as an element of $\Omega^2(\mathcal{P}; \pi^*\mathfrak{g})$ by

$$F := d^{\pi^*\nabla^{\mathcal{G}}} A \circ (\pi^{\text{H}\mathcal{P}}, \pi^{\text{H}\mathcal{P}}) + \pi^!\zeta,$$

where $\pi^{\text{H}\mathcal{P}} : T\mathcal{P} \rightarrow \text{H}\mathcal{P}$ is the canonical projection onto the associated Ehresmann connection $\text{H}\mathcal{P}$ on $\mathcal{P}$; that is,

$$F(X, Y) = d^{\pi^*\nabla^{\mathcal{G}}} A(\pi^{\text{H}\mathcal{P}}(X), \pi^{\text{H}\mathcal{P}}(Y)) + (\pi^*\zeta)(D\pi(X), D\pi(Y))$$

for all $X, Y \in \mathfrak{X}(\mathcal{P})$.

Of course, we expect certain properties, naturally generalized by our previous discussions:

Proposition 6.92: Properties of the generalized field strength

Let $\mathcal{G} \rightarrow M$ be an LGB over a smooth manifold M and $\mathcal{P} \xrightarrow{\pi} M$ a principal $\mathcal{G}$ -bundle, also let $H\mathcal{G}$ be a multiplicative Yang-Mills connection on $\mathcal{G}$ (w.r.t. a $\zeta \in \Omega^2(M; \mathfrak{g})$) and $A \in \Omega^1(\mathcal{P}; \pi^*\mathfrak{g})$ be a connection 1-form on $\mathcal{P}$. Then we have the following properties of the field strength:

- **(Form of type $\mathcal{A}d$)**

$$\mathcal{r}_\sigma^! F = \mathcal{A}d_{\sigma^{-1}} \circ F$$

for all (local) $\sigma \in \Gamma(\mathcal{G})$.

- **(Horizontal form)**

For $X, Y \in T_p\mathcal{P}$ ($p \in \mathcal{P}$) we have

$$F(X, Y) = 0$$

if either of X and Y is vertical.

However, despite the easy proof in the classical theory, the proof of the first property in Prop. Proposition 6.92 will be rather involved; most of the proof is straight-forward calculation, and then use our discussion about multiplicative Yang-Mills connections; the quintessence will be about using Lemma Lemma 6.79 (for first order terms w.r.t. σ) and Cor. Corollary 6.84 (for second order terms w.r.t. σ).

Proof of Prop. Proposition 6.92.

For this proof we will neglect caring for the base point information in the involved pullback bundles since the base point information is clear by context; recall Subsection Section 1.3. In that sense we also have $\mathcal{A}d_\sigma \cong \pi^* \text{Ad}_\sigma \cong \text{Ad}_\sigma$ for all $\sigma \in \Gamma(\mathcal{G})$ etc.; this will simplify the following notation a bit.

- The second bullet point quickly follows by construction: The first summand of F (recall Def. Definition 6.91) is clearly zero if either of X and Y is vertical; similar holds for the second summand because the vertical bundle is the kernel of $D\pi$.

- Thence, let us focus on proving the first bullet point, and denote with $\pi_h : T\mathcal{P} \rightarrow H\mathcal{P}$ the canonical projection onto the associated Ehresmann connection $H\mathcal{P}$ on $\mathcal{P}$, so that we write

$$F = d^{\pi^* \nabla^{\mathcal{G}}} A \circ (\pi_h, \pi_h) + \pi^! \zeta.$$

We have for $\sigma \in \Gamma(\mathcal{G})$

$$\begin{aligned} (\mathcal{r}_\sigma^! \pi^! \zeta)_p(X, Y) &= \zeta_x(D_{p \cdot \sigma_x} \pi(\mathcal{r}_{\sigma_x^*}(X)), D_{p \cdot \sigma_x} \pi(\mathcal{r}_{\sigma_x^*}(Y))), \\ &= \zeta_x(\underbrace{D_{p \cdot \sigma_x} \pi(D_p r_\sigma(X))}_{=D_p(\pi \circ r_\sigma) = D_p \pi}, D_{p \cdot \sigma_x} \pi(D_p r_\sigma(Y))) \\ &= (\pi^! \zeta)_p(X, Y) \end{aligned}$$

for all $p \in \mathcal{P}_x$ ($x \in M$) and $X, Y \in T_p\mathcal{P}$, making use of that fundamental vector fields are in the kernel of $D\pi$ so that $D\pi \circ \mathcal{r}_{\sigma_x^*} = D\pi \circ r_{\sigma_x^*}$. For the first term in the field strength use Cor. Corollary 6.53 and Remark Theorem 6.54 so that

$$\begin{aligned} \mathcal{r}_\sigma^! (d^{\pi^* \nabla^{\mathcal{G}}} A \circ (\pi_h, \pi_h)) &= (r_\sigma^* d^{\pi^* \nabla^{\mathcal{G}}} A) \circ (r_\sigma^* \pi_h \circ \mathcal{r}_{\sigma^*}, r_\sigma^* \pi_h \circ \mathcal{r}_{\sigma^*}) \\ &= (r_\sigma^* d^{\pi^* \nabla^{\mathcal{G}}} A) \circ (\mathcal{r}_{\sigma^*} \circ \pi_h, \mathcal{r}_{\sigma^*} \circ \pi_h) \\ &= \mathcal{r}_\sigma^! (d^{\pi^* \nabla^{\mathcal{G}}} A) \circ (\pi_h, \pi_h). \end{aligned}$$

Let us denote $X^h := \pi_h(X)$ and $Y^h := \pi_h(Y)$, where X and Y are now elements of $\mathfrak{X}(\mathcal{P})$; since we are looking at tensorial equations we can w.l.o.g. assume that X^h and Y^h are the horizontal lifts of vector fields ω^X and ω^Y , respectively, on M , especially $D\pi(X^h) = D\pi(X) = \pi^* \omega^X$ and

$D\pi(Y^h) = D\pi(Y) = \pi^*\omega^Y$. We also rewrite

$$\mathcal{r}_{\sigma*}(X) = Dr_{\sigma}(X) - r_{\sigma}^*\left(\overline{(\pi^!\Delta\sigma)(X)}\right) \in \Gamma(r_{\sigma}^*\mathcal{T}\mathcal{P}) \cong \mathfrak{X}(\mathcal{P}).$$

In the following we will make use of the canonical identification $\mathfrak{X}(\mathcal{P}) \cong \Gamma(r_{\sigma}^*\mathcal{T}\mathcal{P})$, $X \mapsto r_{\sigma}^*X$, especially in order to calculate Lie brackets.

Then make use of Def. [Definition 6.44](#) and Thm. [Theorem 6.52](#) to show

$$\begin{aligned} & r_{\sigma}^*\left(\left(d^{\pi^*\nabla^{\mathcal{G}}}A\right)(\mathcal{r}_{\sigma*}(X^h), \mathcal{r}_{\sigma*}(Y^h))\right) \\ &= r_{\sigma}^*\left(\left(\pi^*\nabla^{\mathcal{G}}\right)_{\mathcal{r}_{\sigma*}(X^h)}\underbrace{\left((A \circ \mathcal{r}_{\sigma*})(Y^h)\right)}_{=0} \right. \\ &\quad \left. - \left(\pi^*\nabla^{\mathcal{G}}\right)_{\mathcal{r}_{\sigma*}(Y^h)}\left((A \circ \mathcal{r}_{\sigma*})(X^h)\right) \right. \\ &\quad \left. - A([\mathcal{r}_{\sigma*}(X^h), \mathcal{r}_{\sigma*}(Y^h)])\right) \\ &= r_{\sigma}^*\left(-A([\mathcal{r}_{\sigma*}(X^h), \mathcal{r}_{\sigma*}(Y^h)])\right) \\ &= r_{\sigma}^*\left(-A([Dr_{\sigma}(X^h), Dr_{\sigma}(Y^h)]) \right. \\ &\quad \left. + A([Dr_{\sigma}(X^h), r_{\sigma}^*\left(\overline{(\pi^!\Delta\sigma)(Y^h)}\right)]) \right. \\ &\quad \left. + A\left([r_{\sigma}^*\left(\overline{(\pi^!\Delta\sigma)(X^h)}\right), Dr_{\sigma}(Y^h)]\right) \right. \\ &\quad \left. - A\left([r_{\sigma}^*\left(\overline{(\pi^!\Delta\sigma)(X^h)}\right), r_{\sigma}^*\left(\overline{(\pi^!\Delta\sigma)(Y^h)}\right)]\right)\right). \end{aligned} \quad (31)$$

Let us look at each summand individually; for the first summand we use the very well-known fact (see *e.g.* [\[17, §A.1.11, Cor. A.1.51, page 615\]](#)) that

$$[Dr_{\sigma}(X^h), Dr_{\sigma}(Y^h)] = Dr_{\sigma}([X^h, Y^h]),$$

making use of that r_{σ} is an isomorphism, and thus we get for the first summand by using Def. [Definition 6.50](#)

$$\begin{aligned} & r_{\sigma}^*\left(-A([Dr_{\sigma}(X^h), Dr_{\sigma}(Y^h)])\right) \\ &= -(r_{\sigma}^*A)(\mathcal{r}_{\sigma*}([X^h, Y^h])) - (r_{\sigma}^*A)\left(r_{\sigma}^*\left(\overline{(\pi^!\Delta\sigma)([X^h, Y^h])}\right)\right) \\ &= -(\mathcal{r}_{\sigma}^!A)([X^h, Y^h]) - r_{\sigma}^*((\pi^!\Delta\sigma)([X^h, Y^h])) \\ &= -(\mathcal{A}d_{\sigma^{-1}} \circ A)([X^h, Y^h]) - r_{\sigma}^*((\pi^!\Delta\sigma)([X^h, Y^h])) \\ &= \mathcal{A}d_{\sigma^{-1}}\left(\left(d^{\pi^*\nabla^{\mathcal{G}}}A\right)(X^h, Y^h)\right) - r_{\sigma}^*((\pi^!\Delta\sigma)([X^h, Y^h])), \end{aligned}$$

where we used a similar argument for the last equality as for the beginning of the calculation for Eq. (31). We know by Remark [Remark 5.14](#) that the map to fundamental vector fields is a homomorphism of Lie algebras, so that

$$\left[r_{\sigma}^*\left(\overline{(\pi^!\Delta\sigma)(X^h)}\right), r_{\sigma}^*\left(\overline{(\pi^!\Delta\sigma)(Y^h)}\right)\right] = r_{\sigma}^*\left(\overline{[(\pi^!\Delta\sigma)(X^h), (\pi^!\Delta\sigma)(Y^h)]_{\pi^*\mathcal{G}}}\right),$$

and therefore we can derive for the fourth term in Eq. (31) that

$$(r_{\sigma}^*A)\left(\left[r_{\sigma}^*\left(\overline{(\pi^!\Delta\sigma)(X^h)}\right), r_{\sigma}^*\left(\overline{(\pi^!\Delta\sigma)(Y^h)}\right)\right]\right) = r_{\sigma}^*\left([\overline{(\pi^!\Delta\sigma)(X^h)}, \overline{(\pi^!\Delta\sigma)(Y^h)}]_{\pi^*\mathcal{G}}\right).$$

For the second and third summands in Eq. (31) we want to use Lemma 6.79 and our result of the fourth summand to show

$$\begin{aligned}
& (r_\sigma^* A) \left(\left[Dr_\sigma(X^h), r_\sigma^* \left(\overline{(\pi^! \Delta \sigma)(Y^h)} \right) \right] \right) \\
&= (r_\sigma^* A) \left(\left[\mathcal{P}_{\sigma^*}(X^h), r_\sigma^* \left(\overline{(\pi^! \Delta \sigma)(Y^h)} \right) \right] \right) \\
&\quad + (r_\sigma^* A) \left(\left[r_\sigma^* \left(\overline{(\pi^! \Delta \sigma)(X^h)} \right), r_\sigma^* \left(\overline{(\pi^! \Delta \sigma)(Y^h)} \right) \right] \right) \\
&= r_\sigma^* \left((\pi^* \nabla^\mathcal{G})_{\mathcal{P}_{\sigma^*}(X^h)} \left((\pi^! \Delta \sigma)(Y^h) \right) \right) + r_\sigma^* \left([(\pi^! \Delta \sigma)(X^h), (\pi^! \Delta \sigma)(Y^h)]_{\pi^* \mathcal{G}} \right)
\end{aligned}$$

using that $\mathcal{P}_{\sigma^*}(X^h)$ is horizontal due to the fact that X^h is horizontal, and, last but not least, we can write for terms like

$$(\pi^! \Delta \sigma)(Y^h) = (\pi^* \Delta \sigma)(D\pi(Y^h)) = (\pi^* \Delta \sigma)(\pi^* \omega^Y) = \pi^* \left((\Delta \sigma)(\omega^Y) \right),$$

and

$$D\pi(\mathcal{P}_{\sigma^*}(X^h)) = D\pi \left(Dr_\sigma(X^h) - r_\sigma^* \left(\overline{(\pi^! \Delta \sigma)(X^h)} \right) \right) = D \underbrace{(\pi \circ r_\sigma)}_{=\pi}(X^h) = \pi^* \omega^X,$$

using that fundamental vector fields are in the kernel of $D\pi$, and therefore

$$(\pi^* \nabla^\mathcal{G})_{\mathcal{P}_{\sigma^*}(X^h)} \left((\pi^! \Delta \sigma)(Y^h) \right) = \pi^* \left(\nabla_{\omega^X}^\mathcal{G} \left((\Delta \sigma)(\omega^Y) \right) \right).$$

Collecting all these results and using Cor. 6.84 we can write Eq. (31) as

$$\begin{aligned}
& r_\sigma^* \left(\left(d^{\pi^* \nabla^\mathcal{G}} A \right) (\mathcal{P}_{\sigma^*}(X^h), \mathcal{P}_{\sigma^*}(Y^h)) \right) \\
&= \mathcal{A}d_{\sigma^{-1}} \left(\left(d^{\pi^* \nabla^\mathcal{G}} A \right) (X^h, Y^h) \right) \\
&\quad + r_\sigma^* \left(\pi^* \left(\nabla_{\omega^X}^\mathcal{G} \left((\Delta \sigma)(\omega^Y) \right) \right) - \pi^* \left(\nabla_{\omega^Y}^\mathcal{G} \left((\Delta \sigma)(\omega^X) \right) \right) \right. \\
&\quad \left. - \underbrace{(\pi^! \Delta \sigma)([X^h, Y^h])}_{=\pi^* \left((\Delta \sigma)([\omega^X, \omega^Y]) \right)} + \left[\pi^* \left((\Delta \sigma)(\omega^X) \right), \pi^* \left((\Delta \sigma)(\omega^Y) \right) \right]_{\pi^* \mathcal{G}} \right) \\
&= \mathcal{A}d_{\sigma^{-1}} \left(\left(d^{\pi^* \nabla^\mathcal{G}} A \right) (X^h, Y^h) \right) + \underbrace{r_\sigma^* \pi^*}_{=\pi^*} \left(\left(d^{\nabla^\mathcal{G}} \Delta \sigma + \frac{1}{2} [\Delta \sigma \wedge \Delta \sigma]_\mathcal{G} \right) (\omega^X, \omega^Y) \right) \\
&= \mathcal{A}d_{\sigma^{-1}} \left(\left(d^{\pi^* \nabla^\mathcal{G}} A \right) (X^h, Y^h) \right) + (\pi^* (\mathcal{A}d_{\sigma^{-1}} \circ \zeta - \zeta)) (D\pi(X), D\pi(Y)) \\
&= \mathcal{A}d_{\sigma^{-1}} \left(\left(d^{\pi^* \nabla^\mathcal{G}} A \right) (X^h, Y^h) \right) + ((\pi^! (\mathcal{A}d_{\sigma^{-1}} \circ \zeta - \zeta))(X, Y)) \\
&= \left(\mathcal{A}d_{\sigma^{-1}} \circ \left(d^{\pi^* \nabla^\mathcal{G}} A \circ (\pi_h, \pi_h) + \pi^! \zeta \right) - \pi^! \zeta \right) (X, Y)
\end{aligned}$$

where we used that $D\pi(X) = \pi^* \omega^X$ and $D\pi(Y) = \pi^* \omega^Y$, which immediately implies

$$D\pi([X, Y]) = \pi^*([\omega^X, \omega^Y]),$$

which is a well-known fact, as also given in [17, Proposition A.1.49; page 615], but also straightforward to check.

Finally, we can conclude this proof by combining the previous result with our first calculation of this proof, that is,

$$\mathcal{P}_\sigma^! F = \mathcal{A}d_{\sigma^{-1}} \circ \left(d^{\pi^* \nabla^\mathcal{G}} A \circ (\pi_h, \pi_h) + \pi^! \zeta \right) - \pi^! \zeta + \pi^! \zeta = \mathcal{A}d_{\sigma^{-1}} \circ F.$$

■

We achieve of course also a structure equation.

Proposition 6.93: Structure equation of the generalized field strength

Let $\mathcal{G} \rightarrow M$ be an LGB over a smooth manifold M and $\mathcal{P} \xrightarrow{\pi} M$ a principal $\mathcal{G}$ -bundle, also let $H\mathcal{G}$ be a multiplicative Yang-Mills connection on $\mathcal{G}$ (w.r.t. a $\zeta \in \Omega^2(M; \mathfrak{g})$) and $A \in \Omega^1(\mathcal{P}; \pi^*\mathfrak{g})$ be a connection 1-form on $\mathcal{P}$. Then we have the **structure equation**

$$F = d\pi^*\nabla^{\mathcal{G}} A + \frac{1}{2}[A \wedge A]_{\pi^*\mathfrak{g}} + \pi^!\zeta.$$

Remark 6.94: The generalized field strength of the total Maurer-Cartan form

Again viewing $\mathcal{G} \xrightarrow{\pi} M$ as the principal bundle itself with $H\mathcal{P} = H\mathcal{G}$, we know by Cor. [Corollary 6.71](#) that the total Maurer-Cartan form $\mu_{\mathcal{G}}$ is the connection 1-form corresponding to $H\mathcal{G}$. Denoting the associated generalized field strength by $F(\mu_{\mathcal{G}})$, the structure equation and the generalized Maurer-Cartan equation, Thm. [Theorem 6.81](#) (also recall Rem. [Theorem 6.82](#)), imply

$$F_g(\mu_{\mathcal{G}}) = \mathcal{A}_{g^{-1}} \circ \pi_{\mathcal{G}}^! \zeta|_g$$

for all $g \in \mathcal{G}_x$ ($x \in M$). This implies that $F(\mu_{\mathcal{G}})$ is fully encoded by ζ , especially we have

$$e^!(F(\mu_{\mathcal{G}})) = \zeta, \quad (32)$$

where e is the neutral section of $\mathcal{G}$. Thus, ζ is the curvature of $\mu_{\mathcal{G}}$ restricted on M , and

$$F = d\pi^*\nabla^{\mathcal{G}} A + \frac{1}{2}[A \wedge A]_{\pi^*\mathfrak{g}} + (e \circ \pi)^!(F(\mu_{\mathcal{G}})).$$

Following similar calculations as in the proof of Prop. [Proposition 6.92](#), it is trivial to show that the first two summands measure the failure of $H\mathcal{P}$ being a foliation, hence this meaning carries over from the classical theory as also argued in [\[13, §2.5, Cor. 2.23\]](#) for LGBs. Due to the third summand, $H\mathcal{P}$ is involutive if and only if

$$F = (e \circ \pi)^!(F(\mu_{\mathcal{G}})).$$

In that sense Eq. (32) reflects the fact that M is a horizontal leave of $H\mathcal{G}$. Furthermore, F can be non-zero while $H\mathcal{P}$ is involutive, if $F(\mu_{\mathcal{G}})$ is non-zero along M .

Proof of Thm. [Proposition 6.93](#).

The idea of the proof is similar to the proof of the "classical" statement; following the structure of [\[17, §5.5, Lemma 5.5.5, page 276\]](#), see also works like [\[12, §4.6, Lemma 4.25\]](#) and [\[13, §2.5, Prop. 2.24\]](#) where something similar is shown on Lie groupoids instead of principal bundles, but with no ζ . That is, we will look at vertical and horizontal tangent vectors and their mixed terms; recall Def. [Definition 6.50](#) and Thm. [Theorem 6.52](#).

- We have by Prop. [Proposition 6.92](#)

$$F(\bar{\nu}, \bar{\mu}) = 0$$

for all $\mu, \nu \in \Gamma(\mathfrak{g})$. Regarding the right hand side of the structure equation, this trivially also holds for the third summand $(\pi^!\zeta)(\bar{\nu}, \bar{\mu}) = 0$. We also have

$$\left(\frac{1}{2}[A \wedge A]_{\pi^*\mathfrak{g}}\right)(\bar{\nu}, \bar{\mu}) = [A(\bar{\nu}), A(\bar{\mu})]_{\pi^*\mathfrak{g}} = [\pi^*\nu, \pi^*\mu]_{\pi^*\mathfrak{g}} = \pi^*([\nu, \mu]_{\mathfrak{g}}),$$

and

$$\begin{aligned} \left(d^{\pi^* \nabla^{\mathcal{G}}} A \right) (\bar{\nu}, \bar{\mu}) &= \underbrace{(\pi^* \nabla^{\mathcal{G}})_{\bar{\nu}} (A(\bar{\mu}))}_{=\pi^* \left(\nabla_{D\pi(\bar{\nu})}^{\mathcal{G}} \mu \right) = 0} - \underbrace{(\pi^* \nabla^{\mathcal{G}})_{\bar{\mu}} (A(\bar{\nu})) - A([\bar{\nu}, \bar{\mu}])}_{= [\nu, \mu]} = -\pi^* ([\nu, \mu]_{\mathcal{G}}). \end{aligned}$$

Thus,

$$\left(d^{\pi^* \nabla^{\mathcal{G}}} A + \frac{1}{2} [A \wedge A]_{\pi^* \mathcal{G}} + \pi^! \zeta \right) (\bar{\nu}, \bar{\mu}) = 0 = F(\bar{\nu}, \bar{\mu}).$$

- Let X and Y be horizontal vector fields of $\mathcal{P}$, then clearly

$$\left(\frac{1}{2} [A \wedge A]_{\pi^* \mathcal{G}} \right) (X, Y) = [A(X), A(Y)]_{\pi^* \mathcal{G}} = 0,$$

then clearly

$$\left(d^{\pi^* \nabla^{\mathcal{G}}} A + \frac{1}{2} [A \wedge A]_{\pi^* \mathcal{G}} + \pi^! \zeta \right) (X, Y) = \left(d^{\pi^* \nabla^{\mathcal{G}}} A \circ (\pi_h, \pi_h) + \pi^! \zeta \right) (X, Y) = F(X, Y),$$

where $\pi_h : T\mathcal{P} \rightarrow H\mathcal{P}$ denotes the canonical projection onto the horizontal bundle $H\mathcal{P}$ (especially, $\pi_h(X) = X$, $\pi_h(Y) = Y$).

- Now let $X \in \mathfrak{X}(\mathcal{P})$ be again horizontal and $\nu \in \Gamma(\mathcal{G})$; then again by Prop. [Proposition 6.92](#)

$$F(X, \bar{\nu}) = 0,$$

as also clearly $(\pi^! \zeta)(X, \bar{\nu}) = 0$, furthermore

$$\left(\frac{1}{2} [A \wedge A]_{\pi^* \mathcal{G}} \right) (X, \bar{\nu}) = [A(X), A(\bar{\nu})]_{\pi^* \mathcal{G}} = 0,$$

and by Lemma [Lemma 6.79](#)

$$\left(d^{\pi^* \nabla^{\mathcal{G}}} A \right) (X, \bar{\nu}) = (\pi^* \nabla^{\mathcal{G}})_X (\pi^* \nu) - A([X, \bar{\nu}]) \stackrel{\text{Lemma 6.79}}{=} 0,$$

which concludes the proof with

$$F(X, \bar{\nu}) = \left(d^{\pi^* \nabla^{\mathcal{G}}} A + \frac{1}{2} [A \wedge A]_{\pi^* \mathcal{G}} + \pi^! \zeta \right) (X, \bar{\nu}).$$

■

Concluding this subsubsection, we also achieve a generalized Bianchi identity, which we already have proven in [\[5, §7, Thm. 7.3\]](#) and [\[6, §5, Thm. 5.1.42\]](#); the proof is straightforward to show, making use of the infinitesimal compatibility conditions, Def. [Definition 6.67](#).

Lemma 6.95: Generalized Bianchi identity, [\[6, §5, Thm. 5.1.42\]](#)

Let $\mathcal{G} \rightarrow M$ be an LGB over a smooth manifold M and $\mathcal{P} \xrightarrow{\pi} M$ a principal $\mathcal{G}$ -bundle, also let $H\mathcal{G}$ be a multiplicative Yang-Mills connection on $\mathcal{G}$ (w.r.t. a $\zeta \in \Omega^2(M; \mathfrak{g})$) and $A \in \Omega^1(\mathcal{P}; \pi^* \mathfrak{g})$ be a connection 1-form on $\mathcal{P}$. Then we have the **(generalized) Bianchi identity**

$$d^{\pi^* \nabla^{\mathcal{G}}} F + [A \wedge F]_{\pi^* \mathcal{G}} = \pi^! d^{\nabla^{\mathcal{G}}} \zeta.$$

6.4.3. Gauge transformation of the generalized field strength. Similar to Subsection [Section 6.3](#) we will now discuss the gauge transformation of the generalized field strength F ; recall Thm. [Proposition 6.58](#).

Proposition 6.96: Gauge transformation of the generalized field strength

Let $\mathcal{G} \rightarrow M$ be an LGB over a smooth manifold M and $\mathcal{P} \xrightarrow{\pi} M$ a principal $\mathcal{G}$ -bundle, also let $H\mathcal{G}$ be a multiplicative Yang-Mills connection on $\mathcal{G}$ and $A \in \Omega^1(\mathcal{P}; \pi^*\mathcal{G})$ be a connection 1-form on $\mathcal{P}$. Furthermore, let $H \in \mathcal{Aut}(\mathcal{P})$. We then have that $H^!F$ is the field strength related to $H^!A$ and

$$H^!F = \mathcal{Ad}_{\text{pr}_2 \circ (\sigma^H)^{-1}} \circ F,$$

where $\sigma^H \in C^\infty(\mathcal{P}; \mathcal{G})^\mathcal{G}$ is defined as in Prop. Proposition 6.57 and $\text{pr}_2 : \pi^*\mathcal{G} \rightarrow \mathcal{G}$ is the projection onto the second component.

Similar to Def. (21) we may shortly just write

$$H^!F = \mathcal{Ad}_{(\sigma^H)^{-1}} \circ F.$$

Proof.

That $H^!F$ is the field strength related to $H^!A$ follows quickly by the same calculation as for proving Eq. (29), using $\pi \circ H = \pi$,

$$\begin{aligned} H^!F &= d^{(\pi \circ H)^*\nabla^\mathcal{G}}(H^!A) + \frac{1}{2}[H^!A \wedge H^!A]_{\pi^*\mathcal{G}} + (\pi \circ H)^!\zeta \\ &= d^{\pi^*\nabla^\mathcal{G}}(H^!A) + \frac{1}{2}[H^!A \wedge H^!A]_{\pi^*\mathcal{G}} + \pi^!\zeta, \end{aligned}$$

which is the field strength of $H^!A$. Now recall Eq. (23), that is,

$$D_p H(X) = \mathcal{r}_{\bar{\sigma}_p^*}(X) + \overline{((\pi^*\Delta)\sigma^H)_p(X)} \Big|_{H(p)}$$

for all $X \in T_p\mathcal{P}$ ($p \in \mathcal{P}$), where $\bar{\sigma}_p := \text{pr}_2(\sigma_p^H)$. Thus, by Prop. Proposition 6.92,

$$\begin{aligned} (H^!F)_p(X, Y) &= F_{H(p)}(D_p H(X), D_p H(Y)) \\ &= F_{p, \bar{\sigma}_p}(\mathcal{r}_{\bar{\sigma}_p^*}(X), \mathcal{r}_{\bar{\sigma}_p^*}(Y)) \\ &= (\mathcal{r}_{\bar{\sigma}_p^*}^! F)_p(X, Y) \\ &= (\mathcal{Ad}_{\bar{\sigma}_p^{-1}} \circ F)_p(X, Y) \end{aligned}$$

for all $X, Y \in T_p\mathcal{P}$; this finishes the proof. ■

We will now extend this again to a local change of gauges (sections of $\mathcal{P}$).

Definition 6.97: Local field strength

Let $\mathcal{G} \rightarrow M$ be an LGB over a smooth manifold M and $\mathcal{P} \xrightarrow{\pi} M$ a principal $\mathcal{G}$ -bundle, also let $H\mathcal{G}$ be a multiplicative Yang-Mills connection on $\mathcal{G}$ and $A \in \Omega^1(\mathcal{P}; \pi^*\mathcal{G})$ be a connection 1-form on $\mathcal{P}$. Furthermore, let $s \in \Gamma(\mathcal{P}|_U)$ be a (local) gauge over an open subset $U \subset M$. Then we define the **local curvature** of **local field strength** $F_s \in \Omega^2(U; \mathcal{G}|_U)$ (**w.r.t.** s) by

$$F_s := s^!F.$$

In previous calculations for the total Maurer-Cartan form and the generalized Maurer-Cartan equation we have basically already shown the pullback of the structure equation.

Corollary 6.98: Pullback of the structure equation

Let $\mathcal{G} \rightarrow M$ be an LGB over a smooth manifold M and $\mathcal{P} \xrightarrow{\pi} M$ a principal $\mathcal{G}$ -bundle, also let $H\mathcal{G}$ be a multiplicative Yang-Mills connection on $\mathcal{G}$ and $A \in \Omega^1(\mathcal{P}; \pi^*\mathcal{G})$ be a connection

1-form on $\mathcal{P}$. Furthermore, let $s \in \Gamma(\mathcal{P}|_U)$ be a (local) gauge over an open subset $U \subset M$. Then we can express the local field strength as the pullback of the structure equation, that is,

$$F_s = d^{\nabla^{\mathcal{G}}} A_s + \frac{1}{2}[A_s \wedge A_s]_{\mathcal{G}} + \zeta.$$

Proof.

This follows quickly by the same calculation as for proving Eq. (29). $\blacksquare$

The gauge transformation of the field strength F describes the behaviour of F under a change of gauge.

Proposition 6.99: Gauge transformations again as a change of gauge

Let $\mathcal{G} \rightarrow M$ be an LGB over a smooth manifold M and $\mathcal{P} \xrightarrow{\pi} M$ a principal $\mathcal{G}$ -bundle, also let $H\mathcal{G}$ be a multiplicative Yang-Mills connection on $\mathcal{G}$ and $A \in \Omega^1(\mathcal{P}; \pi^*\mathcal{G})$ be a connection 1-form on $\mathcal{P}$. Also let U_i and U_j be two open subsets of M so that $U_i \cap U_j \neq \emptyset$, two gauges $s_i \in \Gamma(\mathcal{P}|_{U_i})$ and $s_j \in \Gamma(\mathcal{P}|_{U_j})$, and the unique $\sigma_{ji} \in \Gamma(\mathcal{G}|_{U_i \cap U_j})$ with $s_i = s_j \cdot \sigma_{ji}$ on $U_i \cap U_j$.

Then we have for the fields strength of A over $U_i \cap U_j$ that

$$F_{s_i} = \text{Ad}_{\sigma_{ji}^{-1}} \circ F_{s_j}.$$

Proof.

The proof is precisely as for Thm. Proposition 6.64, just without catering for the Darboux derivative in the gauge transformation due to that one uses Thm. Proposition 6.96 instead of Thm. Proposition 6.58. $\blacksquare$

Remark 6.100: Integrating curved Yang-Mills gauge theories, part II

Similar to Rem. Remark 6.65, Cor. Corollary 6.98 and Thm. Proposition 6.99 show that we recover the field strength and its gauge transformation as developed by Alexei Kotov and Thomas Strobl, see [1] for a concise summary or [6] for an extended introduction. These references are for the very general situation using Lie algebroids, hence see alternatively [5] for this type of gauge theory restricted to Lie algebra bundles.

7. CURVED YANG-MILLS GAUGE THEORY

7.1. Definition and gauge invariance. We eventually are able to conclude this paper with the definition of what we will call **curved Yang-Mills gauge theory**, highlighting that this theory is an integral of the infinitesimal gauge theory originally developed by Alexei Kotov and Thomas Strobl. For the following we say that a fibre metric κ of $\mathcal{G}$ is **Ad-invariant** if

$$\kappa(\text{Ad}_g(\nu), \text{Ad}_g(\mu)) = \kappa(\nu, \mu)$$

for all $g \in \mathcal{G}_x$ ($x \in M$) and $\mu, \nu \in \mathcal{G}_x$.

Corollary 7.1: Contraction of local field strength with Ad-invariant fibre metric is well-defined

Let $\mathcal{G} \rightarrow M$ be an LGB over a spacetime M so that its LAB $\mathcal{G}$ admits an Ad-invariant fibre metric κ , and let $\mathcal{P} \xrightarrow{\pi} M$ be a principal $\mathcal{G}$ -bundle; also let $H\mathcal{G}$ be a multiplicative Yang-Mills connection on $\mathcal{G}$ and $A \in \Omega^1(\mathcal{P}; \pi^*\mathcal{G})$ be a connection 1-form on $\mathcal{P}$. Furthermore, let $(U_i)_i$ be an open covering of M so that there are subordinate gauges $s_i \in \Gamma(\mathcal{P}|_{U_i})$.

Then the top-degree form $\mathfrak{L}_{\text{CYM}}[A] \in \Omega^{\dim(M)}(M; \mathbb{R})$, defined locally by

$$(\mathfrak{L}_{\text{CYM}}[A])|_{U_i} := -\frac{1}{2}\kappa(F_{s_i} \wedge *F_{s_i}),$$

is well-defined and independent of the choice of gauge, where $*$ is the Hodge star operator w.r.t. the spacetime metric of M .

Proof.

Let s_j be another gauge corresponding to U_j so that $U_i \cap U_j \neq \emptyset$, and we have a unique $\sigma_{ji} \in \Gamma(\mathcal{G}|_{U_i \cap U_j})$ with $s_i = s_j \cdot \sigma_{ji}$ on $U_i \cap U_j$. Then it follows by Thm. [Proposition 6.99](#) and the definition of the Hodge star operator (which is just a certain contraction w.r.t. M) that

$$\kappa(F_{s_i} \wedge *F_{s_i}) = \kappa\left(\left(\text{Ad}_{\sigma_{ji}^{-1}} \circ F_{s_j}\right) \wedge * \left(\text{Ad}_{\sigma_{ji}^{-1}} \circ F_{s_j}\right)\right) = \kappa(F_{s_j} \wedge *F_{s_j}),$$

using the Ad-invariance of κ . This concludes the proof. $\blacksquare$

Definition 7.2: Curved Yang-Mills gauge theory

Let $\mathcal{G} \rightarrow M$ be an LGB over a spacetime M so that its LAB $\mathcal{g}$ admits an Ad-invariant fibre metric κ , and let $\mathcal{P} \xrightarrow{\pi} M$ be a principal $\mathcal{G}$ -bundle; furthermore let $\text{H}\mathcal{G}$ be a multiplicative Yang-Mills connection on $\mathcal{G}$ and $A \in \Omega^1(\mathcal{P}; \pi^*\mathcal{g})$ be a connection 1-form on $\mathcal{P}$. Then the functional $A \mapsto \mathfrak{L}_{\text{CYM}}[A]$ is called the **curved Yang-Mills Lagrangian**, where $\mathfrak{L}_{\text{CYM}}[A]$ is defined in Cor. [Corollary 7.1](#).

By construction, this Lagrangian is gauge-invariant.

Theorem 7.3: Gauge invariance of the curved Yang-Mills Lagrangian

Let $\mathcal{G} \rightarrow M$ be an LGB over a spacetime M so that its LAB $\mathcal{g}$ admits an Ad-invariant fibre metric κ , and let $\mathcal{P} \xrightarrow{\pi} M$ be a principal $\mathcal{G}$ -bundle; also let $\text{H}\mathcal{G}$ be a multiplicative Yang-Mills connection on $\mathcal{G}$ and $A \in \Omega^1(\mathcal{P}; \pi^*\mathcal{g})$ be a connection 1-form on $\mathcal{P}$. Then we have

$$\mathfrak{L}_{\text{CYM}}[H^!A] = \mathfrak{L}_{\text{CYM}}[A]$$

for all $H \in \text{Aut}(\mathcal{P})$.

Proof.

By Thm. [Proposition 6.96](#) we have

$$H^!F = \text{Ad}_{\text{pr}_2 \circ (\sigma^H)^{-1}} \circ F,$$

where $\sigma^H \in C^\infty(\mathcal{P}; \mathcal{G})^{\mathcal{G}}$ is defined as in Prop. [Proposition 6.57](#) and $\text{pr}_2 : \pi^*\mathcal{G} \rightarrow \mathcal{G}$ is the projection onto the second component. Similar to the argument in the proof of Thm. [Proposition 6.64](#) we have

$$(H^!F)_s = \text{Ad}_{\text{pr}_2 \circ (\sigma^H)^{-1} \circ s} \circ F_s$$

for a given gauge $s \in \Gamma(\mathcal{P}|_U)$, where U is an open subset of M . With the same argument as in Cor. [Corollary 7.1](#) it follows that

$$\mathfrak{L}_{\text{CYM}}[H^!A] = \mathfrak{L}_{\text{CYM}}[A].$$

$\blacksquare$

Remark 7.4: Infinitesimal gauge invariance

As we have seen previously, the compatibility conditions (Def. Definition 6.85) were important to derive how F behaves under modified right push-forwards. Recall Rem. Remark 6.65, we have a corresponding formulation of infinitesimal gauge transformation and it is straight-forward to show that we achieve **infinitesimal gauge invariance** as in the classical theory. However, as argued by Alexei Kotov and Thomas Strobl in [1] (for a concise summary; [6] for an extended introduction, or alternatively [5] for a simplified language) the corresponding infinitesimal gauge invariance just needs the infinitesimal compatibility conditions (Def. Definition 6.67).

7.2. Field redefinitions. As developed in [5, 6, the latter reference developed the following for Lie algebroids and groupoids as structure]: There is an equivalence relation of curved Yang-Mills gauge theories, preserving the dynamics and kinematics of this theory. Let us recapitulate and integrate these:

Definition 7.5: Field redefinitions on the LGB

Let $\pi_{\mathcal{G}}: \mathcal{G} \rightarrow M$ be an LGB over a smooth manifold M with LAB $\mathcal{G}$; furthermore let $\mu_{\mathcal{G}}$ be a multiplicative Yang-Mills connection on $\mathcal{G}$ (w.r.t. a $\zeta \in \Omega^2(M; \mathcal{G})$). We then define **field redefinitions w.r.t. $\lambda \in \Omega^1(M; \mathcal{G})$** of $\mu_{\mathcal{G}}$ by

$$\mu_{\mathcal{G}}^{\lambda}|_g := \mu_{\mathcal{G}}|_g + \text{Ad}_{g^{-1}} \circ \pi_{\mathcal{G}}^! \lambda|_g - \pi_{\mathcal{G}}^! \lambda|_g$$

for all $g \in \mathcal{G}$ and of ζ by

$$\zeta^{\lambda} := \zeta + d^{\nabla^{\mathcal{G}}} \lambda + \frac{1}{2} [\lambda \wedge \lambda]_{\mathcal{G}}.$$

Remarks 7.6. Equivalently, since we introduced $\mu_{\mathcal{G}}$ as a composition of the vertical Maurer-Cartan form with the projection π^{vert} onto the vertical part w.r.t. the horizontal distribution on $\mathcal{G}$, we can define the field redefinition of π^{vert} by

$$\pi^{\text{vert}, \lambda} := \pi^{\text{vert}} + \overline{\pi_{\mathcal{G}}^! \lambda}^l + \overline{\pi_{\mathcal{G}}^! \lambda}^r,$$

where $\overline{\pi_{\mathcal{G}}^! \lambda}^r$ denotes the fundamental vector field of $\pi_{\mathcal{G}}^! \lambda$ via right-multiplication, that is,

$$\overline{\pi_{\mathcal{G}}^! \lambda}^r|_g = D_{e_x} R_g \circ \lambda_x \circ D_x \pi_{\mathcal{G}}$$

for all $g \in \mathcal{G}_x$, and $\overline{\pi_{\mathcal{G}}^! \lambda}^l$ is just the typical definition for fundamental vector fields, with an emphasis on that it is via left-multiplication.

Proposition 7.7: Field redefined multiplicative Yang-Mills connection a multiplicative Yang-Mills connection

Given the context of Def. Definition 7.5, the field redefinition of $\mu_{\mathcal{G}}$ is a multiplicative Yang-Mills connection w.r.t. ζ^{λ} .

Proof.

It is a straight-forward check to see that $\mu_{\mathcal{G}}^{\lambda}$ is still multiplicative, and well-known in the mentioned literature regarding multiplicative connections; a quick way to see this is by following Rem. Remark 6.87: We actually add an exact term,

$$\mu_{\mathcal{G}}^{\lambda} = \mu_{\mathcal{G}} + \delta \lambda.$$

Thus, $\mu_{\mathcal{G}}^{\lambda}$ stays closed, and hence it is multiplicative. That it satisfies the generalised Maurer-Cartan equation w.r.t. ζ^{λ} is a straight-forward but a bit lengthy calculation, which is why we keep this as an exercise for the reader; for the infinitesimal, i.e. in the connected case by Thm. Theorem 6.81, see [5, 6]. ■

In fact, a simple field redefinition of an Ehresmann connection preserves the compatibilities with the new multiplicative connection, and this is precisely one of the ideas of the field redefinitions. As usual, the difference of two connection 1-forms $A - A'$ should be a one-form which is horizontal and transforms via the Adjoint representations under the modified push-forward. However, we have seen here that the modified push-forward depends on a connection on the group bundle. Thus, is it possible to vary A with a 1-form which is horizontal *and* does not transform via the Adjoint transformation such that A' is a connection 1-form but w.r.t. a different connection on the group bundle? The answer to this are the field redefinitions; for the following observe that the field redefinition of $\mu_{\mathcal{G}}$ induces a field redefinition of $\nabla^{\mathcal{G}}$ given by

$$\nabla^{\text{YM},\lambda} = \nabla^{\mathcal{G}} + \text{ad} \circ \lambda .$$

Theorem 7.8: Field redefinition of Ehresmann connections

Let $\mathcal{G} \rightarrow M$ be an LGB over a smooth manifold M so that its LAB $\mathcal{g}$ admits an Ad-invariant fibre metric κ , and let $\mathcal{P} \xrightarrow{\pi} M$ be a principal $\mathcal{G}$ -bundle; furthermore let $\mu_{\mathcal{G}}$ be a multiplicative Yang-Mills connection on $\mathcal{G}$ (w.r.t. a $\zeta \in \Omega^2(M; \mathcal{g})$) and $A \in \Omega^1(\mathcal{P}; \pi^* \mathcal{g})$ be a connection 1-form on $\mathcal{P}$ w.r.t. $\mu_{\mathcal{G}}$. We define the **field redefinitions w.r.t. $\lambda \in \Omega^1(M; \mathcal{g})$ of A** by

$$A^\lambda := A - \pi^! \lambda .$$

Then A^λ is a connection 1-form w.r.t. μ^λ and

$$F^\lambda := d^{\pi^* \nabla^{\text{YM},\lambda}} A^\lambda + \frac{1}{2} [A^\lambda \wedge A^\lambda]_{\pi^* \mathcal{g}} + \pi^! \zeta^\lambda \equiv F$$

for all $\lambda \in \Omega^1(M; \mathcal{G})$.

Proof.

That $F^\lambda = F$ follows by similar calculations as in [5, 6]. Therefore let us only prove that A^λ is still a connection 1-form. A^λ is clearly still the identity on the vertical structure, and henceforth let us check its behaviour w.r.t. the modified pushforward induced by $\mu_{\mathcal{G}}^\lambda$, denoted by $\mathcal{r}_\sigma^\lambda$ for $\sigma \in \Gamma(\mathcal{G})$: We have to study

$$\left((\mathcal{r}_\sigma^\lambda)^! A^\lambda \right)_p (Y)$$

for all $p \in \mathcal{P}_x$ and $Y \in T_p \mathcal{P}$. First of all, similar to the beginning of the proof of Prop. [Proposition 6.92](#) we have

$$(\mathcal{r}_\sigma^\lambda)^! (\pi^! \lambda) = \pi^! \lambda ,$$

and thus we essentially only need to study the field redefined Darboux derivative:

$$\Delta^\lambda \sigma := \sigma^! \mu_{\mathcal{G}}^\lambda = \Delta \sigma + \text{Ad}_{\sigma^{-1}} \circ \lambda - \lambda .$$

In total,

$$\begin{aligned} \left((\mathcal{r}_\sigma^\lambda)^! A^\lambda \right)_p (Y) &= A_{p \cdot \sigma_x} (\mathcal{r}_{\sigma*} (Y)) - (\pi^! (\text{Ad}_{\sigma^{-1}} \circ \lambda))_p (Y) + \lambda_x (X) - \lambda_x (X) \\ &= (\text{Ad}_{\sigma^{-1}} \circ A - \text{Ad}_{\sigma^{-1}} \circ \pi^! \lambda)_p (Y) \\ &= (\text{Ad}_{\sigma^{-1}} \circ A^\lambda)_p (Y) , \end{aligned}$$

where $X := D_p \pi(Y)$, $\mathcal{r}_\sigma$ denotes the modified pushforward via $\mu_{\mathcal{G}}$ as usual, and we made use of A being a connection 1-form w.r.t. $\mu_{\mathcal{G}}$. ■

We see there is a transformation keeping the Lagrangian invariant such that the dynamics are preserved; $\mathcal{P}$ itself does not change and thus the gauge transformations in form of (local) principal bundle automorphisms also stay the same, and so the kinematics are preserved, too. However, field redefinitions non-trivially transform data like $\nabla^{\mathcal{G}}$ so that it may become flat, and we may end up at a pre-classical theory describing the same physics. ζ itself might survive as shown in [5, 6]. Let us summarize these results from the point of view as in Rem. [Remark 6.87](#).

Remark 7.9: Meaning of Field redefinitions

Recall Rem. [Remark 6.87](#) and its notation, *i.e.* there is a simplicial differential δ so that

$$\begin{aligned}\delta\mu_{\mathcal{G}} &= 0, \\ F_{\mathcal{G}} &= \delta\zeta,\end{aligned}$$

where $F_{\mathcal{G}}$ denotes the curvature of $\mu_{\mathcal{G}}$ in a classical approach, that is,

$$F_{\mathcal{G}} = d^{\pi_{\mathcal{G}}^* \nabla^{\mathcal{G}}} \mu_{\mathcal{G}} + \frac{1}{2} [\mu_{\mathcal{G}} \wedge \mu_{\mathcal{G}}]_{\pi_{\mathcal{G}}^* \mathcal{G}},$$

where $\pi_{\mathcal{G}}$ is the projection of $\mathcal{G}$. The field redefinition of $\mu_{\mathcal{G}}$ can be written as

$$\mu_{\mathcal{G}}^{\lambda} = \mu_{\mathcal{G}} + \delta\lambda.$$

Prop. [Proposition 7.7](#) and Thm. [Theorem 7.8](#) has now several interpretations and implications, where we denote with $[\mu_{\mathcal{G}}]_{\delta}$ the equivalence class of $\mu_{\mathcal{G}}$ inherited by δ :

- (1) The field redefinition of $\mu_{\mathcal{G}}$ is about picking any other representative for $[\mu_{\mathcal{G}}]_{\delta}$.
- (2) Every other representative of $[\mu_{\mathcal{G}}]_{\delta}$ is not just multiplicative, but also a multiplicative Yang-Mills connection, that is,

$$\begin{aligned}\delta\mu_{\mathcal{G}}^{\lambda} &= 0, \\ F_{\mathcal{G}}^{\lambda} &= \delta\zeta^{\lambda},\end{aligned}$$

where $F_{\mathcal{G}}^{\lambda}$ is defined like $F_{\mathcal{G}}$ but using $\mu_{\mathcal{G}}^{\lambda}$ instead of $\mu_{\mathcal{G}}$.

- (3) Regarding the choice of $\mu_{\mathcal{G}}$, the Lagrangian only depends on the choice of class $[\mu_{\mathcal{G}}]_{\delta}$. However, the explicit choice of ζ may still matter: One could vary ζ with centre-valued forms while keeping $\mu_{\mathcal{G}}$ fixed, this is in general not totally covered by the field redefinitions and may affect the Lagrangian as discussed in [\[6\]](#) (or [\[5\]](#)).

Last but not least, as also mentioned in the introduction, these field redefinitions are not just any transformation preserving the physics. Curved Yang-Mills gauge theory is a reformulation of classical gauge theory in such a way that gauge theory is form-invariant w.r.t. the field redefinitions. This reformulation introduces extra terms, ζ and connection 1-forms of ∇ , that is, every classical gauge theory is equivalent to a possibly curved theory with a non-zero ζ contributing to the field strength. From that point of view it is tempting to ask whether there are examples which cannot be flattened by field redefinitions:

7.3. Examples. First examples which cannot be flattened by field redefinitions were found in [\[5, 6\]](#); but many more were found after the upload of this work: It turned out that connections like $\mu_{\mathcal{G}}$ help classifying singular foliations ([\[8\]](#)), and in return singular foliation theory provides a full theory of examples and even a classification of such examples as long as the group bundle is centerless ([\[8, the appendix\]](#)); motivated by Ex. [Example 6.90](#) (recall also [Example 6.89](#)) such a classification is via viewing the LGB and its connection as associated to a “suitable” principal bundle.

Let us first point out the obvious:

Example 7.10: Classical examples

By Ex. [Example 6.88](#), we just recover the classical formalism of gauge theory if having a classical principal bundle P on which a trivial LGB $\mathcal{G} = M \times G$ acts on the right, where G is a Lie group with Lie algebra $\mathfrak{g}$. $H\mathcal{G}$ is the canonical flat multiplicative Yang-Mills connection, $\nabla^{\mathcal{G}} = \nabla^{\text{cYM}}$ is the canonical flat connection on $\mathcal{G} = M \times \mathfrak{g}$, and $\zeta \equiv 0$. Then every example known in the classical formalism carries over.

Furthermore, starting with such a classical example, by [\[5, §6, Cor. 6.2\]](#) we know that if $\mathfrak{g}$ has a non-zero centre (and if M is at least three-dimensional), then one can always add a

centre-valued ζ to this gauge theory, transforming it into a flat pre-classical gauge theory, $\nabla^{\mathcal{G}} = \nabla^{\text{pYM}}$, with non-trivial ζ . If $d^{\nabla^{\text{pYM}}} \zeta \neq 0$, then there is no $\lambda \in \Omega^1(M; \mathfrak{g})$ with $\bar{\zeta}^\lambda = 0$.

This shows that there are examples with non-trivial ζ , stable w.r.t. field redefinitions. However, as shown in [5, §5.2, Thm. 5.16] (or alternatively [6, §5.1.5, Thm. 5.1.33]), if M is contractible, then there is $\lambda \in \Omega^1(M; \mathfrak{g})$ so that $\nabla^{\text{YM}, \lambda}$ is flat. Thus, if M is contractible, then every curved Yang-Mills Lagrangian is equivalent to a Lagrangian of a pre-classical gauge theory (ζ may be non-vanishing). This may be due to that $\mathcal{G}$ is a trivial fibre bundle, and then the LGB action breaks down to Lie group action; recall all our discussions related to classical principal bundles equipped with an LGB action of a trivial LGB as in the previous example. Hence we conjecture:

Conjecture 7.11: Trivial LGBs are associated with pre-classical theories

Let $\mathcal{G} \rightarrow M$ be a trivial LGB over a smooth manifold M ; furthermore let $\mu_{\mathcal{G}}$ be a multiplicative Yang-Mills connection on $\mathcal{G}$. Then there is a $\lambda \in \Omega^1(M; \mathfrak{g})$ so that $\mu_{\mathcal{G}}^\lambda$ is flat.

For centerless simple Lie group bundles over connected M this turns out to be true, see [7]. Obviously, non-trivial LGBs are not used in the classical formalism, and by the previous discussion one may ask if there is a (global) example with a curved multiplicative Yang-Mills connection starting with a non-trivial LGB. Let us describe these examples, already presented in [6, 7].

There is a natural class of multiplicative Yang-Mills connections which cannot be flattened via field redefinitions. In order to understand this, we follow and use results of [8, in particular the appendix]: Assume an ordinary principal H -bundle P , where H is a Lie group, and assume that M is connected. Also assume that we have a manifold T on which H acts (from the left).¹² Then we have two associated bundles:

$$\begin{aligned} c_H(P) &:= (P \times H) / H, \\ \mathcal{T} &:= (P \times T) / H. \end{aligned} \tag{33}$$

As discussed, we have a canonical left-action, $c_H(P) \curvearrowright \mathcal{T}$, and an Ehresmann connection on P (in the usual sense) induces associated connections on $c_H(P)$ and $\mathcal{T}$. The one on $c_H(P)$ is a multiplicative Yang-Mills connection on $c_H(P)$ whose primitive is given by the corresponding curvature on P , and the one on $\mathcal{T}$ is a connection that has been called *compatible Yang-Mills connection* [8]. That is, this is an Ehresmann connection on $\mathcal{T}$ w.r.t. the $c_H(P)$ -action and the adjustment on $c_H(P)$, which satisfies a certain curvature equation.

The following remark provides further motivation for considering Lie group bundles like $c_H(P)$ beyond the existence of natural multiplicative Yang-Mills connections.

Remarks 7.12. As argued in the appendix of [8] every Lie group bundle $\mathcal{H}$ with a covariant adjustment and whose structural Lie group H has a trivial center is associated to some principal bundle P (whose structural Lie group K is the extension of a Galois group by H); see also [22] for similar arguments. That is, $\mathcal{H} \cong (P \times H) / K$, and the covariant adjustment is given as an associated connection.

In particular [8] provides the following (a modification of [23]):

Lemma 7.13: Simple Lie group bundles, [10]

If H is simple and centerless, then all covariant adjustments on $\mathcal{H}$ are associated to an Ehresmann connection of a unique principal H -bundle P , and their primitive is the corresponding curvature on P .

¹² T and its related constructions and assumptions are only needed to provide an extra geometric interpretation as in Table 5.

Simplicity also implies that the correspondence in Lemma 7.13 is one-to-one. Slightly generalized, assuming that H has a trivial center implies that the associated connection on $c_H(P)$ is in one-to-one correspondence with the Ehresmann connection on P . Thus, let us now assume that H and hence $c_H(P)$ have a trivial center; furthermore assume that $T = \mathbb{R}^d$, and that H acts faithfully on T with 0 as fixed point. Then there is a canonical singular foliation $\mathcal{F}$ on $\mathcal{T}$ with M as a leaf generated by vector fields of the form

$$\chi_{\mathcal{T}}(X) + \bar{\nu} \quad (34)$$

for all $X \in \mathfrak{X}(M)$ and $\nu \in \Gamma(\text{ad}(P))$, where $\chi_{\mathcal{T}}$ is the projectable horizontal lift of the associated connection on $\mathcal{T}$, and $\nu \mapsto \bar{\nu}$ denotes the inherited Lie algebra bundle action. The connection on $\mathcal{T}$ is then also what one calls an $\mathcal{F}$ -connection.

Because the action is faithful, there is the following one-to-one correspondence between:

- the multiplicative Yang-Mills connection on $c_H(P)$;
- the Ehresmann connection on P ;
- the $\mathcal{F}$ -connection on $\mathcal{T}$.

The major result of [8] implies that (in a formal setting, and in a neighborhood around a leaf) every foliation can be modeled like this; however P is then in general infinite-dimensional.¹³ Altogether, we can study multiplicative Yang-Mills connections on such $c_H(P)$ as structure group bundle by looking at $\mathcal{F}$ and P . In particular, $c_H(P)$ admits a flat multiplicative Yang-Mills connection if and only if $\mathcal{F}$ admits a flat $\mathcal{F}$ -connection, which is the case if and only if P is flat. Furthermore, [8] provides a clear geometric interpretation of the field redefinitions, extending the discussion of [5] and [6, §5.1].

TABLE 5. Comparing the geometry of multiplicative Yang-Mills connections (YM conn.) with the geometry of singular foliations. Here, $\mathcal{F}$ -connections only serve to highlight a geometric interpretation, but we actually provided a notion only defined as associated to P ; however, the general notion of $\mathcal{F}$ -connections allows for more than such connections if H is not simple; this will not matter here, see [8] for the general classification of $\mathcal{F}$ -connections and their relation to multiplicative Yang-Mills connections.

Lie group bundle $c_H(P)$	Singular foliations $\mathcal{F}$	principal H -bundle P
multiplicative YM conn. μ	$\mathcal{F}$ -connection	Ehresmann connection
$c_H(P)$ admits flat μ	$\mathcal{F}$ is flat	P is flat
field redefinition of μ	change of $\mathcal{F}$ -connection	change of Ehresmann connection

Thus, in order to find non-trivial multiplicative Yang-Mills connections which cannot be flattened by field redefinitions one only needs to apply the previous construction on principal bundles P which do not admit flat connections; alternatively search for foliations $\mathcal{F}$ which do not admit flat $\mathcal{F}$ -connections. There is plenty of literature on both cases.

Example 7.14: Cooking recipe

Using non-flat principal bundles, for example a non-trivial principal bundle P over a simply connected base manifold, such as the Hopf fibration $S^7 \rightarrow S^4$, leads to curved multiplicative Yang-Mills connections on $c_H(P)$ in the above construction which cannot be flattened by field redefinitions.

What about the full theory? For this we need to require the existence of an Ad-invariant connection on $c_H(P)$ which is simply given by the assumption that H is simple what is covered by the previous discussion; recall Rem. Theorem 7.12. Thus, it is only left to find a principal $c_H(P)$ -bundle $\mathcal{P}$; for this we have several canonical choices:

¹³In our context we simply assume a finite-dimensional P so that the arguments of [8] carry over to a smooth setting.

- $\mathcal{P} = c_H(P)$ itself;
- $\mathcal{P} = P$ via the action of gauge transformations; recall Rem. ??;
- $\mathcal{P} = (P \times P)/H$, the Atiyah groupoid of P , where H acts diagonally on $P \times P$.

About the last bullet point, for the experienced reader: We have a short exact sequence of Lie groupoids

$$c_H(P) \hookrightarrow (P \times P)/H \twoheadrightarrow \mathcal{P}\text{air}(M),$$

where $\mathcal{P}\text{air}(M)$ is the pair groupoid of M . It is a straightforward task to show that the Atiyah groupoid is a principal $c_H(P)$ -bundle,

$$\begin{array}{ccc} & \curvearrowright & \\ \mathcal{P} = (P \times P)/H & & c_H(P) \\ \downarrow \pi_{\mathcal{P}} & \searrow s_{\mathcal{P}} & \downarrow \\ \mathcal{P}\text{air}(M) & \xrightarrow{s_{\mathcal{P}\text{air}(M)}} & M \end{array}$$

where $s_{\mathcal{P}}$ and $s_{\mathcal{P}\text{air}(M)}$ are the source arrows of $(P \times P)/H$ and $\mathcal{P}\text{air}(M)$, respectively. Here one needs the notion of LGB-actions along a moment map, here $s_{\mathcal{P}}$, which we did not discuss here, but it is straight-forward to extend the previous constructions to such setups; see [7]. Observe however, that the base manifold of the Atiyah groupoid as such a principal bundle is the pair groupoid.

Using such principal bundles thence lead to a curved Yang-Mills gauge theory which cannot be described in a classical way.

7.4. Future prospects. As a next step one could now define minimal couplings. Given a vector bundle $V \rightarrow M$, equipped with a LGB representation $\rho: \mathcal{G} \rightarrow \text{Aut}(V)$, we should be able to define associated vector bundles $\mathcal{V} := (\mathcal{P} \times_M V)/\mathcal{G}$, as we already mentioned before. W.r.t. a local gauge s of $\mathcal{P}$ every section $\Psi \in \Gamma(\mathcal{V})$ has locally the form

$$[s, \psi_s],$$

where ψ is a local section of V and $[\cdot, \cdot]$ denotes the equivalence class related to the construction in $\mathcal{V}$. Then one can define the **minimal coupling** $\nabla^A \Psi$ locally by

$$[s, \nabla \psi_s + \rho_*(A_s)(\psi_s)]$$

where ρ_* is the LAB representation inherited by ρ , and ∇ is a vector bundle connection with

$$\begin{aligned} \rho(\sigma^{-1}) \circ \nabla \circ \rho(\sigma) &= \nabla + \rho_*(\Delta\sigma), \\ R_{\nabla} &= \rho_* \circ \zeta \end{aligned}$$

for all $\sigma \in \Gamma(\mathcal{G})$. Observe the similarity with the infinitesimal compatibility conditions Def. Definition 6.67 and Lemma Lemma 6.78; the first condition should lead to that that $\nabla^A \Psi$ is well-defined and independent of the choice of s . The second condition should imply that the curvature of ∇^A is given by the generalised field strength of A . After showing this one could try to construct all the other obvious analogues of the classical formalism and discuss their applications. Essentially, one achieves a special type of curved Yang-Mills-Higgs theory as introduced in [7].

Acknowledgements: I want to thank Siye Wu, Camille Laurent-Gengoux, Mark John David Hamilton and Alessandra Frabetti for their great help and support in making this paper. Also big thanks to my mother, father, Dennis, Gregor, Marco, Nico, Jakob, Kathi, Konstantin, Lukas, Locki, Gareth, Philipp, Ramona, Annerose, Michael, Maxim, Anna, and my girlfriend Rachel for all their love.

Funding: The paper is part of my post-doc fellowship at the National Center for Theoretical Sciences (NCTS, 國家理論科學研究中心), which is why I also want to thank the NCTS.

LIST OF REFERENCES

- [1] Alexei Kotov and Thomas Strobl. Curving Yang-Mills-Higgs gauge theories. *Physical Review D*, 92(8):085032, 2015. <https://doi.org/10.1103/PhysRevD.92.085032>.
- [2] Ana Cannas Da Silva and Alan Weinstein. *Geometric models for noncommutative algebras*, volume 10. American Mathematical Soc., 1999.
- [3] Thomas Strobl. Algebroid Yang-Mills theories. *Physical review letters*, 93(21):211601, 2004. <https://doi.org/10.1103/PhysRevLett.93.211601>.
- [4] Christoph Mayer and Thomas Strobl. Lie algebroid Yang-Mills with matter fields. *Journal of Geometry and Physics*, 59(12):1613–1623, 2009. <https://doi.org/10.1016/j.geomphys.2009.07.018>.
- [5] Simon-Raphael Fischer. Curved Yang-Mills-Higgs gauge theories in the case of massless gauge bosons. *Journal of Geometry and Physics*, 162:104104, 2021. <https://doi.org/10.1016/j.geomphys.2021.104104>.
- [6] Simon-Raphael Fischer. *Geometry of curved Yang-Mills-Higgs gauge theories*. PhD thesis, 05/31 2021. ID: unige:152555, <https://doi.org/10.13097/archive-ouverte/unige:152555>.
- [7] Simon-Raphael Fischer, Mehran Jalali Farahani, Hyungrok Kim, and Christian Saemann. Adjusted connections I: Differential cocycles for principal groupoid bundles with connection. 2024. arXiv:2406.16755, <https://doi.org/10.48550/arXiv.2406.16755>.
- [8] Simon-Raphael Fischer and Camille Laurent-Gengoux. A classification of neighborhoods around leaves of a singular foliation. 2024. arXiv:2401.05966, <https://doi.org/10.48550/arXiv.2401.05966>.
- [9] Marco Castrillón López and Álvaro Rodríguez Abella. Principal bundles and connections modelled by Lie group bundles. *Geometriae Dedicata*, 217(2):32, 2023. <https://doi.org/10.1007/s10711-023-00766-2>.
- [10] Kirill C. H. Mackenzie. General theory of Lie groupoids and Lie algebroids. *London Mathematical Society Lecture Note Series*, 213:xxxviii+501, 2005. <https://doi.org/10.1017/CBO9781107325883>.
- [11] Ieke Moerdijk and Janez Mrcun. *Introduction to foliations and Lie groupoids*, volume 91. Cambridge University Press, 2003. <https://doi.org/10.1017/CBO9780511615450>.
- [12] Camille Laurent-Gengoux, Mathieu Stiénon, and Ping Xu. Non-abelian differentiable gerbes. *Advances in Mathematics*, 220(5):1357–1427, 2009. <https://doi.org/10.1016/j.aim.2008.10.018>.
- [13] Rui Loja Fernandes and Ioan Mărcuț. Multiplicative Ehresmann connections. *Advances in Mathematics*, 427:109124, 2023. <https://doi.org/10.1016/j.aim.2023.109124>.
- [14] Marius Crainic, Maria Amelia Salazar, and Ivan Struchiner. Multiplicative forms and Spencer operators. *Mathematische Zeitschrift*, 279(3):939–979, 2015. <https://doi.org/10.1007/s00209-014-1398-z>.
- [15] Marius Crainic. Differentiable and algebroid cohomology, van Est isomorphisms, and characteristic classes. *Commentarii Mathematici Helvetici*, 78(4):681–721, 2003. <https://doi.org/10.1007/s00014-001-0766-9>.
- [16] Simon-Raphael Fischer, Mehran Jalali Farahani, Hyungrok Kim, and Christian Saemann. Topological classification of symmetry breaking and vacuum degeneracy. 2024. arXiv:2406.17634, <https://doi.org/10.48550/arXiv.2406.17634>.
- [17] Mark JD Hamilton. *Mathematical Gauge Theory*. Springer, 2017. <https://doi.org/10.1007/978-3-319-68439-0>.
- [18] K Ajaykumar, B. S. Kiranagi, and R Rangarajan. Pullback of Lie algebra and Lie group bundles, and their homotopy invariance. *Journal of Algebra and Related Topics*, 8(1):15–26, 2020. <https://doi.org/10.22124/jart.2020.13988.1156>.
- [19] Martin Bojowald, Alexei Kotov, and Thomas Strobl. Lie algebroid morphisms, poisson sigma models, and off-shell closed gauge symmetries. *Journal of Geometry and Physics*, 54(4):400–426, 2005. <https://doi.org/10.1016/j.geomphys.2004.11.002>.
- [20] Christoph Mayer and Thomas Strobl. Lie algebroid Yang-Mills with matter fields. *Journal of Geometry and Physics*, 59(12):1613–1623, 2009. <https://doi.org/10.1016/j.geomphys.2009.07.018>.
- [21] Henrique Bursztyn, Marius Crainic, Alan Weinstein, and Chenchang Zhu. Integration of twisted Dirac brackets. *Duke Mathematical Journal*, 123(3):549–607, 2004. <https://doi.org/10.1215/S0012-7094-04-12335-8>.
- [22] Eckhard Meinrenken. On the integration of transitive Lie algebroids. *L'Enseignement Mathématique*, 67(3):423–454, 2021.
- [23] DB Blázquez-Sanz, Carlos A Marín Arango, and Sedney Suárez Gordon. Group bundles and group connections. *Lobachevskii Journal of Mathematics*, 43(1):49–70, 2022.
- [24] Janusz Grabowski and Mikołaj Rotkiewicz. Higher vector bundles and multi-graded symplectic manifolds. *Journal of Geometry and Physics*, 59(9):1285–1305, 2009. <https://doi.org/10.1016/j.geomphys.2009.06.009>.
- [25] Henrique Bursztyn, Hudson Lima, and Eckhard Meinrenken. Splitting theorems for Poisson and related structures. *Journal für die reine und angewandte Mathematik (Crelles Journal)*, 2017. <https://doi.org/10.1515/crelle-2017-0014>.

APPENDIX A. DOUBLE TANGENT BUNDLE AND ITS CANONICAL FLIP MAP

We will follow [10, §9.6, page 363]. For a smooth manifold M we denote the projection of its tangent bundle by $\pi_{TM} : TM \rightarrow M$; similarly we have the projection of the **double tangent bundle** $\pi_{TTM} : TTM \rightarrow TM$, the tangent bundle of TM . However, there is also $D\pi_{TM} : TTM \rightarrow TM$,

and in fact there is another vector bundle structure on $\text{T}TM$ rendering $D\pi_{TM}$ a projection; see *e.g.* [10, §3.4 *et seq.*; page 110ff.]. Let us give a very rough sketch:

Let $\xi, \eta \in \text{T}TM$ with

$$D_{X_0}\pi_{TM}(\xi) = D_{Y_0}\pi_{TM}(\eta) =: \omega,$$

where $X_0 := \pi_{\text{T}TM}(\xi)$ and $Y_0 := \pi_{\text{T}TM}(\eta)$, and due to the fact that $D\pi_{TM}$ is a vector bundle morphism over π_{TM} we get

$$p = \pi_{TM}(X_0) = \pi_{TM}(Y_0),$$

where $\pi_{TM}(\omega) =: p$. Thus, one can take curves $X, Y : I \rightarrow TM$ ($I \subset \mathbb{R}$ an open interval around 0) with

$$\begin{aligned} X(0) &= X_0, & \left. \frac{d}{dt} \right|_{t=0} X &= \xi, \\ Y(0) &= Y_0, & \left. \frac{d}{dt} \right|_{t=0} Y &= \eta, \end{aligned}$$

such that

$$\pi_{TM} \circ X = \pi_{TM} \circ Y,$$

because the condition on ξ and η imply on the base paths $\pi_{TM} \circ X, \pi_{TM} \circ Y : I \rightarrow M$ that

$$\begin{aligned} (\pi_{TM} \circ X)(0) &= p = (\pi_{TM} \circ Y)(0), \\ \left. \frac{d}{dt} \right|_{t=0} (\pi_{TM} \circ X) &= D_{X_0}\pi_{TM}(\xi) = \omega = D_{Y_0}\pi_{TM}(\eta) = \left. \frac{d}{dt} \right|_{t=0} (\pi_{TM} \circ Y). \end{aligned}$$

Then the addition and scalar multiplication with $\lambda \in \mathbb{R}$ for $\text{T}TM \xrightarrow{D\pi_{TM}} TM$ is defined by

$$\begin{aligned} \xi \oplus \eta &:= \left. \frac{d}{dt} \right|_{t=0} (X + Y), \\ \lambda \cdot \xi &:= \left. \frac{d}{dt} \right|_{t=0} (\lambda X), \end{aligned}$$

where the addition of curves is well-defined because of $\pi_{TM} \circ X = \pi_{TM} \circ Y$ which implies $\pi_{TM} \circ (X + Y) = \pi_{TM} \circ X = \pi_{TM} \circ Y$; so, one can take the sum of the curves and

$$D\pi_{TM}(\xi \oplus \eta) = \left. \frac{d}{dt} \right|_{t=0} \underbrace{(\pi_{TM} \circ (X + Y))}_{=\pi_{TM} \circ X} = D_{X_0}\pi_{TM}(\xi) = \omega.$$

The operations of the linear structure in $\text{T}TM \xrightarrow{\pi_{\text{T}TM}} TM$ is still denoted in the same manner as usual, and by definition one also gets

$$\begin{aligned} \pi_{\text{T}TM}(\xi \oplus \eta) &= \pi_{\text{T}TM}(\xi) + \pi_{\text{T}TM}(\eta), \\ \pi_{\text{T}TM}(\lambda \cdot \xi) &= \lambda \pi_{\text{T}TM}(\xi). \end{aligned}$$

In total, we have a double vector bundle given by the following commuting diagram

$$\begin{array}{ccc} \text{T}TM & \xrightarrow{D\pi_{TM}} & TM \\ \downarrow \pi_{\text{T}TM} & & \downarrow \pi_{TM} \\ TM & \xrightarrow{\pi_{TM}} & M \end{array} \quad (\text{A.1})$$

i.e. each horizontal and vertical line is a vector bundle so that the horizontal and vertical scalar multiplications on $\text{T}TM$ commute; see *e.g.* [24, §3ff.] or [10, §9.1, page 340ff.] for a definition on double vector bundles in general.

Now observe that the flow X of ξ has values in TM , that is, for all $t \in I$ we have a curve $\alpha_t : J \rightarrow M$ (J another open interval containing 0), $s \mapsto \alpha_t(s)$, so that

$$\alpha_t(0) = \pi_{TM}(X(t)),$$

$$\left. \frac{d}{ds} \right|_{s=0} \alpha_t = X(t),$$

and the first equation implies

$$\left. \frac{d}{dt} \right|_{t=0} \alpha_t(0) = D_{X_0} \pi_{TM}(\xi) = \omega.$$

So, in total we have for all $\xi \in TTM$ a smooth map $\alpha : I \times J \rightarrow M$, $(t, s) \mapsto \alpha(t, s) = \alpha_t(s)$, such that

$$\alpha(0, 0) = p = (\pi_{TM} \circ \pi_{TTM})(\xi) = (\pi_{TM} \circ D\pi_{TM})(\xi),$$

$$\left. \frac{d}{dt} \right|_{t=0} \alpha(t, 0) = \omega = D_{X_0} \pi_{TM}(\xi), \quad \left. \frac{d}{ds} \right|_{s=0} \alpha(0, s) = X_0 = \pi_{TTM}(\xi),$$

$$\left. \frac{d}{dt} \right|_{t=0} \left. \frac{d}{ds} \right|_{s=0} \alpha = \xi.$$

As for tangent vectors, the class $[\alpha]$ of all such α uniquely describes ξ , in fact giving rise to an equivalence relation so that $[\alpha]$ is an equivalence class.

In the context of Schwarz's Theorem one may find it natural to define the **canonical involution (or flip)** on TTM as a map $S : TTM \rightarrow TTM$ by

$$S(\xi) := \left. \frac{d}{ds} \right|_{s=0} \left. \frac{d}{dt} \right|_{t=0} \alpha. \quad (\text{A.2})$$

By [10, §9.6, Thm. 9.6.1, page 363; but without proof] one has that S is an isomorphism of double vector bundles with certain "special" properties; we will not need the general definition of these. What we need of this, is that S is a base-preserving vector bundle isomorphism as both maps, from $TTM \xrightarrow{D\pi_{TM}} TM$ to $TTM \xrightarrow{\pi_{TTM}} TM$ and vice versa. Let us prove this, by starting with showing the well-definedness of S :

W.l.o.g. we can put $J = I$ for simplicity. By construction, the equivalence class $[\beta]$ of $S(\xi)$ is represented by $\beta : I^2 \rightarrow M$ given by $\beta(t, s) := \alpha(s, t)$ with

$$\begin{aligned} \beta(0, 0) &= p, \\ D\pi_{TM}(S(\xi)) &= \left. \frac{d}{dt} \right|_{t=0} \beta(t, 0) = \left. \frac{d}{dt} \right|_{t=0} \alpha(0, t) = \pi_{TTM}(\xi) = X_0, \\ \pi_{TTM}(S(\xi)) &= \left. \frac{d}{ds} \right|_{s=0} \beta(0, s) = \left. \frac{d}{ds} \right|_{s=0} \alpha(s, 0) = D\pi_{TM}(\xi) = \omega, \\ S(\xi) &= \left. \frac{d}{dt} \right|_{t=0} \left. \frac{d}{ds} \right|_{s=0} \beta. \end{aligned}$$

With this, we can express $S(\xi)$ in coordinates: Fix local coordinates $(x^i)_i$ on M , then $(\pi_{TM}^* x^i)_i$ and $(dx^i)_i$ are local coordinates on TM . Then by chain rule

$$\xi(\pi_{TM}^* x^i) = \xi(x^i \circ \pi_{TM}) = d_p x^i (D\pi_{TM}(\xi)) = \omega^i,$$

and

$$\xi(dx^i) = \left(\left. \frac{d}{dt} \right|_{t=0} \left. \frac{d}{ds} \right|_{s=0} \alpha \right) (dx^i) = \left. \frac{d}{dt} \right|_{t=0} \underbrace{\left(dx^i \circ \left. \frac{d}{ds} \right|_{s=0} \alpha \right)}_{= \left(\left. \frac{d}{ds} \right|_{s=0} \alpha \right)(x^i)} = \left. \frac{d}{dt} \right|_{t=0} \left. \frac{d}{ds} \right|_{s=0} \alpha^i,$$

thus we have in total

$$\xi = \omega^i \frac{\partial}{\partial(\pi_{TM}^* x^i)} \Big|_{X_0} + \left(\left. \frac{d}{dt} \right|_{t=0} \left. \frac{d}{ds} \right|_{s=0} \alpha^i \right) \frac{\partial}{\partial(dx^i)} \Big|_{X_0}. \quad (\text{A.3})$$

Similarly we can proceed with $S(\xi)$ and then apply the "classical" Schwarz' Theorem on the derivatives of α^i (recall the properties of β), so that we get

$$\begin{aligned} S(\xi) &= X_0^i \frac{\partial}{\partial(\pi_{TM}^* x^i)} \Big|_{\omega} + \left(\frac{d}{dt} \Big|_{t=0} \frac{d}{ds} \Big|_{s=0} \beta^i \right) \frac{\partial}{\partial(dx^i)} \Big|_{\omega} \\ &= X_0^i \frac{\partial}{\partial(\pi_{TM}^* x^i)} \Big|_{\omega} + \left(\frac{d}{ds} \Big|_{s=0} \frac{d}{dt} \Big|_{t=0} \alpha^i \right) \frac{\partial}{\partial(dx^i)} \Big|_{\omega} \\ &= X_0^i \frac{\partial}{\partial(\pi_{TM}^* x^i)} \Big|_{\omega} + \left(\frac{d}{dt} \Big|_{t=0} \frac{d}{ds} \Big|_{s=0} \alpha^i \right) \frac{\partial}{\partial(dx^i)} \Big|_{\omega}. \end{aligned} \quad (\text{A.4})$$

It is now clear that $S(\xi)$ is independent of the choice of α and just depends on $[\alpha]$. Furthermore, due to what we have shown for β , it is also clear that S is a base-preserving map either from $\text{TTM} \xrightarrow{D\pi_{TM}} TM$ to $\text{TTM} \xrightarrow{\pi_{TM}} TM$ or vice versa. Bijectivity follows by $S \circ S = \text{id}_{\text{TTM}}$.

It is only left to show that we have linearity for S as a map from $\text{TTM} \xrightarrow{D\pi_{TM}} TM$ to $\text{TTM} \xrightarrow{\pi_{TM}} TM$ and vice versa. For this we use the derived coordinate expressions. Let $\eta \in \text{TTM}$ be defined as before with associated area function γ such that $\eta = [\gamma]$. Then one can derive by using what we have shown in the discussion about $\text{TTM} \xrightarrow{D\pi_{TM}} TM$ as vector bundle that

$$\lambda \cdot \xi \oplus \kappa \cdot \eta = \omega^i \frac{\partial}{\partial(\pi_{TM}^* x^i)} \Big|_{\lambda X_0 + \kappa Y_0} + \left(\frac{d}{dt} \Big|_{t=0} \frac{d}{ds} \Big|_{s=0} (\lambda \alpha^i + \kappa \gamma^i) \right) \frac{\partial}{\partial(dx^i)} \Big|_{\lambda X_0 + \kappa Y_0} \quad (\text{A.5})$$

for all $\lambda, \kappa \in \mathbb{R}$, and so

$$\begin{aligned} S(\lambda \cdot \xi \oplus \kappa \cdot \eta) &= (\lambda X_0^i + \kappa Y_0^i) \frac{\partial}{\partial(\pi_{TM}^* x^i)} \Big|_{\omega} + \left(\frac{d}{dt} \Big|_{t=0} \frac{d}{ds} \Big|_{s=0} (\lambda \alpha^i + \kappa \gamma^i) \right) \frac{\partial}{\partial(dx^i)} \Big|_{\omega} \\ &= \lambda S(\xi) + \kappa S(\eta). \end{aligned}$$

In the same fashion, let $\zeta = [\delta] \in T_{X_0} TM$ with $D_{X_0} \pi_{TM}(\zeta) =: \varphi$, then $\lambda \xi + \kappa \zeta$ is just the typical sum of tangent vectors and we get

$$\begin{aligned} S(\lambda \xi + \kappa \zeta) &= X_0^i \frac{\partial}{\partial(\pi_{TM}^* x^i)} \Big|_{\lambda \omega + \kappa \varphi} + \left(\frac{d}{dt} \Big|_{t=0} \frac{d}{ds} \Big|_{s=0} (\lambda \alpha^i + \kappa \delta^i) \right) \frac{\partial}{\partial(dx^i)} \Big|_{\lambda \omega + \kappa \varphi} \\ &= \lambda \cdot \left(X_0^i \frac{\partial}{\partial(\pi_{TM}^* x^i)} \Big|_{\omega} + \left(\frac{d}{dt} \Big|_{t=0} \frac{d}{ds} \Big|_{s=0} \alpha^i \right) \frac{\partial}{\partial(dx^i)} \Big|_{\omega} \right) \\ &\quad \oplus \kappa \cdot \left(X_0^i \frac{\partial}{\partial(\pi_{TM}^* x^i)} \Big|_{\varphi} + \left(\frac{d}{dt} \Big|_{t=0} \frac{d}{ds} \Big|_{s=0} \delta^i \right) \frac{\partial}{\partial(dx^i)} \Big|_{\varphi} \right) \\ &= \lambda \cdot S(\xi) \oplus \kappa \cdot S(\zeta). \end{aligned}$$

This finishes the proof, so we have:

Remark A.1: The canonical involution/flip map an isomorphism, [10, §9.6, Thm. 9.6.1, page 363; but without proof]

The canonical involution/flip S is a base-preserving vector bundle isomorphism from $\text{TTM} \xrightarrow{D\pi_{TM}} TM$ to $\text{TTM} \xrightarrow{\pi_{TM}} TM$, and similarly vice versa. We also write $S_M := S$ to give an accentuation on M .

This isomorphism is basically now Schwarz's Theorem:

Remark A.2: Revisit: Schwarz's Theorem

By definition we have

$$\frac{d}{ds} \Big|_{s=0} \frac{d}{dt} \Big|_{t=0} \alpha = S_M \left(\frac{d}{dt} \Big|_{t=0} \frac{d}{ds} \Big|_{s=0} \alpha \right)$$

for all smooth $\alpha : I \times J \rightarrow M$, where I and J are open intervals containing 0. Similarly this extends to smooth maps $F : M \rightarrow N$, where N is another smooth manifold. So let $\xi = [\alpha] \in \text{TTM}$, and then we calculate for $\text{DDF} : \text{TTM} \rightarrow \text{TTN}$ that

$$\begin{aligned} \text{DDF}(\xi) &= \left. \frac{d}{dt} \right|_{t=0} \left. \frac{d}{ds} \right|_{s=0} (F \circ \alpha) \\ &= S_N \left(\left. \frac{d}{ds} \right|_{s=0} \left. \frac{d}{dt} \right|_{t=0} (F \circ \alpha) \right) \\ &= S_N(\text{DDF}(S_M(\xi))) \\ &= (S_N \circ \text{DDF} \circ S_M)(\xi). \end{aligned}$$

Since the canonical involutions are clearly self-inverse, we can also write

$$S_N \circ \text{DDF} = \text{DDF} \circ S_M.$$

For a notation with base points in M , consider a special case with $M = M_1 \times M_2$, where M_i ($i \in \{1, 2\}$) are smooth manifolds. Thus, $\text{TM} \cong \pi_1^* \text{TM}_1 \oplus \pi_2^* \text{TM}_2$, where $\pi_i : M_1 \times M_2 \rightarrow M_i$ are the projections onto the i -th component. Then let $p_i \in M_i$, $Y_i \in \text{T}_{p_i} M_i$ and $\xi := (Y_1, Y_2)$. Denoting with $\gamma_1 : I \rightarrow M_1$ and $\gamma_2 : J \rightarrow M_2$ the curves with velocities Y_1 and Y_2 at 0, respectively, and so $\alpha(t, s) = (\gamma_1(s), \gamma_2(t))$; then we define $\text{D}_{p_1} F(Y_1)$ as a map

$$\begin{aligned} M_2 &\rightarrow \text{TN}, \\ p_2 &\mapsto \text{D}_{p_1} F(Y_1)|_{p_2} := \text{D}_{(p_1, p_2)} F(Y_1, 0_{p_2}), \end{aligned}$$

where 0_{p_2} is the zero tangent vector at $\text{T}_{p_2} M_2$, in a similar fashion for $\text{D}_{p_2} F(Y_2)$. Then

$$\begin{aligned} \text{D}_{p_2} (\text{D}_{p_1} F(Y_1))(Y_2) &= \left. \frac{d}{dt} \right|_{t=0} \text{D}_{(p_1, \gamma_2(t))} F(Y_1, 0_{\gamma_2(t)}) \\ &= \left. \frac{d}{dt} \right|_{t=0} \left. \frac{d}{ds} \right|_{s=0} (F \circ \alpha(t, s)) \\ &= \text{DDF}(\xi), \end{aligned}$$

similarly

$$\begin{aligned} \text{D}_{p_1} (\text{D}_{p_2} F(Y_2))(Y_1) &= \left. \frac{d}{ds} \right|_{s=0} \text{D}_{(\gamma_1(s), p_2)} F(0_{\gamma_1(s)}, Y_2) \\ &= \left. \frac{d}{ds} \right|_{s=0} \left. \frac{d}{dt} \right|_{t=0} (F \circ \alpha(t, s)) \\ &= S_N(\text{DDF}(\xi)), \end{aligned}$$

in total

$$\text{D}_{p_1} (\text{D}_{p_2} F(Y_2))(Y_1) = S_N(\text{D}_{p_2} (\text{D}_{p_1} F(Y_1))(Y_2)).$$

In fact, using the canonical flip/involution, we can construct and state other properties which can be useful for calculations related to second derivatives.

Remark A.3: Total derivatives of tangent bundle morphisms linear with respect to prolonged vertical structure

Consider a vector bundle morphism $L : \text{TM} \rightarrow \text{TN}$ (over some map $M \rightarrow N$), where N is another smooth manifold. Then for $\xi, \eta \in \text{TTM}$ with their approximating curves X and Y , respectively, as previously,

$$DL(\lambda \cdot \xi \oplus \kappa \cdot \eta) = \left. \frac{d}{dt} \right|_{t=0} (L(\lambda X + \kappa Y))$$

$$\begin{aligned}
&= \frac{d}{dt} \Big|_{t=0} (\lambda L(X) + \kappa L(Y)) \\
&= \lambda \cdot DL(\xi) \mathbf{+} \kappa \cdot DL(\eta)
\end{aligned}$$

for all $\lambda, \kappa \in \mathbb{R}$. Hence we achieve linearity with respect to both vector bundle structures of TTM . If we denote the base points as before, then this reads

$$D_{\lambda X_0 + \kappa Y_0} L(\lambda \cdot \xi \mathbf{+} \kappa \cdot \eta) = \lambda \cdot D_{X_0} L(\xi) \mathbf{+} \kappa \cdot D_{Y_0} L(\eta).$$

Remark A.4: Alignment of both vector bundle structures on TTM on the restricted vertical bundle

By definition, the vertical bundle VTM of TM is a subbundle of TTM . The zero section of TM is a natural embedding of M into TM ; this embedded submanifold will be denoted by $\overline{M}$. Then Diagram (A.1) restricts onto

$$\begin{array}{ccc}
VTM|_{\overline{M}} & \xrightarrow{D\pi_{TM}} & \overline{M} \\
\downarrow \pi_{TTM} & & \downarrow \cong \\
\overline{M} & \xrightarrow{\cong} & M
\end{array}$$

and $S = S_M$ does not only restrict onto that, S is actually the identity on $VTM|_{\overline{M}}$, as also stated in [10, §9.6, Thm. 9.6.1, page 363; but without proof]. This follows simply by the coordinate expressions Eq. (A.3) and (A.4) (recall the involved notation). $\xi \in VTM|_{\overline{M}}$ is in the kernel of both projections, $D\pi_{TM}$ and π_{TTM} , thus $\omega = X_0 = 0 \in T_p M$, therefore $S(\xi) = \xi$ by Eq. (A.3) and (A.4). By Remark A.1 the addition $\mathbf{+}$ and scalar multiplication $\cdot$ then also align with the typical addition $+$ and scalar multiplication $\cdot$, respectively, of TTM as tangent bundle.

Motivated by this, we can actually recover something similar for VTM . In this case

$$\begin{array}{ccc}
VTM & \xrightarrow{D\pi_{TM}} & \overline{M} \\
\downarrow \pi_{TTM} & & \downarrow \cong \\
TM & \xrightarrow{\pi_{TM}} & M
\end{array}$$

and, as already mentioned earlier, the vertical bundle of vector bundles is trivially the pullback of the bundle along itself, here $VTM \cong \pi_{TM}^* TM$. Thus, we have a canonical projection $\text{pr}_2 : VTM \rightarrow TM$ onto the second component, and we can write for $\xi = [\alpha] \in V_{X_0} TM$

$$\xi = \left(\frac{d}{dt} \Big|_{t=0} \frac{d}{ds} \Big|_{s=0} \alpha^i \right) \frac{\partial}{\partial(dx^i)} \Big|_{X_0} \cong \left(X_0, \left(\frac{d}{dt} \Big|_{t=0} \frac{d}{ds} \Big|_{s=0} \alpha^i \right) \frac{\partial}{\partial x^i} \Big|_p \right)$$

using the notation of Eq. (A.3) (especially $X_0 \in T_p M$, $p \in M$). Similarly for $\eta = [\beta] \in V_{Y_0} TM$ (same notation as previously in this appendix, i.e. $Y_0 \in T_p M$). Then by Eq. (A.5)

$$\lambda \cdot \xi \mathbf{+} \kappa \cdot \eta \cong \left(\lambda X_0 + \kappa Y_0, \left(\frac{d}{dt} \Big|_{t=0} \frac{d}{ds} \Big|_{s=0} (\lambda \alpha^i + \kappa \beta^i) \right) \frac{\partial}{\partial x^i} \Big|_p \right)$$

for all $\lambda, \kappa \in \mathbb{R}$. Therefore

$$\text{pr}_2(\lambda \cdot \xi \mathbf{+} \kappa \cdot \eta) = \lambda \text{pr}_2(\xi) + \kappa \text{pr}_2(\eta).$$

For $\zeta \in V_{X_0} TM$ we clearly get $\text{pr}_2(\lambda \xi + \kappa \zeta) = \lambda \text{pr}_2(\xi) + \kappa \text{pr}_2(\zeta)$, and so both linear structures on VTM align under pr_2 .

Remark A.5: Tangent lift, [25, §2.2, last paragraph in Subsection 2.2]

Let $X \in \mathfrak{X}(M)$, then its total derivative is a map $DX : TM \rightarrow TTM$ satisfying

$$D\pi_{TM} \circ DX = D \underbrace{(\pi_{TM} \circ X)}_{=\mathcal{K}_M} = \mathcal{K}_{TM},$$

so that DX is a section of $TTM \xrightarrow{D\pi_{TM}} TM$. Due to the fact that $S = S_M$ is a base-preserving vector bundle isomorphism from $TTM \xrightarrow{D\pi_{TM}} TM$ to $TTM \xrightarrow{\pi_{TM}} TM$ (and vice versa), we have a vector field $X_T \in \mathfrak{X}(TM)$ given by

$$X_T := S \circ DX,$$

called the **tangent lift** of X . We then also have

$$D\pi_{TM} \circ X_T = \pi_{TTM} \circ DX = X,$$

thus the label as tangent lift. We also have linearity properties (sort of) and a Leibniz rule, that is, we clearly get

$$X_T(\lambda Y + \kappa Z) = \lambda \cdot X_T(Y) \mathbf{+} \kappa \cdot X_T(Z).$$

for all $Y, Z \in T_p M$ and $\kappa, \lambda \in \mathbb{R}$, and we also have something similar w.r.t. X : Recall Eq. (A.3) and (A.4), including their notation w.r.t. to local coordinates $(x^i)_i$. That is, by definition of DX we can derive

$$D_p X(Y) = Y^i \frac{\partial}{\partial(\pi_{TM}^* x^i)} \Big|_{X_p} + Y^j \frac{\partial X^i}{\partial x^j} \Big|_p \frac{\partial}{\partial(dx^i)} \Big|_{X_p},$$

thus,

$$\begin{aligned} D_p(\kappa X + \lambda W)(Y) &= Y^i \frac{\partial}{\partial(\pi_{TM}^* x^i)} \Big|_{\lambda X_p + \kappa W_p} + Y^j \frac{\partial(\lambda X^i + \kappa W^i)}{\partial x^j} \Big|_p \frac{\partial}{\partial(dx^i)} \Big|_{\lambda X_p + \kappa W_p} \\ &= \lambda \cdot D_p X(Y) \mathbf{+} \kappa \cdot D_p W(Y) \end{aligned}$$

for all $X, W \in \mathfrak{X}(M)$ and $\lambda, \kappa \in \mathbb{R}$, and

$$\begin{aligned} D_p(fX)(Y) &= Y^i \frac{\partial}{\partial(\pi_{TM}^* x^i)} \Big|_{f(p)X_p} + f(p) Y^j \frac{\partial X^i}{\partial x^j} \Big|_p \frac{\partial}{\partial(dx^i)} \Big|_{f(p)X_p} + Y^j X^i(p) \frac{\partial f}{\partial x^j} \Big|_p \frac{\partial}{\partial(dx^i)} \Big|_{f(p)X_p} \end{aligned} \quad (\text{A.6})$$

for all $f \in C^\infty(M)$. Therefore

$$(\lambda X + \kappa W)_T = \lambda X_T + \kappa W_T,$$

and by Eq. (A.4)

$$(fX)_T(Y) = f(p) X_p^i \frac{\partial}{\partial(\pi_{TM}^* x^i)} \Big|_Y + f(p) Y^j \frac{\partial X^i}{\partial x^j} \Big|_p \frac{\partial}{\partial(dx^i)} \Big|_Y + Y(f) X^i(p) \frac{\partial}{\partial(dx^i)} \Big|_Y$$

so

$$(fX)_T = f X_T + df \otimes X^i \frac{\partial}{\partial(dx^i)}.$$

Similar to Remark A.4, the vertical bundle of TM is canonically isomorphic to $\pi_{TM}^* TM$, one can think of the second term as $df \otimes \pi_{TM}^* X$.