

Fourier Analysis Stein: Chapter 8. Problems.

Kelvin Hong
kh.boon2@gmail.com

Xiamen University Malaysia, Asia Pacific University Malaysia — August 17, 2024

1 Problems

1. Prove that there are infinitely many primes by observing that if there were only finitely many, p_1, \dots, p_N , then

$$\prod_{j=1}^N \frac{1}{1 - 1/p_j} \geq \sum_{n=1}^{\infty} \frac{1}{n}.$$

Solution: Assumes there are only finitely many primes p_1, \dots, p_N . Given a positive integer M , each positive integer $n \leq M$ can be expressed as a product of primes $p_1^{k_1} \cdots p_N^{k_N}$ for some integers k_1, \dots, k_N . We let K_1, \dots, K_N be the maximum values of k_1, \dots, k_N across all $n \leq M$. Thus, we have

$$\prod_{j=1}^N \frac{1}{1 - 1/p_j} \geq \prod_{j=1}^N \left(\sum_{k=0}^{K_j} \frac{1}{p_j^k} \right) \geq \sum_{n=1}^M \frac{1}{n}.$$

Taking the limit as $M \rightarrow \infty$, we see that the RHS diverges, which is a contradiction to our assumption. \square

2. In the text we showed that there are infinitely many primes of the form $4k + 3$ by a modification of Euclid's original argument. Adapt this technique to prove the similar result for primes of the form $3k + 2$, and for those of the form $6k + 5$.

Solution: Assume there are only finitely many primes of the form $3k + 2$, and let p_1, \dots, p_N be all of them in increasing order and $p_1 = 5$. Consider the number $n = 3p_1 \cdots p_N + 2$. This is a number of the form $3k + 2$ and $n > p_N$, so it must be composite by our assumption. Since it is not divisible by 3, it must have prime factors of the form $3k + 1$ and $3k + 2$. If it only has prime factors of the form $3k + 1$, the product of these primes would be still of the form $3k + 1$, so it will at least have a prime factor of the form $3k + 2$. But this is a contradiction since n is not divisible by any of the primes p_1, \dots, p_N .

For another solution, we assume there are only finitely many primes of the form $6k + 5$, q_1, \dots, q_M , in increasing order, where $q_1 = 11$. We can let $m = 6q_1 \cdots q_M + 5$, we can argue similarly by noting that if m is composite, it must have prime factors of the form $6k + 1$ and $6k + 5$, and also it must have at least one prime factor of the form $6k + 5$, which create a contradiction. \square

3. Prove that if p and q are relatively prime, then $\mathbb{Z}^*(p) \times \mathbb{Z}^*(q)$ is isomorphic to $\mathbb{Z}^*(pq)$.

Solution: Let $\phi : \mathbb{Z}^*(p) \times \mathbb{Z}^*(q) \rightarrow \mathbb{Z}^*(pq)$ be defined by $\phi(a, b) = aq + bp \pmod{pq}$. Obviously both sets are groups, so we only need to show the mapping is one-to-one and surjective.

To show ϕ is one-to-one, suppose $\phi(a, b) = \phi(c, d)$, then $aq + bp \equiv cq + dp \pmod{pq}$. There is an integer k such that $(a - c)q + (b - d)p = kpq$. Rearrange to get $(a - c)q = p(kq - b + d)$. Since p and q are relatively prime, p divides $a - c$, hence $a \equiv c \pmod{p}$. Similarly $b \equiv d \pmod{q}$.

To show ϕ is surjective, we first let x, y be integers such that $py + qx \equiv 1 \pmod{pq}$ with $x \in \mathbb{Z}^*(p)$ and $y \in \mathbb{Z}^*(q)$. For any $z \in \mathbb{Z}^*(pq)$, we can write $z = z \cdot 1 = z(py + qx) = zpy + zqx$. We find that $\phi(zx, zy) = z$ just because z is both relatively prime to p and q . \square

4. (Skipped)

5. If n is a positive integer, show that

$$n = \sum_{d|n} \varphi(d),$$

where φ is the Euler phi-function.

Solution: For any $1 \leq d \leq n$, if d is a divisor of n , then $\varphi(n/d)$ is the number of integers $1 \leq k \leq n/d$ such that $\gcd(k, n/d) = 1$. Let $m = kd$, then we have $\gcd(m, n) = d$. We see there are exactly $\varphi(n/d)$ integers m satisfying $\gcd(m, n) = d$.

Obviously for each $1 \leq m \leq n$, $\gcd(m, n)$ is a divisor of n , so we actually have

$$n = \sum_{d|n} \varphi(n/d) = \sum_{d|n} \varphi(d).$$

\square

6. (Skipped)

7. Recall that for $|z| < 1$,

$$\log_1 \left(\frac{1}{1-z} \right) = \sum_{k \geq 1} \frac{z^k}{k}.$$

We have seen that

$$e^{\log_1 \left(\frac{1}{1-z} \right)} = \frac{1}{1-z}.$$

(a) Show that if $w = 1/(1-z)$, then $|z| < 1$ if and only if $\Re(w) > 1/2$.

Solution: We rearrange to see $z = 1 - 1/w$, so $|z| < 1$ iff $1 - \frac{1}{w} - \frac{1}{\bar{w}} + \frac{1}{|w|^2} < 1$, we then have

$$\frac{1}{|w|^2} < \frac{1}{w} + \frac{1}{\bar{w}} = \frac{w + \bar{w}}{|w|^2} = \frac{2\Re(w)}{|w|^2}.$$

Since the above derivation are all equivalent, we have $|z| < 1$ iff $\Re(w) > 1/2$. \square

(b) Show that if $\Re(w) > 1/2$ and $w = \rho e^{i\varphi}$ with $\rho > 0, |\varphi| < \pi$, then

$$\log_1 w = \log \rho + i\varphi.$$

Solution: We have $e^{\log_1 w} = w$, so there is an integer k such that $\log_1 w = \log \rho + i(2\pi + \varphi)$. Now view $\log_1 w$ as a function of φ , when $\varphi = 0$ we have that $\log_1 w$ is real, and is exactly $\log \rho$, since k is an integer, it must be zero. \square

8. Let ζ denote the zeta function defined for $s > 1$.

(a) Compare $\zeta(s)$ with $\int_1^\infty x^{-s} dx$ to show that

$$\zeta(s) = \frac{1}{s-1} + O(1) \quad \text{as } s \rightarrow 1^+.$$

Solution: For each integer $N \geq 2$ we let $\zeta_N(s) = \sum_{n=1}^N n^{-s}$, we have

$$\begin{aligned}\zeta_N(s) &\leq 1 + \sum_{n=2}^N \int_{n-1}^n x^{-s} dx \\ &= 1 + \int_1^N x^{-s} dx \\ &= 1 + \frac{1}{s-1} - \frac{1}{(s-1)N^{s-1}} \\ &< 1 + \frac{1}{s-1}.\end{aligned}$$

By letting $N \rightarrow \infty$, we see $\zeta(s) \leq \frac{1}{s-1} + 1$ for every $s > 1$. Similarly, if we fixed $s > 1$, we have

$$\zeta_N(s) \geq \int_1^{N+1} x^{-s} dx = \frac{1}{s-1} - \frac{1}{(s-1)(N+1)^{s-1}} \implies \zeta(s) \geq \frac{1}{s-1}.$$

Hence this also hold for every $s > 1$. Note that we actually proved something stronger:

$$\frac{1}{s-1} \leq \zeta(s) \leq \frac{1}{s-1} + 1.$$

□

(b) Prove as a consequence that

$$\sum_p \frac{1}{p^s} = \log\left(\frac{1}{s-1}\right) + O(1) \quad \text{as } s \rightarrow 1^+.$$

Solution: Note that $\zeta(s) = \prod_p \frac{1}{1-p^{-s}}$, taking logarithm both sides give us

$$\begin{aligned}\log \zeta(s) &= - \sum_p \log\left(1 - \frac{1}{p^s}\right) \\ &= \sum_p \left(\frac{1}{p^s} + \sum_{k \geq 2} \frac{1}{k p^{ks}} \right)\end{aligned}$$

The second term can be bounded as below:

$$0 \leq \sum_{k \geq 2} \frac{1}{k p^{ks}} \leq \frac{1}{2} \sum_{k \geq 2} \frac{1}{p^{ks}} = \frac{1}{2} \cdot \frac{p^{-2s}}{1-p^{-s}} \leq p^{-2s}$$

which gives us

$$\log \zeta(s) = \sum_p \frac{1}{p^s} + \sum_p O\left(\frac{1}{p^{2s}}\right) = \sum_p \frac{1}{p^s} + O(1).$$

When taking $s \rightarrow 1^+$, we have proved the result.

□