

MAT195 TextBook Notes

Aman Bhargava

January 2019

Contents

0.1	Review: Memorizy Stuff	3
0.1.1	Trig Function Derivatives	3
0.1.2	Inverse Trig Derivatives	3
0.1.3	How to complete the Square	3
0.1.4	Trig Angle Sums	3
0.1.5	Hyperbolic Trig Functions	4
0.1.6	Inverse Hyperbolic Trig Function	4
0.2	Introduction and Course Description	4
1	Techniques of Integration (Chapter 7 in Textbook)	5
1.1	Integration by Parts	5
1.1.1	Tips for Integration by Parts:	6
1.2	Trigonometric Integrals	6
1.2.1	Strategy for $\int \sin^m(x)\cos^n(x)$	6
1.2.2	Strategy for $\int \tan^m(x)\sec^n(x)$	7
1.2.3	Strategy for $\int \sin(mx)\cos(nx)dx$	7
1.2.4	Strategy for $\int \csc^m(x)\cot^n(x)dx$	8
1.3	Trig Sub	8
1.3.1	Inverse Substitution	8
1.3.2	List of Trig Subs	8
1.3.3	General Layout for Trig Sub	9
1.4	Partial Fractions	9
1.4.1	Denominator is only distinct Linear Factors	9
1.4.2	Denominator is only Linear Factors but some are Repeated	9
1.4.3	Denominator has non-repeated Quadratic Factors	10
1.4.4	Denominator has repeated Quadratic Factors	10
1.4.5	Rationalizing Substitutions	10
1.5	Strategy for Solving Integrals	10
1.6	Improper Integrals (Involving Infinity)	11
1.6.1	Definition of Improper Integral	11
1.6.2	Comparison Theorem	11

2	Further Applications of Integration	12
2.1	Arc Length	12
2.1.1	Definition of Arc Length	12
2.1.2	Arc Length Formula	12
2.2	Surface Area of Revolution	12
2.2.1	Surface Area of Revolution Formulae	12
2.3	Applications to Physics and Engineering	13
2.3.1	Hydrostatic Force	13
2.3.2	Center of Mass	13
2.3.3	Rotating cross sections around line	13
3	Parametric Equations and Polar Coordinates	14
3.1	Parametric Equations	14
3.1.1	Tangents	14
3.1.2	Integrals	14
3.2	Polar Coordinates	15
3.2.1	Quick Info:	15
3.2.2	Symmetry	15
3.2.3	Tangents to Polar Curves	15
3.3	Areas and Lengths in Polar Coordinates	15
3.3.1	Area	15
3.3.2	Arc Length	16
4	Infinite Sequences and Series	17
4.1	Sequences	17
4.1.1	Limits of Sequences	17
4.1.2	Definitions	17
4.2	Series	18
4.2.1	Partial Sums	18
4.2.2	Infinite Series	18
4.2.3	Geometric Series	18
4.2.4	Test for Divergence	18
4.2.5	Working with Series	18
4.2.6	Comparison Test	18
4.2.7	P-Series	19
4.3	The Integral Test and Estimates of Sums	19
4.3.1	Integral Test	19
4.3.2	Estimating The Sum of a Series	19

0.1 Review: Memorizy Stuff

0.1.1 Trig Function Derivatives

$$\frac{d}{dx}\sin(x) = \cos(x) \quad \frac{d}{dx}\csc(x) = -\csc(x)\cot(x)$$

$$\frac{d}{dx}\cos(x) = -\sin(x) \quad \frac{d}{dx}\sec(x) = \sec(x)\tan(x)$$

$$\frac{d}{dx}\tan(x) = \sec^2(x) \quad \frac{d}{dx}\cot(x) = -\csc^2(x)$$

0.1.2 Inverse Trig Derivatives

$$\frac{d}{dx}\sin^{-1}(x) = \frac{1}{\sqrt{1-x^2}}$$

$$\frac{d}{dx}\cos^{-1}(x) = \frac{-1}{\sqrt{1-x^2}}$$

$$\frac{d}{dx}\tan^{-1}(x) = \frac{1}{1+x^2}$$

0.1.3 How to complete the Square

1. Put $ax^2 + bx$ in brackets and forcefully factor out the a
2. Add $(\frac{b}{2})^2$ to the inside of the brackets and subtract it from the outside (you got it)
3. Factor and be happy that you've completed the square;

0.1.4 Trig Angle Sums

1. $\sin(A + B) = \sin(A)\cos(B) + \cos(A)\sin(B)$
2. $\cos(A + B) = \cos(A)\cos(B) - \sin(A)\sin(B)$
3. $\sin(A - B) = \sin(A)\cos(B) - \cos(A)\sin(B)$
4. $\cos(A - B) = \cos(A)\cos(B) + \sin(A)\sin(B)$

0.1.5 Hyperbolic Trig Functions

0.1.6 Inverse Hyperbolic Trig Function

0.2 Introduction and Course Description

Chapter 1

Techniques of Integration (Chapter 7 in Textbook)

1.1 Integration by Parts

Integration by parts is basically just the reverse product rule.

$$\text{Product rule: } d/dx[f(x)g(x)] = f'(x)g(x) + f(x)g'(x)$$

You could reverse this simply, but it wouldn't be that useful. The more useful form that *is* the integration by parts formula looks like this:

$$\int [f(x)g'(x)] = f(x)g(x) - \int [f'(x)g(x)]$$

You can think through that one pretty easily — you are just splitting up the initial integral, then moving half of it to the other side.

That's pretty useful, but there's an even more useful way to write the formula, and it looks like this:

$$\int u dv = uv - \int v du$$

This works because we let $u = f(x)$ and $v = g(x)$. $g'(x) = v' = dv/dx$,

and same with u' . So to get to that formula we go like this:

$$\int uv' dx = uv - \int u'v dx \quad (1.1)$$

$$\int u \frac{dv}{dx} dx = uv - \int v \frac{du}{dx} dx \quad (1.2)$$

$$\int u dv = uv - \int v du \quad (1.3)$$

1.1.1 Tips for Integration by Parts:

- When using the uv equation, it's useful to define things in this order:
 - $u = ?$ $dv = ?$
 - $du = ?$ $v = ?$
 - keeping in mind that $u' dx = du$ and $\int \frac{dv}{dx} dx = \int v' = v$
- Choose your u so that it becomes simpler when differentiated, and let your v be the thing that gets a little hairier.
- Practice a lot from the textbook, ya dingus

1.2 Trigonometric Integrals

There are a bunch of configurations of trig functions for which we need to learn the steps necessary to take the integral. It's important to practice this because recognizing the form of the integral is the most difficult thing to do here.

1.2.1 Strategy for $\int \sin^m(x) \cos^n(x)$

If the power of cosine is odd: "Save" one of the cosine terms, and then express it as $\int \sin^m(x) \cos^{2k+1}(x) * \cos(x)$. Then turn the $\cos^{2k+1}(x)$ into sin terms with pythagorean identity. Then substitute $u = \sin(x)$ and solve.

If the power of sine is odd: Do the same thing but reverse \sin and \cos (save one $\sin(x)$ and sub u for $\cos(x)$)

If both powers are even: Use the following identities to help you solve it:

$$\sin^2(x) = 1/2(1 - \cos(2x)) \quad (1.4)$$

$$\cos^2(x) = 1/2(1 + \cos(2x)) \quad (1.5)$$

$$\sin x \cos x = 1/2 \sin 2x \quad (1.6)$$

1.2.2 Strategy for $\int \tan^m(x) \sec^n(x)$

If the power of $\sec(x)$ is even, "save" a factor of $\sec^2(x)$ and use identity $\sec^2(x) = 1 + \tan^2(x)$ to express the rest in terms of $\tan(x)$. Then substitute $u = \tan(x)$

If the power of tangent is odd, save a factor of $\sec(x)\tan(x)$ and convert the rest of the $\tan(x)$'s using $\tan^2(x) = \sec^2(x) - 1$

Also note the following:

$$\int \tan(x) dx = \ln|\sec(x)| + C$$

$$\int \sec(x) dx = \ln|\sec(x) + \tan(x)| + C$$

Remember this as well:

$$\frac{d}{dx} \tan(x) = \sec^2(x)$$

$$\frac{d}{dx} \sec(x) = \sec(x)\tan(x)$$

1.2.3 Strategy for $\int \sin(mx)\cos(nx)dx$

Use the following identities:

$$\sin A \cos B = 1/2[\sin(A - B) + \sin(A + B)] \quad (1.7)$$

$$\sin A \sin B = 1/2[\cos(A - B) - \cos(A + B)] \quad (1.8)$$

$$\cos A \cos B = 1/2[\cos(A - B) + \cos(A + B)] \quad (1.9)$$

1.2.4 Strategy for $\int \csc^m(x) \cot^n(x) dx$

Know the following things:

$$\frac{d}{dx} \csc(x) = -\csc(x) \cot(x) \quad (1.10)$$

$$\cot^2(x) = \csc^2(x) - 1 \quad (1.11)$$

$$\frac{d}{dx} \cot(x) = -\csc^2(x) \quad (1.12)$$

1.3 Trig Sub

What is trig sub? Trig sub is when you use the *inverse substitution* rule in conjunction with useful trigonometric identities and trigonometric integrals to solve integrals that you wouldn't otherwise be able to solve.

1.3.1 Inverse Substitution

Unlike u-substitution, you are substituting in a non-equivalent function ($g(x)$) for x instead of substituting a variable like u for the actual value of x . Hence, the following result arises:

$$\int f(x) dx = \int f(g(x)) g'(x) dx$$

Qualifications: g must have an inverse function, and g must be one-to-one.

1.3.2 List of Trig Subs

The main use of trig subs is to get rid of irritating radical signs that make integration hard. The following is a table of types (from the textbook):

Expression	Substitution	Identity
$\sqrt{a^2 - x^2}$	$x = a \sin \theta, -\pi/2 \leq \theta \leq \pi/2$	$1 - \sin^2 \theta = \cos^2 \theta$
$\sqrt{a^2 + x^2}$	$x = a \tan \theta, -\pi/2 < \theta < \pi/2$	$1 + \tan^2 \theta = \sec^2 \theta$
$\sqrt{x^2 - a^2}$	$x = a \sec \theta, -\pi/2 \leq \theta \leq \pi/2$	$\sec^2 \theta - 1 = \tan^2 \theta$

1.3.3 General Layout for Trig Sub

1. Make sure there is no other way (e.g. u-sub, etc.)
2. If there is a quadratic in the root, complete the square
3. Recognize the stuff in the root as one of the three.
4. Set $x = a * \text{trig}(\theta)$ and find dx in terms of θ
5. Solve the stuff.

1.4 Partial Fractions

This is just a way to break up rational functions in to little pieces that we can actually deal with. A proper rational function is one where the power on the top polynomial is lower than that of the bottom. Improper rationals are the other way around.

In order to use partial fractions, you need to make the rational function a proper one.

There are a few cases to consider for splitting things up into partial fractions:

1.4.1 Denominator is only distinct Linear Factors

1. Set an equality between the original rational and $\frac{A}{(\text{root}_1)} + \frac{B}{(\text{root}_2)} + \dots$
2. Multiply both sides by the denominator of the rational
3. Expand and solve for A, B, \dots

1.4.2 Denominator is only Linear Factors but some are Repeated

It's roughly the same as last time, EXCEPT:

1. Suppose (root_1) is repeated k times so $(\text{root}_1)^k$ is a factor
2. Then you need to use $\frac{A_1}{(\text{root}_1)} + \frac{A_2}{(\text{root}_1)^2} + \dots + \frac{A_k}{(\text{root}_1)^k}$
3. Now solve as you did last time.

1.4.3 Denominator has non-repeated Quadratic Factors

Basically just have a linear term on top in the expansion, so $\frac{A_1x+B}{ax^2+bx+c}$ would be a term in the thing.

Also, $\int \frac{dx}{x^2+a^2} = \frac{1}{a} \tan^{-1}\left(\frac{x}{a}\right) + C$

1.4.4 Denominator has repeated Quadratic Factors

Roughly the same as when you have linear repeated factors. if you have $(root_1)^k$, then you get

$$\frac{Ax+B}{(root_1)} + \frac{Cx+D}{(root_1)^2} + \dots + \frac{Yx+Z}{(root_1)^k}$$

Then just solve as before.

1.4.5 Rationalizing Substitutions

Use substitutions to make annoying functions into rational functions and solve from there. For instance, if you have a radical in the numerator, make the substitution $u^2 =$ whatever was in the radical.

1.5 Strategy for Solving Integrals

1. Simplify the integrand with identities/algebra if possible.
2. Look for an obvious u-sub
3. Classify the integrand according to its form:
 - (a) Trig integral ($\sin^m(x)\cos^n(x)$, etc.)
 - (b) Rational Function
 - (c) Integration by parts - especially with *polynomial*transcendental function*
 - (d) Radicals \rightarrow consider trig sub ($x = a \tan \theta$, etc.)
4. Try again, using several methods, and drawing from your MASSIVE PAST EXPERIENCE

1.6 Improper Integrals (Involving Infinity)

1.6.1 Definition of Improper Integral

$\int_a^\infty f(x)dx = \lim_{t \rightarrow \infty} \int_a^t f(x)dx$ Convergent improper integrals have an actual value. Divergent improper integrals don't. Integrals from negative infinity to positive infinity exist if and only if the integral from -infinity to a converges and the integral from a to +infinity converges (it is the sum of the two).

For $\int_1^\infty \frac{1}{x^p} dx$ is convergent if $p > 1$

For integrals where the y value goes to infinity , then you take limits around where it goes to infinity, ya dingus.

1.6.2 Comparison Theorem

You can prove that an integral is convergent if you can find another one for which the function is strictly of greater magnitude that is convergent (and likewise for non-convergency)

Chapter 2

Further Applications of Integration

2.1 Arc Length

2.1.1 Definition of Arc Length

Arc length is the actual length of a curve. It is defined as:

$$L = \lim_{n \rightarrow \infty} \sum_{i=1}^n |P_{i-1}, P_i|$$

2.1.2 Arc Length Formula

$$L = \int_a^b \sqrt{1 + [f'(x)]^2} dx$$

2.2 Surface Area of Revolution

2.2.1 Surface Area of Revolution Formulae

$$S = \int_a^b 2\pi y \sqrt{1 + \left(\frac{dy}{dx}\right)^2} dx$$

$$S = \int_a^b 2\pi y \sqrt{1 + \left(\frac{dx}{dy}\right)^2} dy$$

2.3 Applications to Physics and Engineering

2.3.1 Hydrostatic Force

Integral of the area of each slice times the pressure on that slice at that depth
(pressure = density * depth)

2.3.2 Center of Mass

For a thin plate on the plane, the centroid is at (\bar{x}, \bar{y}) where:

$$\bar{x} = \frac{1}{A} \int_a^b x(f(x) - g(x))dx$$

$$\bar{y} = \frac{1}{2A} \int_a^b (f(x)^2 - g(x)^2)dx$$

2.3.3 Rotating cross sections around line

If cross section is completely outside of the line, volume = area * distance traveled by cross section.

Chapter 3

Parametric Equations and Polar Coordinates

3.1 Parametric Equations

3.1.1 Tangents

$$\frac{dy}{dx} = \frac{\frac{dy}{dt}}{\frac{dx}{dt}}$$
$$\frac{d^2y}{dx^2} = \frac{\frac{d}{dt}\left(\frac{dy}{dx}\right)}{\frac{dx}{dt}}$$

3.1.2 Integrals

Normal Integral: Area from $\alpha \rightarrow \beta$

$$\int_{\alpha}^{\beta} y(t) * \frac{dx}{dt} dt$$

Arc Length

$$L = \int_{\alpha}^{\beta} \sqrt{\frac{dx^2}{dt} + \frac{dy^2}{dt}} dt$$

Surface Area of Revolution:

$$S = \int_{\alpha}^{\beta} 2\pi y \sqrt{\frac{dx^2}{dt} + \frac{dy^2}{dt}} dt$$

3.2 Polar Coordinates

3.2.1 Quick Info:

1. Points are in form (r, θ)
2. Origin denoted by O or as "pole"
3. For point (x, y) in cartesian space and point (r, θ) :
 - $x = r\cos\theta$
 - $y = r\sin\theta$
 - $\tan\theta = \frac{x}{y}$

3.2.2 Symmetry

If an equation is unchanged when $\theta \rightarrow -\theta$, it is symmetric about line $\theta = 0$
If an equation is unchanged when $r \rightarrow -r$ OR $\theta \rightarrow \theta + \pi$, it is symmetric about the pole. If an equation is unchanged when $\theta \rightarrow \pi - \theta$ then the curve is symmetric about the line $\theta = \pi/2$ (a vertical line in polar coordinates)

3.2.3 Tangents to Polar Curves

Treat as parametric equation. Steps:

$$x = r\cos\theta = f(\theta)\cos(\theta)$$

$$y = r\sin\theta = f(\theta)\sin(\theta)$$

Then $\frac{dy}{dx}$ is just $\frac{dy}{d\theta} / \frac{dx}{d\theta}$

3.3 Areas and Lengths in Polar Coordinates

3.3.1 Area

Area of a Sector of a Circle

$$A = \frac{1}{2}r^2\theta$$

Area Inside Curve:

$$A = \int_a^b \frac{1}{2}(f(\theta))^2 d\theta$$

$$A = \int_a^b \frac{1}{2}r^2 d\theta$$

3.3.2 Arc Length

Review of Parametric Arc Length

$$L = \int_a^b \sqrt{r^2 + \left(\frac{dr}{d\theta}\right)^2} d\theta$$

Chapter 4

Infinite Sequences and Series

4.1 Sequences

Definition: List of numbers written in definite order. Notation is the same as set notation.

4.1.1 Limits of Sequences

A sequence has a limit L if: you can make a_n as close as you want to L by increasing n

You can incorporate $(\delta) \epsilon$ notation if you want to.

Also, if you can find a function that matches the sequence at all integer points, then you can just find the limit of the function using regular limit rules to find the limit of the sequence.

You can also disperse the a limit inside of a function.

$$\lim_{n \rightarrow \infty} \sin(\pi/n) = \sin(\lim_{n \rightarrow \infty} \pi/n)$$

4.1.2 Definitions

Monotonic sequences are sequences that are either strictly increasing or strictly decreasing.

Bounded above if no value of n will make a_n greater than M . Ditto bounded below.

Monotonic Sequence Theorem: Every bounded, monotonic sequence is convergent.

4.2 Series

A series is: a sum of a sequence (often infinite)

4.2.1 Partial Sums

$$s_1 = a_1, s_2 = a_1 + a_2, s_n = a_1 + \dots + a_n$$

$$s_n = \sum_{i=1}^n a_i$$

4.2.2 Infinite Series

$$\sum_{n=1}^{\infty} a_n = \lim_{n \rightarrow \infty} \sum_{i=1}^n a_i$$

4.2.3 Geometric Series

If $a_n = (a_1)r^n$, then $s_n = a + ar + ar^2 + \dots + ar^{n-1}$

$$s_n = \frac{a(1 - r^n)}{1 - r}$$

4.2.4 Test for Divergence

If $\lim_{n \rightarrow \infty} a_n$ does not exist or equals ∞ , then the infinite series of a_n is divergent.

4.2.5 Working with Series

You can add and subtract series normally.

$$\sum (a_n + b_n) = \sum a_n + \sum b_n$$

$$\sum (c * a_n) = c \sum a_n$$

4.2.6 Comparison Test

Let a_n be one series you DON'T know the limit of and b_n be another. Make sure a_n/b_n divides nicely. let $c = \lim_{n \rightarrow \infty} a_n/b_n$

Use c as a comparator - if it tells you that a is bigger than b and b is known to be divergent, then a must be divergent, etc.

4.2.7 P-Series

A p-series is of the form:

$$\sum_{n=1}^{\infty} 1/n^p$$

If $p > 1$, it converges. Otherwise it diverges.

4.3 The Integral Test and Estimates of Sums

4.3.1 Integral Test

It's not easy to find the sum of series except for geometric series and $\sum 1/[n(1+n)]$.

To prove that a sum diverges, take the integral of the continuous function $f(n) = a_n$ from $1 \rightarrow \infty$. If the integral is divergent, then so is the sum. If the integral is convergent, so is the sum. Think about the geometric argument and draw out the boxes on the page if necessary (reimann sums).

4.3.2 Estimating The Sum of a Series

Any partial sum of a series is an approximation of the infinite sum. We can get a good picture of how good the approximation is via the **remainder**.

$$R_n = s - s_n = a_{n+1} + a_{n+2} + \dots$$

By the integral test (and intuition), the remainder is less than or equal to the integral from n to ∞

$$R_n = a_{n+1} + a_{n+2} + \dots \leq \int_n^{\infty} f(x)dx$$

$$R_n \geq \int_{n+1}^{\infty} f(x)dx$$

In other words, the

Remainder Estimate for the Integral Test is:

$$\int_{n+1}^{\infty} f(x)dx \leq R_n \leq \int_n^{\infty} f(x)dx$$

4.4 Comparison Tests

4.4.1 The Limit Comparison Test

Let $\sum a_n$ and $\sum b_n$ are series with positive terms. If

$$\lim_{n \rightarrow \infty} \frac{a_n}{b_n} = c$$

and c is a finite number $\neq 0$, then both either converge or diverge.