
Personal Sensing in Clinical Research


Kendra Wyant

PI: John Curtin

UW-Madison Department of Psychology








Personal Sensing and Mental Health

- Screening
 - Passive
 - Scalable
- Monitoring
 - Intervention prior to relapse




Alcohol Use Disorder (AUD)

- AUD is a chronic relapsing disease
- Lapses are often early signs of relapse
- A temporally dynamic sensing system can capture day-to-day changes in lapse risk



Will people find these measures acceptable?

Sample Demographics



Active Mixed Passive

Mixed

Passive























































Mixed






















Passive









"This was my favorite part of the study. It helped me to set a good intention towards my recovery."



"This was my favorite part of the study. It helped me to set a good intention towards my recovery." "I may not want to always mentally realize how often I feel bad."

"If I was more of a diary type of person I'd probably be more into this kind of thing. I mostly just didn't know what to say."



"This was my favorite part of the study. It helped me to set a good intention towards my recovery."

"Nice to speak rather than type through the check in."

"To be able to verbilize my feelings was the most best therapy."



"I may not want to always mentally realize how often I feel bad."

"If I was more of a diary type of person I'd probably be more into this kind of thing. I mostly just didn't know what to say."

"This was my favorite part of the study. It helped me to set a good intention towards my recovery."

"Nice to speak rather than type through the check in."

"To be able to verbilize my feelings was the most best therapy."



"I may not want to always mentally realize how often I feel bad."

"If I was more of a diary type of person I'd probably be more into this kind of thing. I mostly just didn't know what to say."

"I felt like I needed to seek out a private spot to do the audio message. I would have much preferred to type the message."

"It takes time out of your day where you have to completely switch locations just so you can do it in private. I don't like that people could hear me and the topic wherever and whenever so I stopped using it."

"I went about my days as I normally would and never thought about it."



"I went about my days as I normally would and never thought about it."


"Some things were personal and sometimes I forgot they were being looked at."



"I went about my days as I normally would and never thought about it."

"I don't mind turning this info over and I trust the people running the study."

"Some things were personal and sometimes I forgot they were being looked at."



"I went about my days as I normally would and never thought about it."

"I don't mind turning this info over and I trust the people running the study."

"Some things were personal and sometimes I forgot they were being looked at."

"I trust no one tracking my information. Too much of our privacy and protection is lost already In today's society."

"It freaks me out that you can know everything I say. I sometimes say sensitive things."





"I went about my days as I normally would and never thought about it."

"I don't mind turning this info over and I trust the people running the study."

"It is a huge invasion but my sobriety is paramount to my overall health."

"Loss of privacy for the sake of sobriety, okay I'll give in."





"Some things were personal and sometimes I forgot they were being looked at."

"I trust no one tracking my information. Too much of our privacy and protection is lost already In today's society."

"It freaks me out that you can know everything I say. I sometimes say sensitive things."

"I went about my days as I normally would and never thought about it."

"I don't mind turning this info over and I trust the people running the study."

"It is a huge invasion but my sobriety is paramount to my overall health."

"Loss of privacy for the sake of sobriety, okay I'll give in."





"Some things were personal and sometimes I forgot they were being looked at."

"I trust no one tracking my information. Too much of our privacy and protection is lost already In today's society."

"It freaks me out that you can know everything I say. I sometimes say sensitive things."

"It seems this was more to aid with the study research than to help with my recovery. If I could see a way it would be prioritized to help me, I would be more willing to have them tracked longer. "

Can we predict an alcohol lapse from cellular communication logs?

Cellular Communications



- Passive
- Widespread use
- Understudied personal sensing measure for AUD
- Promising signal of lapse risk









Logs

- Phone number of other party
- Duration of call
- Date and time of message/call
- Type of message/call

How many contacts people communicate with each day.

Ratio of incoming to outgoing SMS messages.

A decline in number of outgoing calls.

Frequent communications with a new number.

An increase in outgoing messages on Friday and Saturday nights.

How many supportive contacts.

How many contacts in recovery.

Average level of pleasantness of interactions with contacts.

Ratio of drinking to non-drinking contacts.

How many drinkers in family.

Context

- Type of relationship
- Drank with contact in past
- Drinking status of contact
- Recovery status of contact
- Would drink in front of them
- Pleasantness of interactions
- Supportiveness










Prediction Aims

Aim 1: Train and evaluate the best performing machine learning model to predict alcohol lapse from cellular communication logs and contextual information about the interactions



Prediction Aims


Aim 1: Train and evaluate the best performing machine learning model to predict alcohol lapse from cellular communication data and contextual information about the interactions

Aim 2: Employ a model comparison approach to compare models that use all available features with models that are restricted to only passive signals

Explanation Aim

Aim 3: Evaluate the importance of feature sets within the top performing model to inform treatment





Clinical Significance

References

- Bae S, Ferreira D, Suffoletto B, Puyana JC, Kurtz R, Chung T, Dey AK. Detecting Drinking Episodes in Young Adults Using Smartphone-based Sensors. Proc ACM Interact Mob Wearable Ubiquitous Technol 2017 Jun 30;1(2):1–36.
- Bae S, Chung T, Ferreira D, Dey AK, Suffoletto B. Mobile phone sensors and supervised machine learning to identify alcohol use events in young adults: Implications for just-in-time adaptive interventions. Addict Behav 2018 Aug 1;83:42–47.
- Bai R, Xiao L, Guo Y, Zhu X, Li N, Wang Y, Chen Q, Feng L, Wang Y, Yu X, Xie H. Tracking and Monitoring Mood Stability of Patients With Major Depressive Disorder by Machine Learning Models Using Passive Digital Data: Prospective Naturalistic Multicenter Study. JMIR MHEALTH UHEALTH 2021;15.
- Barnett I, Torous J, Staples P, Sandoval L, Keshavan M, Onnela J-P. Relapse prediction in schizophrenia through digital phenotyping: a pilot study. Neuropsychopharmacology 2018 Jul;43(8):1660–1666.
- De Choudhury M, Counts S, Horvitz E. Predicting postpartum changes in emotion and behavior via social media. Proc SIGCHI Conf Hum Factors Comput Syst CHI 13 [Internet] Paris, France: ACM Press; 2013 [cited 2018 Aug 30]. p. 3267.
- Eichstaedt JC, Smith RJ, Merchant RM, Ungar LH, Crutchley P, Preoţiuc-Pietro D, Asch DA, Schwartz HA. Facebook language predicts depression in medical records. Proc Natl Acad Sci National Academy of Sciences; 2018 Oct 30;115(44):11203–11208.
- Epstein DH, Tyburski M, Kowalczyk WJ, Burgess-Hull AJ, Phillips KA, Curtis BL, Preston KL. Prediction of stress and drug craving ninety minutes in the future with passively collected GPS data. Npj Digit Med 2020 Dec;3(1):26.
- Grünerbl A, Muaremi A, Osmani V, Bahle G, Ohler S, Troester G, Mayora O, Haring C, Lukowicz P. Smart-Phone Based Recognition of States and State Changes in Bipolar Disorder Patients. IEEE J Biomed Health Inform 2014 Jul 25;19.
- Mcmanus K, Mallory E, Goldfeder R, Haynes W, Tatum J. Mining Twitter Data to Improve Detection of Schizophrenia. AMIA Jt Summits Transl Sci Proc AMIA Summit Transl Sci 2015 Aug 26;2015:122–6.
- Palmius N, Tsanas A, Saunders K, Bilderbeck A, Geddes J, Goodwin G, de Vos M. Detecting Bipolar Depression From Geographic Location Data. IEEE Trans Biomed Eng 2016 Oct 25:PP:1–1.
- Razavi R, Gharipour A, Gharipour M. Depression screening using mobile phone usage metadata: a machine learning approach. J Am Med Inform Assoc 2020 Apr 1;27(4):522–530.
- Reece AG, Reagan AJ, Lix KLM, Dodds PS, Danforth CM, Langer EJ. Forecasting the onset and course of mental illness with Twitter data. Sci Rep 2017 Oct 11;7(1):1–11.
- Sekulić I, Gjurković M, Šnajder J. Not Just Depressed: Bipolar Disorder Prediction on Reddit. ArXiv181104655 Cs [Internet] 2019 Mar 27.
- Wang Y, Ren X, Liu X, Zhu T. Examining the Correlation Between Depression and Social Behavior on Smartphones Through Usage Metadata: Empirical Study. JMIR MHealth UHealth 2021 Jan 6;9(1):e19046.
- Wolf M, Theis F, Kordy H. Language Use in Eating Disorder Blogs: Psychological Implications of Social Online Activity. J Lang Soc Psychol SAGE Publications Inc; 2013 Jun 1;32(2):212–226.


Images


- https://www.google.com/url?sa=i&url=https%3A%2F%2Feconsultancy.com%2Fwhat-is-social-media-here-are-34-definitions%2F&psig=AOvVaw2CTHW2qy4qeCwi6an-GkXJ&ust=1617384698085000&source=images&cd=vfe&ved=0CA0QjhxqFwoTCMC-3tfJ3e8CFQAAAAAAAAAAAAAAA
- Vecteezy images
- https://srilankamirror.com/news/16621-don-t-call-back-on-international-missed-calls-trc
- https://www.iup.edu/counselingcenter/services/groups-for-iup-students/

Thank You!

Questions?

Thoughts?





Demographics for enrolled participants (N = 169)

	n	%	М	SD
Age			40.5	12.1
Sex				
Female	83	49.1		
Male	86	50.9		
Race				
American Indian/Alaska Native	3	1.8		
Asian	2	1.2		
Black/African American	8	4.7		
White	146	86.4		
Multiracial or not listed	10	5.9		
Hispanic, Latino, or Spanish Origin				
Yes	5	3.0		
No	164	97.0		
Income			33855	31311
Marital Status				
Never married	78	46.2		
Married	35	20.7		
Divorced	48	28.4		
Separated	6	3.6		
Widowed	2	1.2		

Why Machine Learning?

- Well suited for handling the complexities of AUD and diversity across individuals
- Can fit many models without risk of overfitting because the models are evaluated on new data (i.e., validation and test set)
- Feature engineering
- Can be used for both prediction and explanation

Log Type	Measure	
SMS	Phone number of the other party	
	Date message was received	
	Date message was sent	
	Message type (incoming, outgoing, draft)	
	Contact saved in phone	
	Read status (read, unread)	
Call	Phone number of the other party	
	Date of call	
	Call type (incoming, outgoing, missed, voicemail, rejected, blocked)	
	Facetime (iPhone only)	
	Call duration (in seconds)	
	Contact saved in phone	
	Caller ID (allowed, restricted, unknown, payphone)	