

## 1. Problems

**Problema 1.** Let  $x_1, x_2, \dots, x_n > 0$  such that  $\frac{1}{1+x_1} + \dots + \frac{1}{1+x_n} = 1$ . Prove that

$$x_1 x_2 \dots x_n \geq (n-1)^n.$$

**Solución.** We see that

$$1 - \frac{1}{1+x_1} = \frac{1}{1+x_2} + \dots + \frac{1}{1+x_n} \geq (n-1) \sqrt[n-1]{\frac{1}{1+x_2} \times \dots \times \frac{1}{1+x_n}} = \frac{n-1}{\sqrt[n-1]{(1+x_2) \dots (1+x_n)}}$$

So

$$\frac{x_1}{1+x_1} \geq \frac{n-1}{\sqrt[n-1]{(1+x_2) \dots (1+x_n)}}$$

Generally

$$\frac{x_k}{1+x_k} \geq \frac{n-1}{\sqrt[n-1]{\prod_{i \neq k} (1+x_i)}}$$

Multiply this result for  $k = 1, 2, \dots, n$  we find that

$$\begin{aligned} \frac{x_1}{1+x_1} \cdot \dots \cdot \frac{x_n}{1+x_n} &\geq \frac{(n-1)^{n-1}}{\sqrt[n-1]{(1+x_1)^{n-1} \cdot (1+x_2)^{n-1} \dots (1+x_n)^{n-1}}} \\ \frac{x_1}{1+x_1} \cdot \dots \cdot \frac{x_n}{1+x_n} &\geq \frac{(n-1)^{n-1}}{(1+x_1) \cdot (1+x_2) \dots (1+x_n)} \\ \implies x_1 x_2 \dots x_n &\geq (n-1)^n. \end{aligned}$$

■

**Problema 2.** For positive  $a, b, c$  prove that

$$\sqrt{\frac{a}{b+c}} + \sqrt{\frac{b}{c+a}} + \sqrt{\frac{c}{a+b}} > 2$$

**Solución.** Given real numbers  $x_1, \dots, x_n$  we have

$$\sqrt[n]{x_1 \cdot x_2 \dots x_n} \geq \frac{n}{\frac{1}{x_1} + \frac{1}{x_2} + \dots + \frac{1}{x_n}}.$$

where the case of equality occurs when  $x_1 = x_2 = \dots = x_n$ . So, we find that

$$\sqrt{\frac{a}{b+c}} = \sqrt{1 \cdot \frac{a}{b+c}} \geq \frac{2}{1 + \frac{b+c}{a}} \geq \frac{2a}{a+b+c}$$

Similary  $\sqrt{\frac{b}{c+a}} \geq \frac{2b}{a+b+c}$  and  $\sqrt{\frac{c}{a+b}} \geq \frac{2c}{a+b+c}$  where the equality occurs when  $1 = \frac{a}{b+c} = \frac{b}{c+a} = \frac{c}{a+b}$  it's which no posible. Then  $\sqrt{\frac{a}{b+c}} + \sqrt{\frac{b}{c+a}} + \sqrt{\frac{c}{a+b}} > 2$ . ■

**Problema 3.** For positive  $a, b, c, d$  prove that

$$1 < \frac{a}{a+b+c} + \frac{b}{b+c+d} + \frac{c}{c+d+a} + \frac{d}{d+a+b} < 2.$$

**Solución.** Since  $a, b, c, d$  are positive, we find

$$1 < \frac{a^2}{a^2+ab+ac} + \frac{b^2}{b^2+bc+bd} + \frac{c^2}{c^2+cd+ac} + \frac{d^2}{d^2+ad+bd} < 2.$$

and we take  $A = \frac{a^2}{a^2+ab+ac} + \frac{b^2}{b^2+bc+bd} + \frac{c^2}{c^2+cd+ac} + \frac{d^2}{d^2+ad+bd}$ . By Cauchy-Schwarz we find that

$$(a^2 + b^2 + c^2 + d^2 + ab + 2ac + ad + bc + 2bd + cd) \cdot A \geq (a + b + c + d)^2$$

# GTN EXCL Program

---

$$\begin{aligned} \iff [(a+c)^2 + (a+c)(b+d) + (b+d)^2] \cdot A &\geq (a+c)^2 + 2(a+c)(b+d) + (b+d)^2 \\ \iff A &\geq \frac{(a+c)^2 + 2(a+c)(b+d) + (b+d)^2}{(a+c)^2 + (a+c)(b+d) + (b+d)^2} \end{aligned}$$

Since  $(a+c)^2 + 2(a+c)(b+d) + (b+d)^2 > (a+c)^2 + (a+c)(b+d) + (b+d)^2$ , hence  $A > 1$ .

It's clear that

$$\begin{aligned} -2 &< -\frac{a}{a+b+c} - \frac{b}{b+c+d} - \frac{c}{c+d+a} - \frac{d}{d+a+b} < -1 \\ 2 &< \frac{b+c}{a+b+c} + \frac{c+d}{b+c+d} + \frac{d+a}{c+d+a} + \frac{a+b}{d+a+b} < 3. \end{aligned}$$

Set  $B = \left( \frac{b+c}{a+b+c} + \frac{c+d}{b+c+d} + \frac{d+a}{c+d+a} + \frac{a+b}{d+a+b} \right)$ . By Cauchy-Schwarz we find that

$$\begin{aligned} [(b+c)(a+b+c) + (c+d)(b+c+d) + (d+a)(c+d+a) + (a+b)(d+a+b)] \cdot B &\geq [2(a+b+c+d)]^2 \\ [(a+b+c+d)^2 + a^2 + b^2 + c^2 + d^2 + (a+c)(b+d)] \cdot B &\geq [2(a+b+c+d)]^2 \end{aligned}$$

We know that  $(a+b+c+d)^2 > a^2 + b^2 + c^2 + d^2 + (a+c)(b+d)$  because this inequality is reducible to  $(a+c)(b+d) + 2(ac+bd) > 0$  which is true because all number are positive. Hence

$$2(a+b+c+d)^2 > (a+b+c+d)^2 + a^2 + b^2 + c^2 + d^2 + (a+c)(b+d).$$

Thus

$$B \geq \frac{4(a+b+c+d)^2}{(a+b+c+d)^2 + a^2 + b^2 + c^2 + d^2 + (a+c)(b+d)} > \frac{4(a+b+c+d)^2}{2(a+b+c+d)^2} = 2.$$

Hence  $B > 2$ . ■