

Definitions look more alike through SNOMED CT than through code lists

Overlap between two definitions of one condition (Jaccard), through four vocabularies

● first definition ● second definition ● what they share, over everything either includes

●● 0: nothing shared ●● 0.33: a third ● 1: identical

	ICD-10 codes	phcodeX groups	SNOMED CT concepts	SNOMED CT + parents	pairs
Epilepsy					
any vs focal	●● 0.27	●● 0.25	●● 0.19	●● 0.19	3
any vs generalized	●● 0.09	●● 0.25	●● 0.38	●● 0.50	3
focal vs focal strict	● 1.00	● 1.00	● 1.00	● 1.00	6
focal vs generalized	●● 0.00	●● 0.00	●● 0.00	●● 0.07	9
Sleep apnoea					
any sleep disorder vs hospital	●● 0.06	●● 0.11	●● 0.06	●● 0.20	2
hospital vs + primary care	● 1.00	● 1.00	● 1.00	● 1.00	1
Insomnia					
F51.0 or G47.0 vs all of F51	●● 0.10	●● 0.17	●● 0.18	●● 0.36	1
Constipation					
K59.0 or laxatives vs all of K59	●● 0.12	●● 0.17	●● 0.07	●● 0.06	1
ADHD					
F90.0 vs all of F90	●● 0.20	● 1.00	●● 0.50	● 0.80	1
Intellectual disability					
mild (F70) vs any (F7)	●● 0.17	●● 0.20	●● 0.14	●● 0.35	1
Bladder					
neurogenic (N31) vs other (N32)	●● 0.00	●● 0.00	●● 0.28	●● 0.29	1

READING THIS

Mean overlap over 29 pairs: ICD-10 0.30, phcodeX 0.35, SNOMED CT 0.34, with parents 0.41.
Adding SNOMED CT parents raises the overlap above ICD-10 in 18 of 29 pairs and lowers it in 4.
10 pairs share no ICD-10 code at all, yet overlap once SNOMED CT parents are added.