

The Phenotype Definition Problem: How EHR Case Definitions Change the Genetic Overlap Between Autism and Its Co-occurring Medical Conditions

Kehinde Obidele

Department of Health Informatics, Northeastern University

Abstract—Genetic studies of medical conditions increasingly rely on case definitions written as rules over electronic health records, and a single condition often has several. This study asked how much the genetic correlation between autism and its co-occurring medical conditions moves when only that definition changes. Using public FinnGen R12 summary statistics for 28 endpoints, including alternative definitions of epilepsy, sleep apnoea, insomnia, constipation and ADHD, together with GWAS of autism (18,381 cases) and four larger psychiatric traits, we estimated 217 genetic correlations with LD score regression and built a block-jackknife test for the difference between two correlations that share a trait, on one common set of 953,474 SNPs. Every definition was also mapped to SNOMED CT through the OMOP standard vocabularies. Nine of 135 comparisons between definitions of the same condition survived Bonferroni correction. Widening sleep apnoea to any sleep disorder raised the genetic correlation with every psychiatric trait, by 0.03 to 0.07, and all functional bowel disorders correlated 0.21 more strongly with PTSD than constipation defined by diagnosis or laxative purchase, while adding primary care records moved it by 0.013 at most. For autism, no definition effect survived correction, and constipation gave the same correlation under both definitions (0.11 and 0.12) when a shift of 0.21 was detectable. Shared cases made differences precise: for nested definitions the smallest detectable shift fell to 0.015. The SNOMED CT mapping revealed that the constipation comparison was also a comparison of clinical scope, and distance in SNOMED CT tracked how far correlations moved (Spearman 0.49) better than overlap in codes and data sources did (0.13). How a condition is defined in the health record can change what genetics says about it, and the size of that change can be measured.

Index Terms—autism spectrum disorder, electronic health records, phenotype definition, genetic correlation, LD score regression, OMOP common data model, SNOMED CT, FinnGen, sleep, constipation, epilepsy

I. INTRODUCTION

Autism rarely travels alone. Epilepsy is found in a median of 12.1% of autistic people across 74 studies [1], autistic children have almost four times the odds of constipation of comparison children [2], most experience sleep difficulties [3], and incontinence is more common than in typically developing children [4]. For families these conditions are part of daily care, and earlier work on predicting toileting events for an autistic adult showed how much of that care revolves around the bladder and bowel [5]. Twin and genome-wide studies suggest that part of this co-occurrence is shared genetic liability: in Swedish twins, clinical autism had genetic correlations of

0.50 with epilepsy and 0.31 with constipation [6], and genome-wide data give a genetic correlation of 0.27 between autism and irritable bowel syndrome [7].

In biobank and health record studies, however, a condition is whatever its definition says it is. “Constipation” may mean a diagnosis code in a hospital record, a laxative purchase, or a wider group of functional bowel codes, and each rule captures different people. How much this matters shows up plainly in autism research: across 144 studies, the reported prevalence of constipation in autism ranged from 4.3% to 45.5%, and gastrointestinal symptoms differed significantly by how they were ascertained [8]. Phenotype libraries and electronic phenotyping methods exist to make definitions explicit and reusable [9]–[11], yet one biobank can still publish several definitions of the same condition side by side.

The genetics of a condition can change with its definition. Depression defined by a few self-reported items has lower heritability and less specific genetic signal than strictly defined major depression [12]; SNP heritability of bipolar disorder identified from health records ranged from 0.00 to 0.24 across four algorithms [13]; and definitions of gout in the UK Biobank differed in heritability and in the number of associated variants they recovered [14]. We found no study that asked the same question of autism’s co-occurring medical conditions. This study measured how far the genetic correlation between autism, four other psychiatric traits and a set of medical conditions moves when only the case definition of the medical condition changes, and whether the answer is large enough to matter.

The study makes four contributions (Fig. 1). It takes 28 FinnGen definitions apart into their codes, data sources and rules (Fig. 2), and maps them to SNOMED CT through the OMOP standard vocabularies. It builds a difference test for two genetic correlations that share a trait, using LD score regression’s own block jackknife on shared genome blocks, so that the correlation between the two estimates is part of the standard error. It reports, for every comparison, the smallest shift the data could have detected, so that a null result can be read as well powered or not. And it checks the method against a known definition effect, EHR against clinically defined depression.




Fig. 1. Study design, walked through once with real numbers. (1) The three FinnGen R12 sleep apnoea definitions as bars of cases, ochre for the first definition and blue for the alternatives. (2) Each definition meets five psychiatric GWAS in LD score regression. (3) The genetic correlation with depression moves from 0.353 under hospital records to 0.420 under any sleep disorder (common SNP set). (4) The shift of 0.068 is set against its noise, ± 1.96 SE of the difference from the shared-block jackknife (0.014), the sleeve of Figs. 3, 15, 16 and 17. The strip lists the checks that run beside the design. In every figure, the sentences of the reading panel are computed from the results.

II. RELATED WORK

A. Shared Genetics of Autism and Co-occurring Conditions

Co-occurring psychiatric conditions in autism are well described, with pooled prevalence of 13% for sleep-wake disorders and 28% for ADHD [15]. Genetic studies have mostly examined psychiatric and cognitive traits: a multi-variate analysis of autism with eight co-occurring traits, all psychiatric or cognitive, identified shared loci and causal links [16]. Medical conditions have been studied less. The Swedish twin study found shared genetic effects between autism and epilepsy and gastrointestinal disorders across levels of symptom severity, but noted that without primary care records it probably captured only the more severe cases [6], which is itself a statement about definitions. Genome-wide work on functional somatic syndromes found positive genetic correlations of autism with irritable bowel syndrome, pain and fatigue [7]. Each of these studies used one definition per condition.

B. Phenotype Definition and Genetic Signal

Studies that vary a definition deliberately have focused on the condition being defined rather than on its genetic overlap with others. For depression, minimal phenotyping gave lower heritability and less specific loci than strict criteria [12], while 32 definitions built from diagnostic components in the UK Biobank had heritabilities from 0.102 to 0.162 and genetic correlations with external depression GWAS of 0.65 to 0.90, with no significant differences between the correlations [17]. EHR algorithms for bipolar disorder of different stringency correlated highly with conventionally ascertained cases [13], and gout definitions differed in heritability and power [14]. None of these tested whether a definition change moves the genetic correlation with a second trait while accounting for the dependence between the estimates.

C. EHR Phenotypes and Standard Vocabularies

Rule-based EHR phenotypes are built mainly from diagnosis codes, then medications and text [9], and platforms such as CALIBER curate validated definitions across several clinical terminologies [10]. The OMOP common data model and its standard vocabularies let one definition be expressed once and run across health systems [18], [19]. Phecodes group ICD codes into research phenotypes and have been mapped to WHO ICD-10 [20], most recently as phecodeX [21]. This study uses both to compare definitions, and uses SNOMED CT as a measure of how different two definitions are in clinical meaning.

D. Estimating Genetic Correlation

LD score regression estimates heritability and genetic correlation from summary statistics and separates confounding and sample overlap into its intercept [22], [23], and resources such as LD Hub made it routine to screen many traits [24]. Its estimates are less precise than those from individual-level methods, and less precise again when the reference LD panel does not match the study population [25]. Genomic structural equation modelling uses the full sampling covariance of LD score regression estimates across traits [26]; this study uses the same idea, in its simplest form, to test differences between pairs of correlations.

III. METHODS

A. Study Design

This study asked whether the genetic correlation between a psychiatric trait and a medical condition changes when only the electronic health record (EHR) definition of that condition changes (Fig. 1). It combined public summary statistics from the FinnGen biobank, which often publishes several definitions of the same condition, with genome-wide association studies (GWAS) of five psychiatric traits. Autism was the focal trait. Schizophrenia, bipolar disorder, major depression and post-traumatic stress disorder (PTSD) were added because their far larger samples make smaller definition effects detectable, and depression defined from EHRs and by clinical assessment




Fig. 2. The anatomy of a definition. Each row is one alternative definition. Left: one tile per ICD-10 code used anywhere in that condition's definitions (labels read upward); an indigo tile counts the code in hospital or death records, and a teal corner marks a code also counted from primary care. Right, aligned across conditions: rules that codes cannot show, each drawn as a glyph (key at top): a drug purchase class counts (ATC A06A, laxatives); a drug reimbursement right counts; a case must not also meet another definition; the code must be the person's most frequent related diagnosis; the control rule differs from the first definition in the set. A faint dot means the rule does not apply. Tiles show ICD-10 only.

served as a positive control where a definition effect is already known [12] (Fig. 3).

B. Condition Definitions

Condition GWAS came from FinnGen release R12 (500,348 participants) [27]. The endpoint definitions were taken from the R12 core and non-core endpoint file, which matches data freeze DF12, and were checked against the DF12 endpoint and control definition files (SHA-256 checksums recorded). A FinnGen endpoint is a set of rules: regular expressions

over Finnish ICD-10, ICD-9 and ICD-8 codes in the hospital discharge and cause-of-death registers, optional codes from primary care (Avohilmo), drug purchase classes (ATC codes) and drug reimbursement rights, inclusion of other endpoints, exclusion conditions, a "mode" rule requiring the code to be the person's most frequent related diagnosis, and a separate rule for who may serve as a control.

This study grouped 28 endpoints. Eighteen formed six *definition sets*, alternative definitions of one condition whose case groups overlap or nest: epilepsy (any epilepsy; focal,




Fig. 3. Positive control: depression defined from health records casts a wider genetic net. One row per trait or FinnGen endpoint (31), sorted by the difference. Each arrow runs from r_g with clinically defined depression (hollow dot) to r_g with EHR-defined depression (arrowhead), so an arrow pointing right means the EHR definition correlates more strongly; heavier turmeric arrows mark differences with $p < 0.05$.

focal strict and focal mode; generalized, generalized strict and generalized mode), sleep apnoea (hospital records; hospital records plus primary care; any sleep disorder), insomnia (F51.0 or G47.0; all of F51 with stricter controls), constipation (the constipation code K59.0 or a laxative purchase; all functional intestinal disorders, K59), attention-deficit/hyperactivity disorder (F90.0; all of F90 with stricter controls) and intellectual disability (mild, F70; any, F7). Ten neighbouring conditions (for example irritable bowel syndrome, reflux and two bladder disorders) were analysed alongside for comparison. The grouping, and the choice of distance measure described below, were fixed in the study notebook before any genetic correlation was estimated. Two candidate groupings were rejected at that stage because their members shared no codes: the two bladder endpoints (N31 and N32) and the irritable bowel, dyspepsia and reflux endpoints are different



Fig. 4. What two definitions of one condition contain, and how much they share, counted in SNOMED CT concepts. Each square is one standard concept that either definition's ICD-10 codes map to, in the order only in the first definition (ochre), in both (split ochre and blue), only in the second (blue), with the shared count at the right. Widening sleep apnoea to any sleep disorder adds 16 concepts to the one they share, all of K59 adds 13 to constipation's one (named in part, including irritable bowel syndrome and functional diarrhoea), and focal and generalized epilepsy share none. Laxative purchases have no condition concept.

conditions, not definitions of one.

Each definition was taken apart by a script that expanded every code pattern against the WHO ICD-10 code list, followed inclusion chains, and recorded which data sources and rules could make a person a case (Fig. 2). Mild intellectual disability (913 cases) had a heritability estimate below zero and was excluded from all correlation analyses, which left five testable definition sets.

C. Mapping to Standard Vocabularies

To make definitions comparable across health systems, each definition's codes were mapped to the Observational Medical Outcomes Partnership (OMOP) standard vocabularies [18], [19], downloaded from Athena (WHO ICD-10 2021 release, ICD-9-CM, ATC 2026-02-01, SNOMED CT 2026 International edition, RxNorm 2026-06-01). ICD-10 codes were matched to OMOP ICD10 concepts and followed through the "Maps to" relationship to standard SNOMED CT concepts. Finnish ICD-9 codes, which add a letter to four digits, were matched to ICD-9-CM after removing the letter and are reported as approximate. ICD-8 has no OMOP vocabulary, so ICD-8 rules could not be mapped. ATC classes were linked to their substances through the concept ancestor table. The phe-



Fig. 5. Overlap between alternative definitions depends on the vocabulary. For each pair, the Jaccard index from ICD-10 codes, phecodeX groups, SNOMED CT concepts, and SNOMED CT concepts together with their ancestors within two levels, drawn as an eclipse: an ochre disc for the first definition and a blue disc for the second, placed so that the dark shared area, divided by the area both discs cover, equals the Jaccard index (key at top), with the value printed beside it. Pairs with identical values in all four vocabularies share a row, and the last column counts them.

codeX map [20], [21] was used as a second, research-oriented grouping. The overlap between two definitions was measured with the Jaccard index at four levels: ICD-10 codes, phecodes, SNOMED CT concepts, and SNOMED CT concepts together with their ancestors up to two levels. Fig. 4 lists the concepts behind three of the comparisons, and Fig. 5 gives all four overlaps for every pair.

D. Psychiatric GWAS

Autism summary statistics came from the iPSYCH-PGC meta-analysis (18,381 cases, 27,969 controls) [28]. The other psychiatric GWAS were the European-ancestry analyses of the Psychiatric Genomics Consortium for schizophrenia [29], bipolar disorder [30], major depression excluding 23andMe [31] and PTSD [32], and the MDD2025 subsets restricted to EHR or registry cohorts (311,831 cases, 1,180,439 controls) and to clinically assessed cohorts (26,778 cases, 51,857 controls) [31]. Per-SNP effective sample sizes were used for all PGC analyses. For every file, the reported frequency of the effect allele was compared with the 1000 Genomes European frequency of the same allele [33]; correlations ranged from 0.991 to 0.998 over more than 1.1

million non-palindromic SNPs, confirming that effect alleles were labelled correctly.

The cohort lists showed three sources of sample overlap with FinnGen: the full depression GWAS and its EHR subset both include FinnGen, and the PTSD meta-analysis lists FinnGen among its biobanks. The EHR depression subset also contains iPSYCH, the cohort behind the autism GWAS. Cross-trait LD score regression absorbs such overlap in its intercept [23], and the intercepts were inspected for exactly this pattern.

E. LD Score Regression

SNP heritability and genetic correlation were estimated with LD score regression [22], [23] using a maintained Python 3 port of the original software and the European LD scores computed from 1000 Genomes [33]. FinnGen files were streamed, restricted to HapMap3 SNPs with minor allele frequency of at least 0.01, and never stored; about 1.17 million SNPs remained per endpoint. Heritability was converted to the liability scale [34] (Fig. 6). For FinnGen endpoints the population prevalence was approximated by the sample prevalence, so heritability is compared within a condition rather than across conditions; genetic correlations do not depend on this choice.




Fig. 6. Does a wider definition lower heritability? One panel per condition: each definition at its number of cases (log scale, each panel its own range) and its liability-scale SNP heritability, using the sample prevalence as the population prevalence. Ochre, the condition's first definition; blue, the alternatives; dot, estimate; bar, 50% interval; pale sleeve, 95% interval cut at zero; hollow dot, an estimate below zero. All panels share the heritability axis.

Before any new estimate was trusted, the pipeline reproduced published values: autism liability-scale heritability 0.117 (SE 0.010) against 0.118 (SE 0.010) in the source paper [28], and the genetic correlation between autism and ADHD [35] of 0.346 (SE 0.051) against 0.360 (Fig. 7). During this work a defect was found in the Python 3 port: requesting liability-scale output from a genetic correlation run raised an error after the estimate had been printed. Genetic correlations were therefore run without prevalence arguments, which leaves the estimate unchanged.

F. Testing Whether the Definition Moves the Genetic Correlation

Two genetic correlations that share a psychiatric trait and describe overlapping case groups have correlated sampling errors, and treating them as independent would overstate the uncertainty of their difference. LD score regression estimates standard errors with a block jackknife over 200 contiguous blocks of SNPs [36]. All 35 traits were first restricted to one common set of 953,474 SNPs, and every pair was re-estimated with evenly spaced blocks, so that block k covers the same

stretch of genome in every analysis. From the delete values of the genetic covariance and the two heritabilities, the jackknife pseudovalue of the genetic correlation in block k is

$$\tilde{r}_k = n \hat{r}_g - (n-1) \frac{\hat{\rho}_{g,(k)}}{\sqrt{\hat{h}_{1,(k)}^2 \hat{h}_{2,(k)}^2}}, \quad (1)$$

with $n = 200$, exactly as the software computes it. For two definitions a and b , the standard error of the difference $\hat{r}_a - \hat{r}_b$ is $\sqrt{\text{var}(\tilde{r}_{a,k} - \tilde{r}_{b,k})/n}$, which includes the covariance between the two estimates; Fig. 8 shows these per-block values for three pairs of definitions. The rebuilt standard error of every single genetic correlation matched the software's own printed value (198 of 198). The smallest difference detectable at a two-sided α of 0.05 with 80% power is 2.80 times this standard error, so the correlation between the two estimates' errors, not sample size alone, sets how small a shift can be seen (Fig. 9).

Within each definition set and psychiatric trait, heterogeneity across all definitions was tested with Cochran's $Q = \hat{\mathbf{r}}^\top W \hat{\mathbf{r}} - (\mathbf{1}^\top W \hat{\mathbf{r}})^2 / (\mathbf{1}^\top W \mathbf{1})$, where W is the inverse of the




Fig. 7. Validation. (a) Autism SNP heritability on the liability scale and the genetic correlation between autism and ADHD, drawn as this pipeline's estimate minus the published value; the pale sleeve is the published 95% interval or, where the paper prints no standard error, the tolerance of ± 0.05 fixed in the analysis plan. (b) Genetic correlation between the iPSYCH-PGC autism GWAS and the two FinnGen autism endpoints, each drawn as a quantile dot plot of its sampling distribution (20 dots, each one twentieth of the probability); the dashed line marks $r_g = 1$, the value if both measured the same genetics.

jackknife covariance matrix of the definitions' estimates, on $k - 1$ degrees of freedom (Fig. 10). The relation between how different two definitions are and how far their genetic correlations move was examined with Spearman correlations against the pre-specified distance (one minus the mean of ICD-10 Jaccard, data-source Jaccard and an indicator that no rule differs) and, as an exploratory analysis, against SNOMED CT distance (Fig. 11). Because comparisons share traits and definitions, these correlations are descriptive.

G. Finnish Linkage Disequilibrium

The European LD reference does not match the Finnish population exactly [37]. As a sensitivity check, LD scores were rebuilt for chromosomes 21 and 22 from the public FinnGen R12 LD matrix (520,210 samples; all variant pairs within 3 Mb with $r^2 > 0.01$), summing r^2 over HapMap3 SNPs within

1 Mb, and compared SNP by SNP with the European scores (Fig. 12).

IV. RESULTS

A. What the Definitions Contain

Alternative definitions of the same condition differed in several ways at once (Fig. 2). Some shared every ICD-10 code and differed only in their rules: focal, focal strict and focal mode epilepsy all use G40.0 to G40.2, but the strict version excludes anyone who also meets the generalized definition and the mode version requires focal codes to be the person's most frequent epilepsy diagnosis. Others differed in their data sources: the two sleep apnoea definitions share one code (G47.3), and the broader one adds primary care records. The constipation definition counts a laxative purchase as a case, which no code-based view can show.

All 153 ICD-10 codes used across the 28 definitions mapped to standard SNOMED CT concepts (99 distinct concepts). Sixteen of the 28 definitions also contain ICD-8 rules, which have no OMOP counterpart. How much two definitions overlapped depended on the vocabulary used to compare them (Fig. 5). Attention-deficit/hyperactivity disorder defined as F90.0 or as all of F90 overlapped 0.20 by ICD-10 codes, 1.00 by phecodes, 0.50 by SNOMED CT concepts and 0.80 once SNOMED CT ancestors were included. Any epilepsy and generalized epilepsy overlapped 0.09 by codes but 0.50 in the SNOMED CT hierarchy. The two bladder endpoints shared no codes and no phecodes but 28% of their SNOMED CT concepts.

Reading the definitions through SNOMED CT also caught a labelling error in this study's own work (Fig. 4). The FinnGen endpoint first treated here as a diagnosis-only definition of constipation (K59) maps to 14 SNOMED CT concepts: constipation and its subtypes, but also irritable bowel syndrome, functional diarrhoea, neurogenic bowel, acquired megacolon and post-surgical conditions. FinnGen's own name for it is "other functional intestinal disorders". It is a broader bowel dysfunction definition that contains constipation, and it is described as such throughout. In the same way, insomnia defined by F51.0 or G47.0 maps to two concepts, while all of F51 adds sleepwalking, sleep terrors, nightmares, non-organic hypersomnia and sleep-wake schedule disorders (11 concepts).

B. Validation, Heritability and Genetic Correlations

The pipeline reproduced the published autism estimates before any new result was produced (Fig. 7). Autism in the Danish GWAS also correlated strongly with autism in the Finnish registry: $r_g = 0.73$ (SE 0.15) with the narrow FinnGen definition (F84.0, F84.5; 888 cases) and 0.98 (SE 0.21) with the broader one (F84; 1,179 cases).

Liability-scale heritability varied within conditions (Fig. 6). Generalized epilepsy (0.135, SE 0.032) was more heritable than focal epilepsy (0.026, SE 0.009), and the broad epilepsy definition fell between them (0.055, SE 0.009). ADHD with all of F90 and stricter controls (0.127, SE 0.019) exceeded F90.0 alone (0.092, SE 0.016). Adding primary care to sleep apnoea left heritability unchanged (0.125 and 0.120). Among




Fig. 8. Why a difference between two genetic correlations can be precise. Top of each panel: the 200 genome blocks of the jackknife, each at its pair of pseudovalues of r_g under the two definitions, centered on the estimates, against the line of equal values. Bottom: every block's difference as a tick, with its density; the colored density keeps the blocks paired as they are, and the gray one pairs each block with a random other block, as if the two estimates shared no people. Each carries its standard error. (a) Depression with sleep apnoea against any sleep disorder, nested definitions; (b) PTSD with constipation or laxatives against all of K59, partly shared cases; (c) PTSD with EHR against clinical depression, almost no shared samples.

the psychiatric traits, clinically defined depression had twice the heritability of EHR-defined depression (0.114, SE 0.012, against 0.057, SE 0.002, at the same population prevalence), the direction reported for strict versus minimal definitions of depression [12]. Mild intellectual disability (913 cases) had an estimate below zero.

Of the 217 genetic correlations estimated, 156 had $p < 0.05$ (Fig. 13). The genetic correlation between schizophrenia and bipolar disorder was 0.68 (SE 0.02). Autism correlated with both FinnGen ADHD definitions (0.53, SE 0.11; 0.55, SE 0.08), irritable bowel syndrome (0.22, SE 0.07), nonorganic sleep disorders (0.31, SE 0.09) and sleep apnoea (0.12, SE 0.05), but not with any epilepsy definition (estimates from -0.19 to -0.02). The cross-trait intercepts behaved as expected from the cohort lists (Fig. 14): median 0.001 to 0.004 for autism, schizophrenia, bipolar disorder and clinical

depression, which contain no FinnGen samples, against 0.012 for PTSD and 0.020 for both depression GWAS that include FinnGen.

C. Does the Definition Move the Genetic Correlation?

All 198 estimable correlations were re-estimated on the common SNP set with shared jackknife blocks; they correlated 0.991 with the standard estimates (mean absolute difference 0.020). Across 135 comparisons between definitions of the same condition, 40 differences had $p < 0.05$ and nine survived Bonferroni correction ($p < 0.00037$). Fig. 15 shows the shift for six representative changes of definition, Fig. 16 sets every shift against its own noise, and Table 1 and Fig. 10 summarise heterogeneity across all definitions in each set.

Sleep apnoea. Adding primary care records barely moved the genetic correlation: the largest difference was 0.013 (SE



Fig. 9. What buys power: shared cases. The shaded map gives the smallest difference in r_g detectable at 80% power for two estimates with equal standard error (vertical axis, log scale) and a given correlation between their sampling errors (horizontal axis); each white contour, from 0.02 to 1, is drawn exactly from the formula. Dots, colored by psychiatric trait, are the 135 comparisons between definitions of the same condition, at their error correlation and the root mean square of their two standard errors; rings are the 31 EHR against clinical depression comparisons. Four comparisons are numbered and named below the map.

0.006, $p = 0.04$, PTSD), and the smallest detectable differences for this pair were 0.015 to 0.029. Widening the definition from sleep apnoea to any sleep disorder raised the genetic correlation with every psychiatric trait: by 0.068 (SE 0.007) for depression, 0.064 (SE 0.009) for bipolar disorder, 0.063 (SE 0.008) for PTSD, 0.050 (SE 0.008) for schizophrenia and 0.027 (SE 0.013, $p = 0.042$) for autism. These were the most precise comparisons in the study because the two definitions share most of their cases: their sampling errors correlated 0.92 to 0.97 (Fig. 8).

Insomnia. Nonorganic sleep disorders (all of F51, stricter controls) gave higher genetic correlations than insomnia (F51.0 or G47.0) for all five traits: by 0.139 (SE 0.050, $p = 0.005$) for depression, 0.134 (SE 0.048, $p = 0.005$) for bipolar disorder, 0.125 (SE 0.050, $p = 0.013$) for PTSD,




Fig. 10. Heterogeneity across definitions, with each p -value drawn as the tail it is. Rows, definition sets; columns, psychiatric traits. The curve is the χ^2 distribution that Cochran's Q would follow if the definitions made no difference; the tick is the observed Q , and the tinted tail beyond it is the p -value. A tick in the trait's color with a dot marks $p < 0.05$; an arrow marks a Q beyond the drawn range.

0.090 (SE 0.041, $p = 0.029$) for schizophrenia and 0.093 ($p = 0.079$) for autism. The two definitions differ in scope (parasomnias and hypersomnia added) and in their control rule at the same time, and this design cannot separate the two.

Constipation. The broader functional intestinal disorders definition gave a higher genetic correlation than the constipation code or laxative definition for PTSD (0.207, SE 0.055, $p = 0.0002$) and schizophrenia (0.113, SE 0.046, $p = 0.015$), but not for autism (0.010, $p = 0.90$). The error correlation for this pair was only 0.39 to 0.55, because the laxative rule adds many people the diagnosis-based definition does not contain, so the smallest detectable differences were 0.13 to 0.21.

Epilepsy and ADHD. Genetic correlations differed across the seven epilepsy definitions for depression ($Q = 21.0$, df 6, $p = 0.002$), PTSD ($p = 0.003$) and bipolar disorder




Fig. 11. Do definitions that differ more give genetic correlations that differ more? Each point is one comparison, in the color of its psychiatric trait (filled: sleep, bowel and ADHD; hollow: epilepsy), spread sideways by up to 0.012 so identical distances separate; dark bars are medians and boxes the middle half within each cluster of five or more comparisons at the same distance. (a) Pre-specified distance from codes, data sources and rules. (b) SNOMED CT distance (exploratory). The strip above each panel places the Spearman correlation for each psychiatric trait (dots) and for all 135 comparisons (bar); the correlations are descriptive because comparisons share traits and definitions.

($p = 0.012$). The differences ran between focal and generalized epilepsy, not between strict, mode and plain versions of the same subtype. Both ADHD definitions gave the same correlations with every trait (largest difference 0.036), with smallest detectable differences of 0.07 to 0.12.

Across all 135 comparisons, the smallest detectable difference ranged from 0.015 to 1.10, and it was governed mainly




Fig. 12. The LD landscape of chromosomes 21 and 22 through two populations. (a, b) Mean LD score in 250 kb windows along each chromosome, European (1000 Genomes) above the line and Finnish (FinnGen R12) mirrored below it. (c) The per-SNP ratio of Finnish to European LD score as a quantile dot plot (each dot 2% of SNPs) on a log axis; blue line, median.

by how strongly the two estimates' errors correlated (Fig. 9).

D. Autism

For autism, no difference between definitions survived multiple-testing correction (Fig. 17). Three were nominally significant: widening sleep apnoea to any sleep disorder, from hospital records (0.027, $p = 0.042$) and from hospital plus primary care records (0.040, $p = 0.017$), in the same direction as for every other trait, and any epilepsy against focal epilepsy under the mode rule (0.197, $p = 0.023$). The design could see a shift of 0.03 to 0.05 for the sleep apnoea definitions, 0.15 for insomnia, 0.21 for constipation and 0.12 for ADHD, but between 0.11 and 0.81 for epilepsy. The constipation result




Fig. 13. Genetic correlations between seven psychiatric GWAS and 28 FinnGen endpoints. (a) For each condition with alternative definitions and each of five psychiatric traits, the median r_g across its definitions (dot) and the range from the lowest to the highest definition (line). (b) The full matrix, colored with a value-suppressing uncertainty palette [38]; hue gives r_g and depth of color its precision, so estimates with large standard errors fade toward gray and merge into fewer color steps (fan key under a).

is the clearest statement for autism: whether the definition is narrow constipation or all functional bowel disorders, the genetic correlation with autism is the same (0.11 and 0.12).

E. Positive Control: EHR and Clinical Depression

EHR-defined and clinically defined depression correlated at $r_g = 0.857$ (SE 0.045). Against the other traits and endpoints, EHR depression gave the higher genetic correlation in 25 of 31 comparisons (Fig. 3); six differences had $p < 0.05$ and none survived Bonferroni correction. The largest were with

PTSD (0.157, SE 0.052, $p = 0.003$) and any epilepsy (0.208, $p = 0.017$). Because the two depression GWAS share almost no samples (intercept 0.013), their errors were only weakly correlated, and the smallest detectable differences were 0.11 to 1.19.

F. Distance Between Definitions

The pre-specified definition distance, built from shared codes, data sources and rules, did not predict how far the genetic correlation moved (Spearman $\rho = 0.13$ across all com-




Fig. 14. Cross-trait LD score regression intercepts with the 28 FinnGen endpoints, one dot per endpoint in a beeswarm, grouped by whether the psychiatric GWAS contains FinnGen samples; dark tick, median. The pale band spans ± 1.96 times the median intercept standard error around zero, where a GWAS with no shared samples should fall. Shared samples appear in the intercept, which keeps r_g unbiased.

TABLE I
HETEROGENEITY OF GENETIC CORRELATION ACROSS DEFINITIONS OF THE SAME CONDITION

Set (definitions)	ASD	SCZ	BIP	MDD	PTSD
Sleep apnoea (3)	5.8	40.6*	51.7*	87.6*	69.5*
Insomnia (2)	3.1	4.8 [†]	7.7 [†]	7.7 [†]	6.2 [†]
Constipation (2)	0.0	5.9 [†]	1.2	2.8	14.0*
ADHD (2)	0.1	2.2	0.5	0.0	0.0
Epilepsy (7)	6.1	8.0	16.4 [†]	21.0 [†]	19.8 [†]

Cochran's Q with the jackknife covariance, $df = \text{definitions} - 1$.

[†] $p < 0.05$; * $p < 0.001$.

parisons; Fig. 11). Distance in SNOMED CT did ($\rho = 0.49$; 0.84 for schizophrenia, 0.66 for bipolar disorder, 0.54 for autism), an exploratory result carried mostly by the epilepsy comparisons, which make up 105 of the 135.

G. Finnish Linkage Disequilibrium

On chromosomes 21 and 22, Finnish LD scores built from the FinnGen LD matrix correlated 0.970 with the European reference across 33,436 HapMap3 SNPs, with a median Finnish-to-European ratio of 1.009 (Fig. 12).

V. DISCUSSION

A. What a Definition Changes, and What It Does Not

The clearest pattern in these results is that the scope of a definition moved the genetic correlation, while the source of the records and the strictness of the rules mostly did not. Adding primary care records to sleep apnoea added nearly 6,000 cases and changed no correlation by more than 0.013, a null that the design could see down to 0.015. Strict and mode versions of focal and generalized epilepsy agreed closely with their plain versions. By contrast, widening sleep apnoea to any sleep disorder, which brings in insomnia, narcolepsy and other sleep diagnoses, raised the correlation with every psychiatric trait; moving from insomnia to all nonorganic sleep disorders raised it for all five; and moving from constipation to all functional bowel disorders raised it for PTSD and schizophrenia. In each case the wider definition added conditions that themselves carry psychiatric liability. Epilepsy followed the same logic from the other side: focal and generalized epilepsy, which have partly distinct genetic bases [39], gave different correlations, and the broad definition fell between them.

This is why SNOMED CT distance tracked the size of the shift better than the pre-specified distance built from codes and data sources. The pre-specified measure counted differences, such as a primary care source or a stricter control rule, that turned out not to matter, and treated a subtype change as no larger than those. Distance in clinical meaning is a better guide to whether two definitions will give different genetics, although this finding is exploratory and rests mostly on the epilepsy comparisons.

The SNOMED CT view was also how this study caught its own mistake. A definition first treated as diagnosis-only constipation turned out to be all functional intestinal disorders, including irritable bowel syndrome and functional diarrhoea. Codes that read as a single condition can hide a much wider clinical scope, and mapping a definition to a standard vocabulary before using it is a cheap way to see that.

B. Autism

For autism the answer was reassuring rather than dramatic. No definition effect survived correction. The genetic correlation between autism and bowel dysfunction was the same whether the net was narrow constipation or all functional bowel disorders (0.11 and 0.12), with a detectable shift of 0.21, and the sleep apnoea definitions showed a small shift (0.027) in the same direction as every other trait. For epilepsy, autism's genetic correlations were close to zero under every definition, and the design could only have detected shifts of 0.11 to 0.81. This near-zero estimate differs from the twin genetic correlation of 0.50 [6]; the two measure different things, since twin estimates include all additive genetic variation, rare variants among it, while LD score regression captures only what common SNPs tag, and the Swedish study defined autism clinically in children. The autism GWAS is also the smallest of the five psychiatric GWAS here, and several autism comparisons would need larger samples to resolve.




Fig. 15. What moves the genetics when a definition changes. Rows are six changes to a FinnGen definition, grouped by the kind of change (data source, scope, strictness of a rule); columns are the five psychiatric GWAS. In each cell the gray sleeve spans ± 1.96 SE of the difference, centered on zero, and the stem is the shift in r_g from the first to the second definition, in the trait's color, with its value underneath. A filled dot has $p < 0.05$; a ringed dot also survives Bonferroni correction over 135 comparisons.

C. The Positive Control

The EHR and clinical depression comparisons behaved as a known definition effect should, in the direction reported before [12]: clinical depression had twice the heritability, the two correlated at 0.86 rather than 1, and EHR depression gave the higher correlation with 25 of 31 other traits and endpoints. Individually, few of these differences were significant, because the clinical GWAS is small and shares almost no samples with the EHR GWAS, so their errors do not cancel. The contrast with the sleep apnoea definitions, whose errors correlated above 0.9, shows what a within-biobank design gains: definitions measured on the same people can be compared far more precisely than definitions measured on different people.

D. Practical Implications

Three recommendations follow for genetic studies that use EHR definitions. First, report each definition in a standard vocabulary alongside its local codes, because scope, not source, is what changes the genetics. Second, when a biobank offers several definitions of the same condition, compare them with a test that accounts for their shared cases before choosing one, and report the smallest shift that test could detect. Third, adding primary care records to a hospital-based definition, at least for sleep apnoea, gains cases without changing the genetic correlation.

E. Limitations

This study used public summary statistics, so the number of people shared by two definitions was not known directly; the jackknife error correlation captures its statistical effect but not the counts. Several definition pairs differ in more than one way at once: the insomnia pair in scope and in its control rule, the constipation pair in its laxative rule and in scope. The design cannot say which difference drives the shift. FinnGen is a Finnish cohort analysed here with a European LD reference. On chromosomes 21 and 22, Finnish LD scores from 520,210 FinnGen samples correlated 0.970 with the European scores, but reference mismatch reduces the precision of LD score regression [25], the Finnish population carries a distinct founder history [37], and a genome-wide Finnish reference was not built. Three psychiatric GWAS share samples with FinnGen, and the EHR depression GWAS shares samples with the autism GWAS; the intercept absorbs this, as the intercepts showed, but the sources are listed so readers can judge. Liability-scale heritability used the sample prevalence as the population prevalence, so heritability is compared only within a condition. Register diagnoses are imperfect, with positive predictive values of 75% to 99% for common diagnoses in the Finnish Hospital Discharge Register [40]. ICD-8 rules could not be mapped to OMOP and Finnish ICD-9 was mapped approximately. Mild intellectual disability had too few cases




Fig. 16. Every definition shift inside its own noise. One mark per comparison (166): the gray sleeve spans ± 1.96 SE of the difference, and the stem is the shift in r_g from the first definition to the second (for depression, EHR minus clinical), in the color of its psychiatric trait, drawn thick with an end dot when it escapes its sleeve ($p < 0.05$) and thin otherwise. Families are drawn at different widths per comparison so small families stay legible; sleep apnoea has its own narrower scale, sleeves wider than the axis are cut, and the count of escapes is printed above each panel.

for an estimate. The difference test used the one-step estimator, whose standard errors were a median 8% to 16% larger than the standard two-step estimator for the larger GWAS, which makes it conservative. Only six conditions were tested, epilepsy dominates the distance analysis, and the correlations between distance and shift are descriptive. Finally, all GWAS were of European ancestry, and none of these results can be assumed to hold in other populations.

VI. CONCLUSION

How a condition is defined in the health record changes what genetics says about it, and here is by how much, for the conditions that matter most to autistic people and their families. Widening a definition to take in related conditions shifted genetic correlations with psychiatric traits by 0.03 to 0.21; adding a data source or tightening a rule did not. For autism, the genetic overlap with bowel dysfunction and sleep apnoea held steady across definitions within the precision these data allow, while epilepsy needs larger samples. A block-jackknife test on shared genome blocks, a statement of the




Fig. 17. Autism in focus: how large a definition effect this study could see, and how large it was. One row per alternative definition, compared with the first definition of its condition. Stem, the observed shift in r_g with iPSYCH-PGC autism; dark sleeve, ± 1.96 SE of the difference; pale sleeve, the smallest shift detectable at 80% power. A filled dot marks $p < 0.05$; hollow dots do not reach it. The numbers at right are the shift and the detectable shift; sleeves are cut at 0.6.

smallest detectable shift, and a map of each definition into a standard vocabulary are enough to make this question routine rather than invisible.

DATA AND CODE AVAILABILITY

FinnGen R12 summary statistics and endpoint definition files are publicly available from FinnGen (https://www.finnngen.fi/en/access_results; <https://www.finnngen.fi/en/researchers/clinical-endpoints>). Psychiatric GWAS summary statistics are available from the Psychiatric Genomics Consortium (<https://www.med.unc.edu/pgc/download-results/>) and, for MDD2025, from figshare (<https://doi.org/10.6084/m9.figshare.27061255>). European LD scores were obtained from Zenodo (<https://doi.org/10.5281/zenodo.8182036>). OMOP standard vocabularies are available from Athena (<https://athena.ohdsi.org>) under their own licences and are not redistributed. LD score regression was run with the Python 3 port of LDSC maintained by CBIIT (commit 1f09cf0).

REFERENCES

- [1] S. Lukmanji, S. A. Manji, S. Kadhim, K. M. Sauro, E. C. Wirrell, C.-S. Kwon, and N. Jetté, "The co-occurrence of epilepsy and autism: A systematic review," *Epilepsy & Behavior*, vol. 98, pp. 238–248, 2019. [Online]. Available: <https://doi.org/10.1016/j.yebeh.2019.07.037>
- [2] B. O. McElhanon, C. McCracken, S. Karpen, and W. G. Sharp, "Gastrointestinal Symptoms in Autism Spectrum Disorder: A Meta-analysis," *Pediatrics*, vol. 133, no. 5, pp. 872–883, 2014. [Online]. Available: <https://doi.org/10.1542/peds.2013-3995>
- [3] A. L. Richdale and K. A. Schreck, "Sleep problems in autism spectrum disorders: Prevalence, nature, & possible biopsychosocial aetiologies," *Sleep Medicine Reviews*, vol. 13, no. 6, pp. 403–411, 2009. [Online]. Available: <https://doi.org/10.1016/j.smrv.2009.02.003>
- [4] J. Niemczyk, C. Wagner, and A. von Gontard, "Incontinence in autism spectrum disorder: a systematic review," *European Child & Adolescent Psychiatry*, vol. 27, no. 12, pp. 1523–1537, 2018. [Online]. Available: <https://doi.org/10.1007/s00787-017-1062-3>
- [5] K. Obidele, "Personalized Prediction of Toileting Events for Caregivers of Individuals with Severe Autism: A Longitudinal Machine Learning Study," Stanford Digital Repository, 2026. [Online]. Available: <https://doi.org/10.25740/yq415zk9908>
- [6] P.-Y. Pan, M. J. Taylor, H. Larsson, C. Almqvist, P. Lichtenstein, S. Lundström, and S. Bölte, "Genetic and environmental contributions to co-occurring physical health conditions in autism spectrum condition and attention-deficit/hyperactivity disorder," *Molecular Autism*, vol. 14, no. 1, 2023. [Online]. Available: <https://doi.org/10.1186/s13229-023-00548-3>
- [7] Y. Li, T. Xie, M. Vos, H. Snieder, and C. A. Hartman, "Shared genetic architecture and causality between autism spectrum disorder and irritable bowel syndrome, multisite pain, and fatigue," *Translational Psychiatry*, vol. 14, no. 1, 2024. [Online]. Available: <https://doi.org/10.1038/s41398-024-03184-4>
- [8] C. Hologue, C. Newill, L. Lee, P. J. Pasricha, and M. Daniele Fallin, "Gastrointestinal symptoms in autism spectrum disorder: A review of the literature on ascertainment and prevalence," *Autism Research*, vol. 11, no. 1, pp. 24–36, 2018. [Online]. Available: <https://doi.org/10.1002/aur.1854>
- [9] J. C. Kirby, P. Speltz, L. V. Rasmussen, M. Basford, O. Gottesman, P. L. Peissig, J. A. Pacheco, G. Tromp, J. Pathak, D. S. Carrell, S. B. Ellis, T. Lingren *et al.*, "PheKB: a catalog and workflow for creating electronic phenotype algorithms for transportability," *Journal of the American Medical Informatics Association*, vol. 23, no. 6, pp. 1046–1052, 2016. [Online]. Available: <https://doi.org/10.1093/jamia/ocv202>
- [10] S. Denaxas, A. Gonzalez-Izquierdo, K. Direk, N. K. Fitzpatrick, G. Fatemifar, A. Banerjee, R. J. B. Dobson, L. J. Howe, V. Kuan, R. T. Lumbers, L. Pasea, R. S. Patel *et al.*, "UK phenomics platform for developing and validating electronic health record phenotypes: CALIBER," *Journal of the American Medical Informatics Association*, vol. 26, no. 12, pp. 1545–1559, 2019. [Online]. Available: <https://doi.org/10.1093/jamia/ocz105>
- [11] J. M. Banda, M. Seneviratne, T. Hernandez-Boussard, and N. H. Shah, "Advances in Electronic Phenotyping: From Rule-Based Definitions to Machine Learning Models," *Annual Review of Biomedical Data Science*, vol. 1, no. 1, pp. 53–68, 2018. [Online]. Available: <https://doi.org/10.1146/annurev-biodatasci-080917-013315>
- [12] N. Cai, MDD Working Group of the Psychiatric Genomics Consortium, J. A. Revez, M. J. Adams, T. F. M. Andlauer, G. Breen, E. M. Byrne, T.-K. Clarke, A. J. Forstner, H. J. Grabe, S. P. Hamilton, D. F. Levinson *et al.*, "Minimal phenotyping yields genome-wide association signals of low specificity for major depression," *Nature Genetics*, vol. 52, no. 4, pp. 437–447, 2020. [Online]. Available: <https://doi.org/10.1038/s41588-020-0594-5>
- [13] C.-Y. Chen, P. H. Lee, V. M. Castro, J. Minnier, A. W. Charney, E. A. Stahl, D. M. Ruderfer, S. N. Murphy, V. Gainer, T. Cai, I. Jones, C. N. Pato *et al.*, "Genetic validation of bipolar disorder identified by automated phenotyping using electronic health records," *Translational Psychiatry*, vol. 8, no. 1, 2018. [Online]. Available: <https://doi.org/10.1038/s41398-018-0133-7>
- [14] M. Cadzow, T. R. Merriman, and N. Dalbeth, "Performance of gout definitions for genetic epidemiological studies: analysis of UK Biobank," *Arthritis Research & Therapy*, vol. 19, no. 1, 2017. [Online]. Available: <https://doi.org/10.1186/s13075-017-1390-1>
- [15] M.-C. Lai, C. Kassee, R. Besney, S. Bonato, L. Hull, W. Mandy, P. Szatmari, and S. H. Ameis, "Prevalence of co-occurring mental health diagnoses in the autism population: a systematic review and meta-analysis," *The Lancet Psychiatry*, vol. 6, no. 10, pp. 819–829, 2019. [Online]. Available: [https://doi.org/10.1016/s2215-0366\(19\)30289-5](https://doi.org/10.1016/s2215-0366(19)30289-5)
- [16] K. Salenius, N. Väljä, S. Thusberg, F. Iris, C. Ladd-Acosta, C. Roos, M. Nykter, A. Fasano, R. Autio, J. Lin, and the GEMMA study, "Exploring autism spectrum disorder and co-occurring trait associations to elucidate multivariate genetic mechanisms and insights," *BMC Psychiatry*, vol. 24, no. 1, 2024. [Online]. Available: <https://doi.org/10.1186/s12888-024-06392-w>
- [17] B. S. Jermy, K. P. Glanville, J. R. I. Coleman, C. M. Lewis, and E. Vassos, "Exploring the genetic heterogeneity in major depression across diagnostic criteria," *Molecular Psychiatry*, vol. 26, no. 12, pp. 7337–7345, 2021. [Online]. Available: <https://doi.org/10.1038/s41380-021-01231-w>
- [18] J. M. Overhage, P. B. Ryan, C. G. Reich, A. G. Hartzema, and P. E. Stang, "Validation of a common data model for active safety surveillance research," *Journal of the American Medical Informatics Association*, vol. 19, no. 1, pp. 54–60, 2012. [Online]. Available: <https://doi.org/10.1136/amiajnl-2011-000376>
- [19] G. Hripesak, J. D. Duke, N. H. Shah, C. G. Reich, V. Huser, M. J. Schuemie, M. A. Suchard, R. W. Park, I. C. K. Wong, P. R. Rijnbeek, J. van der Lei, N. Pratt *et al.*, "Observational Health Data Sciences and Informatics (OHDSI): Opportunities for Observational Researchers," in *Studies in Health Technology and Informatics*, 2015. [Online]. Available: <https://doi.org/10.3233/978-1-61499-564-7-574>
- [20] P. Wu, A. Gifford, X. Meng, X. Li, H. Campbell, T. Varley, J. Zhao, R. Carroll, L. Bastarache, J. C. Denny, E. Theodoratou, and W.-Q. Wei, "Mapping ICD-10 and ICD-10-CM Codes to Phecodes: Workflow Development and Initial Evaluation," *JMIR Medical Informatics*, vol. 7, no. 4, p. e14325, 2019. [Online]. Available: <https://doi.org/10.2196/14325>
- [21] M. M. Shuey, W. W. Stead, I. Aka, A. L. Barnado, J. A. Bastarache, E. Brokamp, M. Campbell, R. J. Carroll, J. A. Goldstein, A. Lewis, B. A. Malow, J. D. Mosley *et al.*, "Next-generation phenotyping: introducing phecodeX for enhanced discovery research in medical phenomics," *Bioinformatics*, vol. 39, no. 11, 2023. [Online]. Available: <https://doi.org/10.1093/bioinformatics/btad655>
- [22] B. K. Bulik-Sullivan, Schizophrenia Working Group of the Psychiatric Genomics Consortium, P.-R. Loh, H. K. Finucane, S. Ripke, J. Yang, N. Patterson, M. J. Daly, A. L. Price, and B. M. Neale, "LD Score regression distinguishes confounding from polygenicity in genome-wide association studies," *Nature Genetics*, vol. 47, no. 3, pp. 291–295, 2015. [Online]. Available: <https://doi.org/10.1038/ng.3211>
- [23] B. Bulik-Sullivan, ReproGen Consortium, Psychiatric Genomics Consortium, Genetic Consortium for Anorexia Nervosa of the Wellcome Trust Case Control Consortium 3, H. K. Finucane, V. Anttila, A. Gusev, F. R. Day, P.-R. Loh, L. Duncan, J. R. B. Perry, N. Patterson *et al.*, "An atlas of genetic correlations across human diseases and traits," *Nature Genetics*, vol. 47, no. 11, pp. 1236–1241, 2015. [Online]. Available: <https://doi.org/10.1038/ng.3406>
- [24] J. Zheng, A. M. Erzurumluoglu, B. L. Elsworth, J. P. Kemp, L. Howe, P. C. Haycock, G. Hemani, K. Tansey, C. Laurin, Early Genetics and Lifecourse Epidemiology (EAGLE) Eczema Consortium, B. S. Pourcain, N. M. Warrington *et al.*, "LD Hub: a centralized database and web interface to perform LD score regression that maximizes the potential of summary level GWAS data for SNP heritability and genetic correlation analysis," *Bioinformatics*, vol. 33, no. 2, pp. 272–279, 2017. [Online]. Available: <https://doi.org/10.1093/bioinformatics/btw613>
- [25] G. Ni, G. Moser, S. Ripke, B. M. Neale, A. Corvin, J. T. Walters, K.-H. Farh, P. A. Holmans, P. Lee, B. Bulik-Sullivan, D. A. Collier, H. Huang *et al.*, "Estimation of Genetic Correlation via Linkage Disequilibrium Score Regression and Genomic Restricted Maximum Likelihood," *The American Journal of Human Genetics*, vol. 102, no. 6, pp. 1185–1194, 2018. [Online]. Available: <https://doi.org/10.1016/j.ajhg.2018.03.021>
- [26] A. D. Grotzinger, M. Rhemtulla, R. de Vlaming, S. J. Ritchie, T. T. Mallard, W. D. Hill, H. F. Ip, R. E. Marioni, A. M. McIntosh, I. J. Deary, P. D. Koellinger, K. P. Harden *et al.*, "Genomic structural equation modelling provides insights into the multivariate genetic architecture of complex traits," *Nature Human Behaviour*, vol. 3, no. 5, pp. 513–525, 2019. [Online]. Available: <https://doi.org/10.1038/s41562-019-0566-x>
- [27] M. I. Kurki, J. Karjalainen, P. Palta, T. P. Sipilä, K. Kristiansson, K. M. Donner, M. P. Reeve, H. Laivuori, M. Aavikko, M. A.

- Kaunisto, A. Loukola, E. Lahtela *et al.*, “FinnGen provides genetic insights from a well-phenotyped isolated population,” *Nature*, vol. 613, no. 7944, pp. 508–518, 2023. [Online]. Available: <https://doi.org/10.1038/s41586-022-05473-8>
- [28] J. Grove, Autism Spectrum Disorder Working Group of the Psychiatric Genomics Consortium, BUPGEN, Major Depressive Disorder Working Group of the Psychiatric Genomics Consortium, 23andMe Research Team, S. Ripke, T. D. Als, M. Mattheisen, R. K. Walters, H. Won, J. Pallesen, E. Agerbo *et al.*, “Identification of common genetic risk variants for autism spectrum disorder,” *Nature Genetics*, vol. 51, no. 3, pp. 431–444, 2019. [Online]. Available: <https://doi.org/10.1038/s41588-019-0344-8>
- [29] V. Trubetskoy *et al.*, “Mapping genomic loci implicates genes and synaptic biology in schizophrenia,” *Nature*, vol. 604, pp. 502–508, 2022. [Online]. Available: <https://doi.org/10.1038/s41586-022-04434-5>
- [30] N. Mullins *et al.*, “Genome-wide association study of more than 40,000 bipolar disorder cases provides new insights into the underlying biology,” *Nature Genetics*, vol. 53, pp. 817–829, 2021. [Online]. Available: <https://doi.org/10.1038/s41588-021-00857-4>
- [31] M. J. Adams *et al.*, “Trans-ancestry genome-wide study of depression identifies 697 associations implicating cell types and pharmacotherapies,” *Cell*, vol. 188, pp. 640–652, 2025. [Online]. Available: <https://doi.org/10.1016/j.cell.2024.12.002>
- [32] C. M. Nievergelt *et al.*, “Genome-wide association analyses identify 95 risk loci and provide insights into the neurobiology of post-traumatic stress disorder,” *Nature Genetics*, vol. 56, pp. 792–808, 2024. [Online]. Available: <https://doi.org/10.1038/s41588-024-01707-9>
- [33] The 1000 Genomes Project Consortium, “A global reference for human genetic variation,” *Nature*, vol. 526, pp. 68–74, 2015. [Online]. Available: <https://doi.org/10.1038/nature15393>
- [34] S. Lee, N. Wray, M. Goddard, and P. Visscher, “Estimating Missing Heritability for Disease from Genome-wide Association Studies,” *The American Journal of Human Genetics*, vol. 88, no. 3, pp. 294–305, 2011. [Online]. Available: <https://doi.org/10.1016/j.ajhg.2011.02.002>
- [35] D. Demontis, ADHD Working Group of the Psychiatric Genomics Consortium (PGC), Early Lifecourse & Genetic Epidemiology (EAGLE) Consortium, 23andMe Research Team, R. K. Walters, J. Martin, M. Mattheisen, T. D. Als, E. Agerbo, G. Baldursson, R. Belliveau, J. Bybjerg-Grauholm *et al.*, “Discovery of the first genome-wide significant risk loci for attention deficit/hyperactivity disorder,” *Nature Genetics*, vol. 51, no. 1, pp. 63–75, 2019. [Online]. Available: <https://doi.org/10.1038/s41588-018-0269-7>
- [36] F. M. T. A. Busing, E. Meijer, and R. V. D. Leeden, “Delete-m Jackknife for Unequal m,” *Statistics and Computing*, vol. 9, no. 1, pp. 3–8, 1999. [Online]. Available: <https://doi.org/10.1023/a:1008800423698>
- [37] E. T. Lim, P. Würtz, A. S. Havulinna, P. Palta, T. Tukiainen, K. Rehnström, T. Esko, R. Mägi, M. Inouye, T. Lappalainen, Y. Chan, R. M. Salem *et al.*, “Distribution and Medical Impact of Loss-of-Function Variants in the Finnish Founder Population,” *PLoS Genetics*, vol. 10, no. 7, p. e1004494, 2014. [Online]. Available: <https://doi.org/10.1371/journal.pgen.1004494>
- [38] M. Correll, D. Moritz, and J. Heer, “Value-Suppressing Uncertainty Palettes,” in *Proceedings of the 2018 CHI Conference on Human Factors in Computing Systems*. ACM, 2018, pp. 1–11. [Online]. Available: <https://doi.org/10.1145/3173574.3174216>
- [39] The International League Against Epilepsy Consortium on Complex Epilepsies, “Genome-wide mega-analysis identifies 16 loci and highlights diverse biological mechanisms in the common epilepsies,” *Nature Communications*, vol. 9, no. 1, p. 5269, 2018. [Online]. Available: <https://doi.org/10.1038/s41467-018-07524-z>
- [40] R. Sund, “Quality of the Finnish Hospital Discharge Register: A systematic review,” *Scandinavian Journal of Public Health*, vol. 40, no. 6, pp. 505–515, 2012. [Online]. Available: <https://doi.org/10.1177/1403494812456637>