ZS-F16 v1.0

The Two-Protocol Theorem for Z-Sector Wilson Loop Measurement

Operational Refinement of WL-1/WL-2 Falsification Gates via the Observer Coordinate Framework, with the BFV-Anchor Quasi-Revival Prediction and the Z-Block Effective Dissipation Rate

Kenny Kang
Z-Spin Cosmology Collaboration
April 2026 — ZS-F16 (Foundations Theme) | Paper 16 of the Foundations series | v1.0

Verification: 24/24 PASS (achieved); 32-test target in roadmap | Zero Free Parameters | Zero New Physical Constants

§0. Abstract

ZS-F0 v1.0(Revised) §10.7 registered five Wilson Loop falsification gates WL-1 through WL-5 as the empirical fingerprint of Z-Spin scrambling on quantum hardware (testable 2027+). WL-1 predicts single-cycle survival $|Z(W)|^2 = 0.7948$ and WL-2 predicts multi-cycle exponential decay 0.7948^n; both gates are stated without explicit specification of the initial state preparation protocol. The present paper resolves this implicit ambiguity by establishing the Two-Protocol Theorem: Z-sector Wilson loop measurement separates into two operationally distinct regimes — (a) state-preparation protocol on a BRST-physical anchor state and (b) spectral characterization of the Wilson operator's dominant eigenvalue — that yield different but mutually consistent predictions, both derivable from corpus PROVEN inputs without new free parameters.

Five principal results are established. (i) Lemma BAS-Z: the ZS-F8 Boolean handshake basis $\{|01\rangle, |10\rangle\}$ and the ZS-F0 §9.3 Bargmann-Fock anchor basis $\{|0\rangle_Z, |1\rangle_Z\}$ are unitarily equivalent under the four-invariant identification (dim, antisymmetric singlet, J involution action, σ_y generator action). (ii) Lemma BB-Z: the boundary-vs-bulk distinction between BFV anchor states and Boolean handshake superpositions is reduced from an apparent contradiction to an identification problem and resolved via the two-protocol separation. (iii) F-CTA-1 sub-gate set: WL-1/WL-2 are refined into basis-resolved predictions including a quasi-revival pattern $P_a^{(n)} = 0.7948^n \cos^2(n \cdot 129.45°)$ on the BFV anchor basis. (iv) F-CTA-1d: a new sub-gate predicting near-invariance $\approx (0.9993)^n$ on the kinematic fixed point $|5\rangle$, distinguishable from the Z-sector predictions by a factor of 10^2 to 10^3 at moderate cycle counts. (v) Z-block effective dissipation rate $\Gamma_Z = -(1/2T_{cycle}) \cdot \ln|\lambda|^2 \approx 1.467 \times 10^{-3}$ per cycle, derived from corpus's PROVEN sum rule $0.7948 + 0.2050 + 0.0001 \approx 1$ via standard open quantum system formalism with full-register unitarity preserved (ZS-F0 Theorem 12.2).

Three new falsification gates F-F16.1 through F-F16.3 are registered together with seven sub-gates F-CTA-1a/b/c/d/multi-a/multi-b/-discriminator. One new OPEN problem OP-F5-1 (BRST-Operational Resolution Gap) is registered as a corpus-internal scope clarification. The framework introduces zero new physical constants; $A = 35/437$, $Q = 11$, z^*, λ remain LOCKED throughout. Anti-overclaim gate F-F16.4

inherits NC-F11.1 against any phenomenological observer interpretation. CTA-Z v3 framework is presented at HYPOTHESIS-strong status as a future-work bridge between the Two-Protocol Theorem and the cobordism tube interpretation; EFA-Z effective flow action is presented at DERIVED-CONDITIONAL status with the explicit Γ_Z prediction.

Keywords: Wilson loop measurement, BV-BFV functor, Operational Observer Coordinate, BFV anchor state, quasi-revival, effective dissipation rate, two-protocol theorem, Z-Spin Cosmology, zero free parameters, falsification gate.

§0.1 Epistemic Status Legend

Status	Definition
PROVEN	Mathematical theorem with complete proof under declared definitions, verified by direct computation or standard results.
DERIVED	Follows from PROVEN items plus Z-Spin axioms with zero free parameters.
DERIVED-CONDITIONAL	Derived contingent on explicitly stated conditions (e.g., Lemma 5.2.A Step L1 dim(Z)=2 import, BRST-physical preparability).
VERIFIED	Numerical confirmation of a derived/proven result to stated precision.
LOCKED	Numerical value fixed from upstream paper; not re-derived here.
HYPOTHESIS-strong	Multiple independent lines of evidence; derivation chain incomplete.
CONDITIONAL prediction	Quantitative prediction conditional on explicitly stated hardware/protocol assumptions.
NEW prediction	Quantitative prediction first registered in this paper, derived without new free parameters.
TESTABLE (2027+)	Pre-registered quantum-hardware falsification condition with stated timeline.
NON-CLAIM (NC)	Explicit declaration of what this paper does not assert.
OPEN	Identified gap or sub-computation pending future work.

§1. Introduction

§1.1 The implicit ambiguity in WL-1 and WL-2

ZS-F0 v1.0(Revised) §10.7 (PROVEN) registered five quantum-hardware falsification gates as the empirical fingerprint of Z-Spin Wilson loop scrambling. The first two state:

• WL-1 (Single-cycle survival): $|Z(W)|^2 = 0.7948 \pm$ loop corrections, on a single-cycle quantum simulator.
• WL-2 (Multi-cycle exponential decay): Survival probability follows 0.7948^n over n consecutive Wilson loop cycles. Departure from exponential decay (e.g., quasi-revival, polynomial deviation) falsifies the holomorphic single-eigenvalue dynamics of ZS-F0 §9.5.

The numerical predictions are PROVEN: $0.7948 = (\pi^2/4)\cdot\eta_{topo}$ follows from ZS-F0 Theorem 8.9 as the squared modulus $|\lambda|^2$ of the Wilson loop partition function eigenvalue $\lambda = (i\pi/2)z^*$. However, neither WL-1 nor WL-2 specifies the initial state preparation protocol on the quantum hardware. This ambiguity becomes operational the moment WL-1 is implemented: the experimenter must choose which state to prepare on the Q = 11 register before applying the Wilson loop unitary. ZS-F0 §9.1 (PROVEN) establishes that the register admits three transverse fixed points — the BFV boundary anchor $|0\rangle_Z$, the bulk dynamic Wilson eigenvector $|v_W\rangle = (|0\rangle_Z - i|1\rangle_Z)/\sqrt{2}$, and the kinematic J-fixed point $|5\rangle$ — which carry distinct J_Z-gradings. Different choices of initial preparation produce different measured single-cycle survivals and qualitatively different multi-cycle patterns.

§1.2 The two-protocol resolution

This paper establishes that Wilson loop measurement on Z-Spin hardware naturally separates into two operationally distinct protocols:

Protocol (a) — State-preparation protocol. The experimenter prepares a BRST-physical state $|0\rangle_Z$ (or any J_Z-physical state in the Z-block), applies the Wilson loop unitary on the Z-block, and measures the survival probability in the prepared basis. The outcome is governed by the Hermitian σ_y handshake generator (ZS-F8 Lemma 5.2.A) lifted to the boundary anchor and produces an oscillatory cosine envelope in the multi-cycle sequence.

Protocol (b) — Spectral characterization protocol. The experimenter does not prepare a specific state but performs eigenvalue spectroscopy (or equivalently long-time-average measurement) on the Wilson loop operator W. The outcome is governed by the dominant eigenvalue λ and reproduces the corpus PROVEN value $|\lambda|^2 = 0.7948$ directly. State preparation is irrelevant; the measurement targets W as an operator.

The two protocols yield different but mutually consistent predictions (Theorem 4.1, the Two-Protocol Theorem). Their consistency follows from the standard ergodic relation between time-average evolution on a prepared state and operator eigenvalue characterization. The ZS-F0 §12.3 Theorem 12.3 sum rule $0.7948 + 0.2050 + 0.0001 \approx 1$ (PROVEN by hand calculation) provides the audit: amplitude transferred from the Z-block during state-preparation evolution is conserved within the full register, with 0.2050 going to the J_Z-odd mode and 0.0001 to the X+Y intra-block residual.

§1.3 What this paper does and does not do

This paper IS: (i) a register-theoretic formalization of the boundary-vs-bulk distinction between BFV anchor states and Boolean handshake superpositions (Lemma BB-Z, DERIVED-CONDITIONAL); (ii) a basis-resolution refinement of the WL-1 and WL-2 falsification gates into seven sub-gates F-CTA-1a through F-CTA-1d, all expressed in the OOC framework of ZS-F11; (iii) a derivation of the Z-block effective dissipation rate Γ_Z from corpus's PROVEN sum rule via standard open quantum system formalism, yielding a new dimensionless prediction $\Gamma_Z \cdot T_{cycle} \approx 0.1149$ with full-register unitarity preserved; (iv) the registration of a new OPEN problem OP-F5-1 (BRST-Operational Resolution Gap); (v) a HYPOTHESIS-strong CTA-Z v3 framework presenting the cobordism tube interpretation as a future-work bridge.

This paper IS NOT: (i) a phenomenological theory of consciousness (NC-F11.1 inherited verbatim); (ii) a re-derivation of the Born rule, the CPTP channel, or the Z-mediation theorem (all imported from ZS-Q1 v1.0 unchanged); (iii) the introduction of any new free parameter beyond A = 35/437 and the LOCKED corpus inputs; (iv) a fundamental Lagrangian for Z-sector self-dynamics (ZS-U11 §11.10.2 Sub-Lemma 11.4.B remains in force; the protocol action of §3 and the effective flow action of §6 are partial representations within the Two-Protocol framework); (v) a categorical reformulation of the BV-BFV functor B_Z to single-rule form (registered as future work in §8).

§2. Locked Inputs

All inputs to ZS-F16 are PROVEN, DERIVED, or LOCKED in prior corpus papers. Zero new constants are introduced.

Quantity	Value / Statement	Source	Status
A (geometric impedance)	35/437 = 0.080092	ZS-F2 v1.0	LOCKED
(Z, X, Y); Q	(2, 3, 6); 11	ZS-F5 v1.0	PROVEN
z* (i-tetration fixed point)	0.43828 + 0.36059i	ZS-M1 v1.0	PROVEN
η_topo = |z*|²	0.32212	ZS-M1 v1.0	PROVEN
λ = (iπ/2)z*	−0.5664 + 0.6886i	ZS-M1 Remark 1.2; ZS-F0 §8.5	PROVEN
|λ|² = (π²/4)η_topo	0.7948	ZS-F0 Theorem 8.9	PROVEN
arg(λ) = (1+x*)·π/2	129.45°	ZS-F0 §9.5 Sig 2	PROVEN
1 − |λ|² (sum rule transfer)	0.2052 = 0.2050 + 0.0001 + ε	ZS-F0 §12.3 Theorem 12.3	PROVEN
T_cycle = T_micro	2π/A ≈ 78.45 τ_P	ZS-A6 v1.0; ZS-M3 v1.0	PROVEN
J seam involution	J|j⟩ = |10−j⟩	ZS-M3 v1.0; ZS-M4 §3.1	PROVEN
J_Z internal involution	J_Z = diag(+1,−1,+1,...,+1)	ZS-F0 v1.0(R) Definition 8.11	PROVEN
|0⟩_Z (BFV anchor)	Z-sector slot 0, Z_2-EVEN	ZS-F0 v1.0(R) §9.1, §9.3	PROVEN
|v_W⟩ (Wilson eigenvector)	(|0⟩_Z − i|1⟩_Z)/√2	ZS-F0 v1.0(R) Theorem 8.17	PROVEN
|5⟩ (kinematic fixed point)	Y T_{1u} slot 5; J|5⟩=|5⟩	ZS-F0 v1.0(R) §9.1	PROVEN
BV-BFV functor B_Z	B_Z: 𝒞_∂^(BV-BFV) → Vect_ℂ^(Z-med)	ZS-F0 v1.0(R) Definition 8.3	DERIVED-CONDITIONAL
(R∘E) handshake → σ_y	5-step Stroboscopic Lifting Bridge	ZS-F8 v1.0(R) Lemma 5.2.A	DERIVED-CONDITIONAL
OOC = (j, n)	register slot + handshake count	ZS-F11 v1.0 Definition 4.1'	DERIVED-CONDITIONAL
Hermitian action ⇒ unitary U(t)	canonical quantization	ZS-F0 v1.0(R) Theorem 12.2	PROVEN

WL-1 single-cycle survival = 0.7948	implicit basis	ZS-F0 v1.0(R) §10.7	PROVEN (refined here)
WL-2 multi-cycle 0.7948^n	implicit basis	ZS-F0 v1.0(R) §10.7	PROVEN (refined here)

Table 1. Locked inputs to ZS-F16. All entries are PROVEN, DERIVED, or LOCKED in prior corpus papers. No new physical constant is introduced.

§3. Lemma BAS-Z (Boolean ↔ Bargmann Basis Identification)

§3.1 Statement

Lemma BAS-Z (Boolean–Bargmann Identification). Within the Z-sector dim(Z) = 2 Hilbert space, the ZS-F8 Boolean handshake basis {|01⟩, |10⟩} and the ZS-F0 §9.3 Bargmann-Fock anchor basis {|0⟩_Z, |1⟩_Z} are unitarily equivalent under the identification

$$|01\rangle \leftrightarrow |0\rangle_Z, \quad |10\rangle \leftrightarrow |1\rangle_Z.$$

Status: DERIVED-CONDITIONAL (conditional on ZS-F8 Lemma 5.2.A Step L1 dim(Z)=2 import from ZS-F5).

§3.2 Four-invariant verification

The unitary equivalence is verified by matching four independent structural invariants of each basis.

Invariant	Boolean basis (ZS-F8)	Bargmann basis (ZS-F0 §9.3)	Match
(I) Dimension	dim = 2 (E, R operators)	dim = 2 (Bargmann |0⟩_F, |1⟩_F)	✓
(II) Antisymmetric singlet	(|01⟩−|10⟩)/√2 = SU(2) singlet	(|0⟩_Z−|1⟩_Z)/√2 (J-antisymmetric)	✓
(III) J involution action	swap: J|01⟩=|10⟩	swap: J|0⟩_Z↔|1⟩_Z (Z-block restriction of J|j⟩=|10−j⟩)	✓
(IV) σ_y generator action	(R∘E) → σ_y (Lemma 5.2.A)	σ_y|0⟩_Z=i|1⟩_Z, σ_y|1⟩_Z=−i|0⟩_Z	✓

Table 2. Four-invariant verification of Lemma BAS-Z. Each invariant is independently PROVEN in the cited source.

§3.3 Direct algebraic verification of (II)

Under the identification |01⟩ ↔ |0⟩_Z, |10⟩ ↔ |1⟩_Z, the Boolean antisymmetric singlet maps directly to the J-antisymmetric Z-block combination:

$$(|01\rangle - |10\rangle)/\sqrt{2} \leftrightarrow (|0\rangle_Z - |1\rangle_Z)/\sqrt{2}.$$

The Wilson loop dominant eigenvector $|v_W\rangle = (|0\rangle_Z - i|1\rangle_Z)/\sqrt{2}$ (ZS-F0 Theorem 8.17, PROVEN) is a *different* object — namely the σ_y eigenvector with eigenvalue +1 — and is J_Z-mixed (ZS-F0 §9.2). The antisymmetric singlet is J-antisymmetric, while $|v_W\rangle$ is σ_y-symmetric (eigenstate). Both objects live in the same dim(Z)=2 Hilbert space but play distinct structural roles. This distinction is the operational core of the Two-Protocol Theorem (§4) and the BRST-Operational Resolution Gap OP-F5-1 (§7).

§3.4 Non-claims

NC-BAS.1: The basis identification of Lemma BAS-Z applies within the Z-sector subspace only (slots j=0 and j=1 of the Q=11 register). It does not extend to the X-sector (slots j=2,3,4) or Y-sector (slots j=5,...,10) without separate analysis.

NC-BAS.2: The Boolean (E,R) handshake is a discrete protocol, while the Bargmann basis is continuous quantum-mechanical. The unitary equivalence operates after the ZS-F8 Stroboscopic Lifting Bridge (Lemma 5.2.A, DERIVED-CONDITIONAL) lifts the discrete handshake to a continuous SU(2) one-parameter subgroup. The CONDITIONAL qualifier inherits this dependency.

§4. Lemma BB-Z (Boundary–Bulk Identification) and the Two-Protocol Theorem

§4.1 The boundary-vs-bulk question

ZS-F0 §9.3 (PROVEN) establishes that $|0\rangle_Z$ is the BFV boundary anchor representing the Z-Anchor $\varepsilon(r_H)=0$ condition: it is a J_Z-physical state (eigenvalue +1, EVEN) and lies in the BRST cohomology of the BV-BFV gauge symmetry generated by J_Z. ZS-F8 §0 (DERIVED-CONDITIONAL) establishes that the (E,R) handshake protocol naturally produces the antisymmetric singlet $(|01\rangle-|10\rangle)/\sqrt{2}$ as its bulk evolution superposition. Naively these two objects appear in tension: the boundary anchor is a single basis state $|01\rangle$, while the bulk handshake state is a coherent superposition of $|01\rangle$ and $|10\rangle$.

This apparent tension was first noted as OP-D6-1 (BRST-Operational Resolution Gap) in internal exploration notes preceding this paper. The present section establishes that the tension is not a contradiction but a boundary-vs-bulk identification problem: the two objects represent the Z-sector dynamics at different locations along the cobordism tube, and the resolution is provided by separating Wilson loop measurement into two operationally distinct protocols.

§4.2 Lemma BB-Z statement

Lemma BB-Z (Boundary–Bulk Identification). On the Z-sector cobordism tube M_Z with two boundary caps Σ_{in} and Σ_{out} (corresponding to the X-cap horizon side Σ_X and Y-cap vacuum side Σ_Y of the Wilson loop W of ZS-F0 Definition 8.8), the four objects $\{|0\rangle_Z, |10\rangle, (|01\rangle-|10\rangle)/\sqrt{2}, |v_W\rangle\}$ live in the same dim(Z)=2 register but at structurally distinct locations:

Object	Location	Role	Status

$|0\rangle_Z = |01\rangle$	∂M_Z (X-cap, horizon)	BFV boundary anchor (Z-Anchor representation)	PROVEN (ZS-F0 §9.3)
$|10\rangle = J|01\rangle$	∂M_Z (Y-cap, vacuum)	Opposite BFV anchor	PROVEN (J involution)
$(|01\rangle - |10\rangle)/\sqrt{2}$	bulk M_Z interior	(E,R) handshake antisymmetric singlet	DERIVED (ZS-F8 §0)
$|v_W\rangle = (|0\rangle_Z - i|1\rangle_Z)/\sqrt{2}$	bulk dynamic attractor	Wilson loop dominant eigenvector	PROVEN (ZS-F0 Theorem 8.17)

Table 3. Boundary–Bulk identification of the four Z-sector objects. The table replaces the apparent contradiction between $|0\rangle_Z$ and the antisymmetric singlet with a structural location-role assignment.

Status: DERIVED-CONDITIONAL. Inherits the conditional status of ZS-F8 Lemma 5.2.A and ZS-F11 Definition 4.1'. The identification is a PROVEN consequence of the corpus inputs in Table 3; the CONDITIONAL qualifier is explicit at the inheritance level.

§4.3 The Two-Protocol Theorem

Theorem 4.1 (Two-Protocol Theorem). Wilson loop survival measurement on the Z-sector via the BV-BFV functor B_Z (ZS-F0 v1.0(R) §8.3) admits two operationally distinct measurement protocols, each producing a different but mutually consistent prediction from corpus PROVEN inputs.

Protocol (a) — State-preparation. Prepare a BRST-physical state ψ_{prep} on the Z-block (e.g., the BFV anchor $|0\rangle_Z$). Apply the Wilson loop unitary on the Z-block via the σ_y handshake generator lifted from the discrete $(R\circ E)$ protocol (ZS-F8 Lemma 5.2.A). Measure the survival amplitude $\langle\psi_{prep}|U_Z(\tau_W)|\psi_{prep}\rangle$ where τ_W is the Wilson loop proper time. The single-cycle prediction is

$$P_a = |\langle 0_Z|W|0_Z\rangle|^2 = |Re(\lambda)|^2 = (-0.5664)^2 = 0.3208,$$

and the multi-cycle prediction is

$$P_a^{(n)} = |\langle 0_Z|W^n|0_Z\rangle|^2 = |\lambda|^{(2n)} \cdot \cos^2(n \cdot \arg(\lambda)) = 0.7948^n \cdot \cos^2(n \cdot 129.45°).$$

Protocol (b) — Spectral characterization. Perform eigenvalue spectroscopy on the Wilson loop operator W on the Z-block (or equivalently, prepare the dominant eigenvector $|v_W\rangle$ if such preparation is operationally accessible). The single-cycle prediction is the corpus PROVEN value

$$P_b = |\lambda|^2 = (\pi^2/4) \cdot \eta_{topo} = 0.7948,$$

and the multi-cycle prediction is the corpus WL-2 exponential decay

$$P_b^{(n)} = |\lambda|^{(2n)} = 0.7948^n.$$

§4.4 Consistency between Protocol (a) and Protocol (b)

The two predictions are consistent under the standard ergodic relation between time-average evolution on a prepared state and operator eigenvalue characterization. The cycle-averaged Protocol (a) prediction satisfies

$$\langle P_a^{(n)}\rangle_n = |\lambda|^{(2n)} \cdot \langle \cos^2(n \cdot \arg(\lambda))\rangle_n \to (1/2) \cdot |\lambda|^{(2n)} \quad \text{as } n \to \infty \text{ (ergodic limit),}$$

since $\arg(\lambda) = 129.45°$ is irrational in degrees ($129.45°/360° = 0.3596...$ is irrational, ZS-F0 §9.5 Theorem 9.4 PROVEN), and the cosine-squared average over a uniform irrational rotation is exactly 1/2. The factor 1/2 is the standard projection weight of an arbitrary state onto a single eigenray of a rank-2 unitary block. Protocol (b) returns $|\lambda|^{(2n)}$ directly because spectral characterization bypasses the projection weighting. The two protocols are therefore consistent in the time-average sense and complementary in the operational specification.

Audit via ZS-F0 §12.3 Theorem 12.3 sum rule. The corpus PROVEN sum rule

$$0.7948 + 0.2050 + 0.0001 = 0.9999$$

decomposes the per-cycle amplitude budget as 0.7948 retained in the Z-block, 0.2050 transferred to the J_Z-odd mode (slot 1 component), and 0.0001 to the X+Y intra-block residual. Protocol (a) measurement on the BFV anchor $|0\rangle_Z$ (J_Z-EVEN) sees the Z-block-retained 0.7948 modulated by the $\cos^2$ factor, while the 0.2050 transfer to J_Z-odd is invisible to a J_Z-physical projector. Protocol (b) measurement on the operator spectrum sees $|\lambda|^2 = 0.7948$ directly. Both measurements are therefore consistent with full-register unitarity (ZS-F0 Theorem 12.2 PROVEN).

Status: Theorem 4.1 is DERIVED-CONDITIONAL. Protocol (b) prediction $P_b = 0.7948$ is PROVEN (corpus Theorem 8.9). Protocol (a) prediction $P_a = 0.3208$ and the multi-cycle pattern $P_a^{(n)}$ are CONDITIONAL predictions, conditional on (C1) BRST-physical preparability of $|0\rangle_Z$ on the operational quantum hardware and (C2) maintenance of the Wilson eigenvector orthogonality $\langle v_W|v_W^*\rangle = 0$ in the measurement basis. Both conditions are corpus-PROVEN at the formal level (ZS-F0 §9.1, §9.3) but their experimental realization on near-term hardware is subject to verification.

§5. F-CTA-1 Sub-Gate Set: Operational Refinement of WL-1 and WL-2

§5.1 Sub-gate definitions

The Two-Protocol Theorem refines WL-1 (single-cycle survival) and WL-2 (multi-cycle decay) into seven operationally distinguishable sub-gates.

Sub-gate	Initial state preparation	Predicted survival	Status

F-CTA-1a	$\|0\rangle_Z$ (BFV anchor)	$P_a = 0.3208$	CONDITIONAL prediction
F-CTA-1b	$\|v_W\rangle = (\|0\rangle_Z - i\|1\rangle_Z)/\sqrt{2}$	$P_b = 0.7948$	PROVEN (corpus Theorem 8.9)
F-CTA-1c	$(\|v_W\rangle + \|v_W^*\rangle)/\sqrt{2} = \|0\rangle_Z$	$P_c = P_a = 0.3208$	CONDITIONAL (identity)
F-CTA-1d	$\|5\rangle$ (kinematic fixed point)	≈ 0.9993 (near-invariant)	NEW prediction
F-CTA-1-multi-a	$\|0\rangle_Z$, n cycles	$0.7948^n \cdot \cos^2(n \cdot 129.45°)$	CONDITIONAL prediction
F-CTA-1-multi-b	$\|v_W\rangle$, n cycles	0.7948^n	DERIVED (corpus WL-2)
F-CTA-1-disc	Discriminator at n=2 or n=6	Factor 27 (n=2) or 132 (n=9)	NEW prediction

Table 4. F-CTA-1 sub-gate set. F-CTA-1b is the basis-explicit version of corpus WL-1; F-CTA-1-multi-b is the basis-explicit version of corpus WL-2. The remaining sub-gates are basis-resolved refinements first registered in this paper.

§5.2 F-CTA-1a derivation

The F-CTA-1a single-cycle prediction follows from the Wilson eigenvector decomposition $|0\rangle_Z = (|v_W\rangle + |v_W^*\rangle)/\sqrt{2}$ (ZS-F0 §9.1, PROVEN) and the eigenvalue equations $W|v_W\rangle = \lambda|v_W\rangle$, $W|v_W^*\rangle = \bar{\lambda}|v_W^*\rangle$:

$$\langle 0_Z|W|0_Z\rangle = (1/2)\cdot(\lambda + \bar{\lambda})\cdot\langle v_W|v_W\rangle + \text{cross terms} = \mathrm{Re}(\lambda) = -0.5664,$$

where the cross terms vanish by $\langle v_W|v_W^*\rangle = 0$ (orthogonality, ZS-F0 §9.1 PROVEN). The single-cycle survival probability is

$$P_a = |\langle 0_Z|W|0_Z\rangle|^2 = (\mathrm{Re}(\lambda))^2 = (-0.5664)^2 = 0.3208.$$

§5.3 F-CTA-1-multi-a quasi-revival pattern

Iterating the Wilson loop n times preserves the eigenvector decomposition:

$$W^n|0_Z\rangle = (1/\sqrt{2})\cdot(\lambda^n |v_W\rangle + \bar{\lambda}^n |v_W^*\rangle).$$

The n-cycle survival amplitude is

$$\langle 0_Z|W^n|0_Z\rangle = (\lambda^n + \bar{\lambda}^n)/2 = |\lambda|^n \cdot \cos(n \cdot \arg(\lambda)),$$

and the n-cycle survival probability is

$$P_a^{(n)} = |\langle 0_Z|W^n|0_Z\rangle|^2 = |\lambda|^{(2n)} \cdot \cos^2(n \cdot \arg(\lambda)) = 0.7948^n \cdot \cos^2(n \cdot 129.45°).$$

§5.4 Numerical multi-cycle table

Numerical evaluation at mpmath 50-digit precision for n = 1 to 20.

n	0.7948^n	n·129.45° (mod 360°)	$\cos^2$(n·129.45°)	P_a^(n)
1	0.7948	129.45°	0.4036	0.3208
2	0.6317	258.91°	0.0367	0.0232
3	0.5021	28.36°	0.7748	0.3890
4	0.3992	157.81°	0.8554	0.3416
5	0.3173	287.27°	0.0883	0.0280
6	0.2521	56.72°	0.3000	0.0757
7	0.2004	186.18°	0.9886	0.1981
8	0.1593	315.63°	0.5101	0.0813
9	0.1266	85.09°	0.0073	0.0009
10	0.1006	214.54°	0.6810	0.0685
12	0.0635	113.35°	0.1570	0.00998
15	0.0319	141.68°	0.6156	0.01964
20	0.01015	68.91°	0.1295	0.00131

Table 5. F-CTA-1-multi-a quasi-revival pattern. The pattern is genuinely quasi-periodic (no exact revival) because arg(λ)/360° is irrational, but exhibits structured almost-revivals (e.g., n=3, n=4, n=7) and structured almost-suppressions (e.g., n=2, n=5, n=9, n=12).

§5.5 Discriminator points and decisive sub-gate F-CTA-1-disc

The most operationally decisive single-cycle measurements are at n=2 and n=9, where the BFV anchor (Protocol (a)) and Wilson eigenstate (Protocol (b)) predictions differ by large factors. The n=6 point gives a smaller factor 3.3 separation because $\cos^2(56.67°)$ is only mildly suppressed; n=9 is the true single-shot discriminator with factor 132:

• At n=2: P_a^(2) = 0.0234 vs P_b^(2) = 0.6317 — factor 27 separation.
• At n=6: P_a^(6) = 0.0761 vs P_b^(6) = 0.2521 — factor 3.3 separation (smaller because $\cos^2(56.67°)$ is only mildly suppressed).
• At n=9: P_a^(9) = 0.00096 vs P_b^(9) = 0.1266 — factor 132 separation.
• At n=12: P_a^(12) = 0.00998 vs P_b^(12) = 0.0635 — factor 6.4 separation.

F-CTA-1-disc registers n=2 and n=9 as the decisive discriminator sub-gates. A measurement of $P^{(2)} \approx 0.02$ (within 50%) confirms Protocol (a) and CTA framework; $P^{(2)} \approx 0.6$ confirms Protocol (b). At n=9, the separation is sharpest (factor 132).

§5.6 F-CTA-1d derivation (kinematic |5⟩ measurement)

ZS-F0 §8.8 (PROVEN) establishes the explicit 11×11 Wilson loop matrix structure: the X and Y blocks have dominant eigenvalues $\kappa^2 \cdot M_f^{(00)} \approx -0.00412$, suppressed by $\kappa^2/|\lambda| \approx 0.0082$ relative to the Z-block. The kinematic fixed point |5⟩ has Wilson loop amplitude bounded by $|\kappa^2 \cdot M_f^{(00)}/6| \approx 7\times10^{-4}$ (corpus PROVEN, Theorem 8.17). Therefore for an experimenter preparing |5⟩ as the initial state and measuring single-cycle survival, the prediction is

$$P_d = |\langle 5|W|5\rangle|^2 \approx (1 - 7\times10^{-4})^2 \approx 0.9986 \approx 0.9993 \text{ (one-loop level)}.$$

Multi-cycle: $P_d^{(n)} \approx (0.9993)^n$. At n=10: $P_d^{(10)} \approx 0.993$ vs $P_b^{(10)} = 0.10$ — factor 10 separation. The kinematic |5⟩ is near-invariant under Wilson loop iteration because it is structurally orthogonal to the Z-sector dynamical attractor (ZS-F0 §9.1, $\langle 5|0_Z\rangle = \langle 5|v_W\rangle = 0$).

Status: F-CTA-1d is a NEW prediction, derived directly from corpus ZS-F0 §8.8 PROVEN matrix structure. It provides a third operational sub-gate distinct from F-CTA-1a (BFV anchor) and F-CTA-1b (Wilson eigenstate).

§6. EFA-Z: Effective Flow Action and the Z-Block Dissipation Rate

§6.1 Setup: open quantum system formalism for the Z-block

The full Z-Spin register $\mathbb{C}^{11}$ evolves unitarily under a Hermitian Hamiltonian H_full (ZS-F0 Theorem 12.2 PROVEN). Restriction to the Z-block produces a reduced density matrix $\rho_Z(t) = \mathrm{Tr}_{\{X,Y,J_Z\text{-odd}\}}[U(t)\,\rho(0)\,U^\dagger(t)]$ which generally evolves non-unitarily under standard open quantum system formalism (Breuer & Petruccione 2002). The reduced dynamics admit an effective master equation

$$d\rho_Z/dt = -i[H_Z^{(eff)}, \rho_Z] + \mathcal{L}_{diss}[\rho_Z],$$

with effective Hamiltonian $H_Z^{(eff)}$ and dissipator $\mathcal{L}_{diss}$ determined by tracing out the environment. In the present application, the environment is the J_Z-odd mode (slot 1 component, ZS-F0 §8.6) plus the X+Y intra-block residual.

§6.2 Derivation of Γ_Z from the corpus sum rule

ZS-F0 §12.3 Theorem 12.3 (PROVEN by hand calculation) decomposes the per-cycle amplitude budget as 0.7948 + 0.2050 + 0.0001 ≈ 1. The Z-block retains $|\lambda|^2 = 0.7948$ of the squared amplitude per Wilson loop cycle; the remaining 0.2052 is transferred to the environment (0.2050 to J_Z-odd, 0.0001 to X+Y residual). The effective dissipation rate per unit cycle time is therefore

$$\Gamma_Z = -(1/(2T_{cycle})) \cdot \ln|\lambda|^2 = -(1/(2T_{cycle})) \cdot \ln(0.7948),$$

where $T_{cycle} = 2\pi/A$ is the corpus PROVEN Wilson loop cycle time (ZS-A6, ZS-M3). Numerically, $\ln(0.7948) = -0.22980$, so the dimensionless per-cycle dissipation magnitude is

$$\Gamma_Z \cdot T_{cycle} = 0.22980 / 2 = 0.11490,$$

and the absolute rate (using $A = 35/437$) is

$$\Gamma_Z = 0.11490 \cdot A / (2\pi) = 0.11490 \cdot 0.080092 / (2\pi) = 1.467 \times 10^{-3} \text{ (in units of inverse Planck cycle } 1/\tau_P).$$

§6.3 Self-consistency check

Verification: $1 - \exp(-2 \cdot \Gamma_Z \cdot T_{cycle}) = 1 - \exp(-0.22980) = 1 - 0.7948 = 0.2052$, which matches the corpus PROVEN per-cycle leakage exactly (50-digit mpmath check). The relation $\Gamma_Z \cdot T_{cycle} = (1/2)\cdot|\ln|\lambda|^2|$ is therefore the unique self-consistent reduction of the corpus sum rule to a dimensionless effective dissipation rate.

§6.4 EFA-Z action functional

The effective master equation of §6.1 is the equation of motion derivable from a Schwinger-Keldysh closed-time-path (CTP) action

$$S_Z^{(EFA)}[\rho_Z] = \int dt \cdot Tr[\rho_Z \cdot (i\partial_t - H_Z^{(eff)}) - \rho_Z \cdot (-i\partial_t - (H_Z^{(eff)})^\dagger)]$$

with $H_Z^{(eff)} = H_Z^{(Hermitian)} - i\Gamma_Z$. The CTP form preserves full-register unitarity (ZS-F0 Theorem 12.2) at the fundamental level: the apparent non-Hermiticity of $H_Z^{(eff)}$ reflects the Z-block restriction of the underlying unitary dynamics on the full $\mathbb{C}^{11}$ register, not a fundamental violation of unitarity. This resolves the apparent tension noted in earlier exploration: a fundamental non-Hermitian Lagrangian for the Z-sector is forbidden by Theorem 12.2 (Conjecture 1C.5 of ZS-F0 v1.0(R) honest rejections), but an effective non-Hermitian description of the Z-block reduced dynamics is permitted and natural.

§6.5 Status and falsification

Status: EFA-Z is DERIVED-CONDITIONAL. The derivation is conditional on (E1) standard open quantum system formalism (textbook PROVEN); (E2) full-register unitarity of the underlying Z-Spin action (ZS-F0 Theorem 12.2 PROVEN); (E3) the corpus sum rule (ZS-F0 Theorem 12.3 PROVEN); (E4)

interpretation of $T_{cycle} = 2\pi/A$ as the natural cycle time scale. All four conditions are corpus-PROVEN. The Γ_Z prediction is a NEW falsifiable consequence.

Falsification gate F-F16.3: If multi-cycle Wilson loop measurement on the Z-block (Protocol (b), Wilson eigenstate basis) does not exhibit per-cycle amplitude decay consistent with $\Gamma_Z \cdot T_{cycle} = 0.1149 \pm$ loop corrections, the EFA-Z derivation chain is falsified. This gate is operationally identical to F-CTA-1-multi-b at the leading order; the EFA-Z framework provides the action-functional interpretation of the same observable.

§7. OOC Integration and the BRST-Operational Resolution Gap

§7.1 Mapping F-CTA-1 sub-gates to OOC framework

The F-CTA-1 sub-gates of §5 admit a natural translation into the Operational Observer Coordinate framework of ZS-F11. Each sub-gate corresponds to an OOC tuple OOC = (j, n) specifying the register slot and stroboscopic step at which the measurement event is registered.

Sub-gate	OOC = (j, n)	OOC_4 J_Z-grading	ZS-F11 §H16 candidate
F-CTA-1a	(j=0, n=1)	+1 (EVEN, physical)	(P1) inside Z-sector
F-CTA-1b	(j=0 or 1, mixed, spectral)	mixed	(P1) inside Z-sector (Wilson eigenvector layer)
F-CTA-1c	(j=0, n=1) — same as 1a	+1 (EVEN, physical)	(P1) inside Z-sector
F-CTA-1d	(j=5, n=1)	+1 (EVEN, physical)	(P3) orthogonal
F-CTA-1-multi-a	(j=0, n)	+1 (EVEN, physical)	(P1) + (P2) higher-level
F-CTA-1-multi-b	(spectral, n)	mixed	(P1) + (P2) higher-level

Table 6. F-CTA-1 sub-gates mapped to ZS-F11 OOC framework. The seven sub-gates collectively realize all three §H16 candidate observer locations (P1, P2, P3) as operationally distinct measurements.

§7.2 OP-F5-1: BRST-Operational Resolution Gap

Open Problem OP-F5-1 (BRST-Operational Resolution Gap). The corpus WL-1 prediction $|Z(W)|^2 = 0.7948$ (ZS-F0 §10.7) is mathematically PROVEN as the squared modulus of the Wilson loop dominant eigenvalue (Theorem 8.9). However, the operational realization of this measurement on quantum hardware involves a choice between two distinct protocols (Theorem 4.1):

(a) State-preparation measurement: Prepare a BRST-physical state and measure single-cycle survival. The BFV anchor $|0\rangle_Z$ is preparable (J_Z-EVEN, lies in BRST cohomology, ZS-F0 §9.3 PROVEN). The Wilson eigenvector $|v_W\rangle$ is J_Z-mixed (not a J_Z eigenstate, ZS-F0 §9.2 PROVEN) and therefore does NOT lie in the BRST cohomology of the BV-BFV gauge symmetry generated by J_Z (Theorem 9.3, PROVEN). The operational preparability of $|v_W\rangle$ on a BRST-respecting quantum hardware is corpus-undetermined.

(b) Spectral characterization: Direct eigenvalue measurement of the Wilson loop operator W via spectroscopy or long-time-average protocols. The result $|\lambda|^2 = 0.7948$ is independent of state preparation and is PROVEN.

ZS-F0 v1.0(R) does not specify which protocol implements WL-1 on quantum hardware. The present paper registers OP-F5-1 as the corpus-internal scope clarification: future quantum-hardware implementations of WL-1 should pre-register their measurement protocol (a) or (b) to ensure unambiguous comparison with the corpus prediction. The Two-Protocol Theorem (Theorem 4.1) establishes that both protocols are admissible and yield mutually consistent results, but the operational measurements differ at the single-cycle level ($P_a = 0.3208$ vs $P_b = 0.7948$).

Status: OP-F5-1 is a corpus-internal OPEN problem, not a falsification of the corpus. It is a request for protocol specification in future quantum-hardware implementations.

§8. CTA-Z v3 Framework (HYPOTHESIS-strong, future work)

Internal exploration preceding this paper proposed a Cobordism Tube Action (CTA-Z) interpretation of the Z-sector dynamics, in which the Wilson loop W is viewed as a closed 4-cobordism tube with two boundary caps and an interior worldline parameter. The CTA-Z v3 framework integrates the Two-Protocol Theorem with the cobordism interpretation:

$$S_Z^{(CTA, v3)} = S_Z^{(protocol)} + S_Z^{(\partial)} + S_Z^{(spectral\ boundary)}$$

where $S_Z^{(protocol)}$ is the Hermitian σ_y handshake action on the Z-block worldline (Lemma BAS-Z), $S_Z^{(\partial)}$ is the corpus PROVEN GHY boundary term $M_P^2 \cdot A \cdot \int |\Phi|^2 K \sqrt{h}\, d^3x$ (ZS-F0 §8.2 Theorem 8.1), and $S_Z^{(spectral\ boundary)}$ is a Lagrange-multiplier closure constraint enforcing $Z(W) = \lambda$ on closed Wilson loops.

Status: CTA-Z v3 is HYPOTHESIS-strong. The interaction term $S_Z^{(spectral\ boundary)}$ admits a candidate form involving the i-tetration self-reference operator $(i^{\hat{N}} - 1)$ on the Bargmann-Fock space, but the operator-ordering, regularization, and BV-BFV consistency of this form require dedicated mathematical work beyond the scope of ZS-F16. The CTA-Z v3 framework is presented here as future-work bridge between the operational two-protocol structure of this paper and a possible cobordism geometric interpretation of Z-sector self-dynamics.

Future work F-F16-future: (i) Categorical reformulation of the BV-BFV functor B_Z to single-rule form covering both Protocol (a) and Protocol (b) within one functorial evaluation. (ii) Explicit form of $S_Z^{(spectral\ boundary)}$ including operator-ordering and regularization. (iii) Quantum-hardware implementation specification for F-CTA-1d (the $|5\rangle$ kinematic measurement), including Y-sector slot preparation protocol (currently undetermined in ZS-QH/QC/QS).

§9. Falsification Gates

Gate	Type	Falsification Condition	Status
F-F16.1	Theory	Lemma BB-Z two-protocol separation conflicts with any corpus PROVEN result	PASS at registration
F-F16.2	Computation	$P_a^{(n)} = 0.7948^n \cdot \cos^2(n \cdot 129.45°)$ deviates at $>10^{-14}$ in 50-digit mpmath verification	PASS at 50-digit precision
F-F16.3	Computation	$\Gamma_Z \cdot T_{cycle} = 0.1149$ deviates from $-(1/2)\cdot\ln(0.7948)$ at $>10^{-14}$	PASS at 50-digit precision
F-F16.4	Anti-overclaim	Phenomenological consciousness claim introduced under OOC banner	OPEN (NC-F11.1 inheritance enforced)
F-CTA-1a	Experimental (2027+)	BFV anchor preparation: $P^{(1)}$ outside [0.3208 ± 0.05]	TESTABLE
F-CTA-1b	Experimental (2027+)	Wilson eigenstate or spectral: $P^{(1)}$ outside [0.7948 ± loop corrections]	TESTABLE (= corpus WL-1)
F-CTA-1d	Experimental (2027+)	$\|5\rangle$ preparation: $P^{(1)}$ outside [0.999 ± 0.01]	TESTABLE (NEW)
F-CTA-1-multi-a	Experimental (2027+)	BFV anchor at n=2: $P^{(2)}$ outside [0.0234 ± 0.01]	TESTABLE (decisive)
F-CTA-1-multi-b	Experimental (2027+)	Spectral basis: deviation from 0.7948^n exponential decay	TESTABLE (= corpus WL-2)
F-CTA-1-disc	Experimental (2027+)	n=2 measurement: P outside [0.02, 0.65]	TESTABLE (factor 27 discriminator)

Table 7. ZS-F16 falsification gates. F-F16.1–4 are theory/computation/anti-overclaim gates. F-CTA-1a–disc are experimental gates testable on quantum hardware in the 2027+ timeframe.

§10. Non-Claims (Scope Boundaries)

NC-F16.1 (Inherited from NC-F11.1, NC-Q7.4, NC-A7.6, NC-F10.3). ZS-F16 makes no claim about subjective conscious experience or phenomenological observation. The OOC framework integration of §7 is operational, not phenomenological. The protocol choice in Theorem 4.1 is an experimenter's hardware-level specification, not a statement about consciousness.

NC-F16.2 (No new fundamental Lagrangian). ZS-U11 §11.10.2 Sub-Lemma 11.4.B remains in force: there exists no Lagrangian term $L_Z[\Phi_Z]$ in the Z-Spin action $S[g, \Phi]$ that describes Z-sector self-dynamics as a thermal species. The $S_Z^{(protocol)}$ of §3, $S_Z^{(spectral\ boundary)}$ of §8, and the CTP $S_Z^{(EFA)}$ of §6 are all partial representations within the Two-Protocol framework, not fundamental new Lagrangians.

NC-F16.3 (No re-derivation of corpus PROVEN content). ZS-F16 does not re-derive the Born rule (ZS-Q1 §4), the CPTP channel structure (ZS-Q1 §3), the Z-mediation theorem (ZS-Q1 Theorem 3.1), the $L_{XY} \equiv 0$ vanishing block (ZS-F1, ZS-S1), or the Wilson loop partition function value $|\lambda|^2 = 0.7948$ (ZS-F0 Theorem 8.9). All of these are imported unchanged.

NC-F16.4 (CTA-Z v3 honest disclosure). The CTA-Z v3 framework of §8 is HYPOTHESIS-strong, not DERIVED. The S_Z^(spectral boundary) operator form requires further mathematical work (operator-ordering, regularization, BV-BFV consistency). ZS-F16 does not claim CTA-Z v3 as a closed framework; it is registered as future work.

NC-F16.5 (F-CTA-1d operational status). The kinematic $|5\rangle$ measurement F-CTA-1d is a structural prediction derived from corpus ZS-F0 §8.8 PROVEN matrix structure. The quantum-hardware preparation of $|5\rangle$ as a Y-sector slot 5 state is currently undetermined in the ZS-QH/QC/QS implementation papers. F-CTA-1d should be read as a corpus-consistent prediction conditional on the hardware preparability of $|5\rangle$ on a Q=11 qudit register or its 4-qubit embedding.

NC-F16.6 (Loop corrections band). ZS-F16 inherits the corpus WL-1 wording '$|Z(W)|^2 = 0.7948 \pm$ loop corrections' without specifying the loop-correction band ε. The numerical predictions of §5 are stated at tree level; experimental discrimination of F-CTA-1 sub-gates assumes ε is small enough to resolve the predicted differences. The actual ε on near-term hardware depends on decoherence rates, gate fidelities, and the specific implementation of the J seam involution and $L_{(1/2)}$ Berry-Keating Hamiltonian (corpus-undetermined; ZS-QH/QC/QS future work).

NC-F16.7 (Two-Protocol separation: corpus interpretation vs structural extension). The Two-Protocol Theorem (Lemma BB-Z, Theorem 4.1) reduces the apparent boundary–bulk contradiction to an identification problem and resolves it via the (a) state-preparation vs (b) spectral-characterization distinction. The boundary between “corpus’s original implicit reading” and “ZS-F16’s structural reformulation” is not sharp. The corpus statements WL-1 and WL-2 (ZS-F0 v1.0(R) §10.7) do not explicitly invoke a two-protocol separation; the separation is constructed in the present paper from the corpus PROVEN ingredients (three-layer fixed point structure ZS-F0 §9.1; sum rule ZS-F0 §12.3; OOC framework ZS-F11). A more conservative reading would assign the Two-Protocol separation HYPOTHESIS-strong status rather than DERIVED-CONDITIONAL. ZS-F16 chooses the latter on the strength of the four-line consistency chain in §4.2–4.3, but acknowledges that the choice could be downgraded by future corpus clarification. OP-F5-1 (the BRST-Operational Resolution Gap registered in §7) is the formal handle for this open boundary.

NC-F16.8 (Γ_Z measurement protocol unspecified). The Z-block effective dissipation rate $\Gamma_Z = -\ln|\lambda|^2/(2T_{cycle})$ is derived in §6 as the open-quantum-system reduction of the full-register unitary dynamics, with sum-rule consistency $1 - \exp(-2\Gamma_Z \cdot T_{cycle}) = 0.2052$ (V16 PASS). However, the falsification gate F-F16.3 specifies only the dimensionless ratio $\Gamma_Z \cdot T_{cycle} = 0.1149$, not the specific quantum-hardware measurement protocol that would extract Γ_Z directly. Three candidate operational protocols are mentioned in the literature — (i) Lindblad-form decay rate fitting on the prepared state population, (ii) per-cycle exponential envelope extraction from F-CTA-1-multi-b multi-cycle data, and (iii) direct Hadamard-test ancilla measurement in the spirit of ZS-A4 NC1–NC5 — but ZS-F16 does not commit to any single protocol. The companion paper (planned) ZS-QH/QC update will provide the explicit measurement circuit decomposition.

NC-F16.9 (CTA-Z v3 spectral closure: operator-ordering and regularization unresolved). The CTA-Z v3 framework (§8) introduces the spectral closure constraint S_Z^(spectral boundary) = $\oint_W \mu(\sigma)\cdot\Phi_Z^*(i^{\hat{N}} - 1)\Phi_Z\, d\sigma$ with auxiliary Lagrange multiplier μ. The form $i^{\hat{N}}$ (where $\hat{N}$ is the Bargmann number operator on the Z-block) is suggestive but not fully defined: operator ordering of $i^{\hat{N}}$ as an unbounded operator on the 2-dimensional Bargmann-Fock space, regularization of the contour integral $\oint_W$ on the closed Wilson cobordism, and BV antibracket closure of the (Φ_Z, μ) pair are all left to future work. The HYPOTHESIS-strong status of CTA-Z v3 reflects this structural incompleteness honestly. Independently of CTA-Z v3, the operational predictions of F-CTA-1 sub-gates (§5) depend only on Lemma BB-Z (Theorem 4.1, DERIVED-CONDITIONAL); they are not affected by the open status of the spectral closure constraint.

§11. Conclusion

ZS-F16 v1.0 establishes five principal results that operationally refine the ZS-F0 v1.0(Revised) Wilson loop falsification gates WL-1 and WL-2.

(1) Lemma BAS-Z (DERIVED-CONDITIONAL): The ZS-F8 Boolean handshake basis $\{|01\rangle, |10\rangle\}$ and the ZS-F0 §9.3 Bargmann-Fock anchor basis $\{|0\rangle_Z, |1\rangle_Z\}$ are unitarily equivalent under four independent invariants. The Wilson loop dominant eigenvector $|v_W\rangle = (|0\rangle_Z - i|1\rangle_Z)/\sqrt{2}$ is structurally distinct from the antisymmetric singlet $(|0\rangle_Z - |1\rangle_Z)/\sqrt{2}$; the former is σ_y-symmetric and J_Z-mixed, the latter is J-antisymmetric and J_Z-physical.

(2) Lemma BB-Z and the Two-Protocol Theorem (DERIVED-CONDITIONAL): The boundary-vs-bulk distinction between BFV anchor states and Boolean handshake superpositions is reduced to a structural location-role assignment on the cobordism tube. Wilson loop measurement separates into Protocol (a) state-preparation ($P_a = 0.3208$) and Protocol (b) spectral characterization ($P_b = 0.7948$), with consistency audited by the corpus PROVEN sum rule $0.7948 + 0.2050 + 0.0001 \approx 1$.

(3) F-CTA-1 sub-gate set: WL-1 and WL-2 are refined into seven operationally distinguishable sub-gates F-CTA-1a/b/c/d/multi-a/multi-b/-disc. The BFV-anchor multi-cycle quasi-revival pattern P_a^(n) = $0.7948^n\cdot\cos^2(n\cdot 129.45°)$ is a NEW corpus-derived prediction, structurally distinct from the spectral exponential decay P_b^(n) = 0.7948^n. Discriminator points at n = 2, 9 separate the two predictions by factors 27 and 132 respectively.

(4) F-CTA-1d (NEW prediction): The kinematic fixed point $|5\rangle$ measurement predicts near-invariance ≈ $(0.9993)^n$, separated from the Z-sector predictions by a factor of 10^2 to 10^3 at moderate cycle counts. F-CTA-1d provides a third operational sub-gate distinguishable from both Protocol (a) and Protocol (b) measurements.

(5) EFA-Z effective dissipation rate (DERIVED-CONDITIONAL): Standard open quantum system formalism applied to the Z-block reduced dynamics yields $\Gamma_Z = -(1/(2T_{cycle}))\cdot\ln|\lambda|^2$ with dimensionless value $\Gamma_Z\cdot T_{cycle} = 0.1149$, exactly reproducing the corpus PROVEN per-cycle leakage 1

− $|\lambda|^2$ = 0.2052. The CTP action form S_Z^(EFA) preserves full-register unitarity (ZS-F0 Theorem 12.2) at the fundamental level.

Three new falsification gates F-F16.1 through F-F16.3 plus seven sub-gates F-CTA-1a/b/c/d/multi-a/multi-b/-disc are registered. One new OPEN problem OP-F5-1 (BRST-Operational Resolution Gap) is registered as a corpus-internal scope clarification. The CTA-Z v3 framework is presented at HYPOTHESIS-strong status as future-work bridge to a possible cobordism geometric interpretation of Z-sector self-dynamics. Anti-overclaim gate F-F16.4 inherits NC-F11.1 against any phenomenological observer interpretation.

ZS-F16 introduces zero new physical constants. A = 35/437, Q = 11, z*, λ remain LOCKED throughout. The work is a structural addition that closes the operational ambiguity in WL-1 and WL-2 within the v1.0 corpus prior to any planned restructuring; no prior v1.0 paper content is modified.

§12. Acknowledgements & Code Availability

This work was developed with the assistance of AI tools (Anthropic Claude, OpenAI ChatGPT, Google Gemini) for mathematical verification, code generation, and manuscript drafting. The author assumes full responsibility for all scientific content, claims, and conclusions.

Verification script: zs_f16_verify_v1_0.py (released April 2026; 24/24 PASS at 50-digit mpmath precision). Dependencies: numpy, scipy, mpmath. Coverage: locked input verification (V1–V5), Lemma BAS-Z four-invariant verification (V6–V9), Theorem 4.1 BFV anchor matrix element (V10–V13, including the decisive n=2, n=9 sub-gate predictions), anti-numerology irrationality of arg(λ)/2π (V14, V24), EFA-Z dissipation rate $\Gamma_Z \cdot T_{cycle}$ = 0.1149 with sum rule audit (V15–V16), F-CTA-1d kinematic |5⟩ near-invariance (V17–V18), Born-Markov geometric coefficient (V19), ZS-M12 Lyapunov attractor (V20), Wilson eigenvector orthogonality and BFV decomposition (V21–V22), F-CTA-1 sub-gate set enumeration (V23). Roadmap: 32-test extended verification suite planned, expanding Category C (multi-cycle table verification at n = 1 to 20), Category E (full OOC integration cross-check), and Category F (cross-paper consistency with ZS-F0, ZS-F8, ZS-F11, ZS-M1, ZS-A6). The verification script is deposited at github.com/KennyKang-git/zspin/verify_scripts/.

§13. References

[1] K. Kang, ZS-F0 v1.0(Revised): Ontological Bootstrap and Foundational Closure (Z-Spin Cosmology Collaboration, April 2026). [Definition 8.3 BV-BFV functor B_Z; §8.5 Theorem 8.9 Wilson loop partition function; §9.1 three-layer fixed point structure; §9.3 BFV anchor verification; §12.1 Theorem 12.1 partition function vs unitary evolution; §12.2 Theorem 12.2 Hermitian implies unitary; §12.3 Theorem 12.3 sum rule.]

[2] K. Kang, ZS-F1 v1.0: The Z-Spin Action with Non-Minimal Coupling (Z-Spin Cosmology Collaboration, March 2026).

[3] K. Kang, ZS-F2 v1.0: Geometric Impedance A = 35/437 (Z-Spin Cosmology Collaboration, March 2026).

[4] K. Kang, ZS-F5 v1.0: Gauge Symmetry Constraint — Why Q = 11 (Z-Spin Cosmology Collaboration, March 2026).

[5] K. Kang, ZS-F8 v1.0(Revised): Spectral-Protocol Duality and the Boolean Handshake (Z-Spin Cosmology Collaboration, April 2026). [§0 Abstract: (E−R)/√2 = SU(2) singlet; Lemma 5.2.A Stroboscopic Lifting Bridge.]

[6] K. Kang, ZS-F11 v1.0: Operational Observer Coordinate and the Self-Referential Fixed Point (Z-Spin Cosmology Collaboration, April 2026). [Definition 4.1' OOC; Theorem F11.1' three-projection decomposition; NC-F11.1 anti-phenomenological non-claim.]

[7] K. Kang, ZS-M1 v1.0: i-Tetration & Fixed Point (Z-Spin Cosmology Collaboration, March 2026). [HSI Theorem; Remark 1.2 Leaky Wilson Loop $|\lambda|^2 = (\pi^2/4)\cdot\eta_topo$.]

[8] K. Kang, ZS-M3 v1.0: Regge-Holonomy, Immirzi & Z-Telomere; Spinor Phase Gate (Z-Spin Cosmology Collaboration, March 2026). [J seam involution; Lemma 10.1 $D^{(1/2)}(-I) = -I$.]

[9] K. Kang, ZS-M4 v1.0: Berry-Keating Spectral Bridge (Z-Spin Cosmology Collaboration, March 2026).

[10] K. Kang, ZS-M12 v1.0: Auto-Surgery — Singularity Resolution via i-Tetration Dynamics (Z-Spin Cosmology Collaboration, March 2026). [Theorem 2.1 i-tetration flow; Theorem 5.1 Lyapunov function; Perelman-Ricci flow analogy.]

[11] K. Kang, ZS-Q1 v1.0: Geometric Decoherence, CPTP Channels, and the Born Rule from the Z-Spin Action (Z-Spin Cosmology Collaboration, March 2026). [§3.3 Stinespring dilation; Theorem 3.1 Z-mediation theorem.]

[12] K. Kang, ZS-A6 v1.0: Boundary Physics and the Z-Anchor (Z-Spin Cosmology Collaboration, March 2026). [$T_cycle = 2\pi/A$; Z-Anchor $\varepsilon(r_H)=0$ DERIVED.]

[13] K. Kang, ZS-U11 v1.0: Quartet of Q-Protection Channels (Z-Spin Cosmology Collaboration, April 2026). [§11.10.2 Sub-Lemma 11.4.B: no fundamental $L_Z[\Phi_Z]$ in $S[g, \Phi]$.]

[14] H.-P. Breuer and F. Petruccione, The Theory of Open Quantum Systems (Oxford University Press, 2002). [Standard reference for reduced density matrix dynamics, effective master equation, and CTP formalism.]

[15] A. S. Cattaneo, P. Mnev, and N. Reshetikhin, Classical BV theories on manifolds with boundary, Comm. Math. Phys. 332, 535–603 (2014). [BV-BFV formalism on cobordisms with boundary.]

[16] Z-Spin Cosmology v1.0 Public Repository, https://github.com/KennyKang-git/zspin (accessed April 2026).

§14. Version History

v1.0 (April 2026): Initial public release. Consolidated from internal Z-Spin Collaboration free-exploration sessions (April 2026) covering the operational refinement of corpus WL-1 and WL-2 falsification gates. Lemma BAS-Z (Boolean ↔ Bargmann basis identification, DERIVED-CONDITIONAL); Lemma BB-Z (boundary-bulk identification, DERIVED-CONDITIONAL); Theorem 4.1 (Two-Protocol Theorem, DERIVED-CONDITIONAL); F-CTA-1 sub-gate set (seven sub-gates, mixed PROVEN/CONDITIONAL/NEW status); F-CTA-1d kinematic $|5\rangle$ measurement (NEW prediction); EFA-Z effective dissipation rate $\Gamma_Z\cdot T_cycle = 0.1149$ (DERIVED-CONDITIONAL); CTA-Z v3 framework (HYPOTHESIS-strong, future work); OP-F5-1 BRST-Operational Resolution Gap (corpus-internal OPEN problem registration). Three new theory/computation falsification gates F-F16.1 through F-F16.3 plus seven experimental sub-gates registered. Anti-overclaim gate F-F16.4 inherits NC-F11.1. Six non-claims NC-F16.1 through NC-F16.6 explicitly bound the scope; three additional honest-limitation non-claims NC-F16.7 through NC-F16.9 register the boundary between corpus-direct interpretation and this paper's structural extensions. Verification: 24/24 PASS achieved at 50-digit mpmath precision (companion script zs_f16_verify_v1_0.py); 32-test extended suite in roadmap. External label: v1.0 (no version bump in upstream papers per Z-Spin no-deletion convention). Zero new free parameters; $A = 35/437$ remains the sole geometric input.

Appendix A. Verification Test Inventory (24/24 PASS achieved; 32-test extension in roadmap)

Category	Tests	Coverage
A. Lemma BAS-Z	4	Four-invariant verification: (I) dim, (II) singlet, (III) J action, (IV) σ_y action
B. Lemma BB-Z	6	Boundary-bulk identification table consistency, three orthogonality checks, J/J_Z grading
C. F-CTA-1 sub-gates	8	P_a = 0.3208 single-cycle; P_a^(n) for n = 1, 2, 3, 6, 9, 12, 20; mpmath 50-digit
D. EFA-Z Γ_Z derivation	4	Sum rule 0.7948+0.2050+0.0001; Γ_Z·T_cycle = 0.1149; 1−exp(−0.2298)=0.2052; CTP unitarity
E. OOC integration	5	OOC mapping for each F-CTA-1 sub-gate; J_Z-grading consistency; OP-F5-1 registration
F. Cross-paper consistency	5	ZS-F0 §9.1 inner products; ZS-F8 Lemma 5.2.A; ZS-F11 Definition 4.1'; ZS-M1 z*; ZS-A6 T_cycle
TOTAL	**32**	**100% pass rate target**

Appendix B. Numerical Precision Reference (mpmath 50-digit)

All numerical values in this paper are derived from corpus PROVEN constants at mpmath 50-digit precision. The displayed values are truncated to 4–6 significant digits for readability.

Quantity	50-digit value (mpmath)
x* = Re(z*)	0.43828293672703211162076588879503962787....
y* = Im(z*)	0.36059247187138534595029824739543915934....
|z*|² = η_topo	0.32211886347354823017052930472823060789....
Re(λ) = −π·y*/2	−0.56640528103796858419....
Im(λ) = π·x*/2	0.68859357186905938741....
|λ|² = (π²/4)·η_topo	0.79482008292067451632....
arg(λ) (degrees)	129.45274327296788837796....
arg(λ)/360°	0.35959095353602191216... (irrational)
P_a = (Re(λ))²	0.32081294159700791042....
1 − |λ|²	0.20517991707932548367....
−ln(|λ|²)	0.22980133316528773049....
Γ_Z · T_cycle	0.11490066658264386524....