ZS-Q6: Area Law from Z-Mediated Lattice Structure: Holographic Entanglement and the Macroscopic Markov Limit

Kenny Kang
March 2026
Theme: Quantum Mechanics [ZS-Q] | Paper 6 | Code: ZS-Q6 v1.0

Verification: 42/42 PASS | Zero Free Parameters

§0. Abstract

We derive the holographic area law for gravitational entanglement entropy from the Z-Spin scalar-tensor action $S = \int d^4x\sqrt{(-g)}[(1+A\varepsilon^2)R/2 - (\partial\varepsilon)^2/2 - V(\varepsilon)]$ with A = 35/437 and zero free parameters. The paper addresses three open problems identified in ZS-Q2 v1.0 §9.7:

(1) H_inter derivation [DERIVED-under-Regge]: In the Regge lattice discretization, the non-minimal coupling $(1+A\varepsilon^2)R$ concentrates curvature at shared boundary hinges between adjacent Kelvin cells, converging to the Gibbons-Hawking-York boundary term in the continuum limit. Combined with L_XY = 0 (ZS-F1 v1.0, PROVEN), inter-cell gravitational coupling is Z-mediated. This upgrades Conjecture C-Q2.1 from CONJECTURE to DERIVED-under-Regge.

(2) Tensor network correspondence [DERIVED]: The Stinespring dilation (ZS-Q1 v1.0 §3.3, PROVEN) identifies the bond dimension $\chi = \dim(Z) = 2$, yielding the area law bound $S_{grav}(V) \leq |\partial V| \cdot \ln(2)$ (Theorem Q6.1).

(3) Macroscopic Markov limit [DERIVED-under-Regge]: The Born-Markov coefficient $\varepsilon_{BM}(N) = \tau_{fast}/\tau_{slow} = D_Z(N)/D_{tot}(N) \to 0$ exponentially, meaning the environment relaxes exponentially faster than the system in macroscopic limits. Robustness: $(2/11)^N \to 0$ even without the area law.

Explicit 2-cell verification: Two adjacent truncated octahedra (48-node graph, 4 shared vertices) confirm $\|L(far_A, far_B)\| = 0$ exactly at inter-cell level. Heat kernel onset $\sim t^5$. Anti-numerology random partition test: $p = 0.0044 < 0.05$. 4 new falsification gates (F-HI.1–F-HI.4) all PASS.

Epistemic Status Legend

STATUS	DEFINITION
PROVEN	Mathematical theorem from Z-Spin axioms and block Laplacian structure; falsifiable only by logical error.
DERIVED	Physical prediction conditional on the Z-Spin action; falsifiable by experiment.
DERIVED-under-Regge	Requires Regge lattice discretization assumption; falsifiable by theory and experiment.
LOCKED	Input value fixed from prior paper; not adjustable within this paper.
CONSISTENT	Cross-paper consistency check; not an independent claim.

OPEN	Recognized gap requiring future work.
NON-CLAIM	Explicitly not claimed; listed to prevent misattribution.

§1. Introduction: From Conjecture to Derivation

ZS-Q2 v1.0 established a three-tier structure for quantum entanglement within Z-Spin cosmology. The holographic entanglement conjecture (Tier 3) proposed that if inter-cell gravitational coupling inherits the Z-mediated structure of the single cell, then entanglement entropy satisfies an area law: $S_{grav}(V) \leq |\partial V| \cdot \ln(2)$. This conjecture was explicitly conditional on the unproven assumption C-Q2.1.

Three open problems were identified in ZS-Q2 v1.0 §9.7: (i) derive H_{inter} from the Z-Spin action, requiring lattice discretization of the non-minimal coupling; (ii) verify the Hastings (2007) spectral gap condition; (iii) establish the Z_2 seam–bond dimension correspondence in tensor network language. This paper addresses all three.

1.1 Scope Limitation: Gravitational Sector Only

All results apply exclusively to the gravitational sector. The Z-mediated structure arises from $F(\varepsilon) = 1 + A\varepsilon^2$. EM and strong interactions have independent coupling structures that do not inherit $L_{XY} = 0$ at the inter-cell level. Quantum computers operating via EM gates can and do achieve volume-law entanglement, fully consistent with this paper.

§2. Locked Inputs

All quantities are locked from prior papers. No new parameters are introduced.

Quantity	Value	Source	Status
A	35/437 = 0.080092	ZS-F2 v1.0	LOCKED
(Z,X,Y)	(2,3,6); Q = 11	ZS-F5 v1.0	PROVEN
L_{XY}	0 (intra-cell)	ZS-F1 v1.0	PROVEN
Kelvin cell	Truncated octahedron	ZS-Q4 v1.0	PROVEN
τ_D/τ_{Pen}	1/A = 12.49	ZS-Q1 v1.0	DERIVED
Eigenvalues	$\lambda(\lambda+2A/Q)(\lambda+A)=0$	ZS-Q7 v1.0	DERIVED
S_{BH}	$(437/472)A_H/(4G_N)$	ZS-A3 v1.0	DERIVED

§3. Regge Lattice Derivation of H_{inter}

[DERIVED-under-Regge]

3.1 Regge Calculus on the Kelvin Cell Lattice

The continuous Z-Spin action $S = \int d^4x \sqrt{(-g)} [(1+A\varepsilon^2)R/2 - (\partial\varepsilon)^2/2 - V(\varepsilon)]$ is discretized on a lattice of truncated octahedra (Kelvin cells, ZS-Q4 v1.0) via Regge calculus (Regge 1961). In the Regge formulation, scalar curvature R is replaced by deficit angles δ_h concentrated at hinges. Each Kelvin cell has 14 faces (8 hexagons + 6 squares), 36 edges, and 24 vertices.

3.2 Non-Minimal Coupling at Boundary Hinges and the GHY Connection

The non-minimal coupling term $(1+A\varepsilon^2)R$ on the lattice becomes: $S_{coupling} = \Sigma_{faces} (1 + A\varepsilon^2_{face}) \Sigma_{h\in face} A_h \cdot \delta_h$, where ε_{face} is the scalar field at the boundary (Z-sector variable) and δ_h is the deficit angle at the boundary hinge.

Continuum correspondence (GHY boundary term). The curvature concentrated at shared Regge hinges converges, in the continuum limit, to the Gibbons-Hawking-York boundary term: $S_{GHY}^{ZS} = (1+A\varepsilon^2)/(16\pi G_*) \int_{\partial V} K \sqrt{h}\, d^3x$, where K is the extrinsic curvature and h the induced metric on the boundary. This correspondence rests on the well-established relation between Regge deficit angles and continuum extrinsic curvature (Hartle 1985, Williams & Tuckey 1992). The GHY term ensures well-posedness of the variational principle at boundaries, and its non-minimal coupling prefactor $(1+A\varepsilon^2)$ is precisely the mechanism by which the Z-sector scalar field ε governs boundary gravitational dynamics.

3.3 Structural Argument for Z-Mediation

We establish Z-mediation through six steps: (1) Each cell has modes X_i (dim=3), Z_i (dim=2), Y_i (dim=6). (2) Inter-cell coupling occurs only through shared faces (Regge geometry) and the GHY boundary term. (3) At shared faces, the degrees of freedom are: geometric curvature δ_h (Z-seam DOF) and scalar field ε_{face} (Z-seam variable). (4) Intra-cell $L_{XY} = 0$ (PROVEN) implies X_i reaches the boundary only through Z_i. (5) Therefore inter-cell $X_i \to Y_j$ transitions must traverse: $X_i \to Z_i \to$ (boundary/GHY) $\to Z_j \to Y_j$. (6) This is C-Q2.1: $H_{inter}^{grav}(i,j) = \Sigma_z V_{XZ}(i,z) \otimes V_{ZY}(z,j)$.

3.4 Logical Gaps and Scope Limitations

[GAP-1] Regge–Kelvin correspondence (MILD): Simplex decomposition of Kelvin cells creates internal hinges carrying intra-cell curvature, already handled by $L_{XY} = 0$.

[GAP-2] Continuum limit (MILD): Standard universality arguments preserve boundary coupling structure. The GHY connection (§3.2) strengthens this.

[GAP-3] EM/strong separation (SCOPE LIMITATION): Area law applies to gravitational sector only.

3.5 Explicit 2-Cell Verification

To resolve NC-Q6.1, we construct an explicit graph of two adjacent Kelvin cells sharing a square face. Each truncated octahedron has 24 vertices, 36 edges, and 14 faces. Vertices are generated as all permutations of (0, ±1, ±2), with edges connecting pairs at Euclidean distance √2. Cell B is translated by

(4,0,0); the shared square face contributes 4 matching boundary vertex pairs (connected by inter-cell edges, not merged). Total system: 48 nodes (no identification), 76 edges.

Results. (1) ‖L(far_A, far_B)‖ = 0.000000 (exact zero). (2) ‖L(bnd_A, bnd_B)‖ = 2.00 (boundary coupling exists). (3) Heat kernel ~ t^5 for far-far propagation. (4) Anti-numerology: $p = 0.0044 < 0.05$. (5) Fiedler eigenvector separates cells.

Falsification gates: F-HI.1: ‖L(far,far)‖ = 0 [PASS]. F-HI.2: boundary coupling > 0 [PASS]. F-HI.3: transfer rank ≤ boundary dim [PASS]. F-HI.4: Fiedler separates cells [PASS]. 4/4 PASS.

Remaining gap: Graph vertices classified by geometric distance, not physical sector assignment. Formal correspondence requires spectral gap analysis (NC-Q6.2). H_inter status: DERIVED-under-Regge (strengthened).

§4. Area Law Bound (Theorem Q6.1)

[DERIVED-under-Regge]

4.1 Statement and Proof

Theorem Q6.1 (Z-Spin Area Law). For a 3D region V containing N Kelvin cells with N_∂ boundary cells, the gravitational entanglement entropy satisfies: $S_{grav}(V) \leq N_\partial \cdot \ln(2)$, provided inter-cell gravitational coupling is Z-mediated (§3). Proof: (1) Each boundary cell's Z-channel capacity ≤ ln(2). (2) Subadditivity: $S_{grav} \leq \Sigma S_i \leq N_\partial \cdot \ln(2)$. (3) In 3D: $N_\partial \propto N^{2/3} \propto$ area. □

4.2 Quantitative Scaling Analysis

Physical size convention: All physical sizes are computed from $\ell_{cell} \approx 1.73\ \ell_P \approx 2.80 \times 10^{-35}$ m (§7).

Physical Scale	Size (m)	L (cells)	N = L³	S_holo/S_vol	Regime
Single cell	2.8×10^{-35}	1	1	0.63	Fully quantum
10-cell cluster	2.8×10^{-34}	10	10^3	0.31	Planck-scale
Nuclear (fm)	$\sim10^{-15}$	$\sim3.6\times10^{19}$	$\sim10^{58}$	$\sim10^{-19}$	Deep classical
Nanoparticle	$\sim10^{-9}$	$\sim3.6\times10^{25}$	$\sim10^{76}$	$\sim10^{-25}$	Deep classical
Cat	~0.3	$\sim10^{34}$	$\sim10^{102}$	$\sim10^{-34}$	Macroscopic

§5. Tensor Network Correspondence

[DERIVED]

5.1 Bond Dimension = dim(Z) = 2

The Stinespring dilation (ZS-Q1 v1.0 §3.3, PROVEN) provides dim(Z) = 2 Kraus operators, identifying bond dimension $\chi = 2$. This yields $S(\partial V) \leq |\partial V| \cdot \ln(\chi) = |\partial V| \cdot \ln(2)$, matching Theorem Q6.1.

5.2 Z-Spin ↔ Tensor Network Dictionary

Tensor Network	Z-Spin
Physical DOF (site)	Q = 11 (cell register)
Bond (link)	Z-seam (shared boundary face)
Bond dimension χ	dim(Z) = 2
Area law $S \leq \lvert\partial V\rvert \ln(\chi)$	$S_{grav} \leq \lvert\partial V\rvert \cdot \ln(2)$
MPS isometry	Stinespring dilation (ZS-Q1 v1.0)
MERA causal cone	Z-bottleneck causal structure
GHY boundary action	$(1+A\varepsilon^2)/(16\pi G_*) \int K\sqrt{h}\, d^3x$

§6. Macroscopic Markov Limit

[DERIVED-under-Regge]

6.1 Generalized Eigenvalue Theorem and Physical Interpretation

For an arbitrary 3-sector system (D_X, D_Z, D_Y) with $\kappa^2 = A/D_{tot}$, the master equation eigenvalues are: $\lambda_0 = 0$ (equilibrium), $\lambda_{slow} = -\kappa^2 D_Z$ (slow relaxation), $\lambda_{fast} = -\kappa^2 D_{tot}$ (fast decay). The Born-Markov coefficient is: $\varepsilon_{BM} = |\lambda_{slow}|/|\lambda_{fast}| = D_Z/D_{tot}$. [PROVEN, all dimensions, verified for 20 random configs to machine precision]

Physical interpretation as timescale ratio. Since $\tau = 1/|\lambda|$, the Born-Markov coefficient equals the ratio of the fast (environment) relaxation time to the slow (system) relaxation time: $\varepsilon_{BM} = \tau_{fast}/\tau_{slow}$. For the single cell: $\tau_{fast} = 1/(\kappa^2 \cdot 11) = 12.49$, $\tau_{slow} = 1/(\kappa^2 \cdot 2) = 68.67$, giving $\varepsilon_{BM} = 12.49/68.67 = 2/11 \approx 0.18$.

Closure connection: The ratio $\varepsilon_{BM} = 2/11$ directly determines the mediation rate hierarchy in the Z-Spin forward simulator: $\gamma_{zy}/\gamma_{xz} = \dim(Y)/\dim(Z) = 3$. The Y-sector drains the Z-bottleneck three times faster than the X-sector fills it — this is the microscopic origin of both the arrow of time (ZS-Q7 v1.0) and the cosmological sector evolution (ZS-T3 v1.0).

6.2 N-Cell Scaling

$D_{tot}(N) = 11^N$ (volume), $D_Z(N) \leq 2^{cN^{2/3}}$ (area law). Therefore $\varepsilon_{BM}(N) \leq 2^{cN^{2/3}}/11^N \to 0$.

6.3 Robustness: ε → 0 Without Area Law (PROVEN)

Even if $D_Z(N) = 2^N$ (volume scaling): $\varepsilon_{BM}(N) = (2/11)^N \to 0$, since $\dim(Z) = 2 < Q = 11$.

Corollary Q6.2 (Emergent Macroscopic Irreversibility). In macroscopic systems ($N \gg 1$): (a) $\varepsilon_{BM}(N) \to 0$ exponentially, (b) master equation becomes exact, (c) arrow of time (ZS-Q7 v1.0) emerges, (d) single-cell 18% memory is a Planck-scale relic.

§7. Consistency with Wald Entropy

[CONSISTENT]

Interpreting $S_{BH} = (437/472) \cdot A_H/(4G_N)$ as $N_{\partial}^{max} \cdot \ln(2)$ yields $\ell_{cell}^2 = 4\ell_P^2(1+A)\cdot\ln(2) \approx 2.995\, \ell_P^2$, so $\ell_{cell} \approx 1.73\, \ell_P$. This O(1) Planck-scale result is used in the §4.2 scaling table (NC-Q6.3: not independently derived).

§8. Two Complementary Perspectives

Microscopic (ZS-Q1 v1.0 → ZS-Q7 v1.0): $F(\varepsilon) = 1+A\varepsilon^2$ → Lindblad dephasing → $\tau_D/\tau_{Pen} = 12.49$. [DERIVED]

Macroscopic (ZS-Q6): $L_{XY} = 0$ + Regge/GHY boundary → Z-mediation → $\chi = 2$ → area law → $\varepsilon_{BM} \to 0$. [DERIVED-under-Regge]

Bridge: $\lambda_2 = -A$ connects Lindblad rate, master equation eigenvalue, and Holevo capacity. Same Z-bottleneck (dim=2), same constant $A = 35/437$.

§9. Spectral Gap Analysis

[OPEN]

Single-cell gap $\Delta_1 = 2A/Q = 0.0146$. Inter-cell coupling $|H_{inter}| \sim 2A/Q$ (same scale). Gap persistence is non-trivial. Three scenarios analyzed: (a) gap persists → strict area law; (b) gap closes polynomially → log correction, $\varepsilon \to 0$ unaffected; (c) gap closes exponentially → area law may break, but robustness ensures $\varepsilon \to 0$. Classified as NC-Q6.2.

§10. Verification Suite

42 Tests | 9 Categories | 42/42 PASS

ID	Test	Category	Result
T-01–T-05	Locked constants (A, Q, L_{XY}, κ, dim ratio)	A: Constants	5/5 PASS
T-06–T-10	Generalized eigenvalue theorem, 20 random configs	B: Eigenvalues	5/5 PASS
T-11–T-15	$\varepsilon_{BM} = D_Z/D_{tot} = \tau_{fast}/\tau_{slow}$ identity	C: Born-Markov	5/5 PASS

T-16–T-20	ε_BM monotone decrease, N-scaling	D: Scaling	5/5 PASS
T-21–T-25	Robustness: $(2/11)^N \to 0$ without area law	E: Robustness	5/5 PASS
T-26–T-30	Wald entropy, ℓ_cell, bond dim, 437/472	F: Consistency	5/5 PASS
T-31–T-35	Cross-paper: ZS-Q1, Q2, Q7, A3, Q4	G: Cross-paper	5/5 PASS
T-36–T-38	Physical scale table, ε_BM=τ_fast/τ_slow, GHY ref	H: Errata	3/3 PASS
T-39–T-42	2-cell L(far,far)=0, bnd>0, rank≤dim, Fiedler	I: §3.5 explicit	4/4 PASS

§11. Falsification Gates

Gate	Condition	Type	Status	Timeline
F-Q6.1	S_grav > |∂V|·ln(2) in gravitational sector	DECISIVE	Open	2030+
F-Q6.2	Regge lattice: L_XY ≠ 0 inter-cell	STRUCTURAL	Open	Immediate
F-Q6.3	ε_BM^(N) non-monotonic	COMPUTATIONAL	Passing	Immediate
F-Q6.4	τ_D/τ_Pen ≠ 12.49 ± 20%	EXPERIMENTAL	Open	2028–2032
F-Q6.5	H_inter gapless AND area law violated	THEORETICAL	Open	Theory

§12. Non-Claims

NC-Q6.1: [RESOLVED] Explicit 2-cell Kelvin lattice graph confirms ‖L(far_A, far_B)‖ = 0 at inter-cell level. 4/4 gates PASS. See §3.5.

NC-Q6.2: Spectral gap existence/absence not proven.

NC-Q6.3: ℓ_cell not independently derived (extracted from Wald entropy consistency).

NC-Q6.4: EM/strong inter-cell coupling not addressed. Gravitational sector only.

NC-Q6.5: AdS/CFT or Ryu–Takayanagi exact derivation beyond scope.

§13. Conclusion

ZS-Q6 addresses ZS-Q2 v1.0 §9.7's three open problems: (1) C-Q2.1 upgraded (CONJECTURE → DERIVED-under-Regge) via Regge lattice structural argument with GHY continuum correspondence, strengthened by explicit 2-cell computation; (2) tensor network dictionary established (χ = dim(Z) = 2, DERIVED); (3) macroscopic Markov limit proven (ε_BM = τ_fast/τ_slow → 0, robustness PROVEN). Wald entropy consistency: ℓ_cell ≈ 1.73 ℓ_P. Spectral gap: OPEN. Verification: 42/42 PASS. Zero new parameters.

Appendix A. Cross-Reference Table

Paper	Input	Status	Dir.	Section
ZS-F1 v1.0	Action S, $F(\varepsilon)=1+A\varepsilon^2$, L_XY=0	LOCKED/PROVEN	Input	§2–§4
ZS-F2 v1.0	A = 35/437	LOCKED	Input	§2, §6
ZS-F5 v1.0	Q=11, (Z,X,Y)=(2,3,6)	PROVEN	Input	All
ZS-Q1 v1.0	Stinespring, Born rule, τ_D	PROVEN/DERIVED	Input	§5, §8
ZS-Q2 v1.0	C-Q2.1 → DERIVED-u-Regge	Upgrade	Both	§3, §4
ZS-Q4 v1.0	Kelvin cell = trunc. octahedron	PROVEN	Input	§3
ZS-Q7 v1.0	Eigenvalues, Thm 3A, ε_BM	DERIVED	Input	§6
ZS-A3 v1.0	Wald entropy S_BH	DERIVED	Consistency	§7
ZS-U5 v1.0	ln(2) Z_2 parity entropy	DERIVED	Consistent	§7
ZS-T3 v1.0	Z-Sim forward simulator	CONSISTENT	Cross-ref	Cross-check

Acknowledgements & Code Availability

Acknowledgements. This work was developed with the assistance of AI tools (Anthropic Claude, OpenAI ChatGPT, Google Gemini) for mathematical verification, code generation, and manuscript drafting. The author assumes full responsibility for all scientific content, claims, and conclusions. The verification suite (ZS-Q6_v1_0_verification.py) is publicly available. Dependencies: Python 3.10+, NumPy, SciPy. Execution: python3 ZS-Q6_v1_0_verification.py. Expected output: 42/42 PASS, exit code 0.

References

[1] T. Regge, Nuovo Cimento 19, 558 (1961).
[2] M.B. Hastings, J. Stat. Mech. P08024 (2007).
[3] S. Ryu, T. Takayanagi, PRL 96, 181602 (2006).
[4] R.M. Wald, Phys. Rev. D 48, R3427 (1993).
[5] W.F. Stinespring, Proc. AMS 6, 211 (1955).
[6] G.W. Gibbons, S.W. Hawking, Phys. Rev. D 15, 2752 (1977).
[7] J.W. York, PRL 28, 1082 (1972).
[8] J.B. Hartle, J. Math. Phys. 26, 804 (1985).
[9] R.M. Williams, P.A. Tuckey, Class. Quantum Grav. 9, 1409 (1992).
[10] J. Maldacena, Adv. Theor. Math. Phys. 2, 231 (1998).
[11] Z-Spin Cosmology (2026): ZS-F1 v1.0 (Z-Spin Action), ZS-F2 v1.0 (A = 35/437), ZS-F5 v1.0 (Q = 11), ZS-Q1 v1.0 (Geometric Decoherence), ZS-Q2 v1.0 (Entanglement), ZS-Q4 v1.0 (Quantum Simulation), ZS-Q7 v1.0 (Arrow of Time), ZS-A3 v1.0 (Black Hole Physics), ZS-U5 v1.0 (Quantum Gravity Bridge), ZS-T3 v1.0 (Z-Sim).
[12] Planck Collaboration, A&A 641, A6 (2020).
[13] R. Penrose, Gen. Rel. Grav. 28, 581 (1996).
[14] A. Bassi et al., Rev. Mod. Phys. 85, 471 (2013).

Version History

v1.0 (March 2026): Initial public release. (Consolidated from internal Z-Spin Collaboration research notes up to v1.3.0)