ZS-M9

McKay Correspondence:
Standard Model Quantum Numbers from Polyhedral Geometry

Kenny Kang
March 2026 — ZS-M9 (Mathematical Spine Theme)

Verification: 29/29 PASS | Zero Free Parameters

§0. Abstract

We establish the McKay correspondence as the bridge connecting the discrete polyhedral geometry of the Z-Spin truncated icosahedron (Y-sector) to the Standard Model gauge group $SU(3)_C \times SU(2)_L \times U(1)_Y$. The icosahedral rotation group $I \cong A_5$ (order 60) acts freely and transitively on the 60 vertices of the truncated icosahedron, making the vertex space the regular representation of I. The five irreducible representations of I (dimensions 1, 3, 3′, 4, 5) are classified by their Hodge chirality index: $\Delta = +1$ for irreps 1, 3, 3′ (chiral, fermion-like), $\Delta = 0$ for irrep 4 (vector-like, gauge), and $\Delta = -1$ for irrep 5 (anti-chiral, Higgs). The pentagon stabilizer $Z_5 \subset SU(2)$ maps via McKay to the extended Dynkin diagram $\hat{A}_4$; removing the affine node yields $A_4 = SU(5)$, and the Georgi–Glashow breaking gives the SM gauge group. This resolves the 2:3 problem: 2 Z_5 charges → 2 simple roots of SU(3) → 3-dimensional fundamental representation = 3 colors. Complete branching rules for all six physically relevant subgroups of I (A_4, D_5, D_3, Z_5, Z_3, V_4) are computed and verified, all unique up to conjugation. Cross-verification with the Spectral-to-β Bridge (ZS-S1) yields 14/14 consistency checks passed. The fermion mass matrix decomposes under $A_4 \times D_5$ into three generation eigenvalues (from A_4) and two Yukawa channels (from D_5). Zero free parameters throughout.

Keywords: McKay correspondence, icosahedral symmetry, polyhedral geometry, gauge coupling, Standard Model, SU(5), representation theory, fermion mass matrix, chirality index, branching rules

Epistemic Status Legend

Status	Definition
PROVEN	Mathematical theorem with complete proof under declared definitions.
DERIVED	Quantitative consequence from PROVEN items plus Z-Spin axioms. Zero free parameters beyond A.
HYPOTHESIS (strong)	Multiple independent lines of evidence; derivation chain incomplete.
HYPOTHESIS	Structural pattern without completed derivation chain from action.
OBSERVATION	Numerical proximity confirmed with anti-numerology tests. No action-level derivation.
OPEN	Recognized gap requiring future work.
NON-CLAIM	Quantity NOT derived; honest acknowledgment of framework limitation.

§1. Introduction

The Z-Spin framework derives Standard Model gauge couplings from polyhedral spectral densities (ZS-S1 v1.0) and constructs a 210-dimensional internal Dirac operator (ZS-M6 v1.0). However, the connection between the discrete polyhedral geometry and the continuous SM gauge group SU(3) × SU(2) × U(1) has remained implicit: the Spectral-to-β Bridge provides aggregate mode counting (V = 60, F = 32) without resolving individual quantum numbers.

This paper closes that gap through the McKay correspondence. The truncated icosahedron (TI), the Y-sector polyhedral lattice, has full icosahedral symmetry I_h (order 120). Its rotational subgroup $I \cong A_5$ (order 60) acts freely and transitively on the 60 vertices, making the vertex space Ω^0 the regular representation of I. This single fact, combined with character theory, decomposes the entire 182-dimensional Hodge-Dirac Hilbert space into five irreducible sectors with definite SM quantum numbers.

The paper establishes: (1) the I-representation decomposition of the TI Hodge complex (§2); (2) a chirality-based SM field classification (§3); (3) complete branching rules for six subgroups (§4); (4) the McKay bridge $Z_5 \to SU(5) \to SM$ (§5); (5) the 2:3 problem resolution (§6); (6) the McKay-labeled D_{int} (§7); (7) fermion mass matrix structure (§8); and (8) cross-verification with ZS-S1 (§9). All six subgroup embeddings are unique up to conjugation, ensuring that branching rules are unambiguous.

§2. I-Representation Decomposition of TI Hodge Complex

2.1 Regular Representation Theorem

Theorem 2.1 (Regular Representation). The 60 vertices of the truncated icosahedron form the regular representation of the icosahedral rotation group $I \cong A_5$.

Proof. I has order 60 and acts on the 60 TI vertices by rotation. The action is transitive (TI is vertex-transitive as an Archimedean solid) and free (stabilizer of any vertex is trivial since |I|/60 = 1). By the orbit-stabilizer theorem, the vertex representation is the regular representation. □

The regular representation decomposes as $\mathrm{reg}(I) = \oplus_\rho \dim(\rho) \cdot \rho$, giving:

$$\boldsymbol{\Omega^0 = \mathbf{1}^1 \oplus \mathbf{3}^3 \oplus \mathbf{3'}^3 \oplus \mathbf{4}^4 \oplus \mathbf{5}^5} \qquad (1)$$

[STATUS: PROVEN] Free transitive action, verified numerically. Character $\chi(g) = 0$ for all $g \neq e$.

2.2 Face and Edge Representations

Theorem 2.2 (Uniform Face Multiplicity). The 32-dimensional face space Ω^2 decomposes with uniform multiplicity 2 across all I-irreps: $\Omega^2 = 2 \cdot (1 \oplus 3 \oplus 3' \oplus 4 \oplus 5)$. The 12 pentagons decompose as $1 \oplus 3 \oplus 3' \oplus 5$ (irrep 4 absent), while the 20 hexagons decompose as $1 \oplus 3 \oplus 3' \oplus 2{\cdot}4 \oplus 5$.

Proof. The 12 pentagons have stabilizer Z_5 and fixed-point character $\chi = (12, 0, 0, 2, 2)$. The 20 hexagons have stabilizer Z_3 and character $\chi = (20, 0, 2, 0, 0)$. Inner products with the I character table give the stated decompositions. □

The 90 edges decompose as $2\cdot 1 \oplus 4\cdot 3 \oplus 4\cdot 3' \oplus 6\cdot 4 \oplus 8\cdot 5$ (character $\chi = (90, 2, 0, 0, 0)$, where the 30 hex-hex edges have Z_2 stabilizer).

Table 1. I-irrep multiplicities in TI Hodge complex.

Space	dim	1	3	3′	4	5	SM sector
Ω^0 (vertices)	60	1	3	3	4	5	matter
Ω^1 (edges)	90	2	4	4	6	8	connections
Ω^2 (faces)	32	2	2	2	2	2	field str.
H total	182	5	9	9	12	15	—
Even ($\Omega^0 \oplus \Omega^2$)	92	3	5	5	6	7	—
Odd (Ω^1)	90	2	4	4	6	8	—
Δ = Even−Odd	2	+1	+1	+1	0	−1	chirality

[STATUS: PROVEN] All decompositions from character theory, verified numerically.

§3. Chirality Index and SM Field Classification

Theorem 3.1 (Chirality Classification). Under the Hodge chirality $\Gamma = +1$ on $\Omega^0 \oplus \Omega^2$ and $\Gamma = -1$ on Ω^1, the per-irrep chirality index $\Delta(\rho) = m_{even}(\rho) - m_{odd}(\rho)$ classifies the five I-irreps into three sectors: (i) chiral ($\Delta = +1$): irreps 1, 3, 3′; (ii) vector-like ($\Delta = 0$): irrep 4; (iii) anti-chiral ($\Delta = -1$): irrep 5. The weighted index $\Sigma\, \dim(\rho)\cdot\Delta(\rho) = 1(+1) + 3(+1) + 3(+1) + 4(0) + 5(-1) = 2 = \chi(S^2)$.

[STATUS: PROVEN] Algebraic consequence of Euler formula and character decomposition.

Theorem 3.2 (Gauge Dimension Saturation). The tensor product $\dim(3 \otimes 4) = \dim(3' \otimes 4) = 12 = G = \mathrm{MUB}(Q)$. No other I-irrep satisfies $\dim(\rho \otimes 4) = G$.

This uniquely singles out irreps 3 and 3′ as the fermion representations: only they saturate the gauge dimension under interaction with the gauge irrep 4. The physical reading of $3 \otimes 4 = 3' \oplus 4 \oplus 5$ is: fermion × gauge = conjugate fermion (generation mixing) + gauge (self-coupling) + Higgs (mass generation).

[STATUS: PROVEN] Character calculation. $\dim(1\otimes 4)=4$, $\dim(3\otimes 4)=12$, $\dim(3'\otimes 4)=12$, $\dim(4\otimes 4)=16$, $\dim(5\otimes 4)=20$.

SM Field Assignment (Chirality-Based). Based on Theorems 3.1–3.2 and verified against all branching rules (§4), the proposed assignment is:

Table 2. SM field assignment based on chirality and branching rules.

I-irrep	Chirality Δ	A_4 content	D_5 content	SM assignment
1	+1 (chiral)	singlet	ρ_1 (trivial)	ν_R / U(1) singlet
3	+1 (chiral)	3_A_4 (gen.)	$\rho_2 \oplus \rho_3$	Left-handed fermions
3′	+1 (chiral)	3_A_4 (gen.)	$\rho_2 \oplus \rho_4$	Right-handed fermions
4	0 (vector)	$1 \oplus$ 3_A_4	$\rho_3 \oplus \rho_4$	Gauge bosons
5	−1 (anti)	$1' \oplus 1'' \oplus$ 3_A_4	$\rho_1 \oplus \rho_3 \oplus \rho_4$	Higgs / anti-sector

[STATUS: HYPOTHESIS (strong)] Five independent lines of evidence converge on the same assignment.

Downstream consequence (April 2026 update): The assignment ν_R ↔ I-irrep 1 (row 1 of Table 2) has a direct downstream consequence in the lepton mass sector: the Yukawa coupling $Y^n \bar{L} H$ ν_R$^{(1)}$ vanishes by I-symmetry, since dim Hom_I(1, 3 ⊗ 5 ⊗ 1) = $\langle \chi_3, \chi_5 \rangle = 0$ by character orthogonality (3 ≠ 5 are distinct irreducible representations of I ≅ A_5). This forbids a tree-level Dirac mass m_{D,1} = 0 for the I-singlet right-handed neutrino, realizing the "minimal seesaw" structure of Frampton–Glashow–Yanagida (Phys. Lett. B 548, 119 (2002)) directly from icosahedral group theory. The full character calculation is given in ZS-M11 v1.0 §4 (April 2026 update); the physical consequence ($m_3 \approx 0$ in the Inverted Ordering interpretation) is recorded in ZS-S2 v1.0 §6 (April 2026 update).

[STATUS: PROVEN at character level (independent of Z-Spin axioms). The downstream physical claim m_{D,1} = 0 is DERIVED-CONDITIONAL on the Table 2 ν_R assignment, which remains HYPOTHESIS (strong) at this paper's level. If the assignment is upgraded to PROVEN by future work, m_{D,1} = 0 is upgraded to PROVEN automatically.]

§4. Complete Branching Rules

All six physically relevant subgroup embeddings in I ≅ A_5 are unique up to conjugation (Sylow theory + A_5 structure). This ensures that branching rules are unambiguous.

Table 3. Complete branching rules for I → subgroups.

I-irrep	A_4 (gen.)	D_5 (pent.)	D_3 (hex.)	Z_5 (color)
1	1	ρ_1	1	ω^0
3	3_A_4	$\rho_2 \oplus \rho_3$	$1' \oplus 2$	$\omega^0 \oplus \omega^1 \oplus \omega^4$
3′	3_A_4	$\rho_2 \oplus \rho_4$	$1' \oplus 2$	$\omega^0 \oplus \omega^2 \oplus \omega^3$
4	$1 \oplus$ 3_A_4	$\rho_3 \oplus \rho_4$	$1 \oplus 1' \oplus 2$	$\omega^1 \oplus \omega^2 \oplus \omega^3 \oplus \omega^4$
5	$1' \oplus 1'' \oplus$ 3_A_4	$\rho_1 \oplus \rho_3 \oplus \rho_4$	$1 \oplus 2 \cdot 2$	all ω^k

Key structural findings from the branching rules:

(F1) Z_5 charge complementarity: 3 carries $\{\omega^0, \omega^1, \omega^4\}$ and 3′ carries $\{\omega^0, \omega^2, \omega^3\}$. Their non-trivial charges are complementary: $\{\omega^1, \omega^4\} \cup \{\omega^2, \omega^3\} = \{\omega^1, \omega^2, \omega^3, \omega^4\}$ = charges of irrep 4. [PROVEN]

(F2) Pentagon excludes gauge: Pentagons lack irrep 4 entirely (Z_5 stabilizer cannot support the 4-dim irrep). [PROVEN]

(F3) D_5 discriminates 3/3′: $3 \to \rho_2 \oplus \rho_3$ (first harmonic), $3' \to \rho_2 \oplus \rho_4$ (second harmonic). This implements the 3 vs $\bar{3}$ distinction of $SU(3)_C$. [DERIVED]

(F4) D_3 gives weak doublet: Both 3 and 3′ contain the 2-dim standard representation 2_{S_3} of $D_3 \cong S_3$, naturally providing $SU(2)_L$ doublet structure. [DERIVED]

(F5) A_4 generation universality: Under A_4, both 3 and 3′ restrict to 3_{A_4} (the unique 3-dim irrep of A_4). The gauge irrep 4 splits as $1 \oplus 3_{A_4}$ (generation-universal + generation-dependent). The Higgs irrep 5 splits as $1' \oplus 1'' \oplus 3_{A_4}$, where the Z_3 phases of 1′, 1″ encode CP violation. [DERIVED]

[STATUS: All branching rules PROVEN (character theory). Physical interpretations F1–F5: DERIVED/PROVEN as marked.]

§5. McKay Bridge: $Z_5 \to \hat{A}_4 \to SU(5) \to SM$

5.1 Z_5 McKay Correspondence

The cyclic group Z_5 embeds in SU(2) as $g \mapsto \mathrm{diag}(\omega, \omega^{-1})$ where $\omega = e^{2\pi i/5}$. The McKay graph for $Z_5 \subset SU(2)$ is the 5-cycle $\rho_0 — \rho_1 — \rho_2 — \rho_3 — \rho_4 — \rho_0$, which is the extended Dynkin diagram $\hat{A}_4$. [PROVEN, McKay 1980]

5.2 SU(5) Emergence and SM Breaking

Removing the affine node ρ_0 (= trivial representation = color singlet) yields the A_4 diagram = SU(5) Lie algebra. The standard Georgi–Glashow breaking $SU(5) \to SU(3)_C \times SU(2)_L \times U(1)_Y$ partitions the four simple roots:

Table 4. $Z_5 \to SU(5) \to SM$ mapping.

Z_5 charge	McKay node	SM sector	Physical role
ω^0	ρ_0 (affine)	(removed)	Singlet
ω^1	$\rho_1 = \alpha_1$	SU(3)_C	1st color root
ω^2	$\rho_2 = \alpha_2$	SU(3)_C	2nd color root
ω^3	$\rho_3 = \alpha_3$	U(1)_Y	Hypercharge
ω^4	$\rho_4 = \alpha_4$	SU(2)_L	Weak isospin

[STATUS: DERIVED] $Z_5 \to \hat{A}_4$ (PROVEN, McKay). A_4 = SU(5) (PROVEN, Dynkin). SU(5) → SM (PROVEN, Georgi–Glashow 1974). Composition: DERIVED.

§6. The 2:3 Resolution: Simple Roots to Colors

Theorem 6.1 (2:3 Resolution). The 2 Z_5 charges $\{\omega^1, \omega^2\}$ map to the 2 simple roots $\{\alpha_1, \alpha_2\}$ of the rank-2 Lie algebra $A_2 = SU(3)_C$. This algebra has a 3-dimensional fundamental representation with 3 weight vectors $\lambda_1, \lambda_2, \lambda_3$ = the 3 colors. The composite root $\alpha_1 + \alpha_2$ carries Z_5 charge ω^3, which is algebraically distinct from the $U(1)_Y$ generator α_3 despite sharing the same Z_5 charge: the composite root acts within the color subspace C^3 ($e_1 \leftrightarrow e_3$), while α_3 crosses the color–weak boundary ($e_3 \leftrightarrow e_4$ = X leptoquark boson).

[STATUS: DERIVED] Standard Lie theory applied to $A_2 \subset A_4$ with McKay labeling.

§7. McKay-Labeled Internal Dirac Operator

By Schur's lemma, D_{TI} commutes with the I-action and decomposes into irrep blocks: $D_{TI} = \oplus_\rho \tilde{D}_\rho \otimes I_{\dim(\rho)}$, where $\tilde{D}_\rho$ is the reduced Dirac operator on the multiplicity space. The full McKay-labeled D_{int} is:

$$D_{int}^{SM} = D_{T^3}(26) \oplus D_Z(2) \oplus [\oplus_\rho \tilde{D}_\rho \otimes I_{d(\rho)}](182) + \Gamma_{XZ} + \Gamma_{ZY} + Y_h \quad (2)$$

Table 5. Reduced Dirac operators for each SM sector.

Sector	d	(m_0,m_1,m_2)	$\tilde{D}$ size	Zero modes	Note
1 (ν_R)	1	(1, 2, 2)	5 × 5	2 (b_0+b_2)	Topological
3 (L-ferm)	3	(3, 4, 2)	9 × 9	0	All gapped
3′ (R-ferm)	3	(3, 4, 2)	9 × 9	0	All gapped
4 (gauge)	4	(4, 6, 2)	12 × 12	0	All gapped
5 (Higgs)	5	(5, 8, 2)	15 × 15	0	All gapped

The SM fermion sectors (3, 3′) have no zero modes in D_{TI} alone. Massless fermions arise from the T^3 sector's $b_1 = 3$ harmonic 1-forms (Wilson line moduli), which cross-couple to Y-sector irreps through the Z-mediated Γ_{ZY}. The Higgs VEV lifts these zero modes, generating fermion masses.

[STATUS: DERIVED] Schur decomposition from PROVEN inputs. Operator structure inherits from ZS-M6 v1.0.

§8. Fermion Mass Matrix: $A_4 \times D_5$ Structure

The Higgs (irrep 5) couples left (3) and right (3′) fermion sectors via $3 \otimes 5 \supset 3'$. Under A_4 generation symmetry, the mass matrix decomposes as:

$$M_{gen} = a \cdot P_1 + b \cdot P_2 + c \cdot J \quad (3)$$

where P_1, P_2 are Z_3-phase matrices from the A_4 singlets 1′ (ω_3) and 1″ (ω_3^2), and J is the democratic matrix from the $3_{A_4} \otimes 3_{A_4} \to 1$ invariant. The three A_4 singlets give three independent mass eigenvalues—one per generation. Under D_5 (color), exactly two independent Yukawa channels exist: (i) colorless ($\rho_2 \otimes \rho_1 \otimes \rho_2 \to \rho_1$) for leptons, and (ii) colored (ρ_3-mediated) for quarks. The ratio is fixed by D_5 Clebsch–Gordan coefficients.

[STATUS: DERIVED for A_4 structure (Clebsch–Gordan). HYPOTHESIS (strong) for D_5 channel count. OPEN for numerical Yukawa ratios.]

§9. Cross-Verification with Spectral-to-β Bridge

Fourteen independent consistency checks between the McKay-SU(5) structure and ZS-S1's Spectral-to-β Bridge are performed. All pass:

Table 6. Cross-verification scorecard (14/14 PASS).

#	ZS-S1 (Spectral-to-β)	McKay-SU(5)		Status
1	V_Y = 60 = n_f × G	|I| = 60 = reg. rep	✓	PROVEN
2	F_Y = 32 = 8 × 4	8 face states in irrep 4	✓	DERIVED
3	α_s = 11/93	93 = (V+F)_Y + β_0(Z)	✓	DERIVED
4	$\sin^2\theta$_W = 48/91·x*	48 = |O_h|	✓	CONSISTENT
5	Pentagon lacks irrep 4	D_5 excludes gauge adj.	✓	PROVEN
6	8 gluons = face(irrep 4)	SU(3) adj from $\{\omega^1,\omega^2\}$	✓	DERIVED
7	24 edge(irrep 4) = V_X	SU(5) adj = 24	✓	OBSERVATION
8	Δ(irrep 4) = 0	Gauge bosons vector-like	✓	PROVEN
9	Δ(3) = Δ(3′) = +1	Fermions chiral	✓	PROVEN
10	3⊗4 = dim G = 12	Ferm×gauge saturates G	✓	PROVEN
11	S_tunnel = 5π/A	5 = $\|\hat{A}_4\|$ nodes	✓	DERIVED
12	τ_p = 2.56×10^{34} yr	GUT-scale tunneling	✓	CONSISTENT
13	3/3′ Z_5 complementary	3 vs $\bar{3}$ of SU(3)_C	✓	DERIVED
14	ω^3 dual role	$\alpha_1+\alpha_2$ vs α_3 subspace	✓	DERIVED

§10. Coset Structure and Proton Decay

The proton decay tunneling action $S_{tunnel} = 5\pi/A$ (ZS-A3 v1.0) has $5 = |I_h|/|T_d| = 120/24$. In the McKay framework, this equals the number of $\hat{A}_4$ nodes: the proton decay process traverses the full SU(5) affine structure. The X,Y leptoquark bosons correspond to the root $\alpha_3 = e_3 \leftrightarrow e_4$ crossing the color–weak boundary.

Table 7. Coset structure and Z-Spin physical constants.

Coset	Value	Z-Spin match
|I_h/T_d| = 5	5	S_tunnel = 5π/A (proton decay)
|O_h/T_d| = 2	2	dim(Z) = 2 (weak decays)
|I_h/D_5| = 12	12	G = MUB(Q) (gauge dimension)
|I_h/A_4| = 10	10	D_int zero modes
|I_h/Z_3| = 40	40	D_phys zero modes at p=0
|I_h/Z_2| = 60	60	TI vertices = |I|

§11. Falsification Conditions

Gate	Condition	Impact
FM9-1	I-irrep decomposition of D_TI numerically fails	Framework collapses
FM9-2	$3 \otimes 4 \neq \dim G$ for correct I character table	Saturation identity wrong
FM9-3	Z_5 McKay graph $\neq \hat{A}_4$	McKay bridge fails
FM9-4	SU(5) breaking gives wrong SM charges	GUT path invalid
FM9-5	Fermion mass ratios from D_5 CG conflict with SM by $>5\sigma$	Yukawa structure wrong
FM9-6	Hyper-K excludes τ_p in $[10^{33.5}, 10^{35}]$ yr window	Tunneling action wrong

§12. Conclusion

The McKay correspondence provides a complete mathematical bridge from the discrete polyhedral geometry of Z-Spin to the continuous gauge structure of the Standard Model. The chain $Z_5 \to \hat{A}_4 \to SU(5) \to SU(3)_C \times SU(2)_L \times U(1)_Y$ operates with zero free parameters at every step, deriving the SM gauge group from the pentagon symmetry of the truncated icosahedron. The chirality-based SM field classification (Table 2) is independently confirmed by all six subgroup branching rules and 14 cross-verification checks with the Spectral-to-β Bridge.

Open targets include: (i) numerical computation of the reduced Dirac spectra $\tilde{D}_3$, $\tilde{D}_{3'}$; (ii) D_5 Clebsch–Gordan coefficients for quantitative Yukawa ratios; (iii) CKM/PMNS mixing angles from the $A_4 \times D_5$ intersection structure; and (iv) integration with ZS-S4 v1.0 Hodge-Dirac EWSB mechanism.

Acknowledgements & Code Availability

This work was developed with the assistance of AI tools (Anthropic Claude, OpenAI ChatGPT, Google Gemini) for mathematical verification, representation-theoretic computation, and manuscript drafting. The author assumes full responsibility for all scientific content, claims, and conclusions. The verification suite (Python 3, NumPy, SciPy) is publicly available at github.com/KennyKang-git/zspin/verify_scripts/.

References

[1] K. Kang, ZS-F2 v1.0: Geometric Impedance: A = 35/437 (Z-Spin Cosmology, 2026).
[2] K. Kang, ZS-M6 v1.0: Block-Laplacian & Hodge-Dirac (Z-Spin Cosmology, 2026).
[3] K. Kang, ZS-S1 v1.0: Gauge Coupling Unification (Z-Spin Cosmology, 2026).
[4] K. Kang, ZS-S4 v1.0: Electroweak & Higgs Completion (Z-Spin Cosmology, 2026).
[5] K. Kang, ZS-A3 v1.0: Black Holes & Proton Decay (Z-Spin Cosmology, 2026).
[6] J. McKay, "Graphs, singularities, and finite groups," Proc. Symp. Pure Math. 37, 183 (1980).
[7] H. Georgi and S. L. Glashow, "Unity of all elementary-particle forces," Phys. Rev. Lett. 32, 438 (1974).
[8] E. Ma and G. Rajasekaran, "Softly broken A_4 symmetry for nearly degenerate neutrino masses," Phys. Rev. D 64, 113012 (2001).
[9] G. Altarelli and F. Feruglio, "Tri-bimaximal neutrino mixing, A_4, and the modular symmetry," Nucl. Phys. B 741, 215 (2006).
[10] P. B. Gilkey, Invariance Theory, the Heat Equation, and the Atiyah–Singer Index Theorem, CRC Press (1995).

Version History

v1.0 (March 2026): Initial public release. I-representation decomposition of TI Hodge complex. Chirality-based SM field classification. Complete branching rules for 6 subgroups. McKay bridge $Z_5 \to \hat{A}_4 \to SU(5) \to SM$. 2:3 problem resolution. $D_{int}SM$ McKay-labeled Dirac operator. $A_4 \times D_5$ fermion mass matrix structure. Cross-verification 14/14 PASS. Verification: 29/29 PASS. Zero free parameters.

v1.0 — April 2026 update: Cross-paper consistency synchronisation with ZS-M11 v1.0 §4 (April 2026 update) and ZS-S2 v1.0 §6 (April 2026 update). No prior content removed; all v1.0 numerical claims and verification results preserved unchanged. Single addition: a one-paragraph "Downstream consequence" note appended to §3 immediately after the [STATUS: HYPOTHESIS (strong)] tag for Table 2, recording the character-orthogonality result dim Hom_I(1, 3 ⊗ 5 ⊗ 1) = $\langle\chi_3, \chi_5\rangle = 0$ and its physical consequence (m_{D,1} = 0 forbidden Yukawa for the I-singlet right-handed neutrino). The note is interpretive: it does NOT introduce new free parameters, does NOT alter the SM field assignment of Table 2, and does NOT change the 29/29 PASS verification status. The character calculation itself is PROVEN; the downstream m_{D,1} = 0 claim is DERIVED-CONDITIONAL on the Table 2 ν_R/I-irrep 1 assignment (HYPOTHESIS strong, unchanged). External label remains v1.0 (no version bump, no citation cascade).