

# Aquatic Ecology

## Leaf litter density and decomposition in small man-made ponds

--Manuscript Draft--

<b>Manuscript Number:</b>		
<b>Full Title:</b>	Leaf litter density and decomposition in small man-made ponds	
<b>Article Type:</b>	Original Research	
<b>Keywords:</b>	allochthonous; organic matter; lentic; lake	
<b>Corresponding Author:</b>	Kenneth Fortino Longwood University Farmville, VA UNITED STATES	
<b>Order of Authors:</b>	Kenneth Fortino  Leanna Tacik	
<b>Funding Information:</b>	Longwood University Faculty Development Grant	Dr Kenneth Fortino
<b>Abstract:</b>	<p>The input of terrestrial leaf litter into freshwater ecosystems supports aquatic food webs and fuels microbial metabolism. Although the role of leaf litter subsidies to streams have been studied extensively the effect of leaf litter on ecosystem function in lentic systems has received less attention. In particular the impact of leaf litter on trophic dynamics and biogeochemistry of small man-made ponds is virtually unknown, despite the fact that these systems are extremely common and likely represent a substantial modification to watersheds in the North America. We measured the areal density of leaf litter and the rate of leaf litter decomposition in small man-made ponds in central Virginia to determine the size of the leaf litter pool in these systems, the rate at which leaf litter is decomposed, and the extent to which pond characteristics alter leaf litter abundance or processing. We found that the areal density of leaf litter in the ponds ranged between 3.4 and 1179.0 g AFDM m<sup>-2</sup>. The areal density of leaf litter was significantly greater in the littoral zones, however leaf litter was present in the sediments throughout the pond. There was no relationship between the areal density of leaf litter in the sediments and the percent organic matter of the fine sediments, suggesting that leaf litter input is decoupled from bulk sediment organic matter. The decomposition rate of <i>Liriodendron tulipifera</i> leaves in coarse mesh leaf bags ranged between 0.0025 and 0.0035 d<sup>-1</sup>, which is among the slowest litter decomposition rates recorded in the literature for ponds and was unrelated to pond characteristics. Our results indicate that leaf litter is an abundant and persistent pool of organic matter in the sediments of small man-made ponds and it is likely to have a substantial effect on the trophic dynamics and biogeochemistry of these systems.</p>	
<b>Suggested Reviewers:</b>	<p>Amy Rosemond University of Georgia rosemond@uga.edu</p> <p>Michael Danger Universite de Lorraine michael.danger@univ-lorraine.fr</p> <p>Andrew Mehring University of California San Diego Scripps Institution of Oceanography amehring@ucsd.edu</p> <p>Samuel Fey Reed College Feys@reed.edu</p>	
<b>Author Comments:</b>	<p>Dear Editor,</p> <p>Please consider our manuscript "Leaf litter density and decomposition in small man-made ponds" for publication in Aquatic Ecology. Our paper shows that small man-made ponds contain abundant terrestrial leaf litter and that this organic matter source decomposes slowly. Given that small ponds are a very abundant alteration to</p>	

watersheds in many parts of the world, our results indicate that small man-made ponds could represent an important alteration to organic matter cycling in watershed where they occur. Our findings are especially important because the impact of small ponds in general, and small man-made ponds especially on ecosystem processes have not received much study. Our study is only one of few estimates of the density of leaf litter within ponds and since leaf litter represents a key resource to many freshwater communities, having data on its abundance is important for understanding energy flow in pond ecosystems.

Thank you for considering our manuscript for publication and we look forward to your feedback.

# 1    **Leaf litter density and decomposition in small man-made ponds**

2    Kenneth Fortino ([orcid.org/0000-0002-5173-363X](https://orcid.org/0000-0002-5173-363X))

3    Leanna Tacik

4    Biological and Environmental Sciences, Longwood University, USA

## 5    **Corresponding Author:**

6    Kenneth Fortino

7    [fortinok@longwood.edu](mailto:fortinok@longwood.edu)

8    (434) 395 – 2223

9

## 10    **Abstract**

11    The input of terrestrial leaf litter into freshwater ecosystems supports aquatic food  
12    webs and fuels microbial metabolism. Although the role of leaf litter subsidies to  
13    streams has been studied extensively the effect of leaf litter on ecosystem function  
14    in lentic systems has received less attention. In particular the impact of leaf litter on  
15    trophic dynamics and biogeochemistry of small man-made ponds is virtually  
16    unknown, despite the fact that these systems are extremely common and likely  
17    represent a substantial modification to watersheds in the North America. We  
18    measured the areal density of leaf litter and the rate of leaf litter decomposition in  
19    small man-made ponds in central Virginia to determine the size of the leaf litter  
20    pool in these systems, the rate at which leaf litter is decomposed, and the extent to  
21    which pond characteristics alter leaf litter abundance or processing. We found that  
22    the areal density of leaf litter in the ponds ranged between 3.4 and 1179.0 g AFDM  
23    m<sup>-2</sup>. The areal density of leaf litter was significantly greater in the littoral zones,

however leaf litter was present in the sediments throughout the pond. There was no relationship between the areal density of leaf litter in the sediments and the percent organic matter of the fine sediments, suggesting that leaf litter input is decoupled from bulk sediment organic matter. The decomposition rate of *Liriodendron tulipifera* leaves in coarse mesh leaf bags ranged between 0.0025 and 0.0035 d<sup>-1</sup>, which is among the slowest litter decomposition rates recorded in the literature for ponds and was unrelated to pond characteristics. Our results indicate that leaf litter is an abundant and persistent pool of organic matter in the sediments of small man-made ponds and it is likely to have a substantial effect on the trophic dynamics and biogeochemistry of these systems.

## **Introduction**

Ecosystem subsidies (i.e., the movement of resources across ecosystem boundaries (Polis et al. 1997)) are an important part of organic matter cycling in freshwater systems. The reciprocal transfer of resources between aquatic and terrestrial systems is common (Nakano and Murakami 2001; Baxter et al. 2005), however the input of terrestrial organic matter to aquatic systems is an especially significant flux of material since, this subsidy has been shown to support metabolism and secondary production in a majority of lentic and lotic ecosystems (Marcarelli et al. 2011). Organic matter subsidies from terrestrial to aquatic ecosystems are dominated by detrital plant material either as dissolved (DOC) or particulate (POC) organic carbon, and can substantially augment autochthonous organic matter production (Hodkinson 1975a; Gasith and Hosier 1976; Wetzel 1984; Wetzel 1995;

46 Webster and Meyer 1997; Kobayashi et al. 2011; Mehring et al. 2014). Seasonal leaf  
47 fall dominates the POC input into most temperate aquatic systems (Wallace et al.  
48 1999) and this detrital material serves to stabilize variation in aquatic metabolism  
49 (Wetzel 1984).

50 The effects of terrestrial leaf litter subsidies on freshwaters have received the most  
51 attention in small lotic systems (Webster and Benfield 1986). In undisturbed lotic  
52 systems, leaf mass loss begins with leaching, which is then followed by conditioning  
53 of leaf material by microbial consumers, and finally consumption by shredding  
54 macroinvertebrates (Cummins 1974; Gessner et al. 1999). Shredders can have a  
55 particularly large impact on leaf breakdown rate and leaf litter may contribute  
56 substantial material to stream secondary production (Wallace 1997; Graça 2001;  
57 Eggert and Wallace 2003; Creed et al. 2009). Anthropogenic modifications to  
58 watersheds associated with agricultural and urban land use do not consistently  
59 change leaf litter processing rates in the stream channel (Bird and Kaushik 1992;  
60 Huryn et al. 2002; Walsh et al. 2005; Hagen et al. 2006) but can have profound  
61 impacts on the mechanisms of leaf breakdown (Bird and Kaushik 1992; Paul et al.  
62 2006; Imberger et al. 2008) and thus alter the impact of detrital subsidies.

63 Small impoundments (i.e., man-made ponds) are a common anthropogenic  
64 alteration to watersheds globally (Downing et al. 2006; Downing 2010), but their  
65 impact on leaf litter processing has received limited study. Impoundments have  
66 been shown to alter litter processing rates downstream of dams (Short and Ward  
67 1980; Mendoza-Lera et al. 2010; Tornwall and Creed 2016), but estimates of litter

processing within man-made ponds is limited (Table 1). Impoundment dramatically alters the physical, chemical, and biological characteristics of the system. Not only does the dam eliminate flow within the created pond or lake, but temperate ponds typically stratify, producing heterogeneity in oxygen, and other dissolved components (Wetzel 2001). Further, the reduction in flow produces a depositional environment within the pond favoring the accumulation of soft sediments (Wetzel 2001). These changes to the chemical and physical environment of the pond result in substantial differences in the composition of the pelagic and benthic communities between the pond and the former lotic system (Ogbeibu 2002). Given that chemical, physical, and biological (microbial and animal, consumers) conditions are central to leaf decomposition, it is likely that man-made ponds differ substantially from surrounding lotic habitats with respect to leaf litter processing. The abundance of the smallest ponds ( $< 0.1 \text{ km}^2$ ) is more than 2 orders of magnitude greater than even modest sized lakes ( $1 \text{ km}^2$ ), and the number of small man-made ponds is approaching the number of natural ponds (Downing 2010), indicating that small man-made ponds represent a potentially important but understudied alteration to aquatic organic matter cycling.

Our objectives for this study were to quantify the abundance of leaf litter and leaf litter decomposition rate in small ponds in a moderately urbanized region of central Virginia. We hypothesized that the ponds would contain abundant leaf litter and that leaf mass loss would be slow relative to rates typical for lotic systems due to the alteration of the physical, chemical, and biological conditions within the pond. We further hypothesized that man-made ponds of different construction, even when

geographically close, would differ substantially in leaf processing rate, since leaf litter decomposition is affected by temperature, nutrient availability, invertebrate community composition, and temperature (Webster and Benfield 1986) and these factors should be affected by the design and construction of man-made ponds.

## **Methods**

### **Study Site**

All of the ponds used in the study are located in central Virginia and are small man-made ponds (Table 2). The ponds used for the quantification of leaf litter areal density and sediment organic matter content were Lancer Park Pond, Daulton Pond, Woodland Court Pond, and Wilck's Lake. Lancer Park Pond has an earth dam and a permanent inlet and outlet. The pond is almost completely surrounded by second growth forest. Daulton Pond is a headwater pond with an earth dam that does not have a permanent inlet and is likely partially spring-fed. The riparian zone of Daulton Pond is approximately 50% second growth forest and 50% mowed grass. The littoral zone of Daulton Pond is mostly covered in an unidentified reed and cattails (*Typha sp.*). Woodland Court Pond was created by an earth dam that is drained by a stand-pipe. The pond has a permanent inlet and a riparian zone that is about 30% second growth forest. The remaining portion of the riparian zone is minimally landscaped disturbed land associated with an apartment complex. Approximately 50% of the littoral zone of Woodland Court Pond is a patch of cattail (*Typha sp.*). Wilck's Lake is the largest pond in the study and was created as a borrow pit for the construction of a railroad. Wilck's Lake has no obvious inlet but is

drained by a stand-pipe into a permanent outlet. Wilck's Lake is part of a city park and approximately 90% of the lake shoreline is second growth forest and the remaining area is mowed grass.

The ponds used to determine litter decomposition rate were Lancer Park Pond, Daulton Pond, and Campus Pond. Lancer Park Pond and Daulton Pond are described above. Campus Pond is a stormwater retention pond with a permanent inlet that is drained by a stand-pipe and is surrounded by landscaping that consists of small trees and mowed grass. Campus Pond is enclosed by a vertical concrete wall, so it has no natural littoral zone and is nearly uniform in depth.

#### **Leaf Litter Density and Sediment Organic Matter**

To estimate the areal density of leaf litter in the ponds we used an Ekman dredge to collect sediment samples from the littoral and open water regions of each pond. We collected 6 replicate littoral and 6 replicate open water samples from Daulton Pond, Woodland Court Pond, and Wilck's Lake on 13 May 2013, 14 May 2013, and 14 June 2013 respectively. We collected 3 replicate littoral samples and 3 replicate open water samples from Lancer Park Pond on 20 March 2013. Finally we collected 3 littoral and 6 open water samples from Wilck's Lake on 20 February 2013. In all lakes except Wilck's Lake, littoral samples were collected approximately 5 – 10 m from the shoreline but the actual distance was not recorded. In Wilck's Lake, dense overhanging vegetation along the shoreline prevented sampling and so littoral samples were collected between 10 – 20 m from the shore. The open water samples were collected away from the shoreline of the ponds.



The contents of the Ekman was homogenized in a plastic basin and a 10 ml sample of the fine sediments was collected with a 30 ml plastic syringe with its tip cut off (opening diameter = 1 cm). This sediment slurry was then placed in a pre-weighed 20 ml glass scintillation vial and dried at 50° C for at least 24 h. The remaining material in the basin was sieved through a 250  $\mu$ m mesh in the field and the material retained by the sieve was preserved in 70% ethanol and transported back to the lab. In the lab the preserved material was passed through a 1 mm sieve and macroinvertebrates were removed. All remaining material retained by the sieve was dried at 50 °C for 48 h and homogenized with a mortar and pestle. The dried fine sediments and a subsample of the homogenized leaf litter were each ashed at 550 °C for 4 h to determine the proportion of organic matter in the sample via loss on ignition (LOI). To calculate the ash-free-dry-mass (AFDM) of the total leaf litter of the sample the total dry mass was multiplied by the proportion of organic matter in the sample. The areal density of leaf litter in the pond was then estimated by normalizing the AFDM of the leaf litter to a square meter. We did not estimate the areal mass of organic matter in the fine sediments because we did not have the total dry mass of the sediments collected by the Ekman dredge.

### **Leaf Litter Decomposition**

To determine the leaf litter decomposition rate in the ponds we measured the mass loss rate of tulip poplar (*Liriodendron tulipifera*) leaf bags. Tulip poplar was chosen for the litter species because it is common in the riparian zone of all of the ponds in the study. The litter was collected by gently pulling senescent leaves from the tree.

Only leaves that released without resistance were used. The leaves were all collected and air-dried during the fall of 2013. The leaf bags were assembled by placing 5.0 g of intact leaves into plastic produce bags with approximately 9 mm<sup>2</sup> mesh. The bags were sealed with a zip-tie, attached to a small bag of rocks that served as an anchor, and placed into the littoral zone of Campus Pond and Daulton Pond on 22 October 2013 and into the littoral zone of Lancer Park Pond on 29 October 2013. To determine the mass lost due to handling and deployment, 5 bags were immediately harvested following deployment at each site. Bags were harvested by gently moving the bag into a 250  $\mu$ m mesh net underwater and then gently lifting from the pond. The bag and any material retained in the net were then placed into an 11.4 L resealable plastic bag and returned to the lab. The contents of the bag was gently rinsed over a 1 mm mesh sieve to remove macroinvertebrates and then placed into a pre-weighed paper bag and dried at 50° C for at least 48 h. The dried leaf material was then weighed and homogenized with a mortar and pestle. This homogenized material was then ashed at 550° C to determine the AFDM of the leaves. Following the initial sampling, 5 haphazardly chosen leaf bags were sampled from each pond regularly using the same procedure. The number of days that the remaining leaves were incubated in each pond is shown in Table 2.

### **Statistical Analysis**

Differences in areal leaf litter density among ponds and between the littoral and open water zones of all ponds was determined using ANOVA. The leaf litter density was natural log transformed to homogenize the variance in the test of pond

differences and for the test between the littoral and open water samples. Specific differences among ponds were assessed with a Tukey HSD post-hoc test. The relationship between areal leaf litter density and the percent organic matter of the sediments was assessed using linear regression.

The decay coefficient ( $k$ ) for the leaves in the litter bags in each pond were determined by calculating the slope of the relationship between the natural log of the percent leaf mass remaining by the number of days in the pond (Benfield 2007). All statistical analysis was performed using R (R Core Team 2014).

## **Results**

### **Leaf Litter Density and Sediment Organic Matter**

The areal density of leaf litter in the ponds ranged between 0.00344 and 1.179 kg AFDM  $m^{-2}$ . The greatest areal leaf litter densities were found in Daulton Pond and Lancer Park Pond but in both cases the greatest areal densities were rather exceptional values (Fig. 1). The total areal leaf litter density (i.e., littoral and open water combined) differed significantly among the ponds ( $F_{3, 38} = 3.955$ ,  $p = 0.015$ ). The greatest areal leaf litter density was found in Lancer Park Pond with a mean ( $\pm 1$  SD) density of  $0.399 (\pm 0.436)$  kg AFDM  $m^{-2}$ . However the areal leaf litter density of Lancer Park Pond was only significantly different from Woodland Court pond, which had a mean ( $\pm 1$  SD) areal density of  $0.036 (\pm 0.055)$  kg AFDM  $m^{-2}$ . The mean ( $\pm 1$  SD) areal leaf litter density of Daulton Pond and Wilck's Lake were  $0.175 (\pm 0.344)$  and  $0.148 (\pm 0.194)$  kg AFDM  $m^{-2}$ , respectively and were not significantly different than each other or the other ponds.

In all the ponds, the greatest areal leaf litter densities were found in the littoral portion of the pond (Fig 1). Across all the ponds areal leaf litter density of the littoral portions of the ponds ranged between 0.0097 and 1.179 kg AFDM m<sup>-2</sup> with a mean ( $\pm 1$  SD) areal density of 0.283 ( $\pm 0.347$ ) kg AFDM m<sup>-2</sup>, which is significantly greater ( $F_{1,40} = 28, p < 0.0001$ ) and much more variable than the areal leaf litter density of the open areas of the ponds, which ranged between 0.0034 and 0.215, with a mean ( $\pm 1$  SD) density of 0.030 ( $\pm 0.0479$ ) kg AFDM m<sup>-2</sup> (Fig. 1).

The percent sediment organic matter of the ponds averaged ( $\pm 1$  SD) 10.3 ( $\pm 0.055$ )% across all ponds and ranged between a low of 0.73% in Wilck's lake and a high of 22.3% in Daulton Pond. The mean ( $\pm 1$  SD) percent sediment organic matter of Wilck's Lake was 6.17 ( $\pm 3.65$ )%, which was significantly lower than any of the other ponds ( $F_{3,37} = 6.664, p = 0.001$ ). The percent sediment organic matter of the sediments of Lancer Park Pond and Woodland Court pond were more homogeneous, but not significantly different from the sediments of Daulton Pond (Fig. 2). In all of the ponds, there was no significant difference between the open and littoral sections of the pond ( $F_{1,39} = 0.963, p = 0.333$ ), nor was there a relationship between percent sediment organic matter and the density of leaf litter ( $r^2 = 0.0046, p = 0.714$ )(Fig. 2).

### **Litter Decomposition Rate**

Litter bags were deployed in Daulton Pond, Campus Pond, and Lancer Park Pond for 211, 209, and 204 days respectively. At the end of these incubations the mean ( $\pm 1$  SD) percent of the original 5 g of leaf mass remaining in Daulton Pond, Campus

Pond, and Lancer Park Pond was 45.3 % ( $\pm$  4.7 %), 42.3 % ( $\pm$  8.2 %), 43.2 % ( $\pm$  8.3 %), respectively. The three ponds had similar decay coefficients (k) but Daulton Pond had the lowest rate at 0.0025 d<sup>-1</sup>, followed by Campus pond and Lancer Park Pond with rates of 0.0030 and 0.0035 d<sup>-1</sup>, respectively. All of the litter bags had been colonized by invertebrates but these were not collected quantitatively.

## **Discussion**

Anthropogenic alterations to stream networks can alter leaf litter processing relative to the undisturbed state. Commonly studied anthropogenic disturbances to streams such as urbanization and agriculture have inconsistent impacts on leaf litter processing (Bird and Kaushik 1992; Huryn et al. 2002; Walsh et al. 2005) but little is known about the effect of man-made ponds on leaf litter processing, despite their abundance. Our results show that small man-made ponds collect substantial amounts of terrestrial leaf litter and have some of the slowest leaf litter decomposition recorded in the literature. Thus, small man-made ponds represent an important alteration to organic matter processing and specifically may serve as an organic matter sink within watersheds where they occur.

The areal leaf litter densities measured in the man-made ponds in this study support the observations of other authors that terrestrial detritus represents an important subsidy to lentic systems (Hodkinson 1975b; Gasith and Hosier 1976; Richey et al. 1978; Marcarelli et al. 2011). All of the ponds sampled had measurable leaf litter in their sediments. We are not aware of any other studies that measure leaf litter density in the sediments of man-made ponds in the same size class as we

studied, so it is not clear how representative our measurements are of the leaf litter density of small man-made ponds globally. The only other lentic system for which we were able to find a measure of leaf litter density was for an intermittent swamp (Mehring et al. 2014). In this study the authors report that leaf litter densities range between 1080 g m<sup>-2</sup> following autumn leaf fall to 578 g m<sup>-2</sup> in the summer (Mehring et al. 2014), which is greater than all but the highest littoral values in the ponds we sampled. Although we did not measure the flux of leaf material to the pond, comparisons between the densities we observed and measures of leaf litter inputs also serve to contextualize our observations. Gasith and Hosier (1976) report an input of 1.64 g m<sup>-2</sup> d<sup>-1</sup> of leaf litter into the littoral zone of a Wisconsin lake. In a forested mountain lake Rau (1976) recorded a much lower deposition rate of 0.173 g m<sup>-2</sup> d<sup>-1</sup> and France and Peters (1995) measured an even lower leaf litter flux of approximately 0.04 and 0.02 g m<sup>-2</sup> d<sup>-1</sup> for the littoral zone of 4 lakes in Ontario. The magnitude of these fluxes would not be able to supply the leaf litter densities that we observed in the ponds in our study unless the litter was accumulating over many years. Our litter decomposition rates indicate that 95% of leaf litter mass would be mineralized in between 786 and 1065 days, which indicates that the litter does not persist in these systems for sufficient time for such low deposition rates to be likely. A more likely explanation is that the flux of leaf litter into the ponds in our study is greater than what has been measured in high latitude lakes but not as high as those recorded in the swamp by Mehring et al. (2014).

The greater density of leaf litter in the littoral samples also confirms the findings of other authors that leaf litter accumulates predominantly near the shoreline (Gasith

and Hosier 1976; Rau 1976; France and Peters 1995). Unlike other studies of larger systems (Rau 1976; France and Peters 1995) however, we found measurable leaf litter in the center of the pond. Gasith and Hosier (1976) hypothesize that leaf litter that enters the lake and floats for a period of time before being blown toward the shore and sinking. The presence of measurable leaf litter in the offshore sediments of the lakes in our study may be due to the small surface area of our ponds, which would be insufficiently exposed to wind to exclude floating leaf litter from the open water. This speculation is supported by the observation that the smallest lake in the sample had the most leaf litter in the offshore samples, however the remaining lakes all have a similar amount of offshore leaf litter despite size differences.

The degree to which sediment leaf litter derives from stream inputs in these ponds is unknown but the significance of stream litter inputs is likely a function of stream discharge, litter load, and pond volume. Lancer Park Pond, and Woodland Court Ponds both have permanent first-order stream inlets, which likely serve as a substantial source of litter, especially during high discharge events. Rau (1976) found that litter inputs from intermittent streams around a mountain lake were minor but the system in that study is not likely to be representative of the ponds in our study. Although we know of no other estimation of stream leaf litter input to ponds, the capacity of small streams to transport leaf litter is well known (Bilby and Likens 1980). Despite the mechanisms involved, the presence of leaf litter in the open water sediments of these small ponds indicates that any impact of leaf litter on nutrient cycling and food-web processes extends beyond the littoral zone of small ponds.

291 The degree of variability in leaf litter density within a pond was affected by the  
292 location in the pond. The samples from littoral sediments were much more variable  
293 than those from the open water sediments. The variability of the leaf litter density in  
294 the littoral samples within each pond suggests that the factors affecting leaf litter  
295 accumulation in the sediments are heterogeneous within a lake. Some of this  
296 variation appears to be due to variation in riparian vegetation. France and Peters  
297 (1995) found that riparian vegetation affected litter fall and that litter deposition  
298 increased with the height, girth, and density of riparian trees. Rau (1976) reported  
299 greater litter deposition along forested shorelines, relative to meadow and talus in a  
300 mountain lake. In our study, the samples with the highest littoral leaf litter density  
301 were from in Daulton Pond and Lancer Park Pond. In both lakes these samples came  
302 from regions of the lake with forested riparian zones. Riparian vegetation does not  
303 explain all of the variation in littoral leaf litter density however. The littoral sample  
304 with the lowest leaf litter density in Lancer Park Pond was collected along the same  
305 forested shoreline as the replicates with much greater littoral litter density and  
306 none of the samples collected from a forested shoreline in Woodland Court Pond  
307 had a littoral litter density as high as those found in Daulton Pond or Lancer Park  
308 Pond.

309 Overall leaf litter was a prominent pool of organic matter in all of the small man-  
310 made ponds in the study. Leaf litter alters lentic food webs (Kobayashi et al. 2011;  
311 Cottingham and Narayan 2013; Fey et al. 2015), nutrient cycles (McConnell 1968;  
312 France and Peters 1995), and energy flow (Hodkinson 1975b). The presence of and  
313 variability of leaf litter throughout the sediments of these small man-made ponds is



likely to have profound effects on the ecology and biogeochemistry happening within the pond, and on the role of the pond in the watershed where it occurs.

The fine sediment organic matter content of the pond sediments was strikingly decoupled from the leaf litter density. Overall, the average percent sediment organic matter in the ponds ( $10.3 \pm 0.06 \%$ ) was very similar to the average  $10.7 (\pm 0.05)\%$  sediment organic matter measured in 16 agricultural impoundments in Iowa by (Downing et al. 2008) but less than the more organic rich ( $> 20 \%$  organic matter) *gyttja* typical of productive natural lakes in the temperate zone (Dean and Gorham 1998). The organic matter content of the sediment was not related to the density of leaf litter in the sediments nor did it differ significantly between the littoral and open water samples. These observations suggest that leaf litter inputs may not be an important driver of the variation in percent organic matter in the sediments. We cannot ascertain from our data the degree which leaf litter contributes to total sediment organic matter because the lack of correlation may be due to the redistribution of fine sediment organic matter within the pond obscuring a spatial correlation. Interestingly the two ponds with permanent inlets (Lancer Park Pond and Woodland Court Pond) have the most homogeneous percent sediment organic matter, which may be a result of the higher energy in these systems. Wilck's Lake appears to have a bimodal distribution of sediment organic matter and this is likely due to the fact that this lake was created as a borrow pit, thus the sediments may reflect the historical disturbance of the substrate. The greatest percent sediment organic matter and the greatest variation in sediment organic matter was found in Daulton Pond, which is mainly groundwater fed. This observation may be due to the

337 lack of permanent surface water inputs which would limit the inorganic sediment  
338 load to the lake and maintain higher sediment heterogeneity.

339 The mean ( $\pm$  SD) leaf litter decomposition rate ( $k$ ) measured for all 3 ponds in our  
340 study was 0.0030 ( $\pm$  0.00005) which is lower than the average decay rate of 0.0059  
341  $d^{-1}$  for woody plant litter in lakes in the review by Webster and Benfield (1986) and  
342 lower than what Webster and Benfield (1986) report for Magnoliaceae litter overall.  
343 Our mean decomposition rate was also lower than all but 5 of the 17 observations  
344 made in similar systems collected from the literature (Table 1). All of the studies  
345 with decomposition rates lower than those measured in our ponds came from  
346 boreal systems (Table 1, see Hodkinson 1975a, and Oertli 1993) and of these, 3  
347 were from recalcitrant species (Table 1, see Hodkinson 1975a). Thus the  
348 decomposition rate of *L. tulipifera* litter in our study was among the lowest recorded  
349 rates for woody litter in the literature, and comparable to the litter decomposition  
350 rate high latitude systems.

351 Litter characteristics clearly affect the rate of leaf litter decomposition in aquatic  
352 systems (Webster and Benfield 1986; Gessner 2010), however it is unlikely that the  
353 slow rate of decomposition that we measured was due to the litter choice. Webster  
354 and Benfield (1986) report that Magnoliaceae litter has the second fastest  
355 breakdown rate of the woody plants in their review of breakdown rates, so *L.*  
356 *tulipifera* is not inherently resistant to decomposition.

357 The low rates of decomposition of the leaves in these ponds is likely partially related  
358 to the near absence of shredder activity. Potential shredding taxa (i.e., crayfish)

were observed colonizing the leaf packs in Lancer Park Pond but there was no obvious evidence of shredding on the leaves recovered from any of the ponds (K. Fortino, personal observation). Shredders can dramatically accelerate leaf litter mass loss in streams (Cummins 1974; Webster and Benfield 1986; Wallace et al. 1999) and lakes (Bjelke 2005). The highly limited shredder fauna and the lack of shredder activity may have been due to low oxygen concentration within the leaf packs which could limit shredder colonization and feeding (Bjelke 2005). We did not measure the oxygen availability within the leaf packs but the leaves were mainly black when harvested, which is evidence of decomposition under anoxic conditions (Anderson and Sedell 1979). The soft sediments found in the ponds may have also limited shredder colonization and contributed to the slow decomposition rate of the leaves. Many of the leaf bags became partially buried in the pond sediments during the course of the incubation (K. Fortino, personal observation), which may have reduced the microbial decomposition of the leaf material (Danger et al. 2012).

We used coarse mesh litter bags for our litter incubation, which allowed for the colonization of macroinvertebrates into the leaf bags, however the lack of evidence of shredding activity and the low decomposition rates suggests that the litter mass loss was due mainly the microbial processes. A lack of shredder activity is a common observation in streams that have been affected by urbanization and in these systems leaf litter mass loss is mainly driven by a combination of microbial activity and physical abrasion (Paul et al. 2006). Despite the substantial accumulation of leaf litter resources in these ponds it is possible that, similar to urban streams, they do not provide suitable environmental conditions for shredders. In the ponds that we

studied, physical abrasion would likely be near zero so we expect that virtually all of the leaf litter decomposition is due to microbial activity.

Our hypothesis that leaf litter decomposition would differ among ponds with different construction types was not supported by the data. All three ponds had similarly low decomposition rates despite their construction differences. The similarity in litter decomposition rate between the ponds suggests that pond construction and gross physical conditions are not substantially affecting microbial decomposition rate, which may respond more to local sediment variables that are more similar between the ponds. Another possibility is that interacting differences between the ponds offset their respective effects. For example, Campus Pond typically has the highest chlorophyll, suggesting abundant available nutrients (Table 2), which may have stimulated leaf litter decomposition (Gulis and Suberkropp 2003; Tant et al. 2013). However, Campus Pond also has the largest inlet which could increase sedimentation and offset the impacts of the nutrients.

Taken together our results indicate that leaf litter is being collected and retained by small man-made ponds. Further we found that within these ponds, leaf litter was decaying at among the slowest rates observed for freshwater systems. Given that these ponds are novel, man-made features of the watershed, we suggest that their presence leads to a substantial alteration of organic matter processing within the watershed, and may serve as a sink for detrital organic matter.

## **Acknowledgements**

403 Invaluable field and lab help was provided by Annie Choi, Andreas Gregoriou, DJ  
 404 Lettieri, Julia Marcellus, Carly Martin, and Kasey McCusker. We would like to thank  
 405 the Longwood Real Estate Foundation and Longwood University for access to  
 406 Lancer Park Pond, the Daulton Family for access to Daulton Pond, the Town of  
 407 Farmville for access to Wilck's Lake, Longwood Univeristy for access to Campus  
 408 Pond, and the Woodland Court Apartment Complex for access to Woodland Court  
 409 Pond. We would like to thank Robert Creed for his suggestions to improve a draft of  
 410 this manuscript.

#### 411 **Funding**

412 Funding was provided by a Longwood University Faculty Development Grant and  
 413 the Longwood University PRISM program.

#### 414 **Conflicts of Interest**

415 The authors declare that they have no conflict of interest.

#### 416 **Tables**

417 Table 1. Summary of lentic decomposition coefficients.

Source	System	Region	Litter	Mesh Size	
				(cm)	k (d <sup>-1</sup> )
Alonso et al.	small man-	central Spain	<i>Ailanthus</i>	0.5	0.008
2010	made lake		<i>altissima</i>		

Alonso et al. 2010	small man- made lake	central Spain	<i>Robinia pseudoacacia</i>	0.5	0.005
Alonso et al. 2010	small man- made lake	central Spain	<i>Fraxinus angustifolia</i>	0.5	0.009
Alonso et al. 2010	small man- made lake	central Spain	<i>Ulmus minor</i>	0.5	0.008
Bottollier- Curtet et al. 2011	small floodplain pond	France	mixed exotic species	1	0.0060 – 0.0575
Bottollier- Curtet et al. 2011	small floodplain pond	France	mixed native species	1	0.0066 – 0.0463
Gonçalves et al. 2004	brackish lagoon	Rio de Janeiro State, Brazil	<i>Nymphaea ampla</i>	0.6	4.37
Gonçalves et al. 2004	brackish lagoon	Rio de Janeiro State, Brazil	<i>Typha domingensis</i>	0.6	0.17
Hodkinson 1975a	abandoned beaver pond	Alberta, Canada	<i>Salix</i>	0.35	0.0027

Hodkinson 1975a	abandoned beaver pond	Alberta, Canada	<i>Deschampsia</i>	0.35	0.0018
Hodkinson 1975a	abandoned beaver pond	Alberta, Canada	<i>Juncus</i>	0.35	0.0011
Hodkinson 1975a	abandoned beaver pond	Alberta, Canada	<i>Pinus</i>	0.35	0.0006
Oertli 1993	small man- made pond	Switzerland	<i>Quercus robur</i>	0.5 and 1.25 (data combined)	0.0014
Reed 1979	small natural lake	Ohio, USA	<i>Acer rubrum</i>	0.30	0.015 – 0.03
This study	small man- made ponds	Virginia, USA	<i>Liriodendron tulipifera</i>	0.4	0.0025 – 0.0035

418

419 Table 2. Descriptions of the ponds used in the study. The Secchi depth and  
420 chlorophyll a concentrations are regularly measured in these ponds, so the values  
421 given are representative of growing season conditions. Maximum Z is the maximum  
422 depth ever recorded in the lake. Surface Area is calculated using the digitized outline  
423 of the pond in from google maps with an online tool that calculates surface areas off  
424 of google maps (<https://www.daftlogic.com/projects-google-maps-area-calculator->

425 tool.htm). The latitude and longitude (Lat,Long) of the pond was measured at the  
 426 approximate center of the pond using the “Whats here?” feature of google maps  
 427 (<https://www.google.com/maps/>). Secchi Z is a representative Secchi depth  
 428 recorded during the growing season. Chl a is a representative chlorophyll a  
 429 concentration measured from the surface water during the growing season.  
 430 Chlorophyll was measured with a Turner Designs Trilogy fluorometer using an  
 431 acetone extraction following filtration of the pond water onto a GFF filter. Days  
 432 Litter Bags Incubated is a list of the days the litter bags were in the water before  
 433 being retrieved for those ponds used in the litter decomposition experiment.  
 434 Chlorophyll a was not measured in Woodland Court Pond or Wilck’s Lake.

Pond	Chl a					
	Max Z (m)	Surface Area (ha)	Lat,Long (DD)	Secchi Z (m)	( $\mu\text{g}$ L <sup>-1</sup> )	Days Incubated
Campus Pond	0.5	0.07	37.297, - 78.398	0.2	40.74	0, 3, 7, 15, 21, 28, 42, 57, 82, 105, 127, 209
Daulton Pond	3.4	0.55	37.283, - 78.388	1.75	6.62	0, 3, 10, 15, 22, 30, 43, 60, 106, 128, 211
Lancer Park Pond	1.5	0.10	37.306, - 78.404	0.5	12.00	0, 2, 10, 18, 23, 37, 53, 100, 116,



Woodland	2.0	0.30	37.284, -	0.8	-	NA
Court Pond			78.392			
Wilck's Lake	2.0	13.18	37.304, -	0.6	-	NA
			78.415			

435

436 **Figure Legends**

437

438 Figure 1. Areal leaf litter density in small man-made ponds near Farmville VA. Each  
 439 point represents a single Ekman sample from the lake. Points are randomly offset on  
 440 the x-axis to make all points visible.

441

442 Figure 2 Percent organic matter of the soft sediment determined from loss on  
 443 ignition at 550° C from the surveyed ponds (A) and in relation to the density of leaf  
 444 litter in the sediments (B).

445

446 **References**

447 Alonso A, González-Muñoz N, Castro-Díez P (2010) Comparison of leaf  
 448 decomposition and macroinvertebrate colonization between exotic and native trees

449 in a freshwater ecosystem. *Ecological Research* 25:647–653. doi: 10.1007/s11284-  
 450 010-0698-y

451 Anderson NH, Sedell JR (1979) Detritus Processing by Macroinvertebrates in Stream  
 452 Ecosystems. *Annual Review of Entomology* 24:351–377. doi:  
 453 10.1146/annurev.en.24.010179.002031

454 Baxter CV, Fausch KD, Saunders WC (2005) Tangled webs: reciprocal flows of  
 455 invertebrate prey link streams and riparian zones. *Freshwater Biology* 50:201–220.  
 456 doi: 10.1111/j.1365-2427.2004.01328.x

457 Benfield EF (2007) Decomposition of Leaf Material. In: *Methods in Stream Ecology*.  
 458 Elsevier, pp 711–720

459 Bird GA, Kaushik NK (1992) Invertebrate colonization and processing of maple leaf  
 460 litter in a forested and an agricultural reach of a stream. *Hydrobiologia* 234:65–77.  
 461 doi: 10.1007/bf00010862

462 Bilby RE, Likens GE (1980) Importance of Organic Debris Dams in the Structure and  
 463 Function of Stream Ecosystems. *Ecology* 61:1107–1113. doi: 10.2307/1936830

464 Bjelke U (2005) Processing of leaf matter by lake-dwelling shredders at low oxygen  
 465 concentrations. *Hydrobiologia* 539:93–98. doi: 10.1007/s10750-004-3369-6

466 Bottollier-Curtet MARION, Charcosset J-Y, Planty-Tabacchi A-M, Tabacchi E (2011)  
 467 Degradation of native and exotic riparian plant leaf litter in a floodplain pond.  
 468 *Freshwater Biology* 56:1798–1810. doi: 10.1111/j.1365-2427.2011.02620.x

469 Creed RP, Cherry RP, Pflaum JR, Wood CJ (2009) Dominant species can produce a  
 470 negative relationship between species diversity and ecosystem function. *Oikos*  
 471 118:723–732. doi: 10.1111/j.1600-0706.2008.17212.x

472 Cottingham KL, Narayan L (2013) Subsidy quantity and recipient community  
 473 structure mediate plankton responses to autumn leaf drop. *Ecosphere* 4:art89. doi:  
 474 10.1890/es13-00128.1

475 Cummins KW (1974) Structure and Function of Stream Ecosystems. *BioScience*  
 476 24:631–641. doi: 10.2307/1296676

477 Dean WE, Gorham E (1998) Magnitude and significance of carbon burial in lakes  
 478 reservoirs, and peatlands. *Geology* 26:535. doi: 10.1130/0091-  
 479 7613(1998)026<0535:masocb>2.3.co;2

480 Danger M, Cornut J, Elger A, Chauvet E (2012) Effects of burial on leaf litter quality  
 481 microbial conditioning and palatability to three shredder taxa. *Freshwater Biology*  
 482 57:1017–1030. doi: 10.1111/j.1365-2427.2012.02762.x

483 Downing JA (2010) Emerging global role of small lakes and ponds. *Limnetica*  
 484 29:0009–24.

485 Downing JA, Cole JJ, Middelburg JJ, and others (2008) Sediment organic carbon  
 486 burial in agriculturally eutrophic impoundments over the last century. *Global*  
 487 *Biogeochemical Cycles* 22:n/a–n/a. doi: 10.1029/2006gb002854

488 Downing JA, Prairie YT, Cole JJ, and others (2006) The global abundance and size  
 489 distribution of lakes ponds, and impoundments. *Limnology and Oceanography*  
 490 51:2388–2397. doi: 10.4319/lo.2006.51.5.2388

491 Eggert SL, Wallace JB (2003) Litter breakdown and invertebrate detritivores in a  
 492 resource-depleted Appalachian stream. *Archiv fur Hydrobiologie* 156:315–338. doi:  
 493 10.1127/0003-9136/2003/0156-0315

494 Fey SB, Mertens AN, Cottingham KL (2015) Autumn leaf subsidies influence spring  
 495 dynamics of freshwater plankton communities. *Oecologia* 178:875–885. doi:  
 496 10.1007/s00442-015-3279-5

497 France RL, Peters RH (1995) Predictive Model of the Effects on Lake Metabolism of  
 498 Decreased Airborne Litterfall through Riparian Deforestation. *Conservation Biology*  
 499 9:1578–1586. doi: 10.1046/j.1523-1739.1995.09061578.x

500 Gasith A, Hasler AD (1976) Airborne litterfall as a source of organic matter in lakes.  
 501 *Limnology and Oceanography* 21:253–258. doi: 10.4319/lo.1976.21.2.0253

502 Gessner MO (2010) Functional leaf traits and biodiversity effects on litter  
 503 decomposition in a stream: reply. *Ecology* 91:1869–1871. doi: 10.1890/0012-9658-  
 504 91.6.1869

505 Gessner MO, Chauvet E, Dobson M (1999) A Perspective on Leaf Litter Breakdown in  
 506 Streams. *Oikos* 85:377. doi: 10.2307/3546505

507 Gonçalves Jr. JFG, Santos AM, Esteves FA (2004) The Influence of the Chemical  
 508 Composition of *Typha Domingensis* and *Nymphaea Ampla* Detritus on Invertebrate

509 Colonization During Decomposition in a Brazilian Coastal Lagoon. *Hydrobiologia*  
 510 527:125–137. doi: 10.1023/b:hydr.0000043190.49651.dc

511 Graça MAS (2001) The Role of Invertebrates on Leaf Litter Decomposition in  
 512 Streams - a Review. *International Review of Hydrobiology* 86:383–393. doi:  
 513 10.1002/1522-2632(200107)86:4/5<383::aid-iroh383>3.0.co;2-d

514 Gulis V, Suberkropp K (2003) Leaf litter decomposition and microbial activity in  
 515 nutrient-enriched and unaltered reaches of a headwater stream. *Freshwater Biology*  
 516 48:123–134. doi: 10.1046/j.1365-2427.2003.00985.x

517 Hagen EM, Webster JR, Benfield EF (2006) Are leaf breakdown rates a useful  
 518 measure of stream integrity along an agricultural landuse gradient?. *Journal of the*  
 519 *North American Benthological Society* 25:330–343. doi: 10.1899/0887-  
 520 3593(2006)25[330:albrau]2.0.co;2

521 Hodgkinson ID (1975a) Dry Weight Loss and Chemical Changes in Vascular Plant  
 522 Litter of Terrestrial Origin Occurring in a Beaver Pond Ecosystem. *The Journal of*  
 523 *Ecology* 63:131. doi: 10.2307/2258847

524 Hodgkinson ID (1975b) Energy flow and organic matter decomposition in an  
 525 abandoned beaver pond ecosystem. *Oecologia* 21:131–139. doi:  
 526 10.1007/bf00345556

527 Huryn AD, Huryn VMB, Arbuckle CJ, Tsomides L (2002) Catchment land-use  
 528 macroinvertebrates and detritus processing in headwater streams: taxonomic

529 richness versus function. *Freshwater Biology* 47:401–415. doi: 10.1046/j.1365-  
530 2427.2002.00812.x

531 Imberger SJ, Walsh CJ, Grace MR (2008) More microbial activity not abrasive flow or  
532 shredder abundance, accelerates breakdown of labile leaf litter in urban streams.  
533 *Journal of the North American Benthological Society* 27:549–561. doi: 10.1899/07-  
534 123.1

535 Kobayashi R, Maezono Y, Miyashita T (2011) The importance of allochthonous litter  
536 input on the biomass of an alien crayfish in farm ponds. *Population Ecology* 53:525–  
537 534. doi: 10.1007/s10144-011-0268-3

538 Lennon JT, Pfaff LE (2005) Source and supply of terrestrial organic matter affects  
539 aquatic microbial metabolism. *Aquatic Microbial Ecology* 39:107–119. doi:  
540 10.3354/ame039107

541 Marcarelli AM, Baxter CV, Mineau MM, Hall RO (2011) Quantity and quality: unifying  
542 food web and ecosystem perspectives on the role of resource subsidies in  
543 freshwaters. *Ecology* 92:1215–1225. doi: 10.1890/10-2240.1

544 McConnell WJ (1968) Limnological effects of organic extracts of litter in a  
545 southwestern impoundment. *Limnology and Oceanography* 13:343–349. doi:  
546 10.4319/lo.1968.13.2.0343

547 Mehring AS, Kuehn KA, Tant CJ, and others (2014) Contribution of surface leaf-litter  
548 breakdown and forest composition to benthic oxygen demand and ecosystem

549 respiration in a South Georgia blackwater river. *Freshwater Science* 33:377–389.  
550 doi: 10.1086/675507

551 Mendoza-Lera C, Larrañaga A, Pérez J, and others (2010) Headwater reservoirs  
552 weaken terrestrial-aquatic linkage by slowing leaf-litter processing in downstream  
553 regulated reaches. *River Research and Applications* 28:13–22. doi:  
554 10.1002/rra.1434

555 Nakano S, Murakami M (2001) Reciprocal subsidies: Dynamic interdependence  
556 between terrestrial and aquatic food webs. *Proceedings of the National Academy of*  
557 *Sciences* 98:166–170. doi: 10.1073/pnas.98.1.166

558 Oertli B (1993) Leaf litter processing and energy flow through macroinvertebrates  
559 in a woodland pond (Switzerland). *Oecologia* 96:466–477. doi:  
560 10.1007/bf00320503

561 Ogbeibu A (2002) Ecological impact of river impoundment using benthic macro-  
562 invertebrates as indicators. *Water Research* 36:2427–2436. doi: 10.1016/s0043-  
563 1354(01)00489-4

564 Paul MJ, Meyer JL, Couch CA (2006) Leaf breakdown in streams differing in  
565 catchment land use. *Freshwater Biology* 51:1684–1695. doi: 10.1111/j.1365-  
566 2427.2006.01612.x

567 Polis GA, Anderson WB, Holt RD (1997) Toward an integration of landscape and  
568 food web ecology: The Dynamics of Spatially Subsidized Food Webs. *Annual Review*  
569 *of Ecology and Systematics* 28:289–316. doi: 10.1146/annurev.ecolsys.28.1.289

570 R Core Team (2014) R: A Language and Environment for Statistical Computing. R  
 571 Foundation for Statistical Computing, Vienna, Austria

572 Rau GH (1976) Dispersal of Terrestrial Plant Litter into a Subalpine Lake. *Oikos*  
 573 27:153. doi: 10.2307/3543445

574 Richey JE, Wissmar RC, Devol AH, and others (1978) Carbon Flow in Four Lake  
 575 Ecosystems: A Structural Approach. *Science* 202:1183–1186. doi:  
 576 10.1126/science.202.4373.1183

577 Short RA, Ward JV (1980) Leaf Litter Processing in a Regulated Rocky Mountain  
 578 Stream. *Canadian Journal of Fisheries and Aquatic Sciences* 37:123–127. doi:  
 579 10.1139/f80-015

580 Tant CJ, Rosemond AD, First MR (2013) Stream nutrient enrichment has a greater  
 581 effect on coarse than on fine benthic organic matter. *Freshwater Science* 32:1111–  
 582 1121. doi: 10.1899/12-049.1

583 Tornwall BM, Creed RP (2016) Shifts in shredder communities and leaf breakdown  
 584 along a disrupted stream continuum. *Freshwater Science* 35:1312–1320. doi:  
 585 10.1086/688857

586 Wallace JB (1997) Multiple Trophic Levels of a Forest Stream Linked to Terrestrial  
 587 Litter Inputs. *Science* 277:102–104. doi: 10.1126/science.277.5322.102

588 Wallace JB, Eggert SL, Meyer JL, Webster JR (1999) Effects of Resource Limitation on  
 589 a Detrital-Based Ecosystem. *Ecological Monographs* 69:409. doi: 10.2307/2657224



590 Walsh CJ, Roy AH, Feminella JW, and others (2005) The urban stream syndrome:  
 591 current knowledge and the search for a cure. *Journal of the North American*  
 592 *Benthological Society* 24:706–723. doi: 10.1899/04-028.1

593 Webster JR, Meyer JL (1997) Organic Matter Budgets for Streams: A Synthesis.  
 594 *Journal of the North American Benthological Society* 16:141–161. doi:  
 595 10.2307/1468247

596 Webster JR, Benfield EF (1986) Vascular Plant Breakdown in Freshwater  
 597 Ecosystems. *Annual Review of Ecology and Systematics* 17:567–594. doi:  
 598 10.1146/annurev.es.17.110186.003031

599 Wetzel RG (1984) Detrital dissolved and particulate organic carbon functions in  
 600 aquatic ecosystems. *Bulletin of Marine Science* 3:503–509.

601 Wetzel RG (1995) Death detritus, and energy flow in aquatic ecosystems.  
 602 *Freshwater Biology* 33:83–89. doi: 10.1111/j.1365-2427.1995.tb00388.x

603 Wetzel RG (2001) *Limnology*. Academic Press, New York

604 Reed FC (1979) Decomposition of *Acer Rubrum* Leaves at three depths in a  
 605 eutrophic Ohio lake. *Hydrobiologia* 64:195–197. doi: 10.1007/bf00020519



