

Journal for Oyster Reef Restoration Literature Review

Kenneth Fortino

January 12, 2024

- 1 Summary of Ecosystem Services Model**
- 2 Summary of Foundation Species Model**
- 3 Application of Models to Oyster Restoration
and Aquaculture**

Oysters and oyster reefs are conceptualized using both the ecosystem services model and the foundation species model — sometimes in the same paper (Mercaldo-Allen et al., 2023). There is evidence in the literature to support the use of both of these models for understanding oyster ecology. The foundation species model proposes that an ecosystem contains one or a few species that have a disproportional effect on community structure by altering the physical, chemical, and biological processes of an ecosystem to facilitate and stabilize a specific community (Fields & Silbiger, 2022). The presence of the foundation species creates and stabilizes the physical environment by minimizing fluctuations in temperature, moisture, pH, or other physical parameters (Ellison et al., 2005). Foundation species can also alter ecosystem production through changes in nutrient cycling, in some cases increasing (Fields & Silbiger, 2022) or decreasing (Ellison et al., 2005) nutrient availability. A main effect of the foundation species is to literally build the three-dimensional habitat that other species require (Angelini, Altieri, Silliman, & Bertness, 2011; Mercaldo-Allen et al., 2023). By facilitating the growth and persistence of organisms that would not be able to colonize a patch, the presence of one or more foundation species create a unique, stable, and often more biodiverse community.

Conceptualizing an ecosystem based on the foundation species model, means understanding that system as being structured by the specific facilitation interactions of one or more numerically dominant species (Dayton, 1972; Ellison et al., 2005; Angelini et al., 2011). Because they are capable of building large biogenic structures that can persist for long time periods, sometimes centuries (Lockwood & Mann, 2019), of altering local physical conditions (Lenihan, 1999),

and of processing large volumes of materials via filtration (Newell, 1988) oyster reefs can be conceptualized as foundation species where they occur. The three-dimensional structure of an oyster reef, mainly its height above the otherwise flat sediments and its heterogeneous complex structure creates, a unique environment that facilitates the growth and persistence of the oysters themselves (Lenihan, 1999), sessile producers and consumers (), and mobile consumers (Smith, Lusk, & Castorani, 2022; Searles, Gipson, Walters, & Cook, 2022). This facilitation, in part, relies on the way that the reef alters the physical environment by altering flow around the reef to alter the temperature, dissolved oxygen, and other physiochemical factors (Lenihan, 1999). The reef also alters species trophic interactions by providing cover for both predators and prey organisms (Smith et al., 2022) and aggregating and increasing resources for consumers (Newell, 1988).

For example, Newell (1988) reports that prior to their depletion during the 19th and 20th centuries, oyster reefs could remove 22 — 44% of phytoplankton C production in the Chesapeake Bay, indicating a significant alteration of material cycling in the system (Fields & Silbiger, 2022). Other studies support the idea that the function of oyster reefs is dependent on their structural development indicating that it is the creation of a particular habitat and its associated conditions that drives the effect. Searles et al. (2022) found that macroinvertebrate communities recovered on the interior of restored reefs but not their margins, suggesting that a certain oyster density or reef structure was required to facilitate the colonization of the macroinvertebrates. The alteration of the physical environment is a key factor in the way that oyster reefs create the specific ecological communities associated with them (Lenihan, 1999; Searles et al., 2022). Because the effects of the reef on the structure and function of the reef community are the result of emergent effects that result from the “physical-biological coupling” if the reef and the estuarine environment, the deconstruction of the reef by harvesting practices that destroy the three-dimensional characteristics of the reef, as well as remove individuals will result in the deminishment or elimination of the facultative properties of the reef (Lenihan, 1999). In this way the reef as a foundation species is defined by the emergent properties that arise from the integrated function of the oysters and the other organisms that are facilitated by the effects that the oysters have on the physical-biological coupling of the system (Lenihan, 1999; Ellison et al., 2005; Angelini et al., 2011).

4 Impact of Models on Understanding Oyster Management

References

- Angelini, C., Altieri, A. H., Silliman, B. R., & Bertness, M. D. (2011, October). Interactions among Foundation Species and Their Consequences for Community Organization, Biodiversity, and Conser-

- vation. *BioScience*, 61(10), 782–789. Retrieved 2024-01-11, from <https://academic.oup.com/bioscience/article-lookup/doi/10.1525/bio.2011.61.10.8> doi: 10.1525/bio.2011.61.10.8
- Dayton, P. K. (1972). Toward an understanding of community resilience and the potential effects of enrichments to the benthos at McMurdo Sound, Antarctica. In *Proceedings of the colloquium on conservation problems in Antarctica* (pp. 81–96). Allen Press Lawrence, Kansas, USA.
- Ellison, A. M., Bank, M. S., Clinton, B. D., Colburn, E. A., Elliott, K., Ford, C. R., ... Webster, J. R. (2005, November). Loss of foundation species: consequences for the structure and dynamics of forested ecosystems. *Frontiers in Ecology and the Environment*, 3(9), 479–486. Retrieved 2023-09-29, from [http://doi.wiley.com/10.1890/1540-9295\(2005\)003\[0479:LOFSCF\]2.0.CO;2](http://doi.wiley.com/10.1890/1540-9295(2005)003[0479:LOFSCF]2.0.CO;2) doi: 10.1890/1540-9295(2005)003[0479:LOFSCF]2.0.CO;2
- Fields, J., & Silbiger, N. (2022, February). Foundation species loss alters multiple ecosystem functions within temperate tidepool communities. *Marine Ecology Progress Series*, 683, 1–19. Retrieved 2024-01-10, from <https://www.int-res.com/abstracts/meps/v683/p1-19/> doi: 10.3354/meps13978
- Lenihan, H. S. (1999, August). PHYSICAL–BIOLOGICAL COUPLING ON OYSTER REEFS: HOW HABITAT STRUCTURE INFLUENCES INDIVIDUAL PERFORMANCE. *Ecological Monographs*, 69(3), 251–275. Retrieved 2023-09-29, from [http://doi.wiley.com/10.1890/0012-9615\(1999\)069\[0251:PBCOOR\]2.0.CO;2](http://doi.wiley.com/10.1890/0012-9615(1999)069[0251:PBCOOR]2.0.CO;2) doi: 10.1890/0012-9615(1999)069[0251:PBCOOR]2.0.CO;2
- Lockwood, R., & Mann, R. (2019, December). A conservation palaeobiological perspective on Chesapeake Bay oysters. *Philosophical Transactions of the Royal Society B: Biological Sciences*, 374(1788), 20190209. Retrieved 2022-04-04, from <https://royalsocietypublishing.org/doi/10.1098/rstb.2019.0209> doi: 10.1098/rstb.2019.0209
- Mercaldo-Allen, R., Auster, P. J., Clark, P., Dixon, M. S., Estela, E., Liu, Y., ... Rose, J. M. (2023, April). Oyster aquaculture cages provide fish habitat similar to natural structure with minimal differences based on farm location. *Frontiers in Marine Science*, 10, 1058709. Retrieved 2023-11-20, from <https://www.frontiersin.org/articles/10.3389/fmars.2023.1058709/full> doi: 10.3389/fmars.2023.1058709
- Newell, R. I. (1988). Ecological changes in Chesapeake Bay: are they the result of overharvesting the American oyster, *Crassostrea virginica*. *Understanding the estuary: advances in Chesapeake Bay research*, 129, 536–546. Retrieved from <http://www.oyster-restoration.org/wp-content/uploads/2012/06/Newell-1988-filtering.pdf> (Publisher: Chesapeake Research Consortium Gloucester Point, Virginia)
- Searles, A. R., Gipson, E. E., Walters, L. J., & Cook, G. S. (2022, May). Oyster reef restoration facilitates the recovery of macroin-

vertebrate abundance, diversity, and composition in estuarine communities. *Scientific Reports*, 12(1), 8163. Retrieved 2023-09-13, from <https://www.nature.com/articles/s41598-022-11688-6> doi: 10.1038/s41598-022-11688-6

Smith, R. S., Lusk, B., & Castorani, M. C. N. (2022, July). Restored oyster reefs match multiple functions of natural reefs within a decade. *Conservation Letters*, 15(4), e12883. Retrieved 2023-09-13, from <https://conbio.onlinelibrary.wiley.com/doi/10.1111/conl.12883> doi: 10.1111/conl.12883