

Journal for Oyster Reef Restoration Literature Review

Kenneth Fortino

January 11, 2024

- 1 Summary of Ecosystem Services Model**
- 2 Summary of Foundation Species Model**
- 3 Application of Models to Oyster Restoration
and Aquaculture**

Oysters and oyster reefs are modeled using both the ecosystem services model and the foundation species model. Sometimes both of these models are referenced in the same paper (Mercaldo-Allen et al., 2023). There is evidence in the literature to support the use of both of these models for understanding oyster ecology. For example, Newell (1988) reports that prior to their depletion during the 19th and 20th centuries, oyster reefs could remove 22 — 44% of phytoplankton C production in the Chesapeake Bay, indicating a significant alteration of material cycling in the system consistent with the role of a foundation species (Fields & Silbiger, 2022). Other studies support the idea that the function of oyster reefs is dependent on their structural development indicating that it is the creation of a particular habitat and its associated conditions that drives the effect. ? (?) found that macroinvertebrate communities recovered on the interior of restored reefs but not their margins, suggesting that a certain oyster density or reef structure was required to facilitate the colonization of the macroinvertebrates.

- 4 Impact of Models on Understanding Oyster
Management**

References

Fields, J., & Silbiger, N. (2022, February). Foundation species loss alters multiple ecosystem functions within temperate tidepool communi-

- ties. *Marine Ecology Progress Series*, 683, 1–19. Retrieved 2024-01-10, from <https://www.int-res.com/abstracts/meps/v683/p1-19/> doi: 10.3354/meps13978
- Mercaldo-Allen, R., Auster, P. J., Clark, P., Dixon, M. S., Estela, E., Liu, Y., ... Rose, J. M. (2023, April). Oyster aquaculture cages provide fish habitat similar to natural structure with minimal differences based on farm location. *Frontiers in Marine Science*, 10, 1058709. Retrieved 2023-11-20, from <https://www.frontiersin.org/articles/10.3389/fmars.2023.1058709/full> doi: 10.3389/fmars.2023.1058709
- Newell, R. I. (1988). Ecological changes in Chesapeake Bay: are they the result of overharvesting the American oyster, *Crassostrea virginica*. *Understanding the estuary: advances in Chesapeake Bay research*, 129, 536–546. Retrieved from <http://www.oyster-restoration.org/wp-content/uploads/2012/06/Newell-1988-filtering.pdf> (Publisher: Chesapeake Research Consortium Gloucester Point, Virginia)