

Abstract

- 30 Dor region is a great target for hierarchical star formation studies because the object is an actively star-forming region (Sun et al., 2017).
- Hierarchical star formation is star formation without a preferred spatial scale.
- It appears to be driven by supersonic turbulence in the interstellar medium (ISM)
- Binary star clusters seem to be overrepresented in galaxies; is that a real effect?

Main aim: To investigate the hierarchical star formation of 30 Doradus (30 Dor) in the Large Magellanic Cloud (LMC). This includes identifying stellar and cluster ages and spatial distributions.

Secondary aim: Assess whether binary star clusters are overrepresented compared with standard hierarchical star-formation expectations.

- The fractal dimension (FD) and virial ratio (α_{vir}) are key. Sun et al. (2017) selected Upper Main Sequence (UMS) stars from the color-magnitude diagram (CMD).
- To compare UMS and old stars, a new CMD was produced, including the UMS selection from Sun et al. (2017). Red Giant Branch (RGB) and Red Clump stars were selected as older comparison sample.
- The Orion Nebula Cluster (ONC) and Westerlund 1 (Wd1), projected to the LMC distance, were selected for realistic comparison.

Background

- The FD measures the number of points in a sample; it is represented by a non-integer number. N is the number of samples while r is a scaling factor:

$$D = \frac{\log(N)}{\log(r)}$$

- The virial ratio measures the equilibrium between kinetic energy (T) and gravitational potential energy (Ω): $\alpha_{\text{vir}} = T/|\Omega|$

Targets




Figure 1 (top). 30 Doradus (top left), the most active LMC star-forming region; Wd1 and the ONC, projected to the LMC distance are shown by a red arrow (DSS2 color). **Figure 2 (bottom left):** Orion Nebula (2MASS IR); **Figure 3 (bottom right):** Westerlund 1 (Hubble Space Telescope, F160W)

Detection



Figure 4 (top): Detection of Wd1 stars at original distances. **Figure 5 (bottom):** Detection of Wd1 stars at projected (LMC) distances. **Note: Detections of ONC stars at the LMC distance are too few to form a star cluster.**

Physical Explanation of the 2PCF & MST

- 2PCF works by measuring the probability of two stars being located at a given distance.
- This function helps trace the amplitude of stellar clustering.
- MST connects stars within a cluster with a minimum distance.
- It also represents the basis of the cluster's connectivity, revealing the hierarchical organization and relationships between the stars.

CMD Results





Figure 6 (top): CMD for the 30 Dor region with isochrones $\log(\tau/\text{yr}) = 6.6, 7, 8 \text{ \& } 9$


Figure 7 (bottom left): CMD for Wd1 at the original distance with UMS & RGB stars.

Figure 7 (bottom right): CMD for Wd1 at the LMC's distance with UMS & RGB stars.


Fractal Dimension Results



Dimension (Slope) : 1.74±0.03



Dimension (Slope): 1.43±0.09



Dimension (Slope): 1.47±0.07

Figure 9 (top): Fractal dimension analysis for 30 Dor
Figure 10 (middle): Fractal dimension analysis for the ONC
Figure 11 (bottom): Fractal dimension analysis for Wd1 (Original distance)




Figure 12:




Figure 13:




Figure 14:

2-point correlation for 30 Dor (top, figure 12), the Wd1 Original Distance (middle, figure 12), and Wd1 projected LMC Distance (bottom, figure 13).

Conclusions

- 30 Dor, the ONC and Wd1 have very similar values of FD at their original distances, ranging from 1.43 to 1.74 (highly substructured and clumpy for clusters).
- The ONC projected at the LMC distance has a very low resolution and few stars detected; it would be unidentifiable as a star cluster.
- The Minimum Spanning Tree (MST) helps achieve better results for the FD.
- The 2-point correlation function (2PCF) achieves similar results for 30 Dor, ONC and Wd1 (*work in progress*).
- Wd1 can still be identified as a star cluster projected at the LMC distance.

Contact

Kenny Chan
 Macquarie University
 Email: kingheikenny.chan@hdr.mq.edu.au | kenny.chan@mq.edu.au
 Phone: +61 406 574 967 | +1 (213) 589 7197

References

- Arnold, B., Goodwin, S.P., Griffiths, D.W. and Parker, R.J., 2017. How do binary clusters form?. Monthly Notices of the Royal Astronomical Society, 471(2), pp.2498-2507.
- Akhmetali, A., Ussipov, N., Zaidyn, M., Akhmetali, A., Sakan, A., Akhtanov, S., Kalambay, M. and Shukirgaliyev, B., 2024. Fractal dimension of star clusters. arXiv preprint arXiv:2406.05695.
- Sun, N.C., De Grijs, R., Cioni, M.R.L., Rubele, S., Subramanian, S., Van Loon, J.T., Bekki, K., Bell, C.P., Ivanov, V.D., Marconi, M. and Muraveva, T., 2018. The VMC Survey. XXIX. Turbulence-controlled Hierarchical Star Formation in the Small Magellanic Cloud. The Astrophysical Journal, 858(1), p.31.
- Bisoi, A.K. and Mishra, J., 2001. On calculation of fractal dimension of images. Pattern Recognition Letters, 22(6-7), pp.631-637.
- Pagnini, G., Di Matteo, P., Khoperskov, S., Mastrobuono-Battisti, A., Haywood, M., Renaud, F. and Combes, F., 2023. The distribution of globular clusters in kinematic spaces does not trace the accretion history of the host galaxy. Astronomy & Astrophysics, 673, p.A86.
- Goodwin, S.P. and Whitworth, A.P., 2004. The dynamical evolution of fractal star clusters: The survival of substructure. Astronomy & Astrophysics, 413(3), pp.929-937.
- Grasha, K., Calzetti, D., Adamo, A., Kim, H., Elmegreen, B.G., Gouliermis, D.A., Dale, D.A., Fumagalli, M., Grebel, E.K., Johnson, K.E. and Kahre, L., 2017. The hierarchical distribution of the young stellar clusters in six local star-forming galaxies. The Astrophysical Journal, 840(2), p.113.
- Keihänen, E., Kurki-Suonio, H., Lindholm, V., Viitanen, A., Suur-Uski, A.S., Allevato, V., Branchini, E., Marulli, F., Norberg, P., Tavagnacco, D. and de La Torre, S., 2019. Estimating the galaxy two-point correlation function using a split random catalog. Astronomy & Astrophysics, 631, p.A73.