


Marmara University, 2021

Probability and Statistics

Subject 4
Probability and Probability Distributions


Mujdat Soyuturk, Ph.D.
Associate Professor



Aim

- You have learned to describe a data set,
 - How can you use sample data to draw conclusions about the sampled populations?
 - The technique involves a statistical tool called **probability**.
- The variables that we measured can now be redefined as **random variables**, whose values depend on the chance selection of the elements in the sample.
- Using the probability as a tool, you can create **probability distributions** that serve as models for discrete random variables.

4 - 2 Mujdat Soyuturk, Probability and Statistics, Spring 2021, Marmara University




Contents

- Events and the Sample Space
- Calculating Probabilities
- Counting Rules
- Event Relations and Probability Rules
- Independence and Conditional Probability
- Baye's Rule
- Discrete Random Variables

Most parts of the slides are derived from the textbook: "Mendenhall, Beaver, Beaver, Introduction to Probability and Statistics, 14th Ed., Brooks/Cole, Cengage Learning, 2013"

4 - 3 Mujdat Soyuturk, Probability and Statistics, Spring 2021, Marmara University



What is Probability?


- In Subject 2 and 3, we used graphs and numerical measures to describe data sets which were usually **samples**.
- We measured "how often" using

Relative frequency = f/n

Sample → Population

"How often" = Relative frequency → Probability


4 - 4 Mujdat Soyuturk, Probability and Statistics, Spring 2021, Marmara University



What is Probability?

- When the population is **known**, probability is used to describe the likelihood of observing a particular sample outcome.
 - (subject of next 2-3 weeks)
- When the population is **unknown** and only a sample from that population is available, probability is used in making statements about the makeup of the population, that is, in making statistical inferences.
 - (subject of last 5 weeks)

4 - 5 Mujdat Soyuturk, Probability and Statistics, Spring 2021, Marmara University



Basic Concepts

- An **experiment** is the process by which an observation (or measurement) is obtained.
- Experiment: Record an age**
- Experiment: Toss a die and observe the result**
- Experiment: Record an opinion (yes, no)**
- Experiment: Toss two coins and observe the results**

4 - 6 Mujdat Soyuturk, Probability and Statistics, Spring 2021, Marmara University

Basic Concepts

- A **simple event** is the outcome that is observed on a single repetition of the experiment.
 - The basic element to which probability is applied.
 - One and only one simple event can occur when the experiment is performed.
- A **simple event** is denoted by E with a subscript.

4 - 7

Mujdat Soyuturk, Probability and Statistics, Spring 2021, Marmara University

Basic Concepts


- Each simple event will be assigned a probability, measuring "how often" it occurs.
- The set of all simple events of an experiment is called the **sample space, S** .

4 - 8

Mujdat Soyuturk, Probability and Statistics, Spring 2021, Marmara University


Example

- The die toss:
- Simple events:



Sample space:

$$S = \{E_1, E_2, E_3, E_4, E_5, E_6\}$$



4 - 9

Mujdat Soyuturk, Probability and Statistics, Spring 2021, Marmara University

Basic Concepts


- An **event** is a collection of one or more **simple events**.

•The die toss:

- A: an odd number
- B: a number > 2

$$A = \{E_1, E_3, E_5\}$$

$$B = \{E_3, E_4, E_5, E_6\}$$



4 - 10

Mujdat Soyuturk, Probability and Statistics, Spring 2021, Marmara University

Basic Concepts

- Two events are **mutually exclusive** if, when one event occurs, the other cannot, and vice versa.

•Experiment: Toss a die

- A: observe an odd number
- B: observe a number greater than 2

Not Mutually Exclusive

- C: observe a 6
- D: observe a 3

Mutually Exclusive

B and C?
B and D?

4 - 11

Mujdat Soyuturk, Probability and Statistics, Spring 2021, Marmara University

The Probability of an Event

- The probability of an event A measures "how often" we think A will occur. We write $P(A)$.
- Suppose that an experiment is performed n times. The relative frequency for an event A is

$$\frac{\text{Number of times } A \text{ occurs}}{n} = \frac{f}{n}$$

- If we let n get infinitely large,

$$P(A) = \lim_{n \rightarrow \infty} \frac{f}{n}$$

4 - 12

Mujdat Soyuturk, Probability and Statistics, Spring 2021, Marmara University

The Probability of an Event

- $P(A)$ must be between 0 and 1.
 - If event A can never occur, $P(A) = 0$. If event A always occurs when the experiment is performed, $P(A) = 1$.
- The sum of the probabilities for all simple events in S equals 1.

• The **probability of an event A** is found by adding the probabilities of all the simple events contained in A .

4 - 13

Mujdat Soyuturk, Probability and Statistics, Spring 2021, Marmara University

Finding Probabilities

- Probabilities can be found using
 - Estimates from empirical studies
 - Common sense estimates based on equally likely events.

Examples:

– Toss a fair coin.

$$P(\text{Head}) = 1/2$$

– 10% of the U.S. population has red hair. Select a person at random.

$$P(\text{Red hair}) = .10$$

4 - 14

Mujdat Soyuturk, Probability and Statistics, Spring 2021, Marmara University

Finding Probabilities

How to Calculate the Probability of an Event

1. List all the simple events in the sample space.
2. Assign an appropriate probability to each simple event.
3. Determine which simple events result in the event of interest.
4. Sum the probabilities of the simple events that result in the event of interest.

4 - 15

Mujdat Soyuturk, Probability and Statistics, Spring 2021, Marmara University

Example

- Toss a fair coin twice. What is the probability of observing at least one head?

1st Coin	2nd Coin	E_i	$P(E_i)$
H	H	HH	1/4
H	T	HT	1/4
T	H	TH	1/4
T	T	TT	1/4

$$\begin{aligned}
 P(\text{at least 1 head}) &= P(E_1) + P(E_2) + P(E_3) \\
 &= 1/4 + 1/4 + 1/4 = 3/4
 \end{aligned}$$

4 - 16

Mujdat Soyuturk, Probability and Statistics, Spring 2021, Marmara University

Example

- A bowl contains three candies, one red, one blue and one green. A child selects two candies at random. What is the probability that at least one is red?

1st candy	2nd candy	E_i	$P(E_i)$
R	B	RB	1/6
R	G	RG	1/6
B	R	BR	1/6
B	G	BG	1/6
G	B	GB	1/6
G	R	GR	1/6

$$\begin{aligned}
 P(\text{at least 1 red}) &= P(RB) + P(BR) + P(RG) + P(GR) \\
 &= 4/6 = 2/3
 \end{aligned}$$

4 - 17

Mujdat Soyuturk, Probability and Statistics, Spring 2021, Marmara University

Counting Rules

- If the simple events in an experiment are **equally likely**, you can calculate

$$P(A) = \frac{n_A}{N} = \frac{\text{number of simple events in } A}{\text{total number of simple events}}$$

- You can use **counting rules** to find n_A and N .

4 - 18

Mujdat Soyuturk, Probability and Statistics, Spring 2021, Marmara University

The mn Rule

- If an experiment is performed in two stages, with m ways to accomplish the first stage and n ways to accomplish the second stage, then there are mn ways to accomplish the experiment.
- This rule is easily extended to k stages, with the number of ways equal to

$$n_1 n_2 n_3 \dots n_k$$

Example: Toss two coins. The total number of simple events is:

$$2 \times 2 = 4$$

4 - 19

Mujdat Soyuturk, Probability and Statistics, Spring 2021, Marmara University

Examples

Example: Toss three coins. The total number of simple events is:

$$2 \times 2 \times 2 = 8$$

Example: Toss two dice. The total number of simple events is:

$$6 \times 6 = 36$$

Example: Two candies are drawn from a dish containing two red and two blue candies. The total number of simple events is:

$$4 \times 3 = 12$$

4 - 20

Mujdat Soyuturk, Probability and Statistics, Spring 2021, Marmara University

Permutations

- The number of ways you can arrange n distinct objects, taking them r at a time is

$$P_r^n = \frac{n!}{(n-r)!}$$

where $n! = n(n-1)(n-2)\dots(2)(1)$ and $0! = 1$.

Example: How many 3-digit lock combinations can we make from the numbers 1, 2, 3, and 4?

The order of the choice is important!

$$P_3^4 = \frac{4!}{1!} = 4(3)(2) = 24$$

4 - 21

Mujdat Soyuturk, Probability and Statistics, Spring 2021, Marmara University

Examples

Example: A lock consists of five parts and can be assembled in any order. A quality control engineer wants to test each order for efficiency of assembly. How many orders are there?

The order of the choice is important!

$$P_5^5 = \frac{5!}{0!} = 5(4)(3)(2)(1) = 120$$

4 - 22

Mujdat Soyuturk, Probability and Statistics, Spring 2021, Marmara University

Combinations

- Sometimes the ordering or arrangements of the objects is not important, but only the objects that are chosen.
- In this case, you can use a counting rule for **combinations**.

4 - 23

Mujdat Soyuturk, Probability and Statistics, Spring 2021, Marmara University

Combinations

- The number of distinct combinations of n distinct objects that can be formed, taking them r at a time is

$$C_r^n = \frac{n!}{r!(n-r)!}$$

Example: Three members of a 5-person committee must be chosen to form a subcommittee. How many different subcommittees could be formed?

The order of the choice is not important!

$$C_3^5 = \frac{5!}{3!(5-3)!} = \frac{5(4)(3)(2)1}{3(2)(1)(2)1} = \frac{5(4)}{(2)1} = 10$$

4 - 24

Mujdat Soyuturk, Probability and Statistics, Spring 2021, Marmara University

Example

- A box contains six candies, four red and two green. A child selects two candies at random. What is the probability that exactly one is red?

The order of the choice is not important!

$$C_2^6 = \frac{6!}{2!4!} = \frac{6(5)}{2(1)} = 15$$

ways to choose 2 candies.

$$C_1^2 = \frac{2!}{1!1!} = 2$$

ways to choose 1 green candy.

$$C_1^4 = \frac{4!}{1!3!} = 4$$

ways to choose 1 red candy.

$4 \times 2 = 8$ ways to choose 1 red and 1 green candy.

$$P(\text{exactly one red}) = 8/15$$

Ex 4.15, pp 136

4 - 25

Mujdat Soyuturk, Probability and Statistics, Spring 2021, Marmara University

Event Relations

- Sometimes the event of interest can be formed as a combination of several other events.
- Then, we have to look at the relationships between events.


4 - 26

Mujdat Soyuturk, Probability and Statistics, Spring 2021, Marmara University

Event Relations

- The **union** of two events, A and B, is the event that either A or B or **both** occur when the experiment is performed. We write

$$A \cup B$$




4 - 27

Mujdat Soyuturk, Probability and Statistics, Spring 2021, Marmara University

Event Relations

- The **intersection** of two events, A and B, is the event that both A and B occur when the experiment is performed.

We write $A \cap B$.




- If two events A and B are **mutually exclusive**, then $P(A \cap B) = 0$.

4 - 28

Mujdat Soyuturk, Probability and Statistics, Spring 2021, Marmara University

Event Relations

- The **complement** of an event A consists of all outcomes of the experiment that do not result in event A. We write A^c .



4 - 29

Mujdat Soyuturk, Probability and Statistics, Spring 2021, Marmara University

Example

- Select a student from the classroom and record his/her **hair color** and **gender**.
 - A: student has brown hair
 - B: student is female
 - C: student is male

Mutually exclusive; $B = C^c$

- What is the relationship between events B and C?

• A^c : Student does not have brown hair

• $B \cap C$: Student is both male and female = \emptyset

• $B \cup C$: Student is either male and female = all students = S

Ex 4.16, pp 140

4 - 30

Mujdat Soyuturk, Probability and Statistics, Spring 2021, Marmara University

Calculating Probabilities for Unions and Complements




- There are special rules that will allow you to calculate probabilities for composite events.

The Additive Rule for Unions:

- For any two events, **A** and **B**, the probability of their union, $P(A \cup B)$, is

$$P(A \cup B) = P(A) + P(B) - P(A \cap B)$$



4 - 31 Mujdat Soyuturk, Probability and Statistics, Spring 2021, Marmara University

Example: Additive Rule



Example: Suppose that there were 120 students in the classroom, and that they could be classified as follows:

A: brown hair

$$P(A) = 50/120$$

B: female

$$P(B) = 60/120$$

	Brown	Not Brown
Male	20	40
Female	30	30

$$\begin{aligned} P(A \cup B) &= P(A) + P(B) - P(A \cap B) \\ &= 50/120 + 60/120 - 30/120 \\ &= 80/120 = 2/3 \end{aligned}$$

$$\begin{aligned} \text{Check: } P(A \cup B) &= (20 + 30 + 30)/120 \end{aligned}$$

4 - 32 Mujdat Soyuturk, Probability and Statistics, Spring 2021, Marmara University

A Special Case



When two events **A** and **B** are **mutually exclusive**,

$$P(A \cap B) = 0$$

$$\text{and } P(A \cup B) = P(A) + P(B).$$

A: male with brown hair

$$P(A) = 20/120$$

B: female with brown hair

$$P(B) = 30/120$$

	Brown	Not Brown
Male	20	40
Female	30	30

A and B are mutually exclusive, so that

$$\begin{aligned} P(A \cup B) &= P(A) + P(B) \\ &= 20/120 + 30/120 \\ &= 50/120 \end{aligned}$$

4 - 33 Mujdat Soyuturk, Probability and Statistics, Spring 2021, Marmara University

Calculating Probabilities for Complements




- We know that for any event **A**:

$$P(A \cap A^c) = 0$$

- Since either **A** or A^c must occur,

$$P(A \cup A^c) = 1$$

- so that $P(A \cup A^c) = P(A) + P(A^c) = 1$



$$P(A^c) = 1 - P(A)$$

4 - 34 Mujdat Soyuturk, Probability and Statistics, Spring 2021, Marmara University

Example



Select a student at random from the classroom.

Define:

A: male

$$P(A) = 60/120$$

B: female

	Brown	Not Brown
Male	20	40
Female	30	30

A and B are complementary, so that

$$\begin{aligned} P(B) &= 1 - P(A) \\ &= 1 - 60/120 = 60/120 \end{aligned}$$

4 - 35 Mujdat Soyuturk, Probability and Statistics, Spring 2021, Marmara University

Calculating Probabilities for Intersections



- In the previous example, we found $P(A \cap B)$ directly from the table. Sometimes this is impractical or impossible. The rule for calculating $P(A \cap B)$ depends on the idea of **independent and dependent events**.

Two events, **A** and **B**, are said to be **independent** if and only if the probability that event **A** occurs does not change, depending on whether or not event **B** has occurred.

4 - 36 Mujdat Soyuturk, Probability and Statistics, Spring 2021, Marmara University

Conditional Probabilities

- The probability that **A** occurs, given that event **B** has occurred is called the **conditional probability of A given B** and is defined as

$$P(A|B) = \frac{P(A \cap B)}{P(B)}, \text{ if } P(B) \neq 0$$

"given"

4 - 37

Mujdat Soyuturk, Probability and Statistics, Spring 2021, Marmara University

Example 1

- Toss a fair coin twice. Define
 - A: head on second toss
 - B: head on first toss

HH	1/4
HT	1/4
TH	1/4
TT	1/4

$$P(A|B) = \frac{1}{2}$$

$$P(A|\text{not } B) = \frac{1}{2}$$

P(A) does not change, whether B happens or not...

A and B are independent!

4 - 38

Mujdat Soyuturk, Probability and Statistics, Spring 2021, Marmara University

Example 2

- A bowl contains five candies, two red and three blue. Randomly select two candies, and define
 - A: first candy is blue.
 - B: second candy is red.



$$P(B|A) = P(2^{\text{nd}} \text{ red} | 1^{\text{st}} \text{ blue}) = 2/4 = 1/2$$

$$P(B|\text{not } A) = P(2^{\text{nd}} \text{ red} | 1^{\text{st}} \text{ red}) = 1/4$$

P(B) does change, depending on whether A happens or not...

A and B are dependent!

4 - 39

Mujdat Soyuturk, Probability and Statistics, Spring 2021, Marmara University

Defining Independence

- We can redefine independence in terms of conditional probabilities:

Two events A and B are **independent** if and only if

$$P(A|B) = P(A) \quad \text{or} \quad P(B|A) = P(B)$$

Otherwise, they are **dependent**.

- Once you've decided whether or not two events are independent, you can use the following rule to calculate their intersection.

4 - 40

Mujdat Soyuturk, Probability and Statistics, Spring 2021, Marmara University

The Multiplicative Rule for Intersections

- For any two events, A and B, the probability that both A and B occur is

$$P(A \cap B) = P(A) P(B \text{ given that } A \text{ occurred}) = P(A)P(B|A)$$

- If the events **A** and **B** are independent, then the probability that both A and B occur is

$$P(A \cap B) = P(A) P(B)$$

4 - 41

Mujdat Soyuturk, Probability and Statistics, Spring 2021, Marmara University

Example 1

In a certain population, 10% of the people can be classified as being high risk for a heart attack. Three people are randomly selected from this population. What is the probability that exactly one of the three are high risk?

Define **H**: high risk **N**: not high risk

$$\begin{aligned} P(\text{exactly one high risk}) &= P(HNN) + P(NHN) + P(NNH) \\ &= P(H)P(N)P(N) + P(N)P(H)P(N) + P(N)P(N)P(H) \\ &= (.1)(.9)(.9) + (.9)(.1)(.9) + (.9)(.9)(.1) = 3(.1)(.9)^2 = .243 \end{aligned}$$

4 - 42

Mujdat Soyuturk, Probability and Statistics, Spring 2021, Marmara University

Example 2

Suppose we have additional information in the previous example. We know that only 49% of the population are female. Also, of the female patients, 8% are high risk. A single person is selected at random. What is the probability that it is a high risk female?

Define **H**: high risk **N**: not high risk

From the example, $P(F) = .49$ and $P(H|F) = 0.08$. Use the Multiplicative Rule:

$$\begin{aligned} P(\text{high risk female}) &= P(H \cap F) \\ &= P(F)P(H|F) = .49(.08) = .0392 \end{aligned}$$

4 - 43

Mujdat Soyuturk, Probability and Statistics, Spring 2021, Marmara University

The Law of Total Probability


- Let $S_1, S_2, S_3, \dots, S_k$ be mutually exclusive and exhaustive events (that is, one and only one must happen). Then the probability of another event A can be written as

$$\begin{aligned} P(A) &= P(A \cap S_1) + P(A \cap S_2) + \dots + P(A \cap S_k) \\ &= P(S_1)P(A|S_1) + P(S_2)P(A|S_2) + \dots + P(S_k)P(A|S_k) \end{aligned}$$

4 - 44

Mujdat Soyuturk, Probability and Statistics, Spring 2021, Marmara University

The Law of Total Probability



$$\begin{aligned} P(A) &= P(A \cap S_1) + P(A \cap S_2) + \dots + P(A \cap S_k) \\ &= P(S_1)P(A|S_1) + P(S_2)P(A|S_2) + \dots + P(S_k)P(A|S_k) \end{aligned}$$

4 - 45

Mujdat Soyuturk, Probability and Statistics, Spring 2021, Marmara University

Bayes' Rule

- Let $S_1, S_2, S_3, \dots, S_k$ be mutually exclusive and exhaustive events with prior probabilities $P(S_1), P(S_2), \dots, P(S_k)$.
- If an event A occurs, the posterior probability of S_i , given that A occurred is

$$P(S_i | A) = \frac{P(S_i)P(A | S_i)}{\sum P(S_i)P(A | S_i)} \text{ for } i = 1, 2, \dots, k$$

4 - 46

Mujdat Soyuturk, Probability and Statistics, Spring 2021, Marmara University

Example

From a previous example, we know that 49% of the population are female. Of the female patients, 8% are high risk for heart attack, while 12% of the male patients are high risk. A single person is selected at random and found to be high risk. What is the probability that it is a male?

Define **H**: high risk **F**: female **M**: male

We know:

$$\begin{aligned} P(F) &= .49 \\ P(M) &= .51 \\ P(H|F) &= .08 \\ P(H|M) &= .12 \end{aligned}$$

$$\begin{aligned} P(M | H) &= \frac{P(M)P(H | M)}{P(M)P(H | M) + P(F)P(H | F)} \\ &= \frac{.51(.12)}{.51(.12) + .49(.08)} = .61 \end{aligned}$$

4 - 47

Mujdat Soyuturk, Probability and Statistics, Spring 2021, Marmara University

Random Variables

- A quantitative variable x is a **random variable** if the value that it assumes, corresponding to the outcome of an experiment is a chance or random event.
- Random variables can be **discrete** or **continuous**.

Examples:

- x = SAT score for a randomly selected student
- x = number of people in a room at a randomly selected time of day
- x = number on the upper face of a randomly tossed die

4 - 48

Mujdat Soyuturk, Probability and Statistics, Spring 2021, Marmara University

Probability Distributions for Discrete Random Variables



- The **probability distribution** for a random variable x is the **relative frequency distribution** constructed for the entire population of measurements. It is a **graph, table or formula** that gives the possible values of x and the probability $p(x)$ associated with each value.

We must have

$$0 \leq p(x) \leq 1 \quad \text{and} \quad \sum p(x) = 1$$


4 - 49

Mujdat Soyuturk, Probability and Statistics, Spring 2021, Marmara University

Example



- Toss a fair coin three times and define x = number of heads.



4 - 50

Mujdat Soyuturk, Probability and Statistics, Spring 2021, Marmara University

Probability Distributions



- Probability distributions can be used to describe the population, just as we described samples in Ch.1.
 - Shape:** Symmetric, skewed, mound-shaped...
 - Outliers:** unusual or unlikely measurements
 - Center and spread:** mean and standard deviation. A population mean is called μ and a population standard deviation is called σ .

4 - 51

Mujdat Soyuturk, Probability and Statistics, Spring 2021, Marmara University

The Mean and Standard Deviation



- Let x be a discrete random variable with probability distribution $p(x)$. Then the mean, variance and standard deviation of x are given as

Mean : $\mu = \sum xp(x)$ Expected Value

Variance : $\sigma^2 = \sum (x - \mu)^2 p(x)$

Standard deviation : $\sigma = \sqrt{\sigma^2}$

4 - 52

Mujdat Soyuturk, Probability and Statistics, Spring 2021, Marmara University

The Mean and Standard Deviation



- The **population mean**, which measures the average value of x in the population, is also called the **expected value** of the random variable x is written as $E(x)$.
- It is the value that you would **expect** to observe on **average** if the experiment is repeated over and over again.

4 - 53

Mujdat Soyuturk, Probability and Statistics, Spring 2021, Marmara University

Example



- Toss a fair coin 3 times and record x the number of heads.

x	$p(x)$	$xp(x)$	$(x - \mu)^2 p(x)$
0	1/8	0	$(-1.5)^2(1/8)$
1	3/8	3/8	$(-0.5)^2(3/8)$
2	3/8	6/8	$(0.5)^2(3/8)$
3	1/8	3/8	$(1.5)^2(1/8)$

$$\mu = \sum xp(x) = \frac{12}{8} = 1.5$$

$$\sigma^2 = \sum (x - \mu)^2 p(x)$$

$$\sigma^2 = .28125 + .09375 + .09375 + .28125 = .75$$


$$\sigma = \sqrt{.75} = .688$$

4 - 54

Mujdat Soyuturk, Probability and Statistics, Spring 2021, Marmara University

Example

- The probability distribution for x the number of heads in tossing 3 fair coins.



- Shape? Symmetric; mound-shaped
- Outliers? None
- Center? $\mu = 1.5$
- Spread? $\sigma = .688$

4 - 55

Mujdat Soyuturk, Probability and Statistics, Spring 2021, Marmara University

Key Concepts

I. Experiments and the Sample Space

- Experiments, events, mutually exclusive events, simple events
- The sample space
- Venn diagrams, tree diagrams, probability tables

II. Probabilities

- Relative frequency definition of probability
- Properties of probabilities
 - Each probability lies between 0 and 1.
 - Sum of all simple-event probabilities equals 1.
- $P(A)$, the sum of the probabilities for all simple events in A

4 - 56

Mujdat Soyuturk, Probability and Statistics, Spring 2021, Marmara University

Key Concepts

III. Counting Rules

- mn Rule; extended mn Rule
- Permutations:

$$P_r^n = \frac{n!}{(n-r)!}$$

- Combinations:

$$C_r^n = \frac{n!}{r!(n-r)!}$$

IV. Event Relations

- Unions and intersections
- Events
 - Disjoint or mutually exclusive: $P(A \cap B) = 0$
 - Complementary: $P(A) = 1 - P(A^c)$

4 - 57

Mujdat Soyuturk, Probability and Statistics, Spring 2021, Marmara University

Key Concepts

- Conditional probability:

$$P(A|B) = \frac{P(A \cap B)}{P(B)}$$

- Independent and dependent events

- Additive Rule of Probability:

$$P(A \cup B) = P(A) + P(B) - P(A \cap B)$$

- Multiplicative Rule of Probability:

$$P(A \cap B) = P(A)P(B|A)$$

- Law of Total Probability
- Bayes' Rule

4 - 58

Mujdat Soyuturk, Probability and Statistics, Spring 2021, Marmara University

Key Concepts

V. Discrete Random Variables and Probability Distributions

- Random variables, discrete and continuous
- Properties of probability distributions

$$0 \leq p(x) \leq 1 \text{ and } \sum p(x) = 1$$
- Mean or expected value of a discrete random variable:

$$\text{Mean : } \mu = \sum xp(x)$$
- Variance and standard deviation of a discrete random variable:

$$\text{Variance : } \sigma^2 = \sum (x - \mu)^2 p(x)$$

$$\text{Standard deviation : } \sigma = \sqrt{\sigma^2}$$

4 - 59

Mujdat Soyuturk, Probability and Statistics, Spring 2021, Marmara University