


BACKGROUND

Wildfires are increasing in intensity, severity, size and duration.

Western, mountainous states bear the highest burden.

Increased temperatures and changes in weather patterns will continue contributing to wildfires.

Which climate factors have impact on wildfire?



DATA SOURCE

2.3 Million US Wildfires (1992-2020) 6th Edition
And

Daily Summaries Dataset from National Oceanic and Atmospheric Administration (NOAA)

Process Summary:

- Gathered 280 million weather data points spanning 8 years (2013 to 2020) from NOAA
 - Filtered out Min, Max, Average Temperatures, Rain, Snow, and Wind. Also included elevation and Latitude and Longitude.
 - Added weather readings to fire data by matching location and date

DATA SUMMARY

Final Dataset had 955k weather readings for both Fire and No Fire present

Fire Size	Ave Temp (C)	Max Temp (C)	Min Temp (C)	Precipitation	Snow (mm)	Snow Depth	Wind (10ths	Elevation	Days to
FILE SIZE	Ave remp (c)	iviax reilip (C)	wiiii Teilip (C)	(10ths of mm)	Snow (mm)	(mm)	meters/second)	Elevation	Contain
No Fire	10.8	16.1	5.5	31.4	2.9	24.3	37.5	380.7	N/A
Α	18.7	26.1	11.4	8.7	0.2	0.7	35.0	566.3	0.6
В	17.4	25.0	10.3	8.5	0.1	0.4	35.3	333.3	0.5
С	17.1	25.0	9.7	8.2	0.1	0.2	37.0	361.2	1.1
D	18.8	26.9	10.9	7.9	0.1	0.1	39.6	506.1	3.2
E	19.8	28.0	11.7	6.5	0.0	0.0	40.6	587.0	6.2
F	21.8	30.1	13.4	5.4	0.1	0.0	40.8	674.3	12.4
G	23.5	31.5	14.8	3.9	0.0	0.0	37.4	655.0	31.0



TECHNOLOGY



Pandas



SQLite




Matplotlib Tableau



ML Ski-kit learn

Random forest & Neural Network


ANALYZING WILDFIRE TREND 2013 TO 2020



Wildfire


Wildfire seemed to be higher in 2016

California and Texas seems to has more fires compared to any other states



Count of Fire by Each Month and By State in US

WILDFIRE MAP COMPARISON



In 2013, wildfire incidents in the United States numbered 46,615, consuming a total of 4,307,176 acres. Fast-forward to 2020, the scale of devastation has multiplied, witnessing over double the acreage burned. A staggering 10,274,679 acres were ravaged by wildfires that year, coinciding with a substantial rise in the number of fires, reaching 58,258 incidents.

"Nature is not a place to visit. It is home."


- Gary Snyder

ANALYZING CLIMATE FACTORS AFFECTING WILDFIRE

As of June 1, 2023, around 18,300 wildfires have impacted over half of million (511,000) acres this year.

Wildfires have the potential to harm property, livelihoods, and human health. Fire related threats are increasing, especially as more people live in and around forests, grasslands, and other natural areas. Over the past few decades, the United States has routinely spent more than \$1 billion per year to fight wildfires, including \$2.3 billion in 2020.

ANALYZING CLIMATE TREND 2013 TO 2020




SUMMARY


with higher temperature and higher wind fire count has significantly increased.

It is evident that climate has impact on wildfire

WILDFIRE CLIMATE CONNECTION



Climate change, including increased heat, extended drought, and a thirsty atmosphere, has been a key driver in increasing the risk and extent of wildfires in the western United States during the last two decades. Wildfires require the alignment of a number of factors, including temperature, humidity, and the lack of moisture in fuels, such as trees, shrubs, grasses, and forest debris. All these factors have strong direct or indirect ties to climate variability and climate change.



RANDOM FOREST - MODEL METRICS

Following is the confusion matrix of the model:

	Predicted 0	Predicted 1
Actual 0	79411	13909
Actual 1	10972	134493


Accuracy Score : 0.8958016625834956

Classification Report

CIUSSIIIC	G C I C	ii itcpoi c			
		precision	recall	f1-score	support
	0	0.88	0.85	0.86	93320
	1	0.91	0.92	0.92	145465
accur	асу			0.90	238785
macro	avg	0.89	0.89	0.89	238785
weighted	avg	0.90	0.90	0.90	238785

- The model does a pretty good job for both positive and negative case with an overall accuracy, recall and f1-score of about 90%
- The scores are slightly better for positive cases probably because of the volume of data of positive cases being higher than the negative ones

RANDOM FOREST - FEATURES IMPORTANCES



RANDOM FOREST - OPTIMIZATIONS

Our initial dataset used for the model had the columns populated only for positive (Fire occurred) cases. As a result, it was predicting the Fire occurrence with 100% accuracy. This obviously was a flawed dataset for our purpose.

In the next iteration, we dropped the positive case — only columns. The data set had weather columns - Temperature, Precipitation, latitude, longitude and elevation. The period of the dataset was 1 year.

With this dataset, we had the following confusion matrix:

	Predicted 0	Pred	licted 1	
Actual 0	88647	,	8127	
Actual 1	7840)	10384	
	5	064454	40702704	
-	Score : 0.		10/02/94	ŏ
Classific	cation Repo	rt		
	preci	sion	recall	
	0	0.92	0.92	
	1	0.56	0.57	
accur	racy			
macro	avg	0.74	0.74	
weighted	avg	0.86	0.86	





NEURAL NETWORK MODEL

MODEL I

39% accuracy



Preprocessing:

MinMaxScalar.fit for preprocessing and normalizing the data




Model

- Sequential structure model
- 10 input features
- 4 hidden layers neuron unit ranging between 2 to 5 neurons
- Relu as activation function
- Output layer with one neuron and "Sigmod" as activation function

Model compilation

- Loss function: "mean_squared_logarithmic_error"
- Optimizer: 'adam'
- Metrics: accuracy





NEURAL NETWORK MODEL

MODEL 2

85% accuracy



Preprocessing:

MinMaxScalar.fit for preprocessing and normalizing the data




Model

- Sequential structure model
- 10 input features
- 5 hidden layers neuron unit ranging between 60 to 600 neurons
- Relu as activation function
- Output layer with one neuron and "Sigmod" as activation function

Model compilation

- Loss function: "mean_squared_error"
- Optimizer: 'adam'
- Metrics: accuracy





NEURAL NETWORK MODEL

MODEL 3

86% accuracy



Preprocessing:

MinMaxScalar.fit for preprocessing and normalizing the data



- Sequential structure model
- 11 input features
- bidden layers neuron unit ranging between 400 to 1100 neurons
- Relu as activation function
- Output layer with one neuron and "Sigmod" as activation function

Model compilation

- Loss function: "mean_squared_logarithmic_error"
- Optimizer: 'adam'
- Metrics: accuracy


Using the Machine learning and Neural network, optimized tensor flow model reached accuracy of 86%.

Based on the optimization approach it is recommended

- * To have higher neuron units and minimum of 5 hidden layers column
- * Using mean_squared_logarithmic_error would significantly reduce loss
- * Have a minimum of 10 to 11 features for evaluation as reduced input features results in lower accuracy.

Overall, based on volume of the data collected for wildfire and climate factor, this model performed pretty good and may not need more additional collection of training data. This model would predict reliable results. However,...







FURTHER IMPROVEMENTS OPPORTUNITIES

- * Try improving accuracy of neural network using different preprocessing and/or model and/or compilation
- * Trying random forest regression model and using the probability function of the random forest model
- $\ensuremath{^{*}}$ Using the solution predict the future outcome for the upcoming years

and there could be many more... to be further explored.....



QUESTIONS?

THANKS!

QUESTION?

CREDITS: This presentation template was created by Slidesgo, including icons by Flaticon, and infographics & images by Freepik.

PLEASE KEEP THIS SLIDE FOR ATTRIBUTION.

