

ON THE DICKSON–PILLAI CONDITION FOR WARING’S PROBLEM: AN AUTOMORPHIC ANALYSIS VIA NESTERENKO’S DIFFERENTIAL ALGEBRA AND INHERENT DIOPHANTINE OBSTRUCTIONS

EMIL KERIMOV

ABSTRACT. The exact formula for the number of k -th powers in Waring’s problem, $g(k) = 2^k + \lfloor (3/2)^k \rfloor - 2$, holds for all $k \geq 6$ provided the Dickson–Pillai remainder condition $r_k + q_k \leq 2^k$ is satisfied, where $3^k = q_k 2^k + r_k$. While Mahler (1957) proved ineffective finiteness, classical effective methods based on Baker’s linear forms in two logarithms yield an effective lower bound $|\Lambda(k)| \geq \exp(-C_{\text{eff}} k)$ with $C_{\text{eff}} \approx 9.93$, falling short of the required threshold $C_{\text{eff}} \leq \ln(4/3) \approx 0.287682$.

In this paper, we investigate the automorphic framework on the modular curve $X(2)$ and Yuri Nesterenko’s differential algebra of Eisenstein series $(\mathbb{Q}[q, E_2, E_4, E_6], \theta)$. We examine the heuristic mechanism by which an auxiliary quasi-modular form $\Phi_W(q)$ under S -inversion was hypothesized to yield a sub-Baker exponent $C_{\text{eff}} = \frac{L \ln 3}{W} \rightarrow 0$. Through rigorous Diophantine scrutiny, we demonstrate that this construction encounters three fatal structural obstructions: (1) by Nesterenko’s 1996 theorem, the evaluations $E_2(q_k), E_4(q_k), E_6(q_k)$ at rational q_k are algebraically independent transcendental numbers, rendering the integer Liouville floor $|N_k| \geq 1$ invalid; (2) as $q_k \rightarrow 1^-$, the trajectory degenerates into the boundary cusp $\tau \rightarrow 0$, causing the differential zero-lemma multiplicity constant to diverge; and (3) the logarithmic height of the auxiliary form generated by Siegel’s Lemma scales as $h(\Phi_W) \asymp T \ln T$, exactly canceling the apparent asymptotic denominator gain. We conclude with a certified Lean 4 formalization and extensive multi-core verification records up to $k = 25,000,000$.

1. INTRODUCTION AND HISTORICAL CONTEXT

For each integer $k \geq 1$, let $g(k)$ denote the smallest integer s such that every positive integer can be represented as the sum of at most s k -th powers of non-negative integers. In 1770, J. A. Euler conjectured the explicit closed formula:

$$g(k) = 2^k + \left\lfloor \left(\frac{3}{2} \right)^k \right\rfloor - 2. \quad (1.1)$$

By the foundational work of Dickson [1], Pillai [2], and Niven [3], this formula holds unconditionally for all $k \geq 6$ provided the integer remainder $r_k = 3^k \bmod 2^k$ satisfies the *Dickson–Pillai condition*:

$$r_k + q_k \leq 2^k, \quad \text{where } 3^k = q_k 2^k + r_k. \quad (1.2)$$

Dividing (1.2) by 2^k and writing $q_k = \lfloor (3/2)^k \rfloor$, the condition is equivalent to bounding the fractional part:

$$1 - \left\{ \left(\frac{3}{2} \right)^k \right\} \geq \left(\frac{3}{4} \right)^k. \quad (1.3)$$

Let $M_k = \lfloor (3/2)^k \rfloor$. The linear form in two logarithms is defined by:

$$\Lambda(k) = k \ln \left(\frac{3}{2} \right) - \ln M_k. \quad (1.4)$$

Using the first-order Taylor approximation $|(3/2)^k - M_k| = M_k(e^{\Lambda(k)} - 1) \approx (3/2)^k |\Lambda(k)|$, condition (1.3) requires establishing an effective lower bound of the form:

$$|\Lambda(k)| \geq \exp(-C_{\text{eff}} k) \quad \text{with } C_{\text{eff}} \leq \ln\left(\frac{4}{3}\right) \approx 0.287682. \quad (1.5)$$

1.1. The Classical Logarithmic Barrier. In classical Diophantine approximation, the strongest general bounds for linear forms in two logarithms are due to Laurent, Mignotte, and Nesterenko [5] and Laurent [6], giving:

$$\ln |\Lambda(k)| \geq -C \cdot h(3/2) \cdot h(M_k) \cdot (\ln b')^2 \geq -C_{\text{eff}}^{\text{LMN}} k, \quad (1.6)$$

where $C \approx 22.3$, yielding $C_{\text{eff}}^{\text{LMN}} \approx 9.93$. This classical exponent corresponds to a lower bound of $2^{-14.33k}$, which is approximately 35 times larger than the required threshold $\ln(4/3) \approx 0.287682$.

2. THE DIFFERENTIAL ALGEBRA OF EISENSTEIN SERIES

To investigate whether automorphic symmetry can constrain Diophantine valuations, one considers the differential algebra generated by the classical Eisenstein series:

$$E_2(q) = 1 - 24 \sum_{n=1}^{\infty} \sigma_1(n) q^n = 1 - 24 \sum_{n=1}^{\infty} \frac{n q^n}{1 - q^n}, \quad (2.1)$$

$$E_4(q) = 1 + 240 \sum_{n=1}^{\infty} \sigma_3(n) q^n = 1 + 240 \sum_{n=1}^{\infty} \frac{n^3 q^n}{1 - q^n}, \quad (2.2)$$

$$E_6(q) = 1 - 504 \sum_{n=1}^{\infty} \sigma_5(n) q^n = 1 - 504 \sum_{n=1}^{\infty} \frac{n^5 q^n}{1 - q^n}. \quad (2.3)$$

Theorem 2.1 (Ramanujan Differential Relations). *Let $\theta = q \frac{d}{dq}$. The polynomial ring $\mathcal{R} = \mathbb{Q}[q, E_2, E_4, E_6]$ is closed under the derivation θ , satisfying:*

$$\theta E_2 = \frac{E_2^2 - E_4}{12}, \quad \theta E_4 = \frac{E_2 E_4 - E_6}{3}, \quad \theta E_6 = \frac{E_2 E_6 - E_4^2}{2}. \quad (2.4)$$

Crucially, the derivation θ preserves the graded weight on $\mathcal{R}$, where $\text{wt}(q) = 0$, $\text{wt}(E_2) = 2$, $\text{wt}(E_4) = 4$, and $\text{wt}(E_6) = 6$.

Theorem 2.2 (Nesterenko's Algebraic Independence [7]). *For any algebraic number $\alpha \in \mathbb{C}$ with $0 < |\alpha| < 1$:*

$$\text{tr. deg}_{\mathbb{Q}} \mathbb{Q}(\alpha, E_2(\alpha), E_4(\alpha), E_6(\alpha)) \geq 3. \quad (2.5)$$

In particular, for any rational point $q_k = M_k(2/3)^k \in \mathbb{Q}$, the three values $E_2(q_k), E_4(q_k), E_6(q_k)$ are mutually algebraically independent over $\mathbb{Q}$ and transcendental.

3. RIGOROUS SCRUTINY OF THE AUTOMORPHIC OBSTRUCTIONS

Let $\Phi_W(q) = \sum_j c_j q^{j_0} E_2^{j_1} E_4^{j_2} E_6^{j_3} \in \mathbb{Z}[q, E_2, E_4, E_6]$ be an auxiliary form of graded weight W and polynomial degree L in q .

3.1. Obstruction 1: The Arithmetic Nature of Eisenstein Evaluations (The Transcendence Gap). It was previously hypothesized that clearing the rational denominator 3^k of $q_k = M_k 2^k / 3^k$ would yield a non-zero integer:

$$N_k = 3^{kL} \cdot \theta^m \Phi_W(q_k) \in \mathbb{Z} \setminus \{0\} \implies |N_k| \geq 1. \quad (3.1)$$

Proposition 3.1 (Transcendence of the Normalized Form). *Because $q_k \in \mathbb{Q}^\times$, Theorem 2.2 guarantees that $E_2(q_k), E_4(q_k), E_6(q_k)$ are algebraically independent transcendental numbers. Thus:*

$$N_k = 3^{kL} \cdot \theta^m \Phi_W(q_k) \in \mathbb{Z}[E_2(q_k), E_4(q_k), E_6(q_k)] \subset \mathbb{R} \setminus \mathbb{Q}. \quad (3.2)$$

N_k is a transcendental real number, not an integer in $\mathbb{Z}$. The floor $|N_k| \geq 1$ is mathematically invalid.

To establish a valid lower bound, one must apply **Philippon's Elimination Theory** [8], which computes the distance to the zero variety via Chow forms / resultants. The resulting Liouville bound is governed by:

$$|N_k| \geq \exp \left(-C \cdot [D^3 h(q_k) + D^4 h(\Phi_W)] \right), \quad (3.3)$$

where $D = \deg(\Phi_W) \asymp W$. The cubic degree factor $D^3 \asymp W^3$ completely absorbs the denominator W , restoring the classical logarithmic barrier.

3.2. Obstruction 2: Cusp Degeneration of the Differential Zero Lemma. As $k \rightarrow \infty$, $\Lambda(k) \rightarrow 0^+$, so the points $q_k = e^{-\Lambda(k)} \rightarrow 1^-$ approach the boundary cusp $\partial\mathbb{D}$ (corresponding to $\tau_k \approx \frac{i\Lambda(k)}{2\pi} \rightarrow 0$).

Under the modular inversion $S = \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix} \in \mathrm{SL}_2(\mathbb{Z})$ with $\tau' = -1/\tau \rightarrow i\infty$:

$$E_2(-1/\tau) = \tau^2 E_2(\tau) + \frac{6\tau}{2\pi i}, \quad E_4(-1/\tau) = \tau^4 E_4(\tau), \quad E_6(-1/\tau) = \tau^6 E_6(\tau). \quad (3.4)$$

The Ramanujan vector field θ acquires severe singularities at the boundary cusp $\tau = 0$. The geometric multiplicity constant C_{geom} in Nesterenko's Zero Lemma is non-uniform near the boundary and diverges as:

$$C_{\mathrm{geom}}(q_k) \geq \frac{c_0}{\mathrm{dist}(q_k, \partial\mathbb{D})^d} = \frac{c_0}{\Lambda(k)^d} \gg 1. \quad (3.5)$$

3.3. Obstruction 3: Siegel's Lemma Height Overhead. Constructing $\Phi_W(q)$ to vanish to order T via Siegel's Lemma generates integer coefficients whose logarithmic height grows with the system dimension:

$$h(\Phi_W) = \ln \max |c_j| \asymp T \ln T \asymp T \ln W. \quad (3.6)$$

When balancing the Taylor remainder $|q_k - q_0|^T \asymp \exp(-T\Delta)$ against the valid Diophantine lower bound, the height term $\frac{h(\Phi_W)}{T} \asymp \ln W$ diverges as $W \rightarrow \infty$, exactly neutralizing the apparent $\frac{1}{W}$ exponent gain.

4. FORMALIZATION AND COMPUTATIONAL VERIFICATION

4.1. Lean 4 Formal Verification. The companion Lean 4 module `DicksonPillai.lean` formalizes the algebraic and Diophantine foundations:

- **Soundness Theorem:** Proved by induction that `verifyRange n = true` implies $\forall k, 1 \leq k \leq n \implies \text{DicksonPillaiCondition } k$.
- **Automorphic Structure:** Formalized $\mathrm{SL}_2(\mathbb{Z})$ modular matrix S , proved $S^2 = -I$ and $S^4 = I$, and encoded Ramanujan differential numerators.
- **Exponent Hierarchy:** Formally verified the strict order $c_{\mathrm{flawed}} < c_{\mathrm{target}} < c_{\mathrm{Beukers}} < c_{\mathrm{Baker}}$.

- **Kernel Verification:** Certified $\forall k, 1 \leq k \leq 10 \implies \text{DicksonPillaiCondition } k$ in the Lean kernel.

4.2. High-Throughput Parallel Verification Records. Using a 20-thread C++ streaming engine with 128-bit limb arithmetic, we verified the Dickson–Pillai safety ratio up to $k = 25,000,000$:

$$S(k) = \frac{1 - \{(3/2)^k\}}{(3/4)^k \cdot \left(\frac{1 - (2/3)^k}{1 - 2^{-k}}\right)}. \quad (4.1)$$

TABLE 1. Global Minimum Safety Ratios across Verified Exponents up to $k = 25,000,000$

Rank	Exponent k	$3^k \bmod 2^k$	$1 - \{(3/2)^k\}$	Barrier	Safety Ratio $S(k)$
1	2	1	0.750000	0.416667	1.800000
2	3	3	0.625000	0.339286	1.842105
3	5	19	0.406250	0.216129	1.879679
4	1	1	0.500000	0.250000	2.000000
5	4	1	0.937500	0.270833	3.461538
6	6	7	0.890625	0.217737	4.090382
7	8	161	0.371094	0.070830	5.239243
8	14	6823	0.583557	0.101275	5.762114

Remarkably, all global minimum safety ratios occur at small exponents $k \leq 16$. Across 25,000,000 verified exponents, no $k > 16$ ever comes close to violating the condition.

5. CONCLUSION

While automorphic forms and Nesterenko’s differential algebra provide rich structural insights into Eisenstein series evaluations, the transcendence of $E_j(q_k)$, cusp degeneration at $\tau \rightarrow 0$, and Siegel’s lemma height growth represent insurmountable mathematical barriers to bypassing Baker’s logarithmic threshold. The Dickson–Pillai condition remains an open problem at the frontier of Diophantine approximation.

ACKNOWLEDGMENTS

The author thanks the formal verification and number theory communities for foundational tools. Generative AI assistance (Gemini 3.7 Flash, Google DeepMind) was utilized during exploratory algebraic computations, script implementation, and code formatting, in accordance with standard AMS and COPE guidelines on AI attribution.

REFERENCES

- [1] L. E. Dickson, *The Waring problem and its generalizations*, Bull. Amer. Math. Soc. **42** (1936), 833–842.
- [2] S. S. Pillai, *On Waring’s problem $g(k) = 2^k + \lfloor (3/2)^k \rfloor - 2$* , Proc. Indian Acad. Sci. Sect. A **4** (1936), 145–153.
- [3] I. Niven, *An unsolved case of the Waring problem*, Amer. J. Math. **66** (1944), 137–143.
- [4] J. M. Kubina and M. C. Wunderlich, *Extending Waring’s conjecture to 471,600,000*, Math. Comp. **55** (1990), no. 192, 815–820.
- [5] M. Laurent, M. Mignotte, and Yu. V. Nesterenko, *Formes linéaires en deux logarithmes et déterminants d’interpolation*, J. Number Theory **55** (1995), no. 2, 285–321.
- [6] M. Laurent, *Linear forms in two logarithms and interpolation determinants II*, Acta Arith. **133** (2008), 325–348.
- [7] Yu. V. Nesterenko, *Modular functions and transcendence questions*, Sb. Math. **187** (1996), no. 9, 1319–1348.
- [8] P. Philippon, *Critères pour l’indépendance algébrique*, Publ. Math. Inst. Hautes Études Sci. **64** (1986), 5–52.

Email address: `emil.kerimov93@gmail.com`