# Gamma Function

Emil Kerimov

October 31, 2026

## 1 Definition

**Definition 1.1.** *The Gamma Function is defined as:*

$$\Gamma(z) = \int_0^\infty e^{-t}t^{z-1}dt$$

The first few evaluations of the Gamma function are:

$$\begin{aligned}
\Gamma(1) &= \int_0^\infty e^{-t}\ dt = -e^{-t}\Big|_0^\infty = 0 - (-1) = 1 \\
\Gamma(2) &= \int_0^\infty e^{-t}t\ dt = -e^{-t}t\Big|_0^\infty + \int_0^\infty e^{-t}\ dt = 1 \\
\Gamma(3) &= \int_0^\infty e^{-t}t^2\ dt = -e^{-t}t^2\Big|_0^\infty + 2\int_0^\infty e^{-t}t\ dt = 2 \\
\Gamma(4) &= \int_0^\infty e^{-t}t^3\ dt = -e^{-t}t^3\Big|_0^\infty + 3\int_0^\infty e^{-t}t\ dt = 6
\end{aligned} \tag{1}$$

## 2 Relation to the Factorial Function

Expanding on the pattern we found by evaluating the first few integer values of the Gamma function through integration by parts we obtain:

**Theorem 2.1.** $\Gamma(z+1) = z\Gamma(z)$
*Proof by induction*

*Assume true for k*

$$\Gamma(k+1) = k\Gamma(k)$$

*Proving for k+1*

$$\begin{aligned}
\Gamma(k+1) &= \int_0^\infty e^{-t}t^k\ dt \\
&= -e^{-t}t^k\Big|_0^\infty + k\int_0^\infty e^{-t}t^{k-1}\ dt \\
&= 0 - 0 + k\Gamma(k-1) \\
&= k\Gamma(k-1)
\end{aligned}$$

### 2.1 Integer Values

Therefore since $\Gamma(1) = 1$, then for positive integer values

$$\Gamma(n) = (n-1)! \tag{2}$$

## 3 Equivalent Integral Forms

$$\Gamma(z) = \int_0^\infty e^{-t} t^{z-1} dt \tag{3}$$

**Theorem 3.1.** *Starting from Equation 1.1, we can use the substitution* $t = -ln(u)$ *to obtain*

$$\Gamma(z) = \int_0^1 (-lnu)^{z-1} dt \tag{4}$$

*Proof*

$$t = -ln(u)$$
$$dt = -\frac{du}{u}$$
$$\Gamma(z) = \int_1^0 e^{ln(u)} (-ln(u))^{z-1} \frac{-du}{u}$$
$$\Gamma(z) = -\int_1^0 u(-ln(u))^{z-1} \frac{du}{u}$$
$$\Gamma(z) = -\int_1^0 (-ln(u))^{z-1} du$$
$$\Gamma(z) = \int_0^1 (-ln(u))^{z-1} du$$

## 4 Finite Gamma Expression

Define the finite gamma function (find out its real name)

$$\Gamma_p(x) = \frac{p!p^x}{x(x+1)(x+2)...(x+p)} = \frac{p^x}{x(\frac{x}{1}+1)(\frac{x}{2}+1)...(\frac{x}{p}+1)} \quad x > 0 \tag{5}$$

Note that this function has the following properties:

$$\Gamma_p(1) = \frac{p!p}{1(1+1)(1+2)...(1+p)} = \frac{p!p}{p!(1+p)} = \frac{p}{1+p} \tag{6}$$

$$\begin{aligned} \Gamma_p(x+1) &= \frac{p!pp^x}{(x+1)(x+2)...(x+p)(x+p+1)} \\ &= \frac{xp!pp^x}{x(x+1)(x+2)...(x+p)(x+p+1)} \\ &= \frac{xp}{x+p+1}\Gamma_p(x) \end{aligned} \tag{7}$$

### 4.1 Taking the limit to obtain Gamma

Using Equation 7 and assuming $\Gamma_\infty(x)$ converges we get:

$$\lim_{p\to\infty} \Gamma_p(x+1) = \lim_{p\to\infty} \frac{xp}{x+p+1}\Gamma_p(x) = x \lim_{p\to\infty} \Gamma_p(x) \tag{8}$$

Therefore using the above properties we see that if $\Gamma_\infty(x)$ converges then it must be equal to $\Gamma(x)$.

## 5 Euler Mascheroni constant

The Euler Mascheroni constant is defined as:

$$\gamma = \lim_{p\to\infty} \sum_{i=1}^{p} 1/i - ln(p) \approx 0.5772 \tag{9}$$

This constant can be interpreted as the difference between the sum of $1/x$ and the integral of $1/x$ from 1 to $\infty$. It is unknown whether this number is transcendental or even irrational! One can show numerically that if it is rational then the denominator must be greater than $10^{242080}$.

## 6 Weierstrass Formula

**Theorem 6.1.** *Let $\gamma$ be the Euler Mascheroni constant as defined in section 5, and $\Gamma$ be the Gamma function, then*

$$\frac{1}{\Gamma(x)} = x \cdot e^{x\gamma} \cdot \prod_{k=1}^{\infty} (1 + x/k) \cdot e^{-x/k} \tag{10}$$

*Proof*

*Re-writing* $p^x$

$$p^x = e^{xlnp} = e^{x[lnp - 1 - 1/2 - 1/3 - \cdots - 1/p]} e^{x + x/2 + x/3 + \cdots + x/p}$$

*Plugging* $p^x$ *into equation 5*

$$\Gamma_p(x) = \frac{e^{x[lnp - 1 - 1/2 - 1/3 - \cdots - 1/p]} e^{x + x/2 + x/3 + \cdots + x/p}}{x(\frac{x}{1} + 1)(\frac{x}{2} + 1)...(\frac{x}{p} + 1)}$$

$$\Gamma_p(x) = e^{x[lnp - 1 - 1/2 - 1/3 - \cdots - 1/p]} \cdot \frac{1}{x} \cdot \frac{e^x}{x+1} \cdot \frac{e^{x/2}}{x/2+1} \cdot \frac{e^{x/3}}{x/3+1} \cdot \ldots \cdot \frac{e^{x/p}}{x/p+1}$$

$$\Gamma_p(x) = e^{x[lnp - 1 - 1/2 - 1/3 - \cdots - 1/p]} \cdot \frac{1}{x} \cdot \prod_{k=1}^{p} \frac{e^{x/k}}{1 + x/k}$$

*Taking the limit as* $p \to \infty$

$$\Gamma(x) = \lim_{p \to \infty} \Gamma_p(x) = \lim_{p \to \infty} e^{x[lnp - 1 - 1/2 - 1/3 - \cdots - 1/p]} \cdot \frac{1}{x} \cdot \prod_{k=1}^{p} \frac{e^{x/k}}{1 + x/k}$$

$$\Gamma(x) = e^{-x\gamma} \cdot \frac{1}{x} \cdot \prod_{k=1}^{\infty} \frac{e^{x/k}}{1 + x/k}$$

*Equivalently*

$$\frac{1}{\Gamma(x)} = x \cdot e^{x\gamma} \cdot \prod_{k=1}^{\infty} (1 + x/k) \cdot e^{-x/k}$$

## 6.1 Symmetry around 1/2 relation

Using 6 we derive the following functional form.

**Theorem 6.2.**

$$\boxed{\frac{1}{\Gamma(x)} \cdot \frac{1}{\Gamma(1-x)} = \frac{sin(\pi x)}{\pi}} \tag{11}$$

*Proof*
*Starting from 6 and considering* $\frac{1}{\Gamma(x)} \cdot \frac{1}{\Gamma(-x)}$ *we obtain*

$$\frac{1}{\Gamma(x)} \cdot \frac{1}{\Gamma(-x)} = x \cdot e^{x\gamma} \cdot \prod_{k=1}^{\infty} (1 + x/k) \cdot e^{-x/k} \cdot -x \cdot e^{-x\gamma} \cdot \prod_{k=1}^{\infty} (1 - x/k) \cdot e^{x/k}$$

$$\frac{1}{\Gamma(x)} \cdot \frac{1}{\Gamma(-x)} = -x^2 \cdot \prod_{k=1}^{\infty} (1 - x^2/k^2)$$

*Using the functional form of* $\Gamma$, *we get* $\Gamma(1-x) = \Gamma(-x+1) = -x\Gamma(-x)$

$$\frac{1}{\Gamma(x)} \cdot \frac{1}{\Gamma(1-x)} = x \cdot \prod_{k=1}^{\infty} (1 - x^2/k^2)$$

*Using the product form of* $\frac{\sin(\pi x)}{\pi x} = \prod_{k=1}^{\infty} \left(1 - \frac{x^2}{k^2}\right)$ *, See Euler's Sine Product Form document*

$$\frac{1}{\Gamma(x)} \cdot \frac{1}{\Gamma(1-x)} = \frac{sin(\pi x)}{\pi}$$

## 6.2 Special Examples

Plugging in various values for $x$ we obtain certain values and relations for $\Gamma(x)$

$x = \frac{1}{2}$

$$\Gamma(\frac{1}{2}) = \sqrt{\frac{\pi}{sin(\frac{\pi}{2})}} = \sqrt{\pi} \tag{12}$$

$x = \frac{1}{3}$

$$\Gamma(\frac{1}{3})\Gamma(\frac{2}{3}) = \frac{\pi}{sin(\frac{\pi}{3})} = \frac{2\pi}{\sqrt{3}} \tag{13}$$

$x = \frac{1}{4}$

$$\Gamma(\frac{1}{4})\Gamma(\frac{3}{4}) = \frac{\pi}{sin(\frac{\pi}{4})} = \sqrt{2}\pi \tag{14}$$

**Theorem 6.3.** *$x = -\frac{1}{2}$ with equation 12*

$$\Gamma(-\frac{1}{2}) = -2\sqrt{\pi} \tag{15}$$

*Proof*

$$\Gamma(-\frac{1}{2}) \cdot \Gamma(1 - (-\frac{1}{2})) = \frac{\pi}{sin(-\frac{\pi}{2})}$$

$$\Gamma(-\frac{1}{2}) = -\frac{\pi}{\Gamma(\frac{3}{2})}$$

$$\Gamma(-\frac{1}{2}) = -\frac{\pi}{\frac{1}{2}\Gamma(\frac{1}{2})}$$

$$\Gamma(-\frac{1}{2}) = -\frac{\pi}{\frac{1}{2}\sqrt{\pi}}$$

$$\Gamma(-\frac{1}{2}) = -2\sqrt{\pi}$$

Note that equation 12 implies that

$$\Gamma(\frac{1}{2}) = \int_0^\infty \frac{1}{e^t\sqrt{t}}dt = \sqrt{\pi} \tag{16}$$

Since $\Gamma(x) = (x-1)!$ for integer values of $x$, this is where the idea of $(-\frac{1}{2})! = \sqrt{\pi}$ comes from.

# 7 Other relations

## 7.1 Gauss's Multiplication Formula

The multiplication formula, discovered by Carl Friedrich Gauss in 1812, generalizes Legendre's duplication formula (Section 10.5) to an arbitrary positive integer multiplier $n \geq 1$.

**Theorem 7.1.** *For any integer $n \geq 1$ and any complex number $x \notin \{0, -1/n, -2/n, \dots\}$:*

$$\boxed{\Gamma(x) \cdot \Gamma\left(x + \frac{1}{n}\right) \cdot \Gamma\left(x + \frac{2}{n}\right) \cdots \Gamma\left(x + \frac{n-1}{n}\right) = (2\pi)^{\frac{n-1}{2}} n^{\frac{1}{2} - nx} \Gamma(nx)} \tag{17}$$

*Proof*
*We prove the multiplication formula using Euler's limit product representation of the Gamma function from Section 4:*

$$\Gamma(z) = \lim_{p \to \infty} \frac{p!\, p^z}{z(z+1)(z+2)\cdots(z+p)} = \lim_{p \to \infty} \frac{p!\, p^z}{\prod_{j=0}^{p}(z+j)}$$

*For each $k \in \{0, 1, \dots, n-1\}$, evaluating at $z = x + \frac{k}{n}$:*

$$\begin{aligned}
\Gamma\left(x + \frac{k}{n}\right) &= \lim_{p \to \infty} \frac{p!\, p^{x+k/n}}{\prod_{j=0}^{p}\left(x + \frac{k}{n} + j\right)} \\
&= \lim_{p \to \infty} \frac{p!\, p^{x+k/n}}{\prod_{j=0}^{p} \frac{nx+jn+k}{n}} \\
&= \lim_{p \to \infty} \frac{n^{p+1} p!\, p^{x+k/n}}{\prod_{j=0}^{p}(nx+jn+k)}
\end{aligned}$$

*Multiplying these $n$ limits together for $k = 0, 1, \dots, n-1$:*

$$\prod_{k=0}^{n-1} \Gamma\left(x + \frac{k}{n}\right) = \lim_{p \to \infty} \frac{\prod_{k=0}^{n-1}\left(n^{p+1} p!\, p^{x+k/n}\right)}{\prod_{k=0}^{n-1}\prod_{j=0}^{p}(nx+jn+k)}$$

*Notice that as $k$ ranges from $0$ to $n-1$ and $j$ ranges from $0$ to $p$, the index $jn+k$ covers every integer from $0$ to $n(p+1)-1$ exactly once. Therefore, the denominator is:*

$$\prod_{k=0}^{n-1}\prod_{j=0}^{p}(nx+jn+k) = \prod_{m=0}^{n(p+1)-1}(nx+m)$$

*For the numerator:*

$$\begin{aligned}
\prod_{k=0}^{n-1}\left(n^{p+1} p!\, p^{x+k/n}\right) &= n^{n(p+1)} (p!)^n p^{nx + \sum_{k=0}^{n-1} \frac{k}{n}} \\
&= n^{n(p+1)} (p!)^n p^{nx + \frac{n-1}{2}}
\end{aligned}$$

*Now compare this with Euler's limit formula for $\Gamma(nx)$ with parameter $np$:*

$$\Gamma(nx) = \lim_{p \to \infty} \frac{(np)!\,(np)^{nx}}{\prod_{m=0}^{np}(nx+m)}$$

*Taking the ratio of the two expressions:*

$$\begin{aligned}
\frac{\prod_{k=0}^{n-1}\Gamma\left(x + \frac{k}{n}\right)}{\Gamma(nx)} &= \lim_{p \to \infty} \frac{n^{n(p+1)} (p!)^n p^{nx + \frac{n-1}{2}}}{(np)!\,(np)^{nx}} \cdot \frac{\prod_{m=0}^{np}(nx+m)}{\prod_{m=0}^{n(p+1)-1}(nx+m)} \\
&= n^{-nx} \lim_{p \to \infty} \frac{n^{np+n} (p!)^n p^{\frac{n-1}{2}}}{(np)!} \cdot \frac{1}{\prod_{m=np+1}^{n(p+1)-1}(nx+m)}
\end{aligned}$$

*The remaining product has* $(n(p+1)-1)-np = n-1$ *factors:*

$$\prod_{m=np+1}^{n(p+1)-1}(nx+m) = (nx+np+1)(nx+np+2)\cdots(nx+np+n-1)$$

*As* $p\to\infty$*, this product is asymptotic to* $(np)^{n-1} = n^{n-1}p^{n-1}$*. Substituting this asymptotic behavior:*

$$\begin{aligned}\frac{\prod_{k=0}^{n-1}\Gamma\left(x+\frac{k}{n}\right)}{\Gamma(nx)} &= n^{-nx}\lim_{p\to\infty}\frac{n^{np+n}(p!)^n p^{\frac{n-1}{2}}}{(np)!\cdot n^{n-1}p^{n-1}}\\ &= n^{1-nx}\lim_{p\to\infty}\frac{n^{np}(p!)^n p^{\frac{1-n}{2}}}{(np)!}\end{aligned}$$

*Using Stirling's asymptotic formula for factorials (Section 11):*

$$\begin{aligned}p! &\sim \sqrt{2\pi p}\left(\frac{p}{e}\right)^p \implies (p!)^n \sim (2\pi)^{n/2}p^{n/2}\left(\frac{p}{e}\right)^{np}\\ (np)! &\sim \sqrt{2\pi np}\left(\frac{np}{e}\right)^{np} = \sqrt{2\pi n}\,n^{np}p^{1/2}\left(\frac{p}{e}\right)^{np}\end{aligned}$$

*Substituting these into the limit:*

$$\begin{aligned}\lim_{p\to\infty}\frac{n^{np}(p!)^n p^{\frac{1-n}{2}}}{(np)!} &= \lim_{p\to\infty}\frac{n^{np}\cdot(2\pi)^{n/2}p^{n/2}(p/e)^{np}\cdot p^{\frac{1-n}{2}}}{\sqrt{2\pi n}\,n^{np}p^{1/2}(p/e)^{np}}\\ &= \frac{(2\pi)^{n/2}}{\sqrt{2\pi n}}\lim_{p\to\infty}\frac{p^{n/2+(1-n)/2}}{p^{1/2}}\\ &= (2\pi)^{\frac{n-1}{2}}n^{-1/2}\end{aligned}$$

*Multiplying by* $n^{1-nx}\Gamma(nx)$*:*

$$\prod_{k=0}^{n-1}\Gamma\left(x+\frac{k}{n}\right) = n^{1-nx}\cdot(2\pi)^{\frac{n-1}{2}}n^{-1/2}\Gamma(nx) = (2\pi)^{\frac{n-1}{2}}n^{\frac{1}{2}-nx}\Gamma(nx)$$

*which completes the proof.*

**Corollary 7.1.1.** *Multiplying both sides of Equation (17) by* $x$ *and taking the limit as* $x\to 0$*, we have* $\lim_{x\to 0}x\Gamma(x) = \Gamma(1) = 1$*, and* $\lim_{x\to 0}x\Gamma(nx) = \frac{1}{n}\lim_{x\to 0}(nx)\Gamma(nx) = \frac{1}{n}$*. Therefore:*

$$\boxed{\Gamma\left(\frac{1}{n}\right)\cdot\Gamma\left(\frac{2}{n}\right)\cdots\Gamma\left(\frac{n-1}{n}\right) = \frac{(2\pi)^{\frac{n-1}{2}}}{\sqrt{n}}} \tag{18}$$

**Example 7.2.** *For* $n=2$*, Gauss's multiplication formula reduces directly to Legendre's duplication formula (Theorem 10.5):*

$$\Gamma(x)\Gamma\left(x+\frac{1}{2}\right) = (2\pi)^{1/2}2^{\frac{1}{2}-2x}\Gamma(2x) = \frac{\sqrt{\pi}}{2^{2x-1}}\Gamma(2x)$$

*For* $n=3$*, Equation (18) gives:*

$$\Gamma\left(\frac{1}{3}\right)\Gamma\left(\frac{2}{3}\right) = \frac{(2\pi)^{2/2}}{\sqrt{3}} = \frac{2\pi}{\sqrt{3}}$$

*which perfectly agrees with Euler's reflection formula (11):*

$$\Gamma\left(\frac{1}{3}\right)\Gamma\left(1-\frac{1}{3}\right) = \frac{\pi}{\sin(\pi/3)} = \frac{\pi}{\sqrt{3}/2} = \frac{2\pi}{\sqrt{3}}$$

## 7.2 Relation to the Riemann Zeta Function

**Theorem 7.3.** *For* $\mathrm{Re}(s) > 1$*, the Gamma function is related to the Riemann Zeta function* $\zeta(s) = \sum_{n=1}^{\infty} \frac{1}{n^s}$ *by:*

$$\boxed{\Gamma(s)\zeta(s) = \int_0^\infty \frac{u^{s-1}}{e^u - 1}\,\mathrm{d}u} \tag{19}$$

*Proof*
*Starting from the integral definition of the Gamma function (Equation 1.1):*

$$\Gamma(s) = \int_0^\infty t^{s-1}e^{-t}\,\mathrm{d}t \qquad (\mathrm{Re}(s) > 0)$$

*For any positive integer* $n \geq 1$*, substitute* $t = nu$*, so that* $\mathrm{d}t = n\,\mathrm{d}u$*:*

$$\Gamma(s) = \int_0^\infty (nu)^{s-1}e^{-nu}n\,\mathrm{d}u = n^s \int_0^\infty u^{s-1}e^{-nu}\,\mathrm{d}u$$

*Dividing by* $n^s$*:*

$$\frac{\Gamma(s)}{n^s} = \int_0^\infty u^{s-1}e^{-nu}\,\mathrm{d}u$$

*Summing both sides over all positive integers* $n \geq 1$*:*

$$\Gamma(s)\sum_{n=1}^{\infty}\frac{1}{n^s} = \sum_{n=1}^{\infty}\int_0^\infty u^{s-1}e^{-nu}\,\mathrm{d}u$$

*For* $\mathrm{Re}(s) > 1$*, we can interchange the summation and the integral by the monotone convergence theorem (or dominated convergence):*

$$\begin{aligned}\Gamma(s)\zeta(s) &= \int_0^\infty u^{s-1}\left(\sum_{n=1}^{\infty}e^{-nu}\right)\mathrm{d}u \\ &= \int_0^\infty u^{s-1}\left(\frac{e^{-u}}{1-e^{-u}}\right)\mathrm{d}u \\ &= \int_0^\infty \frac{u^{s-1}}{e^u - 1}\,\mathrm{d}u\end{aligned}$$

*which proves the identity.*

# 8 Digamma function

## 8.1 Definition

Define the Digamma function

$$\Psi(x) = \frac{\partial}{\partial x}ln(\Gamma(x)) = \frac{\Gamma'(x)}{\Gamma(x)} \tag{20}$$

## 8.2 Series Representation

**Theorem 8.1.** *Let $\gamma$ be the Euler Mascheroni constant as defined in section 5 then,*

$$\boxed{\Psi(x) = -\gamma - \frac{1}{x} + \sum_{k=1}^{\infty}\left(\frac{1}{k} - \frac{1}{x+k}\right)} \tag{21}$$

*Proof*

$$\Psi(x) = \frac{\partial}{\partial x} ln(\Gamma(x))$$

*Using equation 10*

$$\Psi(x) = -\frac{\partial}{\partial x} ln(x \cdot e^{x\gamma} \cdot \prod_{k=1}^{\infty}(1 + x/k) \cdot e^{-x/k})$$

$$\Psi(x) = -\frac{\partial}{\partial x}\left[ln(x) + x\gamma + \sum_{k=1}^{\infty} ln(1 + x/k) - \frac{x}{k}\right]$$

$$\Psi(x) = -\frac{1}{x} - \gamma - \sum_{k=1}^{\infty} \frac{1}{k} \cdot \frac{1}{1 + x/k} - \frac{1}{k}$$

$$\Psi(x) = -\frac{1}{x} - \gamma + \sum_{k=1}^{\infty} \frac{1}{k} - \frac{1}{x+k}$$

## 8.3 Derivatives

**Theorem 8.2.**

$$\boxed{\Psi'(x) = \sum_{k=1}^{\infty} \frac{1}{(k + x - 1)^2}} \tag{22}$$

*Proof*

*Using equation 21*

$$\Psi(x) = -\gamma - \frac{1}{x} + \sum_{k=1}^{\infty}\left(\frac{1}{k} - \frac{1}{x+k}\right)$$

$$\Psi'(x) = \frac{1}{x^2} + \sum_{k=1}^{\infty} \frac{1}{(x+k)^2}$$

*Including the first term in the sum*

$$\Psi'(x) = \sum_{k=1}^{\infty} \frac{1}{(x + k - 1)^2}$$

**Theorem 8.3.**

$$\boxed{\Psi^n(x) = \sum_{k=1}^{\infty} \frac{(-1)^{n+1} n!}{(k + x - 1)^{n+1}}} \tag{23}$$

*Proof*

*Take repeated derivatives of equation 22*

## 8.4 Special values

**Theorem 8.4.**

$$\Psi^n(x+1) = \Psi^n(x) + \frac{(-1)^n n!}{x^{n+1}} \tag{24}$$

*Proof*

$$\Psi^n(x+1) = \sum_{k=1}^{\infty} \frac{(-1)^{n+1} n!}{(k+x)^{n+1}}$$

$$\Psi^n(x+1) = \frac{(-1)^n n!}{x^{n+1}} + \sum_{k=2}^{\infty} \frac{(-1)^{n+1} n!}{(k+x)^{n+1}}$$

$$\Psi^n(x+1) = \frac{(-1)^n n!}{x^{n+1}} + \sum_{k=1}^{\infty} \frac{(-1)^{n+1} n!}{(k+x-1)^{n+1}}$$

$$\Psi^n(x+1) = \frac{(-1)^n n!}{x^{n+1}} + \Psi^n(x)$$

**Theorem 8.5.** *Reflection formula*

$$\Psi(1-x) = \Psi(x) + \pi cot(\pi x) \tag{25}$$

*Proof*

$$\Psi(1-x) = -\gamma - \frac{1}{1-x} + \sum_{k=1}^{\infty} \Big( \frac{1}{k} - \frac{1}{k+1-x} \Big)$$

*Adding and subtracting* $\Psi(x)$

$$\Psi(1-x) = -\gamma - \frac{1}{1-x} + \sum_{k=1}^{\infty} \Big( \frac{1}{k} - \frac{1}{k+1-x} \Big) + \Psi(x) - \Psi(x)$$

$$\Psi(1-x) = -\gamma - \frac{1}{1-x} + \sum_{k=1}^{\infty} \Big( \frac{1}{k} - \frac{1}{k+1-x} \Big) + \Psi(x) - \left( -\gamma - \frac{1}{x} + \sum_{k=1}^{\infty} \Big( \frac{1}{k} - \frac{1}{x+k} \Big) \right)$$

*Simplifying*

$$\Psi(1-x) = \Psi(x) + \frac{1}{x} - \frac{1}{1-x} + \sum_{k=1}^{\infty} \Big( \frac{1}{k} - \frac{1}{k+1-x} - \frac{1}{k} + \frac{1}{x+k} \Big)$$

$$\Psi(1-x) = \Psi(x) + \frac{1}{x} - \frac{1}{1-x} + \sum_{k=1}^{\infty} \Big( \frac{1}{x+k} - \frac{1}{k+1-x} \Big)$$

$$\Psi(1-x) = \Psi(x) + \sum_{k=0}^{\infty} \Big( \frac{1}{x+k} + \frac{1}{x-k-1} \Big)$$

$$\Psi(1-x) = \Psi(x) + \sum_{k=-\infty}^{\infty} \Big( \frac{1}{x+k} \Big)$$

*Using the series* [1] *for* $\pi cot(\pi x) = \sum_{k=-\infty}^{\infty} \frac{1}{x+k}$

$$\Psi(1-x) = \Psi(x) + \pi cot(\pi x)$$

**Theorem 8.6.**

$$\Psi(2x) = \frac{1}{2} \cdot \Psi(x) + \frac{1}{2} \cdot \Psi(x + \frac{1}{2}) + ln(2) \tag{26}$$

*Proof*
*TODO*

$$\Psi(1) = \Gamma'(1) = -\gamma \tag{27}$$

$$\Psi^{(n)}(1) = (-1)^{n+1} n! \zeta(n+1) \tag{28}$$

Where $\zeta(z)$ is the Riemann Zeta function defined as $\zeta(z) = \sum_{k=1}^{\infty} \frac{1}{k^z}$. CONFIRM THIS?

$$\Psi(n) = \frac{\Gamma'(n)}{\Gamma(n)} = -\gamma + \sum_{k=1}^{n-1} \frac{1}{k} = -\gamma + H_{n-1} \tag{29}$$

Where $H_n$ is the sum of the $n$ first harmonic numbers. Note that this implies that the derivative of the $\Gamma$ function is diverges to $\infty$.

# 9 Using The Gamma Function for Euler Integrals

**Theorem 9.1.** *Let $p = x + yi$ and $\alpha = tan^{-1}\left(\frac{y}{x}\right)$ then*

$$\boxed{\frac{\Gamma(s)}{n|p|^s} sin(\alpha s) = \int_0^{\infty} u^{ns-1} e^{-xu^n} sin(yu^n) du} \tag{30}$$

$$\boxed{\frac{\Gamma(s)}{n|p|^s} cos(\alpha s) = \int_0^{\infty} u^{ns-1} e^{-xu^n} cos(yu^n) du} \tag{31}$$

*Proof*

$$\Gamma(s) = \int_0^{\infty} e^{-t} t^{s-1} dt$$

*Using substitution $t = p \cdot u^n$ hence $dt = p \cdot n \cdot u^{n-1} \cdot du$*

$$\Gamma(s) = \int_0^{\infty} p^{n-1} u^{ns-n} e^{-pu} npu^{n-1} du = n \int_0^{\infty} p^s u^{ns-1} e^{-pu^n} du$$

$$\frac{\Gamma(s)}{np^s} = \int_0^{\infty} u^{ns-1} e^{-pu^n} du$$

*Assuming $p$ is a complex number with real part $x$, imaginary part $y$*
*Taking the conjugate of both sides, we obtain a second equation.*

$$\frac{\Gamma(s)}{n\bar{p}^s} = \int_0^{\infty} u^{ns-1} e^{-\bar{p}u^n} du$$

*Note that $p = x + yi = |p|e^{i\alpha}$ and $\bar{p} = x - yi = |p|e^{-i\alpha}$ where $\alpha = tan^{-1}\left(\frac{y}{x}\right)$*

[1] See the Euler's Sine Product formula document.

*Adding and subtracting both equations we get*

$$\frac{\Gamma(s)}{n}\Big[\frac{1}{\bar{p}^s} \pm \frac{1}{p^s}\Big] = \int_0^\infty u^{ns-1}\Big[e^{-\bar{p}u^n} \pm e^{-pu^n}\Big]du$$

$$\frac{\Gamma(s)}{n|p|^s}\Big[\frac{1}{e^{-i\alpha s}} \pm \frac{1}{e^{i\alpha s}}\Big] = \int_0^\infty u^{ns-1}\Big[e^{-xu^n+yu^n i} \pm e^{-xu^n-yu^n i}\Big]du$$

$$\frac{\Gamma(s)}{n|p|^s}\Big[e^{i\alpha s} \pm e^{-i\alpha s}\Big] = \int_0^\infty u^{ns-1}e^{-xu^n}\Big[e^{yu^n i} \pm e^{-yu^n i}\Big]du$$

*With addition we get*

$$\frac{\Gamma(s)}{n|p|^s}\Big[2cos(\alpha s)\Big] = \int_0^\infty u^{ns-1}e^{-xu^n}\Big[2cos(yu^n)\Big]du$$

$$\frac{\Gamma(s)}{n|p|^s}cos(\alpha s) = \int_0^\infty u^{ns-1}e^{-xu^n}cos(yu^n)\Big]du$$

*With subtraction we get*

$$\frac{\Gamma(s)}{n|p|^s}\Big[2isin(\alpha s)\Big] = \int_0^\infty u^{ns-1}e^{-xu^n}\Big[2isin(yu^n)\Big]du$$

$$\frac{\Gamma(s)}{n|p|^s}sin(\alpha s) = \int_0^\infty u^{ns-1}e^{-xu^n}sin(yu^n)\Big]du$$

## 9.1 Examples

### 9.1.1 Sinc Integral $\int \frac{\sin(x)}{x}$

**Theorem 9.2.**

$$\boxed{\int_0^\infty \frac{\sin(x)}{x} = \frac{\pi}{2}} \tag{32}$$

*Proof*

*Starting from equation 30*

$$\frac{\Gamma(s)}{n|p|^s}sin(\alpha s) = \int_0^\infty u^{ns-1}e^{-xu^n}sin(yu^n)du$$

*setting* $y = 1, n = 1, x = 0$ *and taking the limit as* $s \to 0$ *we obtain*

$$|p| = 1, \alpha = tan^{-1}(1/0) = \pi/2$$

$$\lim_{s\to 0}\Gamma(s)sin(s\frac{\pi}{2}) = \lim_{s\to 0}\int_0^\infty u^{s-1}sin(u)du = \int_0^\infty \frac{sin(u)}{u}du$$

*To evaluate the RHS limit we use equation 11* $\Gamma(s) = \dfrac{\pi}{\Gamma(1-s)sin(\pi s)}$

$$\lim_{s\to 0}\Gamma(s)sin(s\frac{\pi}{2}) = \lim_{s\to 0}\frac{\pi}{\Gamma(1-s)sin(\pi s)}sin(s\frac{\pi}{2})$$

*We can split this limit into the product of 3 limits since all of these limits exist*

$$\lim_{s\to 0}\Gamma(s)sin(s\frac{\pi}{2}) = \lim_{s\to 0}\frac{\frac{\pi}{2}}{\Gamma(1-s)}\cdot\lim_{s\to 0}\frac{\pi}{sin(\pi s)}\cdot\lim_{s\to 0}\frac{sin(s\frac{\pi}{2})}{\frac{\pi}{2}}$$

$$\lim_{s\to 0}\Gamma(s)sin(s\frac{\pi}{2}) = \frac{\pi}{2}$$

$$\int_0^\infty \frac{sin(u)}{u}du = \frac{\pi}{2}$$

*since* $\dfrac{sin(u)}{u}$ *is an even function*

$$\int_{-\infty}^\infty \frac{sin(u)}{u}du = \pi$$

### 9.1.2 $\int \sin(x^2)$

**Theorem 9.3.**

$$\boxed{\int_0^\infty \sin(x^2) = \frac{\sqrt{2\pi}}{4}} \tag{33}$$

*Proof*

*Starting from equation 30*

$$\frac{\Gamma(s)}{n|p|^s}sin(\alpha s) = \int_0^\infty u^{ns-1}e^{-xu^n}sin(yu^n)du$$

*setting* $y = 1, n = 2, x = 0$ *and* $ns - 1 = 0$ *we obtain*

$$|p| = 1, \alpha = tan^{-1}(1/0) = \pi/2, s = 1/2$$

$$\frac{\Gamma(\frac{1}{2})sin(\frac{\pi}{4})}{2} = \int_0^\infty sin(u^2)du$$

*using equation 12:* $\Gamma(\frac{1}{2}) = \sqrt{\pi}$

$$\int_0^\infty sin(u^2)du = \frac{\sqrt{\pi}\frac{1}{\sqrt{2}}}{2} = \frac{\sqrt{2\pi}}{4}$$

### 9.1.3 $\int \cos(x^2)$

**Theorem 9.4.**

$$\boxed{\int_0^\infty \cos(x^2) = \frac{\sqrt{2\pi}}{4}} \tag{34}$$

*Proof*
*Starting from equation 31*

$$\frac{\Gamma(s)}{n|p|^s} cos(\alpha s) = \int_0^\infty u^{ns-1} e^{-xu^n} cos(yu^n) du$$

*setting* $y = 1, n = 2, x = 0$ *and* $ns - 1 = 0$ *we obtain*

$$|p| = 1, \alpha = tan^{-1}(1/0) = \pi/2, s = 1/2$$

$$\frac{\Gamma(\frac{1}{2}) cos(\frac{\pi}{4})}{2} = \int_0^\infty cos(u^2) du$$

*using equation 12:* $\Gamma(\frac{1}{2}) = \sqrt{\pi}$

$$\int_0^\infty cos(u^2) du = \frac{\sqrt{\pi}\frac{1}{\sqrt{2}}}{2} = \frac{\sqrt{2\pi}}{4}$$

### 9.1.4 $\int \frac{\sin(x^2)}{x^2}$

**Theorem 9.5.**

$$\boxed{\int_0^\infty \frac{\sin(x^2)}{x^2} = \sqrt{\frac{\pi}{2}}} \tag{35}$$

*Proof*
*Starting from equation 30*

$$\frac{\Gamma(s)}{n|p|^s} sin(\alpha s) = \int_0^\infty u^{ns-1} e^{-xu^n} sin(yu^n) du$$

*setting* $y = 1, n = 2, x = 0$ *and* $ns - 1 = -2$ *we obtain*

$$|p| = 1, \alpha = tan^{-1}(1/0) = \pi/2, s = -1/2$$

$$\frac{\Gamma(\frac{-1}{2}) sin(-\frac{\pi}{4})}{2} = \int_0^\infty \frac{sin(u^2)}{u^2} du$$

*using equation 15:* $\Gamma(-\frac{1}{2}) = -2\sqrt{\pi}$

$$\int_0^\infty \frac{sin(u^2)}{u^2} du = -\frac{\Gamma(\frac{-1}{2})}{2\sqrt{2}} = \sqrt{\frac{\pi}{2}}$$

### 9.1.5 $\int \frac{\sin(x^n)}{x^n}$

**Theorem 9.6.**

$$\boxed{\int_0^\infty \frac{\sin(x^n)}{x^n} = \frac{\Gamma(\frac{1}{n})}{n-1} \cdot cos(\frac{\pi}{2n}) \quad , n > 1} \tag{36}$$

*Proof*
*Starting from equation 30*

$$\frac{\Gamma(s)}{n|p|^s}sin(\alpha s) = \int_0^\infty u^{ns-1}e^{-xu^n}sin(yu^n)du$$

*setting* $y = 1, x = 0$ *and* $ns - 1 = -n$ *we obtain*

$$|p| = 1, \alpha = tan^{-1}(1/0) = \pi/2, s = 1/n - 1$$

$$\frac{\Gamma(\frac{1}{n} - 1)sin(\frac{\pi}{2n} - \frac{\pi}{2})}{n} = \int_0^\infty \frac{sin(u^n)}{u^n}du$$

*using equation 2.1 and assuming* $n \neq 1$*:* $\Gamma(k+1) = k\Gamma(k)$

$$\int_0^\infty \frac{sin(u^n)}{u^n}du = \frac{\Gamma(\frac{1}{n})}{n(\frac{1}{n} - 1)}sin(\frac{\pi}{2n} - \frac{\pi}{2})$$

$$\int_0^\infty \frac{sin(u^n)}{u^n}du = \frac{\Gamma(\frac{1}{n})}{1 - n)}sin(\frac{\pi}{2n} - \frac{\pi}{2})$$

$$\textit{using } \sin\left(x - \frac{\pi}{2}\right) = -\cos(x)$$

$$\int_0^\infty \frac{sin(u^n)}{u^n}du = \frac{\Gamma(\frac{1}{n})}{n - 1} \cdot cos(\frac{\pi}{2n})$$

# 10 Relation to the Beta Function

## 10.1 Definition

The Beta Function is defined as:

**Definition 10.1.**

$$\beta(x,y) = \int_0^1 t^{x-1}(1-t)^{y-1}dt$$

## 10.2 Symmetry

**Theorem 10.1.**

$$\beta(x,y) = \beta(y,x) \tag{37}$$

*Proof*
*Starting from the definition 10.1*

$$\beta(x,y) = \int_0^1 t^{x-1}(1-t)^{y-1}dt$$

*setting* $u = 1-t$, *hence* $du = -dt$ *we obtain*

$$\beta(x,y) = \int_1^0 (1-u)^{x-1}u^{y-1} - du$$

*Switching the bounds of integration we get*

$$\beta(x,y) = \int_0^1 (1-u)^{x-1}u^{y-1}du = \beta(y,x)$$

## 10.3 Relation to Gamma

**Theorem 10.2.**

$$\boxed{\beta(x,y) = \frac{\Gamma(x)\Gamma(y)}{\Gamma(x+y)}} \tag{38}$$

*Proof*

*Starting from the definition 10.1*

$$\beta(x,y) = \int_0^1 t^{x-1}(1-t)^{y-1}dt$$

*setting* $t = \frac{1}{1+u}$, *hence* $dt = -\frac{1}{(1+u)^2}du$ *we obtain*

$$\beta(x,y) = \int_\infty^0 (\frac{1}{1+u})^{x-1}(1-\frac{1}{1+u})^{y-1}\frac{1}{(1+u)^2}(-du)$$

*Switching the bounds of integration and simplifying we get*

$$\beta(x,y) = \int_0^\infty (\frac{1}{1+u})^{x-1}(\frac{u}{1+u})^{y-1}\frac{1}{(1+u)^2}du$$

*collecting powers of* $(1+u)$

$$\beta(x,y) = \int_0^\infty \frac{u^{y-1}}{(1+u)^{x+y}}du$$

*Using the Gamma definition 1.1 and applying a change of variables* $\alpha w = t$, *hence* $dt = \alpha dw$

$$\Gamma(z) = \alpha^z \int_0^\infty e^{-\alpha w}w^{z-1}dw$$

*setting* $z = x+y$ *and* $\alpha = 1+u$, *we obtain*

$$\Gamma(x+y) = (1+u)^{x+y}\int_0^\infty e^{-(1+u)w}w^{x+y-1}dw$$

*Combining with the Beta function integral we obtained*

$$\beta(x,y)\Gamma(x+y) = \int_0^\infty \Gamma(x+y)\frac{u^{y-1}}{(1+u)^{x+y}}du$$

$$\beta(x,y)\Gamma(x+y) = \int_0^\infty \Big((1+u)^{x+y}\int_0^\infty e^{-(1+u)w}w^{x+y-1}dw\Big)\frac{u^{y-1}}{(1+u)^{x+y}}du$$

*Simplfying*

$$\beta(x,y)\Gamma(x+y) = \int_0^\infty\int_0^\infty e^{-(1+u)w}w^{x+y-1}u^{y-1}dwdu$$

*Rearranging*

$$\beta(x,y)\Gamma(x+y) = \int_0^\infty \Big(\int_0^\infty e^{-uw}u^{y-1}du\Big)e^{-w}w^{x+y-1}dw$$

*Using* $\Gamma(z) = \alpha^z\int_0^\infty e^{-\alpha u}u^{z-1}du$

$$\beta(x,y)\Gamma(x+y) = \int_0^\infty \frac{\Gamma(y)}{w^y}e^{-w}w^{x+y-1}dw$$

*Simplifying and moving* $\Gamma(y)$ *outside the integral*

$$\beta(x,y)\Gamma(x+y) = \Gamma(y)\int_0^\infty e^{-w}w^{x-1}dw$$

*Using the definition of the* $\Gamma(z)$

$$\beta(x,y)\Gamma(x+y) = \Gamma(y)\Gamma(x)$$

$$\beta(x,y) = \frac{\Gamma(y)\Gamma(x)}{\Gamma(x+y)}$$

## 10.4 Relation to integral of powers of sin(x) and cos(x)

**Theorem 10.3.**

$$\boxed{\int_0^{\pi/2} sin^n(t)cos^m(t)dt = \frac{1}{2}\beta\Big(\frac{n+1}{2}, \frac{m+1}{2}\Big) = \frac{\Gamma(\frac{n+1}{2})\Gamma(\frac{m+1}{2})}{\Gamma(\frac{n+m}{2}+1)}} \tag{39}$$

*Proof*

$$A(m,n) = \int_0^{\pi/2} sin^n(t)cos^m(t)dt$$

*setting* $sin^2(t) = u$, *hence* $2\sin(t)\cos(t)dt = du$ *we obtain*

*note* $cos^2(t) = 1-u$, *and* $sin^2(0) = 0$ *and* $sin^2\left(\frac{\pi}{2}\right) = 1$

$$A(m,n) = \int_0^1 u^{\frac{n}{2}}(1-u)^{\frac{m}{2}}\frac{du}{2\sqrt{u}\sqrt{1-u}}$$

*Simplifying the powers we get*

$$A(m,n) = \frac{1}{2}\int_0^1 u^{\frac{n}{2}-\frac{1}{2}}(1-u)^{\frac{m}{2}-\frac{1}{2}}du$$

*Writing it in the Beta function form we get*

$$A(m,n) = \frac{1}{2}\int_0^1 u^{\frac{n+1}{2}-1}(1-u)^{\frac{m+1}{2}-1}du = \frac{1}{2}\beta(\frac{n+1}{2}, \frac{m+1}{2})$$

## 10.5 Proof of the Legendre duplication formula

**Theorem 10.4.**

$$\boxed{\Gamma(x)\cdot\Gamma(x+\frac{1}{2}) = \frac{\sqrt{\pi}}{2^{2x-1}}\Gamma(2x)} \tag{40}$$

*Proof*

*Using the relation to the Beta function, and the definition of the Beta function*

$$\frac{\Gamma(x)\Gamma(y)}{\Gamma(x+y)} = \int_0^1 t^{x-1}(1-t)^{y-1}dt$$

*let* $x = y = s$

$$\frac{\Gamma(s)\Gamma(s)}{\Gamma(2s)} = \int_0^1 t^{s-1}(1-t)^{s-1}dt$$

*let* $t = \frac{1+x}{2}$, *hence* $dt = \frac{dx}{2}$

*note when* $t = 0$ *then* $x = -1$, *and when* $t = 1$ *then* $x = 1$

$$\frac{\Gamma(s)\Gamma(s)}{\Gamma(2s)} = \int_{-1}^1 (\frac{1+x}{2})^{s-1}(\frac{1-x}{2})^{s-1}\frac{dx}{2}$$

*Simplifying the powers of 2 and rearranging*

$$2^{2s-1}\Gamma(s)\Gamma(s) = \Gamma(2s)\int_{-1}^1 (1+x)^{s-1}(1-x)^{s-1}dx$$

$$2^{2s-1}\Gamma(s)\Gamma(s) = \Gamma(2s)\int_{-1}^1 (1-x^2)^{s-1}dx$$

*Notice that the integral is even around 0, hence*

$$2^{2s-1}\Gamma(s)\Gamma(s) = 2\Gamma(2s)\int_0^1 (1-x^2)^{s-1}dx$$

*To evaluate the integral, notice that after a substitution of* $u = x^2$ *the integral is just* $\beta(1/2, s)$

$$\beta(1/2, s) = \int_0^1 t^{-\frac{1}{2}}(1-t)^{s-1}dt$$

*Let* $t = x^2$, *hence* $dt = 2xdx$

$$\beta(1/2, s) = \int_0^1 \frac{1}{x}(1-x^2)^{s-1}2xdx = 2\int_0^1 (1-x^2)^{s-1}dx$$

*Hence*

$$2^{2s-1}\Gamma(s)\Gamma(s) = \Gamma(2s)\beta(1/2, s)$$

*Using theorem 10.2*

$$2^{2s-1}\Gamma(s)\Gamma(s) = \Gamma(2s)\frac{\Gamma(1/2)\Gamma(s)}{\Gamma(s+1/2)}$$

*Rearranging and using equation 12*

$$\Gamma(s)\Gamma(s+1/2) = \Gamma(2s)\frac{\sqrt{\pi}}{2^{2s-1}}$$

# 11 Approximations

## 11.1 Stirling's Approximation

**Theorem 11.1.** *The basic approximation of the factorial function for large n is*

$$\boxed{n! \approx (\frac{n}{e})^n \sqrt{2\pi n}} \tag{41}$$

*Proof*
*Starting from the definition of the Gamma function 1.1*

$$\Gamma(n+1) = n! = \int_0^\infty e^{-t} t^n dt$$

*the idea is to take the log of the integrand and then slightly perturb it*

$$ln(e^{-t}t^n) = nln(t) - t$$

$$let\ t = n + \epsilon$$

$$\ln(e^{-t}t^n) = nln(n+\epsilon) - (n+\epsilon) = nln(n) + nln(1+\frac{\epsilon}{n}) - (n+\epsilon)$$

$$for\ large\ n,\ \frac{\epsilon}{n} \ll 1$$

*Using the Taylor-Expansion of* $\ln(1+x)$

$$\ln(e^{-t}t^n) = n\Big(ln(n) + \sum_{k=1}^{\infty} \frac{(-1)^{k+1}}{k} (\frac{\epsilon}{n})^k\Big) - (n+\epsilon)$$

$$\ln(e^{-t}t^n) = nln(n) - n + \sum_{k=1}^{\infty} \frac{(-1)^{k+1}}{k} (\frac{\epsilon^k}{n^{k-1}}) - \epsilon$$

$$\ln(e^{-t}t^n) = nln(n) - n - \frac{\epsilon^2}{2n} + \frac{\epsilon^3}{3n^2} - \frac{\epsilon^4}{4n^3} + \dots$$

$$\ln(e^{-t}t^n) \approx nln(n) - n - \frac{\epsilon^2}{2n}$$

$$e^{-t}t^n \approx \frac{n^n}{e^n} e^{-\frac{\epsilon^2}{2n}}$$

$$n! \approx \int_{-n}^{\infty} \frac{n^n}{e^n} e^{-\frac{\epsilon^2}{2n}} d\epsilon$$

$$n! \approx \frac{n^n}{e^n} \int_{-n}^{\infty} e^{\frac{-\epsilon^2}{2n}} d\epsilon$$

*Assuming n is large enough*

$$n! \approx \frac{n^n}{e^n} \int_{-\infty}^{\infty} e^{-\frac{\epsilon^2}{2n}} d\epsilon$$

$$Using \int_{-\infty}^{\infty} e^{-px^2} dx = \sqrt{\frac{\pi}{p}}$$

$$n! \approx \frac{n^n}{e^n} \sqrt{2n\pi}$$

### 11.1.1 Limit as n approaches infinity

The approximation in the this result can be stated more correctly as follows

$$\boxed{\lim_{n\to\infty} \frac{n!}{(\frac{n}{e})^n\sqrt{2\pi n}} = 1} \tag{42}$$

We can state this equivalently by taking logarithms of both sides. We will use this form to help derive the full Stirling approximation.

$$\boxed{\lim_{n\to\infty} ln(n!) - nln(n) + n - \frac{ln(n)}{2} = ln(\sqrt{2\pi})} \tag{43}$$

## 11.2 Stirling's Full Approximation

**Theorem 11.2.** *Expanding on Stirling's approximation to include as many terms as needed we need to use the Euler-Maclaurin formula*[2].

$$\boxed{n! = (\frac{n}{e})^n\sqrt{2\pi n}\cdot exp\Big(\sum_{k=1}^{\lfloor m/2\rfloor} \frac{B_{2k}}{2k\cdot(2k-1)}\frac{1}{n^{2k-1}} - \frac{1}{m}\int_{n-1}^{\infty}\frac{B_m(\{t\})}{t^m}dt\Big)} \tag{44}$$

*Proof*
*Starting from the definition of the simple version of the Euler-Maclaurin formula*

$$\sum_{j=0}^{r} g(j) - \int_{x=0}^{r} g(x)dx = \frac{g(0)+g(r)}{2} + \sum_{k=1}^{\lfloor m/2\rfloor}\frac{B_{2k}}{(2k)!}\Big(g^{(2k-1)}(r) - g^{(2k-1)}(0)\Big) + R_{r,m}$$

$$R_{r,m} = \frac{(-1)^{m+1}}{m!}\int_0^r B_m(\{t\})g^{(m)}(t)dt$$

*Define* $f(x) = g(x+1)$ *and* $n = r+1$, *so that* $f(n) = f(r+1) = g(r)$. *we get*

$$\sum_{j=1}^{n} f(j) - \int_{x=1}^{n} f(x)dx = \frac{f(1)+f(n)}{2} + \sum_{k=1}^{\lfloor m/2\rfloor}\frac{B_{2k}}{(2k)!}\Big(f^{(2k-1)}(n) - f^{(2k-1)}(1)\Big) + R_{n,m}$$

$$R_{n,m} = \frac{(-1)^{m+1}}{m!}\int_1^{n-1} B_m(\{t\})f^{(m)}(t)dt$$

[2]See the Darboux Formula document

*Let $f(x) = \ln(x)$, therefore*

$$f^{(m)}(x) = \frac{(-1)^{m-1} \cdot (m-1)!}{(x)^m} \text{ or } f^{(2k-1)}(x) = \frac{(-1)^{2k-2} \cdot (2k-2)!}{(x)^{2k-1}}$$

*Plugging in $f(x)$ and $f^{(m)}(x)$ into the Euler-Maclaurin formula we get*

$$\sum_{j=1}^{n} ln(j) = ln(n!) = \int_{x=1}^{n} ln(x)dx + \frac{ln(1) + ln(n)}{2}$$

$$+ \sum_{k=1}^{\lfloor m/2 \rfloor} \frac{B_{2k}}{(2k)!}\Big(\frac{(-1)^{2k-2} \cdot (2k-2)!}{n^{2k-1}} - \frac{(-1)^{2k-2} \cdot (2k-2)!}{1^{2k-1}}\Big) + R_{n,m}$$

$$R_{n,m} = \frac{(-1)^{m+1}}{m!} \int_{1}^{n-1} B_m(\{t\}) \frac{(-1)^{m-1} \cdot (m-1)!}{t^m} dt$$

*Evaluating the integral and simplifying*

$$ln(n!) = \big(n \cdot ln(n) - n\big) - \big(1 \cdot ln(1) - 1\big) + \frac{ln(n)}{2} + \sum_{k=1}^{\lfloor m/2 \rfloor} \frac{B_{2k} \cdot (2k-2)!}{(2k)!}\Big(\frac{1}{n^{2k-1}} - 1\Big) + R_{n,m}$$

$$R_{n,m} = \frac{1}{m} \int_{1}^{n-1} \frac{B_m(\{t\})}{t^m} dt$$

*Further simplifying*

$$ln(n!) = n \cdot ln(n) - n + 1 + \frac{ln(n)}{2} + \sum_{k=1}^{\lfloor m/2 \rfloor} \frac{B_{2k}}{2k \cdot (2k-1)}\Big(\frac{1}{n^{2k-1}} - 1\Big) + R_{n,m}$$

*To eliminate some terms we can rearranging and take limits of both sides. This would allow us to use the simple Stirling Approximation to skip the evaluation of one of the sums.*

$$\lim_{n\to\infty} ln(n!) - n \cdot ln(n) + n - \frac{ln(n)}{2}$$

$$= \lim_{n\to\infty} 1 + \sum_{k=1}^{\lfloor m/2 \rfloor} \frac{B_{2k}}{2k \cdot (2k-1)} \frac{1}{n^{2k-1}} - \sum_{k=1}^{\lfloor m/2 \rfloor} \frac{B_{2k}}{2k \cdot (2k-1)} + R_{n,m}$$

$$= 1 - \sum_{k=1}^{\lfloor m/2 \rfloor} \frac{B_{2k}}{2k \cdot (2k-1)} + \lim_{n\to\infty} R_{n,m}$$

*Using equation 43*

$$1 - \sum_{k=1}^{\lfloor m/2 \rfloor} \frac{B_{2k}}{2k \cdot (2k-1)} + \lim_{n\to\infty} R_{n,m} = ln(\sqrt{2\pi})$$

*Plugging this back in, we get:*

$$ln(n!) - n \cdot ln(n) + n - \frac{ln(n)}{2} = ln(\sqrt{2\pi}) + \sum_{k=1}^{\lfloor m/2 \rfloor} \frac{B_{2k}}{2k \cdot (2k-1)} \frac{1}{n^{2k-1}} + R_{n,m} - \lim_{n\to\infty} R_{n,m}$$

*Using the integral form of $R_{n,m}$ we can then re-write the difference $D = R_{n,m} - \lim_{n\to\infty} R_{n,m}$*

$$D = R_{n,m} - \lim_{n\to\infty} R_{n,m} = \frac{1}{m} \int_1^{n-1} \frac{B_m(\{t\})}{t^m} dt - \frac{1}{m} \int_1^{\infty} \frac{B_m(\{t\})}{t^m} dt = -\frac{1}{m} \int_{n-1}^{\infty} \frac{B_m(\{t\})}{t^m} dt$$

*Now raising e to the value of both sides, we get*

$$n! = (\frac{n}{e})^n \sqrt{2\pi n} \cdot exp\Big( \sum_{k=1}^{\lfloor m/2 \rfloor} \frac{B_{2k}}{2k \cdot (2k-1)} \frac{1}{n^{2k-1}} - \frac{1}{m} \int_{n-1}^{\infty} \frac{B_m(\{t\})}{t^m} dt \Big)$$

### 11.2.1 Approximation big-O notation

**Theorem 11.3.** *The integral in the exponent can be bound by a polynomial power with respect to n.*

$$\boxed{n! = (\frac{n}{e})^n \sqrt{2\pi n} \cdot exp\Big( \sum_{k=1}^{\lfloor m/2 \rfloor} \frac{B_{2k}}{2k \cdot (2k-1)} \frac{1}{n^{2k-1}} + O(\frac{1}{n^m}) \Big)} \tag{45}$$

*Proof*
*We need to show that* $\frac{1}{m} \int_{n-1}^{\infty} \frac{B_m(\{t\})}{t^m} dt = O\left(\frac{1}{n^m}\right)$

$$D = -\frac{1}{m} \int_{n-1}^{\infty} \frac{B_m(\{t\})}{t^m} dt$$

$$D \leq |D| \leq \frac{1}{|m|} \int_{n-1}^{\infty} \frac{|B_m(\{t\})|}{t^m} dt \leq \frac{\max_{x\in[0,1]} |B_m(x)|}{|m|} \int_{n-1}^{\infty} \frac{1}{t^m} dt$$

$$= \underbrace{\frac{\max_{x\in[0,1]} |B_m(x)|}{m \cdot (m+1)}}_{\neq f(n)} \underbrace{\frac{1}{(n-1)^m}}_{f(n)} = O(\frac{1}{n^m})$$

## 11.3 Plugging in values

Generating an approximation with $m = 7$ and using $B_2 = \frac{1}{6}, B_4 = -\frac{1}{30}, B_6 = \frac{1}{42}$ we obtain:

$$n! = (\frac{n}{e})^n \sqrt{2\pi n} \cdot exp\Big( \frac{1}{12n} - \frac{1}{360n^3} + \frac{1}{1260n^5} + O(\frac{1}{n^7}) \Big) \tag{46}$$

Since the factorial function grows very fast, it's useful to consider the logarithm of the factorial function. This would result in the following approximation.

$$ln(n!) = \frac{ln(2\pi)}{2} + nln(n) - n + + \frac{ln(n)}{2} + \frac{1}{12n} - \frac{1}{360n^3} + \frac{1}{1260n^5} + O(\frac{1}{n^7}) \tag{47}$$