
Supplementary material

Table of contents

Causal structure 1
Model formulation 2
1. Latent childhood trauma metrics 2
2. CTQ measurement (emission) layer 2
3. Latent dynamics 2
4. Longitudinal process state 2
5. Prior distributions 3
Stan code 3
Model diagnostics and posterior predictive checks 6
Density 7
Statistics 9
Latent trauma metric stratification 10
Sensitivity analysis 12
Prior predictive check (priors only) 12
Density 16
Statistics 18
Posterior predictive check (without CTQ emissions) 21
Density 25
Statistics 27
Posterior predictive check (without HCC emissions) 30
Density 34
Statistics 36


```
         used (Mb) gc trigger (Mb) max used (Mb)
Ncells  615695 32.9    1312034 70.1   898818 48.1
Vcells 1168803  9.0    8388608 64.0  1935467 14.8
```

Causal structure



Model formulation

1. Latent childhood trauma metrics

For each subject i, the joint distribution of their underlying exposure to abuse ($c_{i,1}$) and neglect ($c_{i,2}$) is drawn from a centered multivariate normal distribution capturing population variance and structural trait correlation:

$$\begin{bmatrix} c_{i,1} \\ c_{i,2} \end{bmatrix} \sim \text{Multivariate-Normal}\left(\begin{bmatrix} 0 \\ 0 \end{bmatrix}, \boldsymbol{\Sigma}_c \right)$$

$$\boldsymbol{\Sigma}_c = \text{diag}\left(\boldsymbol{\sigma}_c\right) \boldsymbol{\Omega}_c \, \text{diag}\left(\boldsymbol{\sigma}_c\right)$$

Where $\boldsymbol{\sigma}_c = [\sigma_{\text{abuse}}, \sigma_{\text{neglect}}]^T$ maps the population scales, and $\boldsymbol{\Omega}_c$ represents the trait correlation matrix.

2. CTQ measurement (emission) layer

The individual's true latent trauma states directly manifest across the five observed subscales of the Childhood Trauma Questionnaire. The Abuse factor ($c_{i,1}$) drives emotional, physical, and sexual dimensions, while the Neglect factor ($c_{i,2}$) drives emotional and physical neglect indicators:

$$y_m^{(\text{EA})} \sim \text{Normal}\left(\alpha_{\text{EA}} + c_{\text{subj }[m],1}, \sigma_{\text{EA}}\right) \quad \text{[Emotional Abuse]}$$

$$y_m^{(\text{PA})} \sim \text{Normal}\left(\alpha_{\text{PA}} + c_{\text{subj }[m],1}, \sigma_{\text{PA}}\right) \quad \text{[Physical Abuse]}$$

$$y_m^{(\text{SA})} \sim \text{Normal}\left(\alpha_{\text{SA}} + c_{\text{subj }[m],1}, \sigma_{\text{SA}}\right) \quad \text{[Sexual Abuse]}$$

$$y_m^{(\text{EN})} \sim \text{Normal}\left(\alpha_{\text{EN}} + c_{\text{subj }[m],2}, \sigma_{\text{EN}}\right) \quad \text{[Emotional Neglect]}$$

$$y_m^{(\text{PN})} \sim \text{Normal}\left(\alpha_{\text{PN}} + c_{\text{subj }[m],2}, \sigma_{\text{PN}}\right) \quad \text{[Physical Neglect]}$$

3. Latent dynamics

The latent dysregulation state $z_i(t)$ transitions over irregular time intervals ($\Delta t = t - t'$) according to a continuous-time (Ornstein–Uhlenbeck) engine. The individual recovery speed λ_i forces a strict positive bound via the structural baseline floor λ_0:

$$z_i(t) \mid z_i(t') \sim \text{Normal}\left(e^{-\lambda_i \Delta t} \, z_i(t'), \sigma_z \sqrt{1 - e^{-2\lambda_i \Delta t}}\right)$$

$$\lambda_i = \lambda_0 + \exp\left(\gamma_{\text{abuse}} \, c_{i,1} + \gamma_{\text{neglect}} \, c_{i,2}\right)$$

4. Longitudinal process state

The state trajectory begins at a unique initial tracking baseline anchored to a matched-pair intercept (u_{pair}) and shifted by diagnostic status (SSD):

$$z_i(t_0) \sim \text{Normal}\left(u_{\text{pair }[i]} + \beta_{\text{SSD}} \, \text{SSD}_i, 1\right)$$

$$u_j \sim \text{Normal}\left(0, \sigma_{\text{pair}}\right)$$

The evolving state $z_i(t)$ projects concurrent signals down into both historical biological archives and acute clinical cross-sections:

$$y_m^{(\text{HCC})} \sim \text{Student-t}\left(\nu = 6, a_{\text{HCC}} + b_{\text{HCC}} \, z_{\text{state }[m]}, \sigma_{\text{HCC}}\right)$$

$$y_m^{(\text{PANSS})} \sim \text{Normal}\left(a_{\text{PANSS}} + b_{\text{PANSS}} \, z_{\text{state }[m]}, \sigma_{\text{PANSS}}\right)$$

$$y_m^{(\text{PSS})} \sim \text{Normal}\left(a_{\text{PSS}} + b_{\text{PSS}} \, z_{\text{state }[m]}, \sigma_{\text{PSS}}\right)$$

5. Prior distributions

$$
\begin{aligned}
\lambda_0, \gamma_{\text{abuse}}, \gamma_{\text{neglect}}, \beta_{\text{SSD}} &\sim \text{Normal}\ (0, 1) \\
\alpha_{\text{EA}}, \alpha_{\text{PA}}, \alpha_{\text{SA}}, \alpha_{\text{EN}}, \alpha_{\text{PN}} &\sim \text{Normal}\ (0, 1) \\
a_{\text{PANSS}}, a_{\text{PSS}}, a_{\text{HCC}} &\sim \text{Normal}\ (0, 0.5) \\
b_{\text{PANSS}}, b_{\text{PSS}}, b_{\text{HCC}} &\sim \text{Normal}\ (0, 1) \\
\sigma_z, \sigma_{\text{pair}}, \sigma_{\text{HCC}}, \sigma_{\text{PANSS}}, \sigma_{\text{PSS}} &\sim \text{Exponential}\ (1) \\
\sigma_{\text{EA}}, \sigma_{\text{PA}}, \sigma_{\text{SA}}, \sigma_{\text{EN}}, \sigma_{\text{PN}} &\sim \text{Exponential}\ (1) \\
\boldsymbol{\sigma}_c &\sim \text{Exponential}\ (1) \\
\boldsymbol{\Omega}_c &\sim \text{LKJ}\ (\eta = 2)
\end{aligned}
$$

Stan code

The mathematical model is rewritten in the Stan language.

```
data{

int <lower=1> N_state;
array[N_state] int <lower=1> subj_id;
int <lower=1> N_pair;
array[N_state] int <lower=1> pair_id;
vector[N_state] time;
array[N_state] int <lower=0, upper=1> is_first;
array[N_state] int <lower=0, upper=1> scz;

int <lower=1> N_hcc;
array[N_hcc] int <lower=1> state_hcc;
vector[N_hcc] y_hcc;

int <lower=1> N_panss;
array[N_panss] int <lower=1> state_panss;
vector[N_panss] y_panss;

int <lower=1> N_pss;
array[N_pss] int <lower = 1> state_pss;
vector[N_pss] y_pss;

int <lower=1> N_ctq;
int <lower=1> N_subj;
array[N_ctq] int <lower=1> subj_ctq;

vector[N_ctq] y_ea;
vector[N_ctq] y_pa;
vector[N_ctq] y_sa;
vector[N_ctq] y_en;
vector[N_ctq] y_pn;


}

parameters{

vector[N_state] z_raw;
real <lower=0> lambda_0;
real gamma_abuse;
real gamma_neglect;
```

```
real <lower=0> sigma_z;
real <lower=0> sigma_pair;

vector[N_pair] u_pair;

real beta_scz;

real a_hcc;
real <lower=0> b_hcc;
real <lower=0> sigma_hcc;

real a_panss;
real b_panss;
real <lower=0> sigma_panss;

real a_pss;
real b_pss;
real <lower=0> sigma_pss;

real <lower=0> sigma_ea;
real <lower=0> sigma_pa;
real <lower=0> sigma_sa;
real <lower=0> sigma_en;
real <lower=0> sigma_pn;


real alpha_ea;
real alpha_pa;
real alpha_sa;
real alpha_en;
real alpha_pn;

matrix[2, N_subj] c_raw;
vector <lower=0> [2] sigma_c;
cholesky_factor_corr[2] L_c;

}

transformed parameters {

// Covariance matrix for CTQ latent factor c1 (abuse) and c2 (neglect)
matrix[2, N_subj] c;
c = diag_pre_multiply(sigma_c, L_c) * c_raw;

// latent subject-level factors c[1, ] and c[2, ]. Implicit Bayesian imputation.
vector[N_subj] lambda_subj;

for (i in 1:N_subj) {
  lambda_subj[i] = lambda_0 + exp(gamma_abuse * c[1, i] + gamma_neglect * c[2, i]);
}

// dynamic latent process
vector[N_state] z;
for (k in 1:N_state) {
    // first point in time aligned between patients and controls
    if (is_first[k] == 1) {
```

```
        z[k] = u_pair[pair_id[k]] + beta_scz * scz[k] + 1 * z_raw[k];
    // Ornstein-Uhlenbeck process (exponential return with stochastic volatility)
    } else {
        real dt = time[k] - time[k-1];
        real phi = exp(-lambda_subj[subj_id[k]] * dt);
        real sigma_dt = sigma_z * sqrt(1 - square(phi));
        z[k] = phi * z[k-1] + sigma_dt * z_raw[k];
    }
  }


}

model{

// priors
lambda_0 ~ normal(0, 1);
gamma_abuse ~ normal(0, 1);
gamma_neglect ~ normal(0, 1);
sigma_z ~ exponential(1);
sigma_pair ~ exponential(1);
sigma_hcc ~ exponential(1);
sigma_panss ~ exponential(1);
sigma_pss ~ exponential(1);
beta_scz ~ normal(0, 1);
a_hcc ~ normal(0, 0.5);
b_hcc ~ normal(0, 1);
a_panss ~ normal(0, 0.5);
b_panss ~ normal(0, 1);
a_pss ~ normal(0, 0.5);
b_pss ~ normal(0, 1);
z_raw ~ std_normal();

u_pair ~ normal(0, sigma_pair);

sigma_ea ~ exponential(1);
sigma_pa ~ exponential(1);
sigma_sa ~ exponential(1);
sigma_en ~ exponential(1);
sigma_pn ~ exponential(1);

alpha_ea ~ normal(0, 1);
alpha_pa ~ normal(0, 1);
alpha_sa ~ normal(0, 1);
alpha_en ~ normal(0, 1);
alpha_pn ~ normal(0, 1);

to_vector(c_raw) ~ std_normal();

L_c ~ lkj_corr_cholesky(2);
sigma_c ~ exponential(1);

// hcc emissions
for (m in 1:N_hcc) {
    y_hcc[m] ~ student_t(6, a_hcc + b_hcc * z[state_hcc[m]], sigma_hcc);
    }
```

```
// panss emissions
for (m in 1:N_panss) {
    y_panss[m] ~ normal(a_panss + b_panss * z[state_panss[m]], sigma_panss);
    }

// pss emissions
for (m in 1:N_pss) {
    y_pss[m] ~ normal(a_pss + b_pss * z[state_pss[m]], sigma_pss);
    }

// CTQ emissions
y_ea ~ normal(alpha_ea + c[1, subj_ctq]', sigma_ea);
y_sa ~ normal(alpha_sa + c[1, subj_ctq]', sigma_sa);
y_pa ~ normal(alpha_pa + c[1, subj_ctq]', sigma_pa);
y_en ~ normal(alpha_en + c[2, subj_ctq]', sigma_en);
y_pn ~ normal(alpha_pn + c[2, subj_ctq]', sigma_pn);


}


generated quantities{


matrix[2, 2] corr = multiply_lower_tri_self_transpose(L_c); // Correlation matrix
matrix[2, 2] cov = quad_form_diag(corr, sigma_c); // (S * R * S) Covariance matrix


}
```

Model diagnostics and posterior predictive checks


The model was rigorously evaluated for convergence using four Markov chains with 4000 iterations each. The Rhat diagnostic was used to assess convergence, with a value below 1.01 suggesting adequate convergence. The effective sample size (ESS) was over 400 for each parameter. The posterior predictive checks follow.




S_Figure 1: Posterior predictive check of HCC. The red line shows the observed values.



S_Figure 2: Posterior predictive check of PANSS. The red line shows the observed values.




S_Figure 3: Posterior predictive check of PSS. The red line shows the observed values.




S_Figure 4: Posterior predictive check of the latent CTQ. The red line shows the observed values.


Statistics



S_Figure 5: Posterior predictive check of selected statistics of HCC. The red line shows the observed values.




S_Figure 6: Posterior predictive check of selected statistics of PANSS. The red line shows the observed values.



S_Figure 7: Posterior predictive check of selected statistics of PSS. The red line shows the observed values.

Latent trauma metric stratification

CTQ subscales exhibit high mutual inter-correlation due to shared response variance. Modeling observed subscale scores as independent predictors introduces multicollinearity and leads to overestimating trauma burden when multiple subscales are elevated simultaneously (analogous to multiple motion sensors triggering from a single event). To resolve this, we employed a Bayesian implicit latent variable framework, modeling five observed CTQ subscale indicators ($\mathbf{y}_{\mathrm{EA}}, \mathbf{y}_{\mathrm{PA}}, \mathbf{y}_{\mathrm{SA}}, \mathbf{y}_{\mathrm{EN}}, \mathbf{y}_{\mathrm{PN}}$) as expressions of two correlated latent factors: Abuse (c_1) and Neglect (c_2).



S_Figure 8: Latent abuse–neglect correlation. The red line shows the posterior median and the dashed lines display the 95% credibility interval.


Mean latent factor score (c_1, c_2) represents standardized, continuous latent trait category, by which the subjects were assigned into trauma severity tiers.

Low trauma: $\frac{1}{2}(c_1 + c_2) \leq -0.75$

Medium trauma: $-0.75 < \frac{1}{2}(c_1 + c_2) < 0.75$

High trauma: $\frac{1}{2}(c_1 + c_2) \geq 0.75$


Because $c_{1,i}$ and $c_{2,i}$ are full posterior draws (not point estimates), each subject gets a posterior distribution over tiers rather than a single label.



S_Figure 9: Posterior trauma-tier posterior classification per subject - stacked posterior probability of low / medium / high, subjects ordered by mean latent severity.

Posterior predictive check of the latent metrics to the CTQ native scale could be displayed by converting the native CTQ using the following procedure:

1. Taking a mean of the 3 abuse native subscales (EA, PA, SA).
2. Taking a mean of the 2 neglect native subscales (EN, PN).
3. Standardize each construct separately across subjects.
4. Averaging the two z-scores.




S_Figure 10: Native CTQ vs. posterior-mean latent metrics.

```
            used    (Mb) gc trigger    (Mb)   max used    (Mb)
Ncells    4550827   243.1    7446355   397.7    7446355   397.7
Vcells 2684373631 20480.2 4599435086 35091.0 3685902231 28121.3
```

Sensitivity analysis

Prior predictive check (priors only)




```
data{

int <lower=1> N_state;
array[N_state] int <lower=1> subj_id;
```

```
int <lower=1> N_pair;
array[N_state] int <lower=1> pair_id;
vector[N_state] time;
array[N_state] int <lower=0, upper=1> is_first;
array[N_state] int <lower=0, upper=1> scz;

int <lower=1> N_hcc;
array[N_hcc] int <lower=1> state_hcc;
vector[N_hcc] y_hcc;

int <lower=1> N_panss;
array[N_panss] int <lower=1> state_panss;
vector[N_panss] y_panss;

int <lower=1> N_pss;
array[N_pss] int <lower = 1> state_pss;
vector[N_pss] y_pss;

int <lower=1> N_ctq;
int <lower=1> N_subj;
array[N_ctq] int <lower=1> subj_ctq;

vector[N_ctq] y_ea;
vector[N_ctq] y_pa;
vector[N_ctq] y_sa;
vector[N_ctq] y_en;
vector[N_ctq] y_pn;


}

parameters{

vector[N_state] z_raw;
real <lower=0> lambda_0;
real gamma_abuse;
real gamma_neglect;
real <lower=0> sigma_z;
real <lower=0> sigma_pair;

vector[N_pair] u_pair;

real beta_scz;

real a_hcc;
real <lower=0> b_hcc;
real <lower=0> sigma_hcc;

real a_panss;
real b_panss;
real <lower=0> sigma_panss;

real a_pss;
real b_pss;
real <lower=0> sigma_pss;
```

```
real <lower=0> sigma_ea;
real <lower=0> sigma_pa;
real <lower=0> sigma_sa;
real <lower=0> sigma_en;
real <lower=0> sigma_pn;


real alpha_ea;
real alpha_pa;
real alpha_sa;
real alpha_en;
real alpha_pn;

matrix[2, N_subj] c_raw;
vector <lower=0> [2] sigma_c;
cholesky_factor_corr[2] L_c;

}

transformed parameters {

// Covariance matrix for CTQ latent factor c1 (abuse) and c2 (neglect)
matrix[2, N_subj] c;
c = diag_pre_multiply(sigma_c, L_c) * c_raw;

// latent subject-level factors c[1, ] and c[2, ]. Implicit Bayesian imputation.
vector[N_subj] lambda_subj;

for (i in 1:N_subj) {
  lambda_subj[i] = lambda_0 + exp(gamma_abuse * c[1, i] + gamma_neglect * c[2, i]);
}

// dynamic latent process
vector[N_state] z;
for (k in 1:N_state) {
    // first point in time aligned between patients and controls
    if (is_first[k] == 1) {
        z[k] = u_pair[pair_id[k]] + beta_scz * scz[k] + 1 * z_raw[k];
    // Ornstein-Uhlenbeck process (exponential return with stochastic volatility)
    } else {
        real dt = time[k] - time[k-1];
        real phi = exp(-lambda_subj[subj_id[k]] * dt);
        real sigma_dt = sigma_z * sqrt(1 - square(phi));
        z[k] = phi * z[k-1] + sigma_dt * z_raw[k];
    }
  }


}

model{

// priors
lambda_0 ~ normal(0, 1);
gamma_abuse ~ normal(0, 1);
gamma_neglect ~ normal(0, 1);
```

```
sigma_z ~ exponential(1);
sigma_pair ~ exponential(1);
sigma_hcc ~ exponential(1);
sigma_panss ~ exponential(1);
sigma_pss ~ exponential(1);
beta_scz ~ normal(0, 1);
a_hcc ~ normal(0, 0.5);
b_hcc ~ normal(0, 1);
a_panss ~ normal(0, 0.5);
b_panss ~ normal(0, 1);
a_pss ~ normal(0, 0.5);
b_pss ~ normal(0, 1);
z_raw ~ std_normal();

u_pair ~ normal(0, sigma_pair);

sigma_ea ~ exponential(1);
sigma_pa ~ exponential(1);
sigma_sa ~ exponential(1);
sigma_en ~ exponential(1);
sigma_pn ~ exponential(1);

alpha_ea ~ normal(0, 1);
alpha_pa ~ normal(0, 1);
alpha_sa ~ normal(0, 1);
alpha_en ~ normal(0, 1);
alpha_pn ~ normal(0, 1);

to_vector(c_raw) ~ std_normal();

L_c ~ lkj_corr_cholesky(2);
sigma_c ~ exponential(1);

// hcc emissions
//for (m in 1:N_hcc) {
//    y_hcc[m] ~ student_t(6, a_hcc + b_hcc * z[state_hcc[m]], sigma_hcc);
//    }

// panss emissions
//for (m in 1:N_panss) {
//    y_panss[m] ~ normal(a_panss + b_panss * z[state_panss[m]], sigma_panss);
//    }

// pss emissions
//for (m in 1:N_pss) {
//    y_pss[m] ~ normal(a_pss + b_pss * z[state_pss[m]], sigma_pss);
//    }

// CTQ emissions
//y_ea ~ normal(alpha_ea + c[1, subj_ctq]', sigma_ea);
//y_sa ~ normal(alpha_sa + c[1, subj_ctq]', sigma_sa);
//y_pa ~ normal(alpha_pa + c[1, subj_ctq]', sigma_pa);
//y_en ~ normal(alpha_en + c[2, subj_ctq]', sigma_en);
//y_pn ~ normal(alpha_pn + c[2, subj_ctq]', sigma_pn);
```


```
}

generated quantities{



matrix[2, 2] corr = multiply_lower_tri_self_transpose(L_c); // Correlation matrix
matrix[2, 2] cov = quad_form_diag(corr, sigma_c); // (S * R * S) Covariance matrix

}
```


Density




S_Figure 11: Prior predictive check of HCC. The red line shows the observed values.



S_Figure 12: Prior predictive check of PANSS. The red line shows the observed values.




S_Figure 13: Prior predictive check of PSS. The red line shows the observed values.




S_Figure 14: Prior predictive check of the latent CTQ. The red line shows the observed values.


Statistics




S_Figure 15: Prior predictive check of selected statistics of HCC. The red line shows the observed values.



S_Figure 16: Prior predictive check of selected statistics of PANSS. The red line shows the observed values.



S_Figure 17: Prior predictive check of selected statistics of PSS. The red line shows the observed values.



S_Figure 18: Latent abuse–neglect correlation. The red line shows the prior median and the dashed lines display the 95% credibility interval.


Mean latent factor score (c_1, c_2) represents standardized, continuous latent trait category, by which the subjects were assigned into trauma severity tiers.

Low trauma: $\frac{1}{2}(c_1 + c_2) \leq -0.75$


Medium trauma: $-0.75 < \frac{1}{2}(c_1 + c_2) < 0.75$

High trauma: $\frac{1}{2}(c_1 + c_2) \geq 0.75$

Because $c_{1,i}$ and $c_{2,i}$ are full posterior draws (not point estimates), each subject gets a posterior distribution over tiers rather than a single label.




S_Figure 19: Prior trauma-tier posterior classification per subject - stacked prior probability of low / medium / high, subjects ordered by mean latent severity.



S_Figure 20: Dynamic latent shared dysregulation of SSD patients marginalized over subjects stratified by latent childhood trauma severity. Shaded areas represent 50%, 80%, and 95% credible intervals. Numbers indicate the probability that the latent state (z) is positive at the corresponding time points associated with each clinical visit, Pr(z(visit) > 0).

```
            used    (Mb) gc trigger    (Mb)   max used    (Mb)
Ncells    4610075   246.3    9589140   512.2    9589140   512.2
Vcells 2684485055 20481.0 5519402103 42109.7 5519250305 42108.6
```

Posterior predictive check (without CTQ emissions)




```
data{

int <lower=1> N_state;
array[N_state] int <lower=1> subj_id;
int <lower=1> N_pair;
array[N_state] int <lower=1> pair_id;
vector[N_state] time;
array[N_state] int <lower=0, upper=1> is_first;
array[N_state] int <lower=0, upper=1> scz;

int <lower=1> N_hcc;
array[N_hcc] int <lower=1> state_hcc;
vector[N_hcc] y_hcc;

int <lower=1> N_panss;
array[N_panss] int <lower=1> state_panss;
vector[N_panss] y_panss;

int <lower=1> N_pss;
array[N_pss] int <lower = 1> state_pss;
vector[N_pss] y_pss;

int <lower=1> N_ctq;
int <lower=1> N_subj;
array[N_ctq] int <lower=1> subj_ctq;

vector[N_ctq] y_ea;
vector[N_ctq] y_pa;
vector[N_ctq] y_sa;
vector[N_ctq] y_en;
vector[N_ctq] y_pn;
```

```
}

parameters{

vector[N_state] z_raw;
real <lower=0> lambda_0;
real gamma_abuse;
real gamma_neglect;
real <lower=0> sigma_z;
real <lower=0> sigma_pair;

vector[N_pair] u_pair;

real beta_scz;

real a_hcc;
real <lower=0> b_hcc;
real <lower=0> sigma_hcc;

real a_panss;
real b_panss;
real <lower=0> sigma_panss;

real a_pss;
real b_pss;
real <lower=0> sigma_pss;

real <lower=0> sigma_ea;
real <lower=0> sigma_pa;
real <lower=0> sigma_sa;
real <lower=0> sigma_en;
real <lower=0> sigma_pn;


real alpha_ea;
real alpha_pa;
real alpha_sa;
real alpha_en;
real alpha_pn;

matrix[2, N_subj] c_raw;
vector <lower=0> [2] sigma_c;
cholesky_factor_corr[2] L_c;

}

transformed parameters {

// Covariance matrix for CTQ latent factor c1 (abuse) and c2 (neglect)
matrix[2, N_subj] c;
c = diag_pre_multiply(sigma_c, L_c) * c_raw;

// latent subject-level factors c[1, ] and c[2, ]. Implicit Bayesian imputation.
vector[N_subj] lambda_subj;

for (i in 1:N_subj) {
```

```
  lambda_subj[i] = lambda_0 + exp(gamma_abuse * c[1, i] + gamma_neglect * c[2, i]);
}

// dynamic latent process
vector[N_state] z;
for (k in 1:N_state) {
    // first point in time aligned between patients and controls
    if (is_first[k] == 1) {
        z[k] = u_pair[pair_id[k]] + beta_scz * scz[k] + 1 * z_raw[k];
    // Ornstein-Uhlenbeck process (exponential return with stochastic volatility)
    } else {
        real dt = time[k] - time[k-1];
        real phi = exp(-lambda_subj[subj_id[k]] * dt);
        real sigma_dt = sigma_z * sqrt(1 - square(phi));
        z[k] = phi * z[k-1] + sigma_dt * z_raw[k];
    }
  }


}

model{

// priors
lambda_0 ~ normal(0, 1);
gamma_abuse ~ normal(0, 1);
gamma_neglect ~ normal(0, 1);
sigma_z ~ exponential(1);
sigma_pair ~ exponential(1);
sigma_hcc ~ exponential(1);
sigma_panss ~ exponential(1);
sigma_pss ~ exponential(1);
beta_scz ~ normal(0, 1);
a_hcc ~ normal(0, 0.5);
b_hcc ~ normal(0, 1);
a_panss ~ normal(0, 0.5);
b_panss ~ normal(0, 1);
a_pss ~ normal(0, 0.5);
b_pss ~ normal(0, 1);
z_raw ~ std_normal();

u_pair ~ normal(0, sigma_pair);

sigma_ea ~ exponential(1);
sigma_pa ~ exponential(1);
sigma_sa ~ exponential(1);
sigma_en ~ exponential(1);
sigma_pn ~ exponential(1);

alpha_ea ~ normal(0, 1);
alpha_pa ~ normal(0, 1);
alpha_sa ~ normal(0, 1);
alpha_en ~ normal(0, 1);
alpha_pn ~ normal(0, 1);

to_vector(c_raw) ~ std_normal();
```

```
L_c ~ lkj_corr_cholesky(2);
sigma_c ~ exponential(1);

// hcc emissions
for (m in 1:N_hcc) {
    y_hcc[m] ~ student_t(6, a_hcc + b_hcc * z[state_hcc[m]], sigma_hcc);
    }

// panss emissions
for (m in 1:N_panss) {
    y_panss[m] ~ normal(a_panss + b_panss * z[state_panss[m]], sigma_panss);
    }

// pss emissions
for (m in 1:N_pss) {
    y_pss[m] ~ normal(a_pss + b_pss * z[state_pss[m]], sigma_pss);
    }


// CTQ emissions
//y_ea ~ normal(alpha_ea + c[1, subj_ctq]', sigma_ea);
//y_sa ~ normal(alpha_sa + c[1, subj_ctq]', sigma_sa);
//y_pa ~ normal(alpha_pa + c[1, subj_ctq]', sigma_pa);
//y_en ~ normal(alpha_en + c[2, subj_ctq]', sigma_en);
//y_pn ~ normal(alpha_pn + c[2, subj_ctq]', sigma_pn);


}


generated quantities{


matrix[2, 2] corr = multiply_lower_tri_self_transpose(L_c); // Correlation matrix
matrix[2, 2] cov = quad_form_diag(corr, sigma_c); // (S * R * S) Covariance matrix


}
```




S_Figure 21: Posterior predictive check of HCC. The red line shows the observed values.



S_Figure 22: Posterior predictive check of PANSS. The red line shows the observed values.




S_Figure 23: Posterior predictive check of PSS. The red line shows the observed values.




S_Figure 24: Posterior predictive check of the latent CTQ. The red line shows the observed values.


Statistics




S_Figure 25: Posterior predictive check of selected statistics of HCC. The red line shows the observed values.



S_Figure 26: Posterior predictive check of selected statistics of PANSS. The red line shows the observed values.



S_Figure 27: Posterior predictive check of selected statistics of PSS. The red line shows the observed values.



S_Figure 28: Latent abuse–neglect correlation. The red line shows the posterior median and the dashed lines display the 95% credibility interval.


Mean latent factor score (c_1, c_2) represents standardized, continuous latent trait category, by which the subjects were assigned into trauma severity tiers.

Low trauma: $\frac{1}{2}(c_1 + c_2) \leq -0.75$


Medium trauma: $-0.75 < \frac{1}{2}(c_1 + c_2) < 0.75$

High trauma: $\frac{1}{2}(c_1 + c_2) \geq 0.75$

Because $c_{1,i}$ and $c_{2,i}$ are full posterior draws (not point estimates), each subject gets a posterior distribution over tiers rather than a single label.




S_Figure 29: Posterior trauma-tier posterior classification per subject - stacked posterior probability of low / medium / high, subjects ordered by mean latent severity.



S_Figure 30: Dynamic latent shared dysregulation of SSD patients marginalized over subjects stratified by latent childhood trauma severity. Shaded areas represent 50%, 80%, and 95% credible intervals. Numbers indicate the probability that the latent state (z) is positive at the corresponding time points associated with each clinical visit, Pr(z(visit) > 0).

```
            used    (Mb) gc trigger    (Mb)   max used    (Mb)
Ncells    4615379   246.5    9664202   516.2    9664202   516.2
Vcells 2684499511 20481.2 5519402103 42109.7 5519306563 42109.0
```

Posterior predictive check (without HCC emissions)




```
data{

int <lower=1> N_state;
array[N_state] int <lower=1> subj_id;
int <lower=1> N_pair;
array[N_state] int <lower=1> pair_id;
vector[N_state] time;
array[N_state] int <lower=0, upper=1> is_first;
array[N_state] int <lower=0, upper=1> scz;

int <lower=1> N_hcc;
array[N_hcc] int <lower=1> state_hcc;
vector[N_hcc] y_hcc;

int <lower=1> N_panss;
array[N_panss] int <lower=1> state_panss;
vector[N_panss] y_panss;

int <lower=1> N_pss;
array[N_pss] int <lower = 1> state_pss;
vector[N_pss] y_pss;

int <lower=1> N_ctq;
int <lower=1> N_subj;
array[N_ctq] int <lower=1> subj_ctq;

vector[N_ctq] y_ea;
vector[N_ctq] y_pa;
vector[N_ctq] y_sa;
vector[N_ctq] y_en;
vector[N_ctq] y_pn;
```

```
}

parameters{

vector[N_state] z_raw;
real <lower=0> lambda_0;
real gamma_abuse;
real gamma_neglect;
real <lower=0> sigma_z;
real <lower=0> sigma_pair;

vector[N_pair] u_pair;

real beta_scz;

real a_hcc;
real <lower=0> b_hcc;
real <lower=0> sigma_hcc;

real a_panss;
real b_panss;
real <lower=0> sigma_panss;

real a_pss;
real b_pss;
real <lower=0> sigma_pss;

real <lower=0> sigma_ea;
real <lower=0> sigma_pa;
real <lower=0> sigma_sa;
real <lower=0> sigma_en;
real <lower=0> sigma_pn;


real alpha_ea;
real alpha_pa;
real alpha_sa;
real alpha_en;
real alpha_pn;

matrix[2, N_subj] c_raw;
vector <lower=0> [2] sigma_c;
cholesky_factor_corr[2] L_c;

}

transformed parameters {

// Covariance matrix for CTQ latent factor c1 (abuse) and c2 (neglect)
matrix[2, N_subj] c;
c = diag_pre_multiply(sigma_c, L_c) * c_raw;

// latent subject-level factors c[1, ] and c[2, ]. Implicit Bayesian imputation.
vector[N_subj] lambda_subj;

for (i in 1:N_subj) {
```

```
  lambda_subj[i] = lambda_0 + exp(gamma_abuse * c[1, i] + gamma_neglect * c[2, i]);
}

// dynamic latent process
vector[N_state] z;
for (k in 1:N_state) {
    // first point in time aligned between patients and controls
    if (is_first[k] == 1) {
        z[k] = u_pair[pair_id[k]] + beta_scz * scz[k] + 1 * z_raw[k];
    // Ornstein-Uhlenbeck process (exponential return with stochastic volatility)
    } else {
        real dt = time[k] - time[k-1];
        real phi = exp(-lambda_subj[subj_id[k]] * dt);
        real sigma_dt = sigma_z * sqrt(1 - square(phi));
        z[k] = phi * z[k-1] + sigma_dt * z_raw[k];
    }
  }


}

model{

// priors
lambda_0 ~ normal(0, 1);
gamma_abuse ~ normal(0, 1);
gamma_neglect ~ normal(0, 1);
sigma_z ~ exponential(1);
sigma_pair ~ exponential(1);
sigma_hcc ~ exponential(1);
sigma_panss ~ exponential(1);
sigma_pss ~ exponential(1);
beta_scz ~ normal(0, 1);
a_hcc ~ normal(0, 0.5);
b_hcc ~ normal(0, 1);
a_panss ~ normal(0, 0.5);
b_panss ~ normal(0, 1);
a_pss ~ normal(0, 0.5);
b_pss ~ normal(0, 1);
z_raw ~ std_normal();

u_pair ~ normal(0, sigma_pair);

sigma_ea ~ exponential(1);
sigma_pa ~ exponential(1);
sigma_sa ~ exponential(1);
sigma_en ~ exponential(1);
sigma_pn ~ exponential(1);

alpha_ea ~ normal(0, 1);
alpha_pa ~ normal(0, 1);
alpha_sa ~ normal(0, 1);
alpha_en ~ normal(0, 1);
alpha_pn ~ normal(0, 1);

to_vector(c_raw) ~ std_normal();
```

```
L_c ~ lkj_corr_cholesky(2);
sigma_c ~ exponential(1);

// hcc emissions
//for (m in 1:N_hcc) {
//    y_hcc[m] ~ student_t(6, a_hcc + b_hcc * z[state_hcc[m]], sigma_hcc);
//    }

// panss emissions
for (m in 1:N_panss) {
    y_panss[m] ~ normal(a_panss + b_panss * z[state_panss[m]], sigma_panss);
    }

// pss emissions
for (m in 1:N_pss) {
    y_pss[m] ~ normal(a_pss + b_pss * z[state_pss[m]], sigma_pss);
    }

// CTQ emissions
y_ea ~ normal(alpha_ea + c[1, subj_ctq]', sigma_ea);
y_sa ~ normal(alpha_sa + c[1, subj_ctq]', sigma_sa);
y_pa ~ normal(alpha_pa + c[1, subj_ctq]', sigma_pa);
y_en ~ normal(alpha_en + c[2, subj_ctq]', sigma_en);
y_pn ~ normal(alpha_pn + c[2, subj_ctq]', sigma_pn);



}

generated quantities{



matrix[2, 2] corr = multiply_lower_tri_self_transpose(L_c); // Correlation matrix
matrix[2, 2] cov = quad_form_diag(corr, sigma_c); // (S * R * S) Covariance matrix

}
```


Density




S_Figure 31: Posterior predictive check of HCC. The red line shows the observed values.



S_Figure 32: Posterior predictive check of PANSS. The red line shows the observed values.




S_Figure 33: Posterior predictive check of PSS. The red line shows the observed values.




S_Figure 34: Posterior predictive check of the latent CTQ. The red line shows the observed values.


Statistics




S_Figure 35: Posterior predictive check of selected statistics of HCC. The red line shows the observed values.



S_Figure 36: Posterior predictive check of selected statistics of PANSS. The red line shows the observed values.



S_Figure 37: Posterior predictive check of selected statistics of PSS. The red line shows the observed values.



S_Figure 38: Latent abuse–neglect correlation. The red line shows the posterior median and the dashed lines display the 95% credibility interval.


Mean latent factor score (c_1, c_2) represents standardized, continuous latent trait category, by which the subjects were assigned into trauma severity tiers.

Low trauma: $\frac{1}{2}(c_1 + c_2) \leq -0.75$


Medium trauma: $-0.75 < \frac{1}{2}(c_1 + c_2) < 0.75$

High trauma: $\frac{1}{2}(c_1 + c_2) \geq 0.75$

Because $c_{1,i}$ and $c_{2,i}$ are full posterior draws (not point estimates), each subject gets a posterior distribution over tiers rather than a single label.



S_Figure 39: Posterior trauma-tier posterior classification per subject - stacked posterior probability of low / medium / high, subjects ordered by mean latent severity.



S_Figure 40: Dynamic latent shared dysregulation of SSD patients marginalized over subjects stratified by latent childhood trauma severity. Shaded areas represent 50%, 80%, and 95% credible intervals. Numbers indicate the probability that the latent state (z) is positive at the corresponding time points associated with each clinical visit, Pr(z(visit) > 0).

```
            used    (Mb) gc trigger    (Mb)   max used    (Mb)
Ncells    4616349   246.6    9664202   516.2    9664202   516.2
Vcells 2684506828 20481.2 5519402103 42109.7 5519306563 42109.0
```