

PATIENT REFERRAL DOCUMENT

LMU Klinikum Munich

Patient: Friedrich Wagner

Date of Birth: 22.07.1952

Date: 15.05.2026

Referring Physician: Dr. Sabine Richter

General Practice Schwabing, Munich

REASON FOR REFERRAL:

Patient presents with sudden onset of left-sided facial drooping and slurred speech beginning approximately 90 minutes ago. Wife reports patient was speaking normally at breakfast then suddenly was unable to form words and his left side of face appeared to droop. Patient is conscious but confused and unable to communicate clearly.

CLINICAL FINDINGS:

Blood pressure: 195/115 mmHg

Heart rate: 96 bpm

Temperature: 37.1 degrees Celsius

Oxygen saturation: 97% on room air

GCS: 13/15

FAST test: Positive · facial drooping left side, arm weakness left side, speech severely affected

NEUROLOGICAL EXAMINATION:

Left sided facial droop confirmed on examination

Left arm drift present · unable to hold arm raised for 10 seconds

Speech dysarthric and partially aphasic

Pupils equal and reactive bilaterally

No neck stiffness

CURRENT MEDICATIONS:

Ramipril 5mg daily · hypertension

Simvastatin 40mg daily · hypercholesterolaemia

Aspirin 100mg daily · cardiovascular prophylaxis

PAST MEDICAL HISTORY:

Hypertension · diagnosed 2018

Hypercholesterolaemia · diagnosed 2019

Type 2 diabetes mellitus · diagnosed 2020, diet controlled

TIA · possible episode reported 6 months ago, not formally investigated

INVESTIGATIONS PERFORMED:

ECG: Normal sinus rhythm, no acute changes

Random glucose: 8.4 mmol/L

Full blood count: Within normal limits

Renal function: Creatinine 98 umol/L, eGFR 67 mL/min

FAMILY HISTORY:

Father · fatal stroke aged 68

Mother · hypertension, alive aged 84

SOCIAL HISTORY:

Retired engineer, lives with wife

Ex-smoker · stopped 2010, 30 pack year history

Alcohol · occasional, less than 10 units per week

CLINICAL IMPRESSION:

High clinical suspicion of acute ischaemic stroke. Symptom onset within thrombolysis window. Patient requires immediate neurological assessment and urgent CT head with CT angiography to determine eligibility for thrombolysis or thrombectomy. Every minute counts.

Urgency: EMERGENCY · please see immediately.