

CSE402: Optimisation

IRFAN RAHMAN &
MAHMUDUL HASAN TURJOY

Last updated: 8 October 2026

These notes were taken in the course **CSE402: Optimisation** taught at BRAC University in Fall 2026 by Nafis Faisal.

Contents

1	Lecture 01	4
1.1	Review on linear algebra	4
1.1.1	Vectors	4
1.1.2	Linear transformations	6
1.1.3	Basis and dimension	6
1.1.4	Eigenvectors and eigenvalues	6

1 Lecture 01

Date: 06/10/2026

Mathematical language of this course is linear algebra (of real vector spaces) and multivariable calculus. In this lecture, we will give a brief review of linear algebra.

§1.1 Review on linear algebra

Notation. Let $a_1, a_2, \dots, a_n$ be a finite sequence of real numbers. We will denote the sum

$$S = a_1 + a_2 + \dots + a_n \quad (1.1)$$

by

$$S = \sum_{k=1}^n a_k. \quad (1.2)$$

We have the following useful identities:

$$\sum_{k=1}^n k = \frac{n(n+1)}{2}, \quad (1.3)$$

$$\sum_{k=1}^n k^2 = \frac{n(n+1)(2n+1)}{6}. \quad (1.4)$$

§1.1.1 Vectors

Definition 1.1.1 ($\mathbb{R}$ -vector space). Let V be a non-empty set. The set V equipped with two binary operations

$$+ : V \times V \rightarrow V$$

called *vector addition*, and

$$\cdot : \mathbb{R} \times V \rightarrow V$$

called *scalar multiplication* is called a **vector space over $\mathbb{R}$** or **$\mathbb{R}$ -vector space** if for all $u, v, w \in V$ and all $a, b \in \mathbb{F}$, the following axioms hold:

1. **Closure under addition:**

$$u + v \in V \quad (1.5)$$

2. **Commutativity of addition:**

$$u + v = v + u \quad (1.6)$$

3. **Associativity of addition:**

$$(u + v) + w = u + (v + w) \quad (1.7)$$

4. **Existence of zero vector:** There exists a vector $0 \in V$ such that

$$u + 0 = u \quad (1.8)$$

5. **Existence of additive inverse:** For every $u \in V$, there exists $-u \in V$ such that

$$u + (-u) = 0 \quad (1.9)$$

6. **Closure under scalar multiplication:**

$$au \in V \quad (1.10)$$

7. **Distributivity of scalar multiplication over vector addition:**

$$a(u + v) = au + av \quad (1.11)$$

8. **Distributivity of scalar multiplication over field addition:**

$$(a + b)u = au + bu \quad (1.12)$$

9. **Associativity of scalar multiplication:**

$$a(bu) = (ab)u \quad (1.13)$$

10. **Identity element of scalar multiplication:**

$$1u = u \quad (1.14)$$

Hence, $(V, +, \cdot)$ is called a vector space over the field $\mathbb{R}$. Elements of V are called **vectors** and elements of $\mathbb{R}$ are called **scalars**.

Convention. From now on, vector space means $\mathbb{R}$ -vector space, unless stated otherwise.

In this course, we will work with only the following special example of $\mathbb{R}$ -vector space.

Example 1.1.2

Let

$$V := \mathbb{R}^n = \{[x_1, x_2, \dots, x_n]^T \mid x_1, x_2, \dots, x_n \in \mathbb{R}\}. \quad (1.15)$$

For $[x_1, x_2, \dots, x_n]^T, [y_1, y_2, \dots, y_n]^T \in V$ and $\alpha \in \mathbb{R}$ define addition (+) and scalar multiplication ($\cdot$), respectively as follows:

$$\begin{bmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{bmatrix} + \begin{bmatrix} y_1 \\ y_2 \\ \vdots \\ y_n \end{bmatrix} = \begin{bmatrix} x_1 + y_1 \\ x_2 + y_2 \\ \vdots \\ x_n + y_n \end{bmatrix}, \quad (1.16)$$

$$\alpha \cdot \begin{bmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{bmatrix} = \begin{bmatrix} \alpha x_1 \\ \alpha x_2 \\ \vdots \\ \alpha x_n \end{bmatrix}. \quad (1.17)$$

Then it is an easy check that $V = \mathbb{R}^n$ forms an $\mathbb{R}$ -vector space.

§1.1.2 Linear transformations

In this course, we are mostly concerned with following special functions between vector spaces:

Not sure about that

Definition 1.1.3 (Linear transformation). Let V, W be two vector space. A **linear transformation** from V to W is a function $T : V \rightarrow W$ such that for all $\mathbf{x}, \mathbf{y} \in V$ and $\alpha \in \mathbb{R}$ the following holds true:

- **Additivity.**

$$T(\mathbf{x} + \mathbf{y}) = T(\mathbf{x}) + T(\mathbf{y}). \quad (1.18)$$

- **Homogeneity.**

$$T(\alpha \cdot \mathbf{x}) = \alpha \cdot T(\mathbf{x}). \quad (1.19)$$

Need to add visualisation of additivity and homogeneity property of a linear transformation.

Example 1.1.4

Let $V := \mathbb{R}^n, W := \mathbb{R}^m$. An $m \times n$ matrix M then defines a linear transformation $M : \mathbb{R}^n \rightarrow \mathbb{R}^m$ ^a as follows:

For $\mathbf{x} \in \mathbb{R}^n$,

$$M(\mathbf{x}) := M\mathbf{x}. \quad (1.20)$$

Here, M is an $m \times n$ matrix, $\mathbf{x}$ can is an $n \times 1$ matrix. So $M\mathbf{x}$ is an $m \times 1$ matrix. Therefore, M is indeed a function from $\mathbb{R}^n$ to $\mathbb{R}^m$.

Furthermore, for all $\mathbf{x}, \mathbf{y} \in \mathbb{R}^n$ and $\alpha \in \mathbb{R}$, the following holds by the properties of matrix multiplication:

$$M(\mathbf{x} + \mathbf{y}) = M\mathbf{x} + M\mathbf{y}, \quad (1.21)$$

$$M(\alpha\mathbf{x}) = \alpha M\mathbf{x} \quad (1.22)$$

Hence, M is indeed a linear transformation.

^aHere, we are denoting the linear transformation determined by the matrix M by the same letter M using abuse of notation.

Surprising fact is that an arbitrary linear transformation also comes from a matrix. Before that, we need the idea of basis.

§1.1.3 Basis and dimension

Definition 1.1.5 (Linear combination). Let V be a vector space, and $\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_n \in V$. A **linear combination** of $\mathbf{x}_1, \mathbf{x}_2, \dots, \mathbf{x}_n$ is an expression of the form

$$c_1\mathbf{x}_1, c_2\mathbf{x}_2, \dots, c_n\mathbf{x}_n, \quad (1.23)$$

where $c_1, c_2, \dots, c_n \in \mathbb{R}$.

§1.1.4 Eigenvectors and eigenvalues

Suppose we have a linear transformation $L : \mathbb{R}^n \rightarrow \mathbb{R}^m$. We are interested in finding a non-zero vector $\mathbf{x} \in \mathbb{R}^n$ such that

$$L\mathbf{x} = \lambda\mathbf{x}, \quad (1.24)$$

where $\lambda \in \mathbb{R}$.

Definition 1.1.6 (Eigenvector and eigenvalue). $\mathbf{x}, \lambda$ (if exists) are called an **eigenvector** and **eigenvalue** for L . The $(\mathbf{x}, \lambda)$ is called an **eigenpair** for L .

Theorem 1.1.7

Let L be a linear transformation. Then $(\mathbf{x}, \lambda)$ is an eigen pair of L if and only if $\det(L - \lambda I) = 0$.

Proof.

□