CSE402: Optimisation

Irfan Rahman &
Mahmudul Hasan Turjoy

Last updated: 14 October 2026

These notes were taken in the course **CSE402: Optimisation** taught at BRAC University in Fall 2026 by Nafis Faisal. The main text used is [**KW19**]. All errors are our responsibilty. If you find any error, please email us at mh.turjoy@yahoo.com or rasib.rahman.04@gmail.com, even if it is something very small, even if you're not sure.

Contents

1 Lecture 01 **4**

1.1 Review on linear algebra 4

1.1.1 Vectors 4

1.1.2 Linear transformations 6

1.1.3 Basis and Dimension of $\mathbb{R}^n$ 6

1.1.4 Dimension 10

1.1.5 Basis and Dimension of $\mathbb{R}^n$ 10

1.1.6 Eigenvectors and eigenvalues 11

2 Lecture 02 **14**

2.1 Review on linear algebra—continued 14

2.1.1 Inner product and norm 15

2.1.2 Metric 20

3 Lecture 03 **22**

3.1 Review on linear algebra—continued 22

1 Lecture 01

Date: 06/10/2026

Mathematical language of this course is linear algebra (of real vector spaces) and multivariable calculus. In this lecture, we will give a brief review of linear algebra.

§1.1 Review on linear algebra

Notation. Let $a_1, a_2, \ldots, a_n$ be a finite sequence of real numbers. We will denote the sum

$$S = a_1 + a_2 + \cdots + a_n \tag{1.1}$$

by

$$S = \sum_{k=1}^{n} a_k. \tag{1.2}$$

We have the following useful identities:

$$\sum_{k=1}^{n} k = \frac{n(n+1)}{2}, \tag{1.3}$$

$$\sum_{k=1}^{n} k^2 = \frac{n(n+1)(2n+1)}{6}. \tag{1.4}$$

§1.1.1 Vectors

Definition 1.1.1 ($\mathbb{R}$-vector space)**.** Let V be a non-empty set. The set V equipped with two binary operations

$$+ : V \times V \to V$$

called *vector addition*, and

$$\cdot : \mathbb{R} \times V \to V$$

called *scalar multiplication* is called a **vector space over** $\mathbb{R}$ or **$\mathbb{R}$-vector space** if for all $u, v, w \in V$ and all $a, b \in \mathbb{F}$, the following axioms hold:

1. **Closure under addition:**
$$u + v \in V \tag{1.5}$$

2. **Commutativity of addition:**
$$u + v = v + u \tag{1.6}$$

3. **Associativity of addition:**
$$(u + v) + w = u + (v + w) \tag{1.7}$$

4. **Existence of zero vector:** There exists a vector $0 \in V$ such that

$$u + 0 = u \tag{1.8}$$

5. **Existence of additive inverse:** For every $u \in V$, there exists $-u \in V$ such that

$$u + (-u) = 0 \tag{1.9}$$

6. **Closure under scalar multiplication:**

$$au \in V \tag{1.10}$$

7. **Distributivity of scalar multiplication over vector addition:**

$$a(u + v) = au + av \tag{1.11}$$

8. **Distributivity of scalar multiplication over field addition:**

$$(a + b)u = au + bu \tag{1.12}$$

9. **Associativity of scalar multiplication:**

$$a(bu) = (ab)u \tag{1.13}$$

10. **Identity element of scalar multiplication:**

$$1u = u \tag{1.14}$$

Hence, $(V, +, \cdot)$ is called a vector space over the field $\mathbb{R}$. Elements of V are called **vectors** and elements of $\mathbb{R}$ are called **scalars**.

Convention. From now on, vector space means $\mathbb{R}$-vector space, unless stated otherwise.

In this course, we will work with only the following special example of $\mathbb{R}$-vector space.

Example 1.1.2

Let

$$V \coloneqq \mathbb{R}^n = \{[x_1, x_2 \ldots, x_n]^T \mid x_1, x_2, \ldots, x_n \in \mathbb{R}\}. \tag{1.15}$$

For $[x_1, x_2 \ldots, x_n]^T, [y_1, y_2 \ldots, y_n]^T \in V$ and $\alpha \in \mathbb{R}$ define addition $(+)$ and scalar multiplication $(\cdot)$, respectively as follows:

$$\begin{bmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{bmatrix} + \begin{bmatrix} y_1 \\ y_2 \\ \vdots \\ y_n \end{bmatrix} = \begin{bmatrix} x_1 + y_1 \\ x_2 + y_2 \\ \vdots \\ x_n + y_n \end{bmatrix}, \tag{1.16}$$

$$\alpha \cdot \begin{bmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{bmatrix} = \begin{bmatrix} \alpha x_1 \\ \alpha x_2 \\ \vdots \\ \alpha x_n \end{bmatrix}. \tag{1.17}$$

Then it is an easy check that $V = \mathbb{R}^n$ forms an $\mathbb{R}$-vector space.

§1.1.2 Linear transformations

In this course, we are mostly concerned with following special functions between vector spaces:

Not sure about that

Definition 1.1.3 (Linear transformation)**.** Let V, W be two vector space. A **linear transformation** from V to W is a function $T : V \to W$ such that for all $\mathbf{x}, \mathbf{y} \in V$ and $\alpha \in \mathbb{R}$ the following holds true:

- **Additivity.**
$$T(\mathbf{x} + \mathbf{y}) = T(\mathbf{x}) + T(\mathbf{y}). \tag{1.18}$$
- **Homogeneity.**
$$T(\alpha \cdot \mathbf{x}) = \alpha \cdot T(\mathbf{x}). \tag{1.19}$$

Need to add visualisation of additivity and homogeneity property of a linear transformation.

Example 1.1.4

Let $V := \mathbb{R}^n, W := \mathbb{R}^m$. An $m \times n$ matrix M then defines a linear transformation $M : \mathbb{R}^n \to \mathbb{R}^m$[a] as follows:

For $\mathbf{x} \in \mathbb{R}^n$,

$$M(\mathbf{x}) := M\mathbf{x}. \tag{1.20}$$

Here, M is an $m \times n$ matrix, $\mathbf{x}$ can is an $n \times 1$ matrix. So $M\mathbf{x}$ is an $m \times 1$ matrix. Therefore, M is indeed a function from $\mathbb{R}^n$ to $\mathbb{R}^m$.

Furthermore, for all $\mathbf{x}, \mathbf{y} \in \mathbb{R}^n$ and $\alpha \in \mathbb{R}$, the following holds by the properties of matrix multiplication:

$$M(\mathbf{x} + \mathbf{y}) = M\mathbf{x} + M\mathbf{y}, \tag{1.21}$$

$$M(\alpha\mathbf{x}) = \alpha M\mathbf{x} \tag{1.22}$$

Hence, M is indeed a linear transformation.

[a] Here, we are denoting the linear transformation determined by the matrix M by the same letter M using abuse of notation.

Surprising fact is that an arbitrary linear transformation also comes from a matrix. Before that, we need the idea of basis.

§1.1.3 Basis and Dimension of $\mathbb{R}^n$

Consider the vector space

$$\mathbb{R}^2 = \left\{ [x, y]^T : x, y \in \mathbb{R} \right\}. \tag{1.23}$$

A natural question is: how many vectors are required to generate every vector in $\mathbb{R}^2$? Answer to this question is particularly useful in the context of linear transformations.

Suppose

$$L : \mathbb{R}^2 \to V$$

is a linear transformation. Because $\mathbb{R}^2$ contains infinitely many vectors, describing which vector maps to what under L would require an infinite amount of information.

However, suppose every vector of $\mathbb{R}^2$ can be generated by a finite collection of vectors

$$v_1, \ldots, v_n \tag{1.24}$$

by taking "linear combinations". Then, instead of knowing the value of L on every vector, we only need to know the values

$$L(v_1), \ldots, L(v_n). \tag{1.25}$$

Indeed, if

$$v = a_1 v_1 + \cdots + a_n v_n, \tag{1.26}$$

then by the linearity of L,

$$L(v) = L(a_1 v_1 + \cdots + a_n v_n) = a_1 L(v_1) + \cdots + a_n L(v_n). \tag{1.27}$$

Therefore, knowing the images of $v_1, \ldots, v_n$ completely determines the linear transformation L on the whole space.

This motivates the following question: What is the smallest collection of vectors required to generate all vectors of a vector space?

The answer leads us to the concepts of spanning sets, linear independence, basis, and dimension:

Definition 1.1.5 (Linear Independence)**.** A collection of vectors

$$\{v_1, v_2, \ldots, v_k\} \tag{1.28}$$

in a vector space V is called *linearly independent* if

$$a_1 v_1 + a_2 v_2 + \cdots + a_k v_k = 0 \tag{1.29}$$

implies

$$a_1 = a_2 = \cdots = a_k = 0. \tag{1.30}$$

If there exists a non-zero solution of the above equation, then the vectors are called *linearly dependent*.

Intuitively, if a list of vectors are linearly dependent, then at least one of them can be generated by rest of them.

Example 1.1.6

The vectors

$$\begin{bmatrix} 1 \\ 1 \end{bmatrix}, \qquad \begin{bmatrix} 1 \\ -1 \end{bmatrix} \tag{1.31}$$

are linearly independent.

However,

$$\begin{bmatrix} 1 \\ 2 \end{bmatrix}, \qquad \begin{bmatrix} 2 \\ 4 \end{bmatrix} \tag{1.32}$$

are linearly dependent because the second vector is a scalar multiple of the first.

Definition 1.1.7 (Span)**.** Let V be a vector space and let

$$S = \{v_1, v_2, \ldots, v_n\} \tag{1.33}$$

be a collection of vectors in V. The span of S, denoted by

$$\text{span}\{v_1, v_2, \ldots, v_n\}, \tag{1.34}$$

is the set of all possible linear combinations of the vectors in S. That is,

$$\text{span}\{v_1, v_2, \ldots, v_n\} = \{a_1 v_1 + a_2 v_2 + \cdots + a_n v_n : a_1, \ldots, a_n \in \mathbb{R}\}. \tag{1.35}$$

If

$$\text{span}\{v_1, v_2, \ldots, v_n\} = V, \tag{1.36}$$

then we say that $v_1, v_2, \ldots, v_n$ span the vector space V.

A basis is a minimal spanning set, that is, it contains enough vectors to generate the whole space, but no redundant vectors.

Example 1.1.8

Consider the vectors in $\mathbb{R}^2$

$$v_1 = \begin{bmatrix} 1 \\ 1 \end{bmatrix}, \qquad v_2 = \begin{bmatrix} 1 \\ -1 \end{bmatrix}. \tag{1.37}$$

The span of these two vectors is

$$\text{span}\{v_1, v_2\} = \left\{ a \begin{bmatrix} 1 \\ 1 \end{bmatrix} + b \begin{bmatrix} 1 \\ -1 \end{bmatrix} : a, b \in \mathbb{R} \right\}. \tag{1.38}$$

Now,

$$a \begin{bmatrix} 1 \\ 1 \end{bmatrix} + b \begin{bmatrix} 1 \\ -1 \end{bmatrix} = \begin{bmatrix} a+b \\ a-b \end{bmatrix}. \tag{1.39}$$

Given any vector

$$\begin{bmatrix} x \\ y \end{bmatrix} \in \mathbb{R}^2, \tag{1.40}$$

choosing

$$a = \frac{x+y}{2}, \qquad b = \frac{x-y}{2} \tag{1.41}$$

gives

$$\begin{bmatrix} x \\ y \end{bmatrix} = a \begin{bmatrix} 1 \\ 1 \end{bmatrix} + b \begin{bmatrix} 1 \\ -1 \end{bmatrix}. \tag{1.42}$$

Therefore,

$$\text{span}\{v_1, v_2\} = \mathbb{R}^2. \tag{1.43}$$

Hence, the vectors v_1, v_2 span $\mathbb{R}^2$.

Definition 1.1.9 (Basis)**.** Let V be a vector space. A set of vectors

$$\mathcal{B} = \{v_1, v_2, \ldots, v_k\} \tag{1.44}$$

is called a *basis* of V if:

1. $v_1, v_2, \ldots, v_k$ span V, i.e.

$$V = \text{span}\{v_1, v_2, \ldots, v_k\}, \tag{1.45}$$

2. $v_1, v_2, \ldots, v_k$ are linearly independent.

Example 1.1.10

Consider the set of vectors

$$\mathcal{B} = \left\{ \begin{bmatrix} 1 \\ 1 \end{bmatrix}, \begin{bmatrix} 1 \\ -1 \end{bmatrix} \right\}. \tag{1.46}$$

We claim that $\mathcal{B}$ is a basis of $\mathbb{R}^2$.
First, let

$$\begin{bmatrix} x \\ y \end{bmatrix} \in \mathbb{R}^2 \tag{1.47}$$

be an arbitrary vector. We want to express it as a linear combination of the vectors in $\mathcal{B}$. Suppose

$$\begin{bmatrix} x \\ y \end{bmatrix} = a \begin{bmatrix} 1 \\ 1 \end{bmatrix} + b \begin{bmatrix} 1 \\ -1 \end{bmatrix}. \tag{1.48}$$

Then,

$$\begin{bmatrix} x \\ y \end{bmatrix} = \begin{bmatrix} a+b \\ a-b \end{bmatrix}. \tag{1.49}$$

Solving the equations

$$a + b = x, \qquad a - b = y, \tag{1.50}$$

we get

$$a = \frac{x+y}{2}, \qquad b = \frac{x-y}{2}. \tag{1.51}$$

Therefore, every vector in $\mathbb{R}^2$ can be written as a linear combination of the vectors in $\mathcal{B}$. Hence,

$$\text{span}(\mathcal{B}) = \mathbb{R}^2. \tag{1.52}$$

Also, the two vectors in $\mathcal{B}$ are linearly independent. Therefore, $\mathcal{B}$ is a basis of $\mathbb{R}^2$.
Moreover, the representation

$$\begin{bmatrix} x \\ y \end{bmatrix} = a \begin{bmatrix} 1 \\ 1 \end{bmatrix} + b \begin{bmatrix} 1 \\ -1 \end{bmatrix} \tag{1.53}$$

is unique. The scalars a and b are called the coordinates of the vector with respect to the basis $\mathcal{B}$.

§1.1.4 Dimension

The number of vectors contained in a basis gives the size of a vector space.

Theorem 1.1.11

Every basis of a finite-dimensional vector space contains the same number of vectors.

Proof. This proof was not shown in the class. Interested students may refer to [Axl24, p. 44]. □

is it examinable?

Definition 1.1.12 (Dimension)**.** The number of vectors in a basis of a vector space V is called the *dimension* of V, denoted by

$$\dim(V).$$

Example 1.1.13

Since every basis of $\mathbb{R}^2$ contains exactly two vectors,

$$\dim(\mathbb{R}^2) = 2.$$

§1.1.5 Basis and Dimension of $\mathbb{R}^n$

The standard basis of $\mathbb{R}^n$ is

$$e_1 = \begin{bmatrix} 1 \\ 0 \\ \vdots \\ 0 \end{bmatrix}, \quad e_2 = \begin{bmatrix} 0 \\ 1 \\ \vdots \\ 0 \end{bmatrix}, \quad \ldots, \quad e_n = \begin{bmatrix} 0 \\ 0 \\ \vdots \\ 1 \end{bmatrix}. \tag{1.54}$$

Every vector

$$x = \begin{bmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{bmatrix} \tag{1.55}$$

can be written as

$$x_1 e_1 + x_2 e_2 + \cdots + x_n e_n. \tag{1.56}$$

Therefore,

$$\{e_1, e_2, \ldots, e_n\} \tag{1.57}$$

is a basis of $\mathbb{R}^n$.

Theorem 1.1.14

The dimension of $\mathbb{R}^n$ is

$$\dim(\mathbb{R}^n) = n. \tag{1.58}$$

§1.1.6 Eigenvectors and eigenvalues

Suppose we have a linear transformation $L : \mathbb{R}^n \to \mathbb{R}^m$. We are interested in finding a non-zero vector $\mathbf{x} \in \mathbb{R}^n$ such that

$$L\mathbf{x} = \lambda\mathbf{x}, \tag{1.59}$$

where $\lambda \in \mathbb{R}$.

Definition 1.1.15 (Eigenvector and eigenvalue)**.** $\mathbf{x}, \lambda$ (if exists) are called an **eigenvector** and **eigenvalue** for L. The $(\mathbf{x}, \lambda)$ is called an **eigenpair** for L.

Theorem 1.1.16

Let L be a linear transformation. Then $(\mathbf{x}, \lambda)$ is an eigen pair of L if and only if $\det(L - \lambda I) = 0$.

Proof. Suppose λ is an eigenvalue of L. Then there exists a non-zero vector $\mathbf{x} \in \mathbb{R}^n$ such that

$$L\mathbf{x} = \lambda\mathbf{x}. \tag{1.60}$$

Since $I\mathbf{x} = \mathbf{x}$, we have

$$L\mathbf{x} - \lambda I\mathbf{x} = 0, \tag{1.61}$$

or equivalently,

$$(L - \lambda I)\mathbf{x} = 0. \tag{1.62}$$

Since $\mathbf{x} \neq 0$, this homogeneous system has a non-trivial solution. Therefore, $L - \lambda I$ is singular. Hence,

$$\det(L - \lambda I) = 0. \tag{1.63}$$

Conversely, suppose $\det(L-\lambda I) = 0$. Then $L-\lambda I$ is singular, so there exists a non-zero vector $\mathbf{x} \in \mathbb{R}^n$ satisfying

$$(L - \lambda I)\mathbf{x} = 0. \tag{1.64}$$

Therefore,

$$L\mathbf{x} = \lambda\mathbf{x}. \tag{1.65}$$

Thus, λ is an eigenvalue of L, with corresponding eigenvector $\mathbf{x}$. □

Definition 1.1.17 (Eigenvalues of a matrix)**.** Let M be an $n \times n$ matrix. The set of all eigenvalues of M is denoted by

$$\text{eigs}(M) = \{\lambda_1, \lambda_2, \ldots, \lambda_n\}. \tag{1.66}$$

Here, the eigenvalues are listed with multiplicity.

Definition 1.1.18 (Definiteness of a matrix)**.** Let M be a real symmetric $n \times n$ matrix with eigenvalues $\lambda_1, \lambda_2, \ldots, \lambda_n$. Then M is called

1. **Positive definite (PD)** if all eigenvalues are strictly positive, i.e.,
$$\lambda_i > 0, \quad \forall i = 1, 2, \ldots, n. \tag{1.67}$$
2. **Negative definite (ND)** if all eigenvalues are strictly negative, i.e.,
$$\lambda_i < 0, \quad \forall i = 1, 2, \ldots, n. \tag{1.68}$$
3. **Positive semidefinite (PSD)** if all eigenvalues are non-negative, i.e.,
$$\lambda_i \geq 0, \quad \forall i = 1, 2, \ldots, n. \tag{1.69}$$
4. **Negative semidefinite (NSD)** if all eigenvalues are non-positive, i.e.,
$$\lambda_i \leq 0, \quad \forall i = 1, 2, \ldots, n. \tag{1.70}$$
5. **Indefinite (ID)** if M has both positive and negative eigenvalues.

Example 1.1.19

Let
$$L := \begin{bmatrix} -2 & 2 \\ 3 & -1 \end{bmatrix}. \tag{1.71}$$
We will find the eigenvalues and eigenvectors for L.

The eigenvalues are obtained by solving $\det(L - \lambda I) = 0$. Here,
$$\begin{aligned} \det(L - \lambda I) &= \begin{vmatrix} -2-\lambda & 2 \\ 3 & -1-\lambda \end{vmatrix} \\ &= (-2-\lambda)(-1-\lambda) - 6 \\ &= \lambda^2 + 3\lambda - 4 \\ &= (\lambda - 1)(\lambda + 4). \end{aligned} \tag{1.72}$$
Hence, the eigenvalues for L are $\lambda = 1$ and $\lambda = -4$.

For each eigenvalue, we find a non-zero vector $\mathbf{x} = [x_1, x_2]^T$ such that $(L - \lambda I)\mathbf{x} = 0$.

For $\lambda = 1$, we have
$$\begin{bmatrix} -3 & 2 \\ 3 & -2 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}. \tag{1.73}$$
Both equations give $3x_1 = 2x_2$, so $x_2 = \frac{3}{2}x_1$. Therefore, the eigenvectors corresponding to $\lambda = 1$ are precisely the vectors
$$\mathbf{x} = \begin{bmatrix} x_1 \\ \frac{3}{2}x_1 \end{bmatrix} = x_1 \begin{bmatrix} 1 \\ \frac{3}{2} \end{bmatrix}, \qquad x_1 \in \mathbb{R} \setminus \{0\}. \tag{1.74}$$
Choosing $x_1 = 2$, we obtain the eigenvector $\mathbf{x} = [2, 3]^T$.

For $\lambda = -4$, we have
$$\begin{bmatrix} 2 & 2 \\ 3 & 3 \end{bmatrix} \begin{bmatrix} x_1 \\ x_2 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \end{bmatrix}. \tag{1.75}$$

Both equations give $x_1 + x_2 = 0$, so $x_2 = -x_1$. Therefore, the eigenvectors corresponding to $\lambda = -4$ are precisely the vectors

$$\mathbf{x} = \begin{bmatrix} x_1 \\ -x_1 \end{bmatrix} = x_1 \begin{bmatrix} 1 \\ -1 \end{bmatrix}, \qquad x_1 \in \mathbb{R} \setminus \{0\}. \tag{1.76}$$

Choosing $x_1 = 1$, we obtain the eigenvector $\mathbf{x} = [1, -1]^T$.
Hence, $([2, 3]^T, 1)$ and $([1, -1]^T, -4)$ are eigenpairs for L.

Remark 1.1.20. Any nonzero scalar multiple of an eigenvector is also an eigenvector corresponding to the same eigenvalue.

Indeed, let $x \neq 0$ be an eigenvector of L corresponding to the eigenvalue λ. Then

$$Lx = \lambda x. \tag{1.77}$$

For any scalar $c \neq 0$, by linearity of L,

$$L(cx) = cLx = c\lambda x = \lambda(cx). \tag{1.78}$$

Hence, cx is also an eigenvector of L with the same eigenvalue λ.

2 Lecture 02

Date: 11/10/2026

§2.1 Review on linear algebra—continued

Exercise 2.1.1. Find eigenvalues of the matrix

$$\begin{bmatrix} 6 & 0 \\ 0 & 7 \end{bmatrix}, \tag{2.1}$$

and classify it in terms of eigenvalues.

Answer 2.1.2. Let M denote the matrix in (2.1). The eigenvalues are obtained by solving $\det(M - \lambda I) = 0$. Here,

$$\begin{aligned} \det(M - \lambda I) &= \begin{vmatrix} 6-\lambda & 0 \\ 0 & 7-\lambda \end{vmatrix} \\ &= (6-\lambda)(7-\lambda). \end{aligned} \tag{2.2}$$

Therefore,

$$(6-\lambda)(7-\lambda) = 0 \iff \lambda = 6 \text{ or } \lambda = 7. \tag{2.3}$$

Hence, $\text{eigs}(M) = \{6, 7\}$. Since M is real symmetric and both eigenvalues are strictly positive, it is *positive definite (PD)* by Definition 1.1.18.

Remark 2.1.3. The eigenvalues of a diagonal matrix are precisely its diagonal entries.

Now consider the following matrix:

$$\begin{bmatrix} 1 & -2 \\ 1 & -1 \end{bmatrix}. \tag{2.4}$$

To find its eigenvalues, we solve

$$\begin{aligned} \det(A - \lambda I) &= \begin{vmatrix} 1-\lambda & -2 \\ 1 & -1-\lambda \end{vmatrix} \\ &= (1-\lambda)(-1-\lambda) + 2 \\ &= -1 - \lambda + \lambda + \lambda^2 + 2 \\ &= \lambda^2 + 1 = 0. \end{aligned}$$

Therefore,

$$\lambda^2 = -1 \implies \lambda = \pm i.$$

Its eigenvalues $\pm i$ are not real. Therefore, this matrix does not fall into any class given in Definition 1.1.18. But there are matrices that guaranteed to have real eigenvalues, which we discuss next. Recall:

Definition 2.1.4 (Transpose of a matrix)**.** Let A be an $m \times n$ matrix. Denote its entry whose row number is i and column number is j by $a_{i,j}$. Then transpose of the matrix A, denoted by A^T, is an $n \times m$ matrix whose entry at j-th row and i-th column is $a_{i,j}$. Symbolically, if

$$A = \begin{bmatrix} a_{1,1} & a_{1,2} & \dots & a_{1,n} \\ a_{2,1} & a_{2,2} & \dots & a_{2,n} \\ \vdots & \vdots & \ddots & \vdots \\ a_{m,1} & a_{m,2} & \dots & a_{m,n} \end{bmatrix}, \tag{2.5}$$

then

$$A^T = \begin{bmatrix} a_{1,1} & a_{2,1} & \dots & a_{m,1} \\ a_{1,2} & a_{2,2} & \dots & a_{m,2} \\ \vdots & \vdots & \ddots & \vdots \\ a_{1,n} & a_{2,n} & \dots & a_{m,n} \end{bmatrix}. \tag{2.6}$$

Definition 2.1.5 (Symmetric matrix)**.** We call an $n \times n$ matrix A to be *symmetric* if

$$A = A^T. \tag{2.7}$$

Theorem 2.1.6

All the eigenvalues of a symmetric matrix A are real.

Proof. Will be done in the next class after developing some new idea. □

To prove this theorem, we now introduce new ideas in the context of vector spaces.

§2.1.1 Inner product and norm

It is generalisation of dot product of vectors that we learnt in the HSC/A-levels.

Example 2.1.7 (Motivating example)

Consider the $\mathbb{R}$-vector space $\mathbb{R}^2$. Define a function

$$\langle _, _ \rangle : \mathbb{R}^2 \times \mathbb{R}^2 \to \mathbb{R}, \tag{2.8}$$

given by

$$\langle _, _ \rangle (\mathbf{x}, \mathbf{y}) \coloneqq \mathbf{x}^T \mathbf{y}. \tag{2.9}$$

We will denote $\langle _, _ \rangle (\mathbf{x}, \mathbf{y})$ by $\langle \mathbf{x}, \mathbf{y} \rangle$.

Therefore, $\langle \mathbf{x}, \mathbf{y} \rangle$ is nothing but dot product of $\mathbf{x}$ and $\mathbf{y}$ in the high school sense, i.e.,

$$\langle \mathbf{x}, \mathbf{y} \rangle = \mathbf{x} \cdot \mathbf{y} \tag{2.10}$$

If $\mathbf{x} = \begin{bmatrix} x_1 & x_2 \end{bmatrix}^T$ and $\mathbf{y} = \begin{bmatrix} y_1 & y_2 \end{bmatrix}^T$, then

$$\langle \mathbf{x}, \mathbf{y} \rangle = \mathbf{x} \cdot \mathbf{y} = \sum_{k=1}^{2} x_k y_k. \tag{2.11}$$

We can then write length of a vector $\mathbf{x} = \begin{bmatrix} x_1 & x_2 \end{bmatrix} \in \mathbb{R}^2$, denoted by $\|\mathbf{x}\|$, in terms of dot product defined as above:

$$\|x\| = \sqrt{x_1^2 + x_2^2} = \sqrt{\langle x_1, x_2 \rangle}. \tag{2.12}$$

Furthermore, distance of $\mathbf{x}, \mathbf{y} \in \mathbb{R}^2$ can also be written in terms of the function $\langle _, _ \rangle$:

$$\begin{aligned}
&\sqrt{(x_1 - y_1)^2 + (x_2 - y_2)^2} && (2.13)\\
=&\sqrt{\langle (x_1 - y_1, x_2 - y_2), (x_1 - y_1, x_2 - y_2) \rangle} && (2.14)\\
=&\sqrt{\langle \mathbf{x} - \mathbf{y}, \mathbf{x} - \mathbf{y} \rangle} && (2.15)\\
=&\|\mathbf{x} - \mathbf{y}\|. && (2.16)
\end{aligned}$$

Notice, we previously did not have the notion of length of a vector or distance between two vectors in $\mathbb{R}^n$, for arbitrary n. But the above definition of dot product can be generalised to any $\mathbb{R}^n$, and using that we can meaningfully introduce the notion of length and distance in arbitrary $\mathbb{R}^n$. This generalised notion of dot product and length will be called *inner product* and *norm*, respectively.

Definition 2.1.8 (Inner product)**.** An *inner product* on a vector space V over $\mathbb{R}$ is a function

$$\langle \cdot, \cdot \rangle : V \times V \longrightarrow \mathbb{R} \tag{2.17}$$

which assigns a real number $\langle u, v \rangle$ to every pair of vectors $u, v \in V$, satisfying the following properties for all $u, v, w \in V$ and $c \in \mathbb{R}$:

1. **Linearity in the first argument:**

$$\langle cu + v, w \rangle = c\langle u, w \rangle + \langle v, w \rangle. \tag{2.18}$$

2. **Symmetry:**

$$\langle u, v \rangle = \langle v, u \rangle. \tag{2.19}$$

3. **Positive definiteness:**

$$\langle u, u \rangle \geq 0, \tag{2.20}$$

with equality if and only if $u = 0$.

A vector space equipped with an inner product is called an *inner product space*.

Example 2.1.9

In the vector space $\mathbb{R}^n$ define

$$\langle \mathbf{x}, \mathbf{y} \rangle = \mathbf{x}^T \mathbf{y}. \tag{2.21}$$

Then it can be checked that the definition above defines an inner product on $\mathbb{R}^n$.

Let $\mathbf{x}, \mathbf{y}, \mathbf{z} \in \mathbb{R}^n$ and $c \in \mathbb{R}$. We verify the properties of an inner product.

1. **Linearity in the first argument:**

$$\begin{aligned}\langle c\mathbf{x} + \mathbf{y}, \mathbf{z}\rangle &= (c\mathbf{x} + \mathbf{y})^T\mathbf{z} \\ &= c\mathbf{x}^T\mathbf{z} + \mathbf{y}^T\mathbf{z} \\ &= c\langle \mathbf{x}, \mathbf{z}\rangle + \langle \mathbf{y}, \mathbf{z}\rangle.\end{aligned}$$

2. **Symmetry:**

$$\begin{aligned}\langle \mathbf{x}, \mathbf{y}\rangle &= \mathbf{x}^T\mathbf{y} \\ &= \sum_{i=1}^{n} x_i y_i \\ &= \sum_{i=1}^{n} y_i x_i \\ &= \mathbf{y}^T\mathbf{x} \\ &= \langle \mathbf{y}, \mathbf{x}\rangle.\end{aligned}$$

3. **Positive definiteness:**

$$\begin{aligned}\langle \mathbf{x}, \mathbf{x}\rangle &= \mathbf{x}^T\mathbf{x} \\ &= \sum_{i=1}^{n} x_i^2 \geq 0.\end{aligned}$$

Since $x_i^2 \geq 0$ for every i, equality holds if and only if $x_i = 0$ for all i, i.e., $\mathbf{x} = \mathbf{0}$.

Hence, the Euclidean dot product defines an inner product on $\mathbb{R}^n$.

Definition 2.1.10 (Norm induced by an inner product)**.** Suppose V is an $\mathbb{R}$-vector space with an inner product $\langle _, _\rangle$ defined on it. Then the function

$$\|.\| : V \to \mathbb{R} \tag{2.22}$$

defined by

$$\|\mathbf{x}\| \coloneqq \sqrt{\langle \mathbf{x}, \mathbf{x}\rangle} \tag{2.23}$$

is called *norm induced by this inner product.*

Example 2.1.11 (Euclidean norm on $\mathbb{R}^n$)

For $\mathbf{x} = (x_1, x_2, \ldots, x_n)^T \in \mathbb{R}^n$, the norm induced by the Euclidean inner product is

$$\begin{aligned}\|\mathbf{x}\|_2 &= \sqrt{\langle \mathbf{x}, \mathbf{x}\rangle} \\ &= \sqrt{\mathbf{x}^T\mathbf{x}}\end{aligned}$$

$$= \sqrt{x_1^2 + x_2^2 + \cdots + x_n^2}.$$

This is called the *Euclidean norm* or *2-norm* on $\mathbb{R}^n$.

Proposition 2.1.12

Norm induced by an inner product has some intrinsic properties:
For all $\mathbf{x}, \mathbf{y} \in V$ and $c \in \mathbb{R}$

1. **Positive definiteness:**
$$\|\mathbf{x}\| \geq 0, \tag{2.24}$$
and
$$\|\mathbf{x}\| = 0 \iff \mathbf{x} = 0. \tag{2.25}$$
2. **Homogeneity:**
$$\|c\mathbf{x}\| = |c|\,\|\mathbf{x}\|. \tag{2.26}$$
3. **Triangle inequality:**
$$\|\mathbf{x} + \mathbf{y}\| \leq \|\mathbf{x}\| + \|\mathbf{y}\|. \tag{2.27}$$

By making these properties definition, we can define general notion of norm that is independent from inner product.

Definition 2.1.13 (Norm on a vector space)**.** Let V be a real vector space. A *norm* on V is a function

$$\|\cdot\| : V \longrightarrow \mathbb{R}$$

that assigns a real number $\|v\|$ to each vector $v \in V$, satisfying the following properties for all $u, v \in V$ and all $c \in \mathbb{R}$:

1. **Non-negativity:**
$$\|v\| \geq 0. \tag{2.28}$$
2. **Definiteness:**
$$\|v\| = 0 \iff v = 0. \tag{2.29}$$
3. **Absolute homogeneity:**
$$\|cv\| = |c|\,\|v\|. \tag{2.30}$$
4. **Triangle inequality:**
$$\|u + v\| \leq \|u\| + \|v\|. \tag{2.31}$$

A real vector space V equipped with a norm is called a *normed vector space*.

Following example is particularly important for this course.

Example 2.1.14 (p-norm)

For $p \in \mathbb{R}$, define
$$\|.\|_p : \mathbb{R}^n \to \mathbb{R} \tag{2.32}$$


by

$$\|\mathbf{x}\|_p = \left(|x_1|^p + |x_2|^p + \cdots + |x_n|^p\right)^{\frac{1}{p}}, \tag{2.33}$$

where $\mathbf{x} = [x_1, \ldots, x_n]^T$.
We are not going to prove how this defines a norm on $\mathbb{R}^n$ though.
Putting $p = 2$ gives us the Euclidean norm of Example 2.1.11.
Putting $p = 1$ gives us

$$\|x\|_1 = |x_1| + |x_2| + \cdots + |x_n|, \tag{2.34}$$

which is also called "Manhattan norm" or "Taxicab" norm.
For example, consider the points $\mathbf{x} = (1, 1)$ and $\mathbf{y} = (4, 3)$ in $\mathbb{R}^2$.



The Manhattan distance between these points is

$$d_1(\mathbf{x}, \mathbf{y}) = \|\mathbf{x} - \mathbf{y}\|_1 = |4 - 1| + |3 - 1| = 5.$$

Letting $p \to \infty$ gives

$$\|\mathbf{x}\|_p = \max(|x_1|, |x_2|, \ldots, |x_n|) \tag{2.35}$$

because if $x_i = \max(|x_1|, |x_2|, \ldots, |x_n|)$, then

$$\begin{aligned}
\lim_{p\to\infty} \|x\|_p &= \lim_{p\to\infty} (|x_i|^p)^{\frac{1}{p}} \left(\left|\frac{x_1}{x_i}\right|^p + \left|\frac{x_2}{x_i}\right|^p + \cdots + \left|\frac{x_i}{x_i}\right|^p + \cdots + \left|\frac{x_n}{x_i}\right|^p \right)^{\frac{1}{p}} \\
&= |x_i|(0 + 0 + \cdots + 1 + \cdots + 0)^0 \\
&= |x_i|.
\end{aligned}$$

Question 2.1.15. Let $\mathbf{x} = \begin{bmatrix} -6 \\ 4 \\ 0 \\ 1 \end{bmatrix}$. Find the 1-norm and 3-norm of $\mathbf{x}$.

Answer 2.1.16. Using the definition of the p-norm, we have
For $p = 1$,

$$\begin{aligned}
\|\mathbf{x}\|_1 &= |-6| + |4| + |0| + |1| \\
&= 6 + 4 + 0 + 1 \\
&= 11.
\end{aligned}$$

For $p = 3$,

$$\|\mathbf{x}\|_3 = \left(|-6|^3 + |4|^3 + |0|^3 + |1|^3\right)^{\frac{1}{3}}$$

$$= (216 + 64 + 0 + 1)^{\frac{1}{3}}$$
$$= \sqrt[3]{281}.$$

§2.1.2 Metric

Definition 2.1.17 (Metric)**.** Let V be a nonempty set. A function

$$d : V \times V \to \mathbb{R} \tag{2.36}$$

is called a *metric* if for all $\mathbf{u}, \mathbf{v}, \mathbf{w} \in V$, the following properties hold:

1. **Positive definiteness:**
$$d(\mathbf{u}, \mathbf{v}) \geq 0, \tag{2.37}$$
with equality if and only if $\mathbf{u} = \mathbf{v}$.

2. **Symmetry:**
$$d(\mathbf{u}, \mathbf{v}) = d(\mathbf{v}, \mathbf{u}). \tag{2.38}$$

3. **Triangle inequality:**
$$d(\mathbf{u}, \mathbf{w}) \leq d(\mathbf{u}, \mathbf{v}) + d(\mathbf{v}, \mathbf{w}). \tag{2.39}$$

A set equipped with a metric is called a *metric space*.

Let V be a normed vector space. Then a metric can be induced by the norm as

$$d(\mathbf{u}, \mathbf{v}) = \|\mathbf{u} - \mathbf{v}\|. \tag{2.40}$$

In particular, if the norm is induced by an inner product, then

$$d(\mathbf{u}, \mathbf{v}) = \sqrt{\langle \mathbf{u} - \mathbf{v}, \mathbf{u} - \mathbf{v} \rangle}. \tag{2.41}$$

Example 2.1.18 (Euclidean distance on $\mathbb{R}^n$)
For $\mathbf{x}, \mathbf{y} \in \mathbb{R}^n$, the Euclidean distance is

$$d_2(\mathbf{x}, \mathbf{y}) = \|\mathbf{x} - \mathbf{y}\|_2 = \sqrt{\sum_{i=1}^{n} (x_i - y_i)^2}. \tag{2.42}$$

For example, if $\mathbf{x} = (1, 1)$ and $\mathbf{y} = (4, 3)$, then

$$d_2(\mathbf{x}, \mathbf{y}) = \sqrt{(4-1)^2 + (3-1)^2}$$
$$= \sqrt{13}.$$

Definition 2.1.19 (Neighbourhood and ball)**.** Let d be a metric on $\mathbb{R}^n$. For $\mathbf{r}_0 \in \mathbb{R}^n$ and $a > 0$, the *open ball* of radius a centred at $\mathbf{r}_0$ is defined by


$$B_a(\mathbf{r}_0) = \{\mathbf{r} \in \mathbb{R}^n \mid d(\mathbf{r}, \mathbf{r}_0) < a\}. \tag{2.43}$$

It is also called the *a-neighbourhood* of $\mathbf{r}_0$.

The *sphere* of radius a centred at $\mathbf{r}_0$ is defined by

$$S_a(\mathbf{r}_0) = \{\mathbf{r} \in \mathbb{R}^n \mid d(\mathbf{r}, \mathbf{r}_0) = a\}. \tag{2.44}$$

In $\mathbb{R}^2$ with the Euclidean metric, the open ball consists of all points inside a circle, while the sphere is the circle itself. If $\mathbf{u} \in B_a(\mathbf{r}_0)$, then $d(\mathbf{u}, \mathbf{r}_0) < a$. If $\mathbf{r} \in S_a(\mathbf{r}_0)$, then $d(\mathbf{r}, \mathbf{r}_0) = a$.



3 Lecture 03

Date: 13/10/2026

§3.1 Review on linear algebra—continued

References

[**Axl24**] Sheldon Axler. *Linear Algebra Done Right*. 4th. Undergraduate Texts in Mathematics. Open Access. Springer, 2024. doi: 10.1007/978-3-031-41026-0. url: https://linear.axler.net/ (cited p. 10)

[**KW19**] Mykel J. Kochenderfer and Tim A. Wheeler. *Algorithms for Optimization*. Cambridge, MA: The MIT Press, 2019. isbn: 978-0-262-03942-0. url: https://algorithmsbook.com/optimization/ (cited p. 2)