

Your Paper Title Here

Your Name(s)

Date

Abstract

Write your abstract here.

Keywords

Keyword 1, Keyword 2, Keyword 3, Keyword 4

Introduction / Background

Provide an introduction with background information.

Statistical Methods

Model

We define V as the proportion of people who were infected with COVID-19 that received the vaccine. n is the total number of test subjects that were infected (170 in our case) and π_0 is the true but unknown probability of being vaccinated and still getting infected.

This true probability has been calculated in terms of two other probabilities, π_v and π_p , which are the probabilities that the subject received the vaccine and a placebo respectively:

$$\pi = \frac{\pi_v}{\pi_v + \pi_p}$$

What we are interested in is the efficacy of the vaccine, ψ . This efficacy, according to the study, is computed with the following formula:

$$\psi = \frac{1 - 2\pi}{1 - \pi}$$

We can transform this variable in terms of ψ as well (which is more helpful for our inferences):

$$\pi = \frac{1 - \psi}{2 - \psi}$$

Likelihood Inference

Likelihood function and MLE

In this section, we take the likelihood approach and break down the problem through a frequentist's lens. We begin by approximating V , the proportion of subjects infected with COVID-19 that were

vaccinated, to the binomial distribution (proof in appendix):

$$V \approx \text{Binom}(n, \pi_0)$$

In this case, we know $n = 170$ and observe $v = 8$. In our lectures, we proved that the MLE of a binomial distribution is the proportion of successes to the number of trials. Hence,

$$\hat{\pi}_0^{mle} = \frac{8}{170}$$

Then, with our formula for ψ ,

$$\hat{\psi}_0^{mle} = \frac{1 - 2(\frac{8}{170})}{1 - (\frac{8}{170})} \approx 0.9506$$

Let us verify this with the transformed likelihood function, $L^*(\psi)$. We begin with the binomial likelihood function, and transform it to be in terms of ψ :

$$\begin{aligned} L(\pi) &= \binom{n}{v} \pi^v (1 - \pi)^{n-v} \\ L^*(\psi) &= \binom{n}{v} \left(\frac{1 - \psi}{2 - \psi}\right)^v \left(1 - \frac{1 - \psi}{2 - \psi}\right)^{n-v} \\ L^*(\psi) &= \binom{n}{v} \left(\frac{1 - \psi}{2 - \psi}\right)^v \left(\frac{1}{2 - \psi}\right)^{n-v} \\ L^*(\psi) &= \binom{n}{v} \frac{(1 - \psi)^v}{(2 - \psi)^n} \\ \ell^*(\psi) &= \ln\left(\binom{n}{v}\right) + v \ln(1 - \psi) - n \ln(2 - \psi) \\ \frac{d}{d\psi} \ell^*(\psi) &= -\frac{v}{1 - \psi} + \frac{n}{2 - \psi} \end{aligned}$$

The above gives us the log-likelihood function $\ell^*(\psi)$ and its derivative. We find a critical point by equating the derivative to zero, but we would still need to verify it to be a maximum (and thus,

our MLE).

Second Derivative test

This is where the second derivative test comes in. We check the value of the second derivative at the critical point to identify the concavity, and whether or not we found a maximum:


$$\frac{d^2}{d\psi^2}\ell^*(\psi) = -\frac{v}{(1-\psi)^2} + \frac{n}{(2-\psi)^2}$$

If the above second derivative is less than 0 at our critical point (or for all values of ψ), then it is concave down and we have verified it to be our MLE.

In that case, we have our MLE: $\hat{\psi}_0^{mle}$

Large Sample Confidence Interval

The next step of our likelihood inference is finding the confidence interval, for which we take the large sample approach. Before we do so, we plot the second-order Taylor approximation against the log-likelihood to ensure our approximations hold:



We see the second-order Taylor approximation is very accurate, entirely aligned with the log-likelihood. Thus, we move forward with the large sample confidence interval. To do so, we need to calculate our observed Fisher Information:

$$\begin{aligned}
I(\psi) &= -E \left[\frac{d^2}{d\psi^2} \ell^*(\psi) \right] \\
I(\psi) &= E \left[\frac{v}{(1-\psi)^2} - \frac{n}{(2-\psi)^2} \right] \\
I(\psi) &= \frac{E[v]}{(1-\psi)^2} - \frac{n}{(2-\psi)^2}
\end{aligned}$$

Note that: $E[v] \approx n * \pi$

$$\begin{aligned}
I(\psi) &= \frac{n * \pi}{(1-\psi)^2} - \frac{n}{(2-\psi)^2} \\
I(\psi) &= \frac{n - n\psi}{(2-\psi)(1-\psi)^2} - \frac{n}{(2-\psi)^2} \\
I(\psi) &= \frac{(n - n\psi)(2-\psi) - n(1-\psi)^2}{(1-\psi)^2(2-\psi)^2} \\
I_{obs}(\hat{\psi}_0^{mle}) &= \frac{(n - n\hat{\psi}_0^{mle})(2 - \hat{\psi}_0^{mle}) - n(1 - \hat{\psi}_0^{mle})^2}{(1 - \hat{\psi}_0^{mle})^2(2 - \hat{\psi}_0^{mle})^2}
\end{aligned}$$

We then compute the 95% large sample confidence interval with the formula:

$$\hat{\psi}_0^{mle} \pm z_{\alpha/2} \sqrt{\frac{1}{I(\hat{\psi}_0^{mle})}}$$

Significance test: Likelihood Ratio Statistic

We conclude our likelihood inference with a significance test. We want to find the likelihood ratio statistic W to test whether or not our efficacy rate passes the FDA's standard. We define our hypotheses as follows:

$$H_0 : \psi_0 = 0.3 \quad H_1 : \psi_0 \neq 0.3$$

We know $W = 2 \ln(\Lambda)$, where $\Lambda = \frac{L(\hat{\psi}_0^{mle})}{L(\psi_0^{null})}$

We compute as follows:

$$\begin{aligned}
W &= 2 \ln \left(\frac{L(\hat{\psi}_0^{mle})}{L(\psi_0^{null})} \right) \\
W &= 2 \ln \left(\frac{\binom{n}{v} \frac{(1-\hat{\psi}_0^{mle})^v}{(2-\hat{\psi}_0^{mle})^n}}{\binom{n}{v} \frac{(1-\psi_0^{null})^v}{(2-\psi_0^{null})^n}} \right) \\
W &= 2 \ln \left(\frac{(1-\hat{\psi}_0^{mle})^v (2-\psi_0^{null})^n}{(2-\hat{\psi}_0^{mle})^n (1-\psi_0^{null})^v} \right) \\
W &= 2(v \ln(1-\hat{\psi}_0^{mle}) + n \ln(2-\psi_0^{null}) - n \ln(2-\hat{\psi}_0^{mle}) - v \ln(1-\psi_0^{null}))
\end{aligned}$$

We then take the observed w and compare it against the chi-squared distribution of W . That is, our p-value will be $P(W \geq w)$, where $W \sim \chi_1^2$

This concludes our likelihood inference approach, and the methods we will employ to make our final conclusion. The results of these procedures will be presented under the Results section.

Bayesian Inference

Detail the Bayesian approach.

Results

Present your findings.

Likelihood Inference Results

Utilizing the log likelihood functions and its derivatives, we identify and critical point and verify it to be the MLE. Unsurprisingly, we find it to be equivalent to the estimate we computed at the beginning using the transformation of the binomial MLE. That is, we find $\hat{\psi}_0^{mle} \approx 0.9506$.

With the observed Fisher Information, we find our large sample confidence interval to be [0.9155,

0.9857]. On the other hand, the parametric bootstrap returns a confidence interval to be [placeholder]

The significance test returns $2.8223132 \times 10^{-28}$, which is [comparable/different] from our empirical p-value, [placeholder].

Discussion / Conclusion

Discuss / conclude here.

Bibliography

Brown, B. (2024). *Lecture Title*. Lecture slides, Course Name, University Name.

Doe, J. (2020). Title of the Paper. *Journal Name*, 12(3), 45-67.

Last, F., & Last, F. (2025). *Book Title*. Publisher.

Smith, A., & Johnson, C. (2023). *Title of the Online Article*. Retrieved from <https://www.example.com>.

Appendix

Code

```
# You can reference your code in the appendix (sample here).
```

Plot second-order taylor approximation against log-likelihood

```
loglik_psi <- function(psi, n, v){  
  c <- choose(n, v)  
  return(log(c) + (v * log(1 - psi)) - (n * log(2 - psi)))  
}
```

```

}

ml_psi <- maxLik2(loglik = loglik_psi,
                  start = 0.9, n = 170, v = 8)

plot(ml_psi) +
  labs(title = "Second order approximation to the psi likelihood",
       x = expression(psi))

```

Compute large sample confidence interval for the MLE

```

z <- qnorm(0.975)
I <- (8/(1 - 0.9506)^2) - (170/(2-0.9506)^2)
upper <- 0.9506 + z * (1/sqrt(I))
lower <- 0.9506 - z * (1/sqrt(I))

```

Compute pvalue using likelihood ratio statistic

```

pval <- pchisq(121.6012, df = 1, lower.tail = F)

```

Proofs

If applicable, include detailed mathematical derivations or additional theoretical explanations.