

Frequentist and Bayesian Reanalysis of the Pfizer BNT162b2 Vaccine Trial

Blake Xu, Kevin Ji, Sangey Rinchen

03/14/2026

Abstract

COVID-19 caused major global health problems and created an urgent need for effective vaccines. In this study, we reviewed the Pfizer–BioNTech BNT162b2 vaccine trial reported by Polack et al. and reanalyze its efficacy using both frequentist and Bayesian methods. Using the observed data of 8 infections in the vaccine group and 162 in the placebo group, we estimate the vaccine efficacy parameter and evaluate whether it exceeds the FDA benchmark of 30

Keywords

Bayesian Inference, Likelihood Inference, COVID-19, Vaccine efficacy

Introduction

Beginning in 2019, the COVID-19 pandemic has had a profound impact on global public health. It has been estimated that approximately 18.2 million people died worldwide as a result of the pandemic (Wang, Haidong et al. 2020-21). Vaccination therefore became a critical strategy for reducing infection and mortality. There are studies showing that “Authorized mRNA COVID-19 vaccines are effective for preventing SARS-CoV-2 infection in real-world conditions.” and COVID-19 vaccination is recommended for all eligible persons.”(Thompson et al., 2021). One of the most widely used vaccines is the BNT162b2 mRNA vaccine developed by Pfizer and BioNTech.

Vaccine efficacy measures the proportional reduction in infection risk among vaccinated individuals relative to those receiving placebo. The U.S. Food and Drug Administration (FDA) recommended that COVID-19 vaccines demonstrate at least 30% efficacy to be considered effective. The efficacy of this vaccine was evaluated in a large randomized, placebo-controlled, double-blind clinical trial (Polack et al., 2020), which randomly assigned participants aged 16 years or older in a 1:1 ratio to receive two doses, 21 days apart, of either placebo or the BNT162b2 vaccine. The primary efficacy

endpoint was the occurrence of laboratory-confirmed COVID-19 infection. A total of 170 infections were targeted(Polack et al., 2020).

Transparent, corroborated methodology is vital to global perceptions of public health institutions. The mixed messaging during the pandemic led to a deterioration in trust of public health (Ortiz-Prado et al., 2025). In our report, we aim to corroborate and validate Pfizer’s analysis using both Frequentist and Bayesian statistical approaches. The Pfizer vaccine trial was notable in its use of Bayesian methods, whereas other firms used frequentist experimental design. Pfizer’s study has been criticized for lacking explanation of study assumptions (most notably the choice of a prior) and pandemic vaccine trials have been criticized for a lack of data transparency (Senn S., 2022). To provide an independent statistical assessment of Pfizer’s BNT162b2 vaccine efficacy, we estimate the vaccine efficacy parameter and test whether the efficacy exceeds the FDA guideline of 30%.

Table 1: vaccine efficacy at least 7 days after dose two

Group	cases	N	rate
BNT162b2	8	17411	0.00046
Placebo	162	17511	0.00925




Figure 1: Observed Infection Rates by Group

Statistical Methods

Model

We define V as the number of people who were infected with COVID-19 that received the vaccine. n is the total number of test subjects that were infected (170 in our case) and π_0 is the true but unknown probability of being vaccinated given that they are infected.

This true probability has been calculated in terms of two other probabilities, π_{v0} and π_{p0} , which are the proportions of COVID infections, for the vaccine and placebo groups respectively, and since the trial is approximately 1:1 randomized, we have

$$\pi_0 = \frac{\pi_{v0}}{\pi_{v0} + \pi_{p0}}$$

What we are interested in is the efficacy of the vaccine, ψ_0 . The estimate of this efficacy, ψ , is computed as the inverse of the estimated relative risk. That is:

$$\psi = 1 - \frac{\pi_v}{\pi_p} = \frac{1 - 2\pi}{1 - \pi}$$

We can transform this variable in terms of ψ (which is more helpful for our inferences):

We can use this to transform π in terms of ψ (which is more helpful for our inferences):

$$\pi = \frac{1 - \psi}{2 - \psi}$$

Likelihood Inference

Likelihood function and MLE

In this section, we take the likelihood approach and break down the problem through a frequentist's lens. We begin by approximating V , the number of subjects infected with COVID-19 that were vaccinated, to the binomial distribution (proof in appendix):

$$V \approx \text{Binom}(n, \pi_0)$$

In this case, from Table 1, there were $n = 170$ and observe $v = 8$. In our lectures, we proved that the MLE of a binomial distribution is the number of successes to the number of trials (Grove, 2026, Lecture 11). Hence,

$$\hat{\pi}_0^{mle} = \frac{8}{170}$$

Then, with our formula for ψ ,

$$\hat{\psi}_0^{mle} = \frac{1 - 2(\frac{8}{170})}{1 - (\frac{8}{170})} \approx 0.9506$$

Let us verify this with the transformed likelihood function, $L^*(\psi)$. We begin with the binomial likelihood function, and transform it to be in terms of ψ :

$$\begin{aligned} L(\pi) &= \binom{n}{v} \pi^v (1 - \pi)^{n-v} \\ L^*(\psi) &= \binom{n}{v} \left(\frac{1 - \psi}{2 - \psi}\right)^v \left(1 - \frac{1 - \psi}{2 - \psi}\right)^{n-v} \\ L^*(\psi) &= \binom{n}{v} \left(\frac{1 - \psi}{2 - \psi}\right)^v \left(\frac{1}{2 - \psi}\right)^{n-v} \\ L^*(\psi) &= \binom{n}{v} \frac{(1 - \psi)^v}{(2 - \psi)^n} \\ \ell^*(\psi) &= \ln\left(\binom{n}{v}\right) + v \ln(1 - \psi) - n \ln(2 - \psi) \\ \frac{d}{d\psi} \ell^*(\psi) &= -\frac{v}{1 - \psi} + \frac{n}{2 - \psi} \end{aligned}$$

The above gives us the log-likelihood function $\ell^*(\psi)$ and its derivative. We find a critical point by equating the derivative to zero, but we would still need to verify it to be a maximum (and thus, our MLE).

Second Derivative test

This is where the second derivative test comes in. We check the value of the second derivative at the critical point to identify the concavity, and whether or not we found a maximum:

$$\frac{d^2}{d\psi^2} \ell^*(\psi) = -\frac{v}{(1 - \psi)^2} + \frac{n}{(2 - \psi)^2}$$

If the above second derivative is less than 0 at our critical point (or for all values of ψ), then it is concave down and we have verified it to be our MLE, $\hat{\psi}_0^{mle}$

Large Sample Confidence Interval

The next step of our likelihood inference is finding the confidence interval, for which we take the large sample approach. Before we do so, we plot the second-order Taylor approximation against the log-likelihood of π to ensure our approximations hold:




Figure 2: Log-likelihood function and second-order approximation for π .

We see the second-order Taylor approximation is accurate, mostly aligned with the log-likelihood. Thus, we move forward with the large sample confidence interval. To do so, we need to calculate our observed Fisher Information:

$$I(\psi) = -E \left[\frac{d^2}{d\psi^2} \ell^*(\psi) \right]$$

$$I(\psi) = E \left[\frac{v}{(1-\psi)^2} - \frac{n}{(2-\psi)^2} \right]$$

$$I(\psi) = \frac{E[v]}{(1-\psi)^2} - \frac{n}{(2-\psi)^2}$$

Note that: $E[v] \approx n * \pi$

$$I(\psi) = \frac{n * \pi}{(1-\psi)^2} - \frac{n}{(2-\psi)^2}$$

$$I_{obs}(\hat{\psi}_0^{mle}) = \frac{(n - n\hat{\psi}_0^{mle})(2 - \hat{\psi}_0^{mle}) - n(1 - \hat{\psi}_0^{mle})^2}{(1 - \hat{\psi}_0^{mle})^2(2 - \hat{\psi}_0^{mle})^2}$$

We then compute the 95% large sample confidence interval with the formula:

$$\hat{\psi}_0^{mle} \pm z_{\alpha/2} \sqrt{\frac{1}{I(\hat{\psi}_0^{mle})}}$$

Significance test: Likelihood Ratio Statistic

We conclude our likelihood inference with a significance test. We want to find the likelihood ratio statistic W to test whether or not our efficacy rate passes the FDA's standard. We define our hypotheses as follows:

$$H_0 : \psi_0 = 0.3 \quad H_1 : \psi_0 \neq 0.3$$

We know $W = 2 \ln(\Lambda)$, where $\Lambda = \frac{L(\hat{\psi}_0^{mle})}{L(\hat{\psi}_0^{null})}$

$$\begin{aligned} W &= 2 \ln\left(\frac{L(\hat{\psi}_0^{mle})}{L(\hat{\psi}_0^{null})}\right) \\ W &= 2 \ln\left(\frac{\binom{n}{v} \frac{(1-\hat{\psi}_0^{mle})^v}{(2-\hat{\psi}_0^{mle})^n}}{\binom{n}{v} \frac{(1-\hat{\psi}_0^{null})^v}{(2-\hat{\psi}_0^{null})^n}}\right) \\ W &= 2 \ln\left(\frac{(1-\hat{\psi}_0^{mle})^v (2-\hat{\psi}_0^{null})^n}{(2-\hat{\psi}_0^{mle})^n (1-\hat{\psi}_0^{null})^v}\right) \\ W &= 2(v \ln(1-\hat{\psi}_0^{mle}) + n \ln(2-\hat{\psi}_0^{null}) - n \ln(2-\hat{\psi}_0^{mle}) - v \ln(1-\hat{\psi}_0^{null})) \end{aligned}$$

We then take the observed w and compare it against the chi-squared distribution of W . That is, our p-value will be $P(W \geq w)$, where $W \sim \chi_1^2$. At the same time, we conduct a parametric bootstrap to find a bootstrapped confidence interval and empirical p-value, for a more robust analysis into our MLE.

This concludes our the methods of our likelihood inference approach, and the results of these procedures will be presented under the Results section.

Bayesian Inference

In this section we describe the Bayesian approach used to estimate the vaccine efficacy parameter ψ , where $-\infty < \psi \leq 1$.

To complete the Bayesian model, we specify a Beta prior for π :

$$\pi \sim \text{Beta}(a, b).$$

The Beta family is a natural prior for a binomial parameter because it is supported on $[0, 1]$ and is conjugate to the binomial likelihood.

We chose the prior through elicitation on the vaccine efficacy parameter ψ rather than directly on π . Our goal was to use a deliberately skeptical prior that would require the observed data to provide strong evidence before concluding that the vaccine was effective. This choice is reasonable because, at the design stage of a clinical trial, substantial uncertainty remains about whether a candidate vaccine will exceed the FDA efficacy benchmark of 30%.

Specifically, we imposed the following prior beliefs:

1. $P(\Psi \geq 0.3) = 0.1$,
2. $P(\Psi \geq 0) = 0.5$.

The first statement assigns only a 10% prior probability to the event that vaccine efficacy exceeds the FDA benchmark, which shows skepticism about strong efficacy before seeing the trial data. The second statement centers prior belief at the boundary between harmful and beneficial effects by assigning probability 0.5 to $\psi \geq 0$.

Using the transformation

$$\psi = \frac{1 - 2\pi}{1 - \pi} \iff \pi = \frac{\psi - 1}{\psi - 2} \quad (1)$$

these prior beliefs become

$$P\left(\Pi \leq \frac{7}{17}\right) = 0.1, \quad P\left(\Pi \leq \frac{1}{2}\right) = 0.5.$$

Solving these equations numerically gives

$$\Pi \sim \text{Beta}(26.2, 26.2).$$

Because the Beta prior is conjugate to the Binomial likelihood, by theorem 13.2 (Grove, 2026, Lecture 13), the posterior distribution of Π is

$$\Pi \mid V = v \sim \text{Beta}(26.2 + 8, 26.2 + 162) = \text{Beta}(34.2, 188.2).$$

$$\begin{aligned} P(\Psi \leq \psi | v) &= P\left(\frac{1 - 2\pi}{1 - \pi} \leq \psi | v\right) \quad -\infty < \psi \leq 1 \\ &= 1 - P\left(\pi \leq \frac{\psi - 1}{\psi - 2} | v\right) \\ F_{\Psi}(\psi | v) &= 1 - F_{\pi}\left(\frac{\psi - 1}{\psi - 2} | v\right) \end{aligned}$$

Differentiating gives the posterior density of ψ :

$$f_{\Psi}(\psi | v) = (1 - F_{\pi}(\frac{\psi - 1}{\psi - 2} | v))' \left(\frac{\psi - 1}{\psi - 2}\right)' \quad (1)$$

$$f_{\Psi}(\psi | v) = f_{\pi}(\frac{\psi - 1}{\psi - 2} | v) \frac{1}{(\psi - 2)^2} \quad (2)$$

$$(3)$$

Substituting the Beta(34.2, 188.2) density gives

$$f_{\Psi|V}(\psi | v) = \frac{(1 - \psi)^{33.2}}{B(34.2, 188.2)(2 - \psi)^{222.4}}, \quad -\infty < \psi \leq 1.$$

Similarly, the prior density of ψ is

$$f_{\Psi}(\psi) = \frac{(1 - \psi)^{25.2}}{B(26.2, 26.2)(2 - \psi)^{52.4}}, \quad -\infty < \psi \leq 1.$$

The prior and posterior densities (as well as the Pfizer study's prior distribution for comparison) for ψ are shown below.




Figure 3: Prior and posterior densities for Psi.

To obtain the posterior distribution of ψ , we use the transformation (1)

For $-\infty < \psi \leq 1$, the cumulative distribution function of Ψ is

$$\begin{aligned}
 P(\Psi \leq \psi | V = v) &= P\left(\frac{1 - 2\Pi}{1 - \Pi} \leq \psi | V = v\right) \quad -\infty < \psi \leq 1 \\
 &= 1 - P\left(\Pi \leq \frac{\psi - 1}{\psi - 2} | V = v\right) \\
 F_{\Psi|V}(\psi|v) &= 1 - F_{\Pi|V}\left(\frac{\psi - 1}{\psi - 2} | v\right)
 \end{aligned}$$

Differentiating gives the posterior density of ψ :

$$\begin{aligned}
 f_{\Psi|V}(\psi|v) &= (1 - F_{\Pi|V}(\frac{\psi - 1}{\psi - 2} | v))' \left(\frac{\psi - 1}{\psi - 2}\right)' \\
 f_{\Psi|V}(\psi|v) &= f_{\Pi|V}(\frac{\psi - 1}{\psi - 2} | v) \frac{1}{(\psi - 2)^2} = \frac{(1 - \psi)^{a-1}}{B(a, b)(2 - \psi)^{a+b}}
 \end{aligned}$$

A similar transformation can be used to obtain the 95% credible interval for ψ . Let $q_{\psi, 0.025}$ and $q_{\psi, 0.975}$ denote the 2.5th and 97.5th percentiles of the posterior distribution of ψ . The corresponding endpoints are

Substituting the Beta(34.2, 188.2) density gives

$$f_{\Psi|V}(\psi | v) = \frac{(1 - \psi)^{33.2}}{B(34.2, 188.2)(2 - \psi)^{222.4}}, \quad -\infty < \psi \leq 1.$$

$$P(\psi \leq 0.3 \mid v),$$

which can be evaluated either directly from the posterior density of ψ or by transforming the equivalent event on π .

Minding the asymptote, the plot suggests that Pfizer's prior of $\Pi \sim B(0.700102, 1)$ is optimistic and assumes efficacy.

Posterior summaries of Ψ can also be obtained from the posterior distribution of π using the transformation $\psi = \frac{1-2\pi}{1-\pi}$.

The posterior median of Ψ is defined by the value $q_{\Psi,0.5}$ satisfying

$$P(\Psi \leq q_{0.5} \mid V = v_{obs}) = 0.5.$$

Using the transformation between ψ and π ,

$$\begin{aligned} P(\Psi \leq q_{\Psi,0.5} \mid V = v_{obs}) &= P\left(\frac{1-2\Pi}{1-\Pi} \leq q_{\Psi,0.5} \mid V = v_{obs}\right) \\ &= P(\Pi(q_{\Psi,0.5} - 2) \leq q_{\Psi,0.5} - 1 \mid V = v_{obs}) \\ &= P\left(\Pi \geq \frac{q_{\Psi,0.5} - 1}{q_{\Psi,0.5} - 2} \mid V = v_{obs}\right) = 0.5 \\ P\left(\Pi \leq \frac{q_{\Psi,0.5} - 1}{q_{\Psi,0.5} - 2} \mid V = v_{obs}\right) &= 1 - 0.5 \end{aligned}$$

A similar transformation can be used to obtain the 95% credible interval for Ψ . Let $q_{\Psi,0.025}$ and $q_{\Psi,0.975}$ denote the 2.5th and 97.5th percentiles of the posterior distribution of Ψ . The corresponding endpoints are

$$q_{\Psi,0.025} = \frac{1 - 2q_{\Pi,1-0.025}}{1 - q_{\Pi,1-0.025}}, \quad q_{\Psi,0.975} = \frac{1 - 2q_{\Pi,1-0.975}}{1 - q_{\Pi,1-0.975}}.$$

To assess whether the vaccine meets the FDA efficacy guideline, we compute the posterior probability

$$P(\Psi \leq 0.3 \mid V = v_{obs}),$$

which can be evaluated either directly from the posterior density of Ψ or by transforming the equivalent event on π .

Results

With the observed Fisher Information, we find our large sample confidence interval to be [0.9155, 0.9857]. On the other hand, the parametric bootstrap returns a confidence interval of [0.9102564, 0.9820359].

The significance test returns $2.8223132 \times 10^{-28}$, which is comparable to our empirical p-value, 10^{-5} . Since none of the bootstrapped values were greater than our observed likelihood ratio statistic, W , our empirical P-value was calculated as $\frac{1}{\text{bootstrap sample size}+1}$.

Bayesian Inference Results

Using the posterior distribution of $\Pi \mid V = v \sim \text{Beta}(34.2, 188.2)$ and the transformation discussed in the Method,

$$q_{\Psi,0.025} = \frac{1 - 2q_{\Pi,0.975}}{1 - q_{\Pi,0.975}}, \quad q_{\Psi,0.5} = \frac{1 - 2q_{\Pi,0.5}}{1 - q_{\Pi,0.5}}, \quad q_{\Psi,0.975} = \frac{1 - 2q_{\Pi,0.025}}{1 - q_{\Pi,0.025}}$$

we obtained a posterior median of 0.82, and a 95% credible interval of (0.744, 0.877).

We can also use the posterior density derived in the previous section and numerical integration to compute summary statistics for the posterior distribution of the vaccine efficacy parameter Ψ . In particular, the posterior mean, posterior median, and a 95% Bayesian credible interval were obtained.

The posterior summaries are reported in Table 2.

Table 2: Posterior summary statistics for Psi.

Mean	Median	Lower.Bound	Upper.Bound
0.817	0.82	0.744	0.877

Calculate a Bayesian P-value

To assess whether the vaccine meets the FDA efficacy guideline, we consider the hypothesis test

$$H_0 : \Psi = 0.3 \quad H_1 : \Psi > 0.3$$

Using the posterior distribution of Ψ , we compute the posterior probability

$$P(\Psi \leq 0.3 \mid V = v_{obs}),$$

which serves as the Bayesian p-value for this one-sided test.

The posterior probability is

$$P(\Psi \leq 0.3 \mid V = v_{obs}) = 2.959 \cdot 10^{-17}$$

Since the posterior probability that $\Psi \leq 0.3$ is essentially zero, we can reject the null hypothesis. That is, the vaccine efficacy is greater than 30% and therefore does meet the FDA criteria.

Discussion / Conclusion

Table 3: ‘95% CI for Psi’

	Lower Bound	Upper Bound
MLE Confidence Interval	0.9155326	0.9856674
Our Credible Interval	0.7438560	0.8769644
Pfizer’s Credible Interval	0.9030000	0.9760000

Both our frequentist and Bayesian analysis corroborate Pfizer’s finding that the BNT162b2 vaccine was 95% effective in preventing COVID-19. Similar to Pfizer’s original report, our significance tests for both Bayesian and frequentist estimates of ψ were very statistically significant, with a p-value well below 0.01 or any other reasonable rejection level. This means we should reject the null hypothesis that the vaccine efficacy does not exceed the FDA guidelines and it is highly probable that the vaccine efficacy is above FDA minimum guidelines. Our empirical tests for both Bayesian and frequentist methods also support the rejection of the null hypothesis.

While all three tests validate that the BNT162b2 vaccine was more efficacious than FDA guidelines ($\psi > 0.3$) at a 95% CI, our results show that Pfizer overestimated the efficacy of their vaccine based on Bayesian methods. The confidence interval for ψ^{mle} is similar to Pfizer’s credible interval for ψ , our posterior lower bound was ~15% smaller and our posterior upperbound was ~10% smaller than Pfizer’s credible interval. This overestimation mainly comes from the difference in choosing the prior. While Pfizer themselves state their prior of $\psi = 0.3$ “can be considered pessimistic”, this makes the assumption that the vaccine is more efficacious before seeing any trial (BioNTech/Pfizer, 2020). They assumed π is centered at 0.4118, which is lower than 0.5. A prior π of 0.5 assumes that the efficacy of a placebo and the vaccine are the same. Our prior does not make this assumption, as the most rigorous standard for clinical trials is to assume no efficacy from any given medical intervention.

In the future, large pharmaceutical companies that use Bayesian methods should be more rigorous and transparent about their selection of a prior. We have demonstrated that through selecting a more rigorous prior, prior selection can impact the magnitude of final estimates significantly- even if the treatment is highly effective. Without fully transparent methodology, fellow researchers and the general can become skeptical of one’s findings which can create material harms in public health.

References

- Grove, R. (2026). *Chapter 13.1: Bayesian Inference*. Lecture 13, Slide 30, STAT 342, University of Washington - Seattle.
- Grove, R. (2025). *Chapter 3*. Lecture 3, Slide 7, STAT 341, University of Washington - Seattle.
- Grove, R. (2026). *Chapter 11: Maximum Likelihood Estimation*. Lecture 11, Slide 33, STAT 342, University of Washington - Seattle.
- Grove, R. (2026). *Chapter 13.1: Bayesian Inference*. Lecture 13, Slide 30, STAT 342, University of Washington - Seattle.
- Polack, F. P., Thomas, S. J., Kitchin, N., Absalon, J., Gurtman, A., Lockhart, S., Perez, J. L., et al. (2020). *Safety and efficacy of the BNT162b2 mRNA Covid-19 vaccine*. **New England Journal of Medicine**, 383(27), 2603–2615. <https://doi.org/10.1056/NEJMoa2034577>
- Thompson, M. G., Burgess, J. L., Naleway, A. L., etour empirical p-value, al. (2021). *Interim estimates of vaccine effectiveness of BNT162b2 and mRNA-1273 COVID-19 vaccines in preventing SARS-CoV-2 infection among health care personnel, first responders, and other essential and frontline workers — Eight U.S. locations, December 2020–March 2021*. *MMWR Morbidity and Mortality Weekly Report*, 70, 495–500.
<http://dx.doi.org/10.15585/mmwr.mm7013e3>
- Wang, Haidong, et al. (2022). *Estimating excess mortality due to the COVID-19 pandemic: A systematic analysis of COVID-19-related mortality, 2020–21*. *The Lancet*, 399(10334), 1513–1536. [https://www.thelancet.com/journals/lancet/article/PIIS0140-6736\(21\)02796-3/fulltext](https://www.thelancet.com/journals/lancet/article/PIIS0140-6736(21)02796-3/fulltext)
- Polack, F. P., Thomas, S. J., Kitchin, N., Absalon, J., Gurtman, A., Lockhart, S., Perez, J. L., et al. (2020). *Safety and efficacy of the BNT162b2 mRNA Covid-19 vaccine*. **New England Journal of Medicine**, 383(27), 2603–2615. <https://doi.org/10.1056/NEJMoa2034577>
- Thompson, M. G., Burgess, J. L., Naleway, A. L., etour empirical p-value, al. (2021). *Interim estimates of vaccine effectiveness of BNT162b2 and mRNA-1273 COVID-19 vaccines in preventing SARS-CoV-2 infection among health care personnel, first responders, and other essential and frontline workers — Eight U.S. locations, December 2020–March 2021*. *MMWR Morbidity and*

Mortality Weekly Report, 70, 495–500.

<http://dx.doi.org/10.15585/mmwr.mm7013e3>

Wang, Haidong, et al. (2022). *Estimating excess mortality due to the COVID-19 pandemic: A systematic analysis of COVID-19-related mortality, 2020–21*. The Lancet, 399(10334), 1513–1536.

[https://www.thelancet.com/journals/lancet/article/PIIS0140-6736\(21\)02796-3/fulltext](https://www.thelancet.com/journals/lancet/article/PIIS0140-6736(21)02796-3/fulltext)

Appendix

Code

Setup

```
#Use this code chunk to include libraries, and set global options.
library(ggplot2)
library(tidyverse)
library(fastR2)
```

Data visualization

```
df <- data.frame(
  Group = c('BNT162b2', 'Placebo'),
  cases = c(8, 162),
  N = c(17411, 17511),
  rate = c(8/17411, 162/17511)
)
knitr::kable(df, digits = 5, caption = "vaccine efficacy at least 7 days after dose two"
)
ggplot(df, aes(x = Group, y = rate, fill = Group))+
  geom_col()+ scale_fill_manual(
    values = c('BNT162b2' = 'steelblue', 'Placebo' = 'firebrick')
  )+labs(x = "Group",y = "Infection Rate",title = "Observed Infection Rates by Group",
    caption = "Figure 1: Observed Infection Rates by Group") +
  theme_classic()+ theme(legend.position = "none", plot.caption = element_text(hjust = 0.5))
```

Plot second-order taylor approximation against log-likelihood

```
loglik_pi <- function(pi, n, v){
  c <- choose(n, v)
  return(log(c) + (v * log(pi)) + ((n-v) * log(1 - pi)))
}
ml_pi <- maxLik2(loglik = loglik_pi,
  start = 0.5, n = 170, v = 8)
plot(ml_pi) +
  labs(title = "Second order approximation to the pi likelihood",
```

```
x = expression(pi), caption = "Figure 2: Log-likelihood function and second-order approximat
")+
theme_classic()+
theme(plot.caption = element_text(hjust = 0.5))
```

Compute large sample confidence interval for the MLE

```
z <- qnorm(0.975)
I <- (8/(1 - 0.9506)^2) - (170/(2-0.9506)^2)
upper <- 0.9506 + z * (1/sqrt(I))
lower <- 0.9506 - z * (1/sqrt(I))
```

Compute pvalue using likelihood ratio statistic

```
pval <- pchisq(121.6012, df = 1, lower.tail = F)
```

Parametric Bootstrap for psi

Find Prior

```
library(LearnBayes)
beta.select(quantile1 = list(p = 0.1, x = 7/17),
            quantile2 = list(p = 0.5, x = 0.5))
```

Compute statistics for psi

```
post_psi <- function(x) {(1-x)^33.2/(beta(34.2, 188.2)*(2-x)^222.4)}
exp_post_psi<- function(x) {x*(1-x)^33.2/(beta(34.2, 188.2)*(2-x)^222.4)}
find_quantile <- function(p){
  uniroot(f = function(q) {
    integrate(post_psi, 0, q)$value - p},
    interval = c(0,1))$root}
q_025 <- find_quantile(0.025)
q_median <- find_quantile(0.5)
q_975<- find_quantile(0.975)
psi_mean <- integrate(exp_post_psi, 0, 1)$value
result_table <- data.frame(Mean = psi_mean, Median = q_median,
                          'Lower Bound' = q_025, 'Upper Bound' = q_975)
knitr::kable(result_table, digits = 3, caption = "Posterior summary statistics for Psi.")
```

Plot of posterior distribution of psi

```
prior_psi <- function(x) {(1-x)^25.2/(beta(26.2,26.2)*(2-x)^52.4)}
post_psi <- function(x) {(1-x)^33.2/(beta(34.2, 188.2)*(2-x)^222.4)}
pfizer_prior <- function(x) {(1-x)^-0.700102/(beta(0.700102,1) * (2-x)^1.700102)}
ggplot() +
```



```
geom_function(fun = post_psi, xlim = c(0,1), aes(color = "Posterior"), linewidth = 1.2) +
geom_function(fun = pfizer_prior, xlim = c(0,1),
aes(color = "Pfizer Prior"), linewidth = 1.2) +
geom_function(fun = prior_psi, xlim = c(0,1),
aes(color = "Prior"), linewidth = 1.2, linetype = 'dashed') +
scale_color_manual(values = c("Posterior" = "firebrick",
"Prior" = "steelblue", "Pfizer Prior" = "darkgreen")) +
labs(x = expression(psi), y = "Density", color = "",
title = "Pfizer's Prior, and Our Prior and\n Posterior Distributions of Psi",
caption = "Figure 3: Prior and posterior densities for Psi.") +
theme_classic()+
theme(plot.caption = element_text(hjust = 0.5))
```

Computing 50th, 2.5th, 0.975th of Psi

```
inverse <- qbeta(0.5, 34.2, 188.2)
q_05 <- (2*inverse - 1)/(inverse - 1)
inverse_025 <- qbeta(0.025, 34.2, 188.2)
inverse_975 <- qbeta(0.975, 34.2, 188.2)
q_025 <- (1- 2*inverse_975)/(1-inverse_975)
q_975 <- (1- 2*inverse_025)/(1-inverse_025)
```

Parametric Bootstrap

Table Code

```
library(knitr)
library(latex2exp)
ci.df <- data.frame(Lower_Bound = c(lower, q_025, 0.903),
Upper_Bound = c(upper, q_975, 0.976))
rownames(ci.df) <- c("MLE Confidence Interval" ,
"Our Credible Interval", "Pfizer's Credible Interval")
colnames(ci.df) <- c("Lower Bound", "Upper Bound")

kable(ci.df,
caption = TeX("95% CI for Psi", output = "character"))
```

Proofs

If applicable, include detailed mathematical derivations or additional theoretical explanations.

Proving that the random variable V , representing the proportion of people who were infected with COVID that were in the vaccine group, approximates to the binomial distribution.

From the above data, we define X as a random variable representing the number of subjects in the vaccine group (of size n_1) who were infected with COVID, and Y as a random variable representing the number of subjects in the placebo group (of size n_2) who were infected with COVID. We assume they take the binomial distribution as follows:

Group	Cases	Sample size
BNT162b2	8	17,411
Placebo	162	17,511
Total	170	34,922

$$X \sim \text{Binom}(n_1, \pi_v) \quad Y \sim \text{Binom}(n_2, \pi_p)$$

Where $\pi_v = P(\text{COVID}|\text{Vaccine})$, $\pi_p = P(\text{COVID}|\text{Placebo})$, and X and Y are independent. We then define V as a random variable that represents the number of subjects that were infected COVID that were in the vaccine group. Then, V is the value of X given the constraint that $X + Y = n$, where n is the sample size of our variable V . (which, in our data, is $n = 170$). We define π as the rate at which a subject in the sample size n is in the vaccine group. As the sample sizes n_1 and n_2 are quite large, we know that the binomial random variables X and Y can be approximated with the Poisson. That is,

$$X \approx \text{Pois}(n_1\pi_v) \quad Y \approx \text{Pois}(n_2\pi_p)$$

Where $\lambda_x = E[X] = n_1\pi_v$, and similarly for $\lambda_y = E[Y]$

Before we proceed with the proof, we will prove two Lemmas, which will help with our main argument.

Lemma I Theorem: The rate of a COVID patient being in the vaccine group, π , is the probability $P(\text{Vaccine}|\text{COVID})$. Then:

$$\pi = \frac{P(\text{COVID}|\text{Vaccine}) \cdot P(\text{Vaccine})}{P(\text{COVID})} = \frac{n_1\pi_v}{\pi_v n_1 + \pi_p n_2}$$

Proof:

Using the data and the distribution definitions above:

$$\begin{aligned} P(\text{COVID}|\text{Vaccine}) &= \pi_v \\ P(\text{Vaccine}) &= \frac{n_1}{n_1 + n_2} \\ P(\text{COVID}) &= \frac{\pi_v n_1 + \pi_p n_2}{n_1 + n_2} \end{aligned}$$

Thus,

$$\begin{aligned} \pi &= \frac{\pi_v \cdot \frac{n_1}{n_1 + n_2}}{\frac{\pi_v n_1 + \pi_p n_2}{n_1 + n_2}} \\ \pi &= \frac{n_1 \pi_v}{\pi_v n_1 + \pi_p n_2} \quad \square \end{aligned}$$

Lemma II Theorem: Let X and Y be two independent Poisson random variables. Then, the sum of X and Y is also a Poisson distribution. That is,

$$S \sim Pois(\lambda_1 + \lambda_2) \quad S = X + Y$$

Proof:

Let $X \sim Pois(\lambda_1)$ and, independently, $Y \sim Pois(\lambda_2)$. Suppose $S = X + Y$

Then:

$$P(X = x) = \frac{e^{-\lambda_1} \lambda_1^x}{x!}$$

$$P(Y = n - x) = \frac{e^{-\lambda_2} \lambda_2^{n-x}}{(n-x)!}$$

By theorem 3.1 (Grove, 2025, Lecture 3):

$$P(S = n) = \sum_{x=0}^n P(X = x)P(Y = n - x)$$

$$P(S = n) = \sum_{x=0}^n \frac{e^{-\lambda_1} \lambda_1^x}{x!} \cdot \frac{e^{-\lambda_2} \lambda_2^{n-x}}{(n-x)!}$$

$$P(S = n) = e^{-\lambda_1 + \lambda_2} \sum_{x=0}^n \frac{\lambda_1^x \lambda_2^{n-x}}{x!(n-x)!}$$

$\sum_{x=0}^n \frac{\lambda_1^x \lambda_2^{n-x}}{x!(n-x)!}$ is the binomial expansion of $\frac{(\lambda_1 + \lambda_2)^n}{n!}$. So,

$$P(S = n) = \frac{e^{-\lambda_1 + \lambda_2} (\lambda_1 + \lambda_2)^n}{n!}$$

Thus, from the PMF above,

$$X + Y = S \sim Pois(\lambda_1 + \lambda_2) \square$$

Main proof Theorem: The random variable V , defined above, approximates to the binomial distribution.

Proof:

We proceed by finding the Probability Mass Function (PMF) of V . We know V to be the value of X given the constraint that $X + Y = n$. Our process, minding the independence of X and Y , thus follows:

$$\begin{aligned}
P(V = v) &= P(X = v | X + Y = n) \\
P(V = v) &= \frac{P(X = v \cap X + Y = n)}{P(X + Y = n)} \\
P(V = v) &= \frac{P(X = v) \times P(Y = n - v)}{P(X + Y = n)}
\end{aligned}$$

As X and Y can be approximated with the Poisson distribution, we will use the Poisson PMF, $\frac{e^{-\lambda} \lambda^x}{x!}$. Remember that $X \approx \text{Pois}(n_1 \pi_v)$ and $Y \approx \text{Pois}(n_2 \pi_p)$. And so, by Lemma II, $X + Y \approx \text{Pois}(n_1 \pi_v + n_2 \pi_p)$

$$\begin{aligned}
P(V = v) &= \frac{\frac{e^{-(n_1 \pi_v)} (n_1 \pi_v)^v}{v!} \times \frac{e^{-(n_2 \pi_p)} (n_2 \pi_p)^{n-v}}{(n-v)!}}{\frac{e^{-(n_1 \pi_v + n_2 \pi_p)} (n_1 \pi_v + n_2 \pi_p)^n}{n!}} \\
P(V = v) &= \frac{n!}{v!(n-v)!} \times \frac{(n_1 \pi_v)^v (n_2 \pi_p)^{n-v} e^{-(n_1 \pi_v + n_2 \pi_p)}}{(n_1 \pi_v + n_2 \pi_p)^n e^{-(n_1 \pi_v + n_2 \pi_p)}} \\
P(V = v) &= \binom{n}{v} \frac{(n_1 \pi_v)^v (n_2 \pi_p)^{n-v}}{(n_1 \pi_v + n_2 \pi_p)^n}
\end{aligned}$$

We apply Lemma I, to simplify the following:

$$\begin{aligned}
\frac{(n_1 \pi_v)^v}{(n_1 \pi_v + n_2 \pi_p)^v} &= \left(\frac{n_1 \pi_v}{\pi_v n_1 + \pi_p n_2} \right)^v = \pi^v \\
\left(\frac{n_2 \pi_p}{n_1 \pi_v + n_2 \pi_p} \right)^{n-v} &= \left(1 - \frac{n_1 \pi_v}{\pi_v n_1 + \pi_p n_2} \right)^{n-v} = (1 - \pi)^{n-v}
\end{aligned}$$

Thus,

$$P(V = v) = \binom{n}{v} \pi^v (1 - \pi)^{n-v}$$

The PMF of the binomial distribution. Therefore:

$$V \approx \text{Binom}(n, \pi) \blacksquare$$