



**INFLUENCE DES CONTRAINTES ENVIRONNEMENTALES, DES  
INTERACTIONS ET DE LA DISTRIBUTION DES TRAITS SUR LA  
RÉPARTITION GÉOGRAPHIQUE DES ESPÈCES**

Thèse présentée

dans le cadre du programme de doctorat en Biologie  
en vue de l'obtention du grade de Philosophiae Doctor

PAR

©KÉVIN CAZELLES

Septembre 2016



**Composition du jury :**

**[Prénom Nom], président du jury, [Université d'attache]**

**Dominique Gravel, directeur de recherche, Université du Québec à Rimouski**

**Nicolas Mouquet, codirecteur de recherche, Université de Montpellier**

**Nicolas Gotelli, examinateur externe, [Université d'attache]**

Dépôt initial le [13 septembre 2016]

Dépôt final le [-]

# UNIVERSITÉ DU QUÉBEC À RIMOUSKI

Service de la bibliothèque

## Avertissement

La diffusion de ce mémoire ou de cette thèse se fait dans le respect des droits de son auteur, qui a signé le formulaire « *Autorisation de reproduire et de diffuser un rapport, un mémoire ou une thèse* ». En signant ce formulaire, l'auteur concède à l'Université du Québec à Rimouski une licence non exclusive d'utilisation et de publication de la totalité ou d'une partie importante de son travail de recherche pour des fins pédagogiques et non commerciales. Plus précisément, l'auteur autorise l'Université du Québec à Rimouski à reproduire, diffuser, prêter, distribuer ou vendre des copies de son travail de recherche à des fins non commerciales sur quelque support que ce soit, y compris l'Internet. Cette licence et cette autorisation n'entraînent pas une renonciation de la part de l'auteur à ses droits moraux ni à ses droits de propriété intellectuelle. Sauf entente contraire, l'auteur conserve la liberté de diffuser et de commercialiser ou non ce travail dont il possède un exemplaire.

à Jascerb

## **REMERCIEMENTS**

Au royaume des idées, nul ne c'est ce qu'il va rencontrer. Chaque rencontre est la promesse d'un échange. Chaque conversation est une caisse de raisonnement possible pour les idées qu'on se fait de notre propre expérience. Chaque échange est alors une possible source d'inspiration pour sa réflexion. De fait, je ne peux faire la liste exhaustive des gens avec qui les échanges ont conduit à la présente thèse. Je me restreint à une portion de ceux avec qui les échanges ont été plus soutenus et ceux qui de longue date me soutiennent. Longueur des présents remerciements demeure ainsi raisonnable, en essayant d'être personnel et aussi avec une certaine pudeur qui m'amène à l'anonymat. Ces remerciements venant du cœur, ils expriment ma gratitude à ceux qui ont été présents dans mon parcours je les remercie pour ce qu'ils sont. Je ne suis pas tellement un homme de poésie, mais pourquoi se refuser le plaisir de contraindre. Ainsi, ai-je choisi d'écrire mes remerciements en alexandrins deux hémistiches égales rimes croisés. Les remerciements plus formels sont dans l'avant-propos.

Longue est la liste des gens que j'ai côtoyé

La liste est longue des personnes qui ont soutenu plus ou moins directement. On vote avec des personnes dont on se nourrit plus ou moins consciemment, ils participent à ce que nous sommes et l'on grandit avec eux Marie, Gigi, Elo, Léo, Alex, Tristan, Laurence, Elissa, Camille.

Des soutiens qui sont de longues dates, je ne sais pas dans quelle mesure ils ont contribué. On a tous des enfances plus ou moins joyeuses mais on porte en nous la trace drôle de statuts en permanent changement. La mienne a été une enfance une adolescence très heureuse avec quelques remous. Dans son sillage on entraîne des gens qui vous supporteront très longtemps, mais ils l'ont Miste, Gros Nico, Gras Nico, Gomar, Ariane, Cendrars, K-wik, Cimer !

La famille, certains disent qu'on ne l'a choisie pas, alors, j'avouerai être très chanceux.

Merci Josette, Jean-Louis, Mireille. merci.

Nico, Benjami, Guy, votre amitié et vos produit...

Le boulot c'est des rencontres de gens passionnés et passionnant avant tout tellement nous et nous sommes tellement uni par le destin particulier de la recherche. Merci Claire, Clara, Marie, Simon, Sonia, Andrei, Florian, Rob, Isabelle, Alain, Sonia. Merci Jaquet, Idaline, Hedou, Kevin, Jo Brasco, Pipin le Solver, Tim.

Legagneux, Casajus, Vissault, plaisir de travailler avec vous encore et encore.

Directeur Nicolas, Dominique merci pour la confiance. David

Merci d'être qui tu es Palette, ça fait du bien de savoir qu'il y en a des comme toi.

Chevrinias, Chevalier, Matoche, noué par le destin ....

Albouy de Montpellier à Rimouski ! Merci à toi et Severine,

Merci les soeurs, je sais que l'frétot est pas souvent là mais bin

Merci Mother

Papi, mami, éternelle soutien, des grands-parents comme vous tous les petits enfants en rêve... Peut pas être autrement, je fais de la science.

## ***AVANT-PROPOS***

Je n'avais pas d'objectif particuliers une envie de m'intéresser à la biologie et plus généralement les écosystèmes. J'ai ainsi envoyé divers mail et j'ai eu une réponse de Nicolas Mouquet qui m'a propulsé au Canada. Là j'ai découvert les travaux entre Nicolas Mouquet et Dominique Gravel dont je ne fais qu'essayer de les prolonger. Alors décider d'aller au doctorat et voilà ce que ça donne

David Mouillot.

Merci Miguel Araujo.

Merci au soutien financier, il rend possible.

Merci France, merci frontenac, merci CSBQ. rendu possible ma thèse.

Merci UQAR / Université de Montpellier. Merci au geek.

Les articles sont en anglais mais chaque section a une partie *résumé en français* dans laquelle je détaille le contexte scientifique, le contexte d'écriture et de publication et je traduit l'abstract.



## ***TABLE DES MATIÈRES***

REMERCIEMENTS . . . . .	vi
AVANT-PROPOS . . . . .	viii
TABLE DES MATIÈRES . . . . .	ix
LISTE DES ABRÉVIATIONS . . . . .	xiv
LISTE DES SYMBOLES . . . . .	xv
INTRODUCTION GÉNÉRALE . . . . .	1
Des îles et des espèces . . . . .	1
En suivant Wallace . . . . .	1
En suivant MacArthur et Wilson . . . . .	3
Quelles informations renferment les distributions d'espèces ? . . . . .	6
Enjeux de la connaissance de la répartition géographique des espèces . . . . .	7
Travail théorique et modélisation . . . . .	9
Les processus qui façonnent les aires de répartition . . . . .	13
Biogéographie historique . . . . .	13
Capacités de dispersion . . . . .	14
Contraintes abiotiques et niche écologique . . . . .	15
Réseaux d'interactions : interdépendance des espèces . . . . .	18
Echanges d'informations génétiques et processus micro-evolutifs . . . . .	20
L'invasion européenne du frelon asiatique . . . . .	21
Cadre théorique de la thèse . . . . .	23
Une vision puissante de la dynamique des distributions d'espèces . . . . .	23
Le modèle mathématique et les prédictions de la TIB . . . . .	26
L'importance de la TIB dans des développements théoriques plus récents . . . . .	29
Le rôle des interactions dans la distribution des espèces . . . . .	32
Importance des interactions dans la distribution . . . . .	32
Un problème d'échelle ? . . . . .	36

Vers une biogéographie énergétique . . . . .	38
ARTICLE 1	
A PROPOS DES INTERACTIONS BIOTIQUES ET DES CONTRAINTES ENVIRON- NEMENTALES À L'ÉCHELLE BIOGÉOGRAPHIQUE . . . . .	40
1.1 Résumé en français . . . . .	40
1.1.1 Contexte scientifique . . . . .	40
1.1.2 Publication associée . . . . .	41
1.1.3 Traduction du résumé de l'article publié . . . . .	41
1.2 Title . . . . .	43
1.3 Authors . . . . .	43
1.4 Abstract . . . . .	43
1.5 Introduction . . . . .	44
1.6 The model . . . . .	47
1.6.1 A simple probabilistic biogeographical model . . . . .	47
1.6.2 Integrating biotic interactions . . . . .	50
1.6.3 Integrating environmental gradients . . . . .	53
1.7 Exploring the model . . . . .	55
1.7.1 Biodiversity distribution over environmental gradients . . . . .	58
1.8 Discussion . . . . .	62
1.9 Acknowledgment . . . . .	68
ARTICLE 2	
THEORIE DE CO-OCCURRENCE DES ESPECES DANS LES RÉSEAUX D'INTERACTION . . . . .	69
2.1 Résumé en français du deuxième article . . . . .	69
2.1.1 Publication associée . . . . .	70
2.1.2 Traduction du résumé de l'article publié . . . . .	71
2.2 Titre . . . . .	73
2.3 Auteurs . . . . .	73
2.4 Résumé en anglais . . . . .	73
2.5 Mot-clef . . . . .	74

2.6	Introduction . . . . .	74
2.7	Definitions . . . . .	78
2.8	ACKNOWLEDGMENTS . . . . .	93
2.9	Context . . . . .	93
2.10	Material and Methods . . . . .	95
2.10.1	Datasets . . . . .	95
2.10.2	Measures of co-occurrence . . . . .	96
2.11	Results . . . . .	96
2.11.1	Discussion . . . . .	99
2.11.2	Box 1 . . . . .	101
2.11.3	Tables . . . . .	103
2.12	Figures . . . . .	104
2.13	Material and methods . . . . .	105
2.13.1	Datasets . . . . .	105
2.13.2	Building metawebs . . . . .	107
2.13.3	Co-occurrence measurement . . . . .	109
2.13.4	Hypergeometric distribution . . . . .	109
2.13.5	GLM and RF . . . . .	110
2.14	Supporting Tables . . . . .	111
2.15	Supporting Figures . . . . .	118
ARTICLE 3		
TOWARDS A METABOLIC THEORY OF BIOGEOGRAPHY . . . . .		129
3.0.1	Résumé en français du troisième article . . . . .	129
3.0.2	Publication envisagée . . . . .	129
3.1	Introduction . . . . .	130
3.2	Model . . . . .	133
3.2.1	Primary producers and habitat heterogeneity . . . . .	134
3.2.2	Metawebs . . . . .	134
3.2.3	Migration of species from the metaweb . . . . .	135

3.2.4	Energetic constraints on local food webs . . . . .	135
3.2.5	Energy fluxes and transfer efficiency . . . . .	136
3.2.6	Extinctions . . . . .	139
3.2.7	Simulations . . . . .	140
3.3	Results . . . . .	140
CONCLUSION GÉNÉRALE . . . . .		141
Introduire davantage d'écologie des communautés dans les modèles de distribution		
	d'espèce . . . . .	141
	L'information des réseaux . . . . .	141
3.3.1	Interaction des différents mécanismes . . . . .	142
	Vers une théorie en intégrative de la biogéographie . . . . .	142
	Des contraintes physiologiques . . . . .	143
3.3.2	L'abstraction des espèces . . . . .	143
	Traits fonctionnels . . . . .	143
	Des données nouvelles . . . . .	145
	Les défis à relever dans un monde en changement . . . . .	146
	Une érosion de la biodiversité affolantes . . . . .	146
3.3.3	Avons-nous des espoirs vains ? . . . . .	146
	Des écosystèmes bouleversés . . . . .	147
	Transients . . . . .	148
	Vers une écologie prédictive ? . . . . .	148
	Un espoir . . . . .	148
	Catégories . . . . .	148
	Les produits de l'évolution . . . . .	149
	Quels types de prédictions pouvons-nous faire ? . . . . .	150
	Les dangers d'aller trop vite . . . . .	151
3.3.4	Contraintes énergétiques . . . . .	151
ANNEXE I		
COMMENT LA BIODIVERSITÉ S'INSTALLE EN TERRITOIRES ISOLÉS . . . . .		152

4.4	Le modèle de MacArthur et Wilson . . . . .	152
4.5	Comment un modèle déterministe émerge-t-il de phénomènes aléatoires . . .	153
4.5.1	Modèle stochastique ? . . . . .	153
4.5.2	Les objets mathématiques requis . . . . .	154
4.5.3	Les briques élémentaires du modèle stochastique . . . . .	154
4.5.4	Assemblons les briques ! . . . . .	155
4.5.5	Retrouvons le modèle classique . . . . .	156
4.6	Applications et perspectives . . . . .	157
RÉFÉRENCES . . . . .		162

## ***LISTE DES ABRÉVIATIONS***

**SDM** Modèle de distribution d'espèce (en référence terme anglais : *Species Distribution Model*)

**TIB** Théorie insulaire de la biogéographie

**TTIB** Théorie trophique de la biogéographie des îles (en référence terme anglais : *Species Distribution Model*)

**RCP** Representative Concentration Pathway

## ***LISTE DES SYMBOLES***

**SYMBOLE 1** Ceci est la définition du symbole 1.

**SYMBOLE 2** Ceci est la définition du symbole 2.

**SYMBOLE 3** Ceci est la définition du symbole 3.

## INTRODUCTION GÉNÉRALE

La biogéographie est l'étude de la répartition géographiques des espèces. Aujourd'hui, ce terme est souvent remplacé par celui de macroécologie. Outre la distinction historique, ce mot met en avant l'importance du rapport des espèces à leur environnement (biotique ou abiotique) plutôt que la dimension évolutive pourtant tout aussi importante. C'est pour garder à l'esprit la richesse des facteurs qui dessinent les aires de répartition que je garde le terme de biogéographie, discipline dont je dresse un portrait dans la présente introduction. J'y aborde aussi bien la complexité de la compréhension de la distribution spatiale des espèces que les cadres théoriques associés. Chemin faisant, je discute de l'importance du lien qu'il existe entre les interactions écologiques et la répartition des espèces ; cette réflexion est l'essence même de ma thèse.

### Des îles et des espèces

#### En suivant Wallace

Dans l'introduction de son livre *Island Life* paru en 1881, le célèbre naturaliste Alfred Russel Wallace nous rapporte deux faits étonnants qui justifient pleinement l'examen attentif de la répartition géographique des espèces ([Wallace, 1881](#)). Premièrement, le biogéographe démontre, à l'aide de nombreux exemples, que l'éloignement entre deux régions du monde n'est pas suffisant pour conclure quand à l'éloignement de leur composition faunistique et floristique. Ainsi, la comparaison des avifaunes de l'île japonaise d'Hokkaido et de l'Angleterre, séparées par des milliers de kilomètres, révèle une proximité des paysages ornithologiques très supérieure à celle constatée dans l'analyse comparée des oiseaux des îles indonésiennes de Bali et de Lombok pourtant distantes de quelques kilomètres seulement. Deuxièmement, en s'appuyant sur les différences des faunes brésiliennes et africaines, Wallace souligne la faiblesse du pouvoir prédictif des variables climatiques pour décrire les compositions fau-



nistiques présentes sous des latitudes similaires. Ces constatations soulignent l'utilité de croiser les informations des distributions à la lumière d'une analyse taxonomique pour y apporter du sens. Dans le cadre de la théorie de l'évolution<sup>1</sup>, encore toute jeune en 1881, cette analyse taxonomique est une analyse historique : Wallace montre que la compréhension d'un problème spatial, celui des aires de répartition de groupes d'espèces, n'est possible que par une compréhension temporelle, celle de l'histoire des espèces. Cette idée est clairement énoncée dans la suite de son introduction :

« Many years study of this class of subjects has convinced me that there is no short and easy method of dealing with them ; because they are, in their very nature, the visible outcome and residual product of the whole past history of the earth. »

Tout au long de son livre, Wallace démontre que la connaissance à l'échelle mondiale de la distribution des êtres vivants permet d'associer les différentes îles aux grands ensembles régionaux biologiques (que nous appelons aujourd'hui écozones) sur la base des ressemblances biologiques des espèces qui témoignent du lien temporel unissant les différentes zones géographiques de la Terre. Ce travail de caractérisation d'ensemble géographiques conduit notamment Wallace, dans un article de 1860 ([Wallace, 1860](#)), à tracer la ligne éponyme séparant l'écozone indomalaise de l'écozone australienne (séparant les îles de Bali et Lombok mentionnée au paragraphe précédent). La connaissance apportée à la géographie par l'histoire est saisissante et les exemples de Wallace deviennent autant d'arguments en faveur de la théorie de l'évolution. Le discours de Wallace porte sur des processus à des échelles spatiales et temporelles très grandes<sup>2</sup>, ce qui apporte certes un éclairage substan-

---

1. Wallace a publié en 1858 un article *On the Tendency of Varieties to Depart Indefinitely From the Original Type* qui témoigne très clairement que ses idées sur les variations temporelles des espèces étaient très proches de celles de Charles Darwin à qui il avait d'ailleurs envoyé son manuscrit ([Wallace, 1858](#)).

2. L'âge de la terre est très débattu à l'époque. Bien que l'ensemble des savants s'accordent pour aller bien au-delà des 6000 ans bibliques, il n'y a alors pas de consensus. Wallace affirme à la page 212 du chapitre 10 de *Island Life* que la vie se développait il y a au moins 500 millions d'années ([Wallace, 1881](#)), ce qui est

tiel qui se double cependant d'un obstacle épistémologique majeur : si l'explication ultime de la présence d'une espèce en un point donné est le produit d'une série de contingences historiques, quelles peuvent être les fondations d'une théorie de la biogéographie ? Ce n'est qu'au XX<sup>ème</sup> siècle que des réponses convaincantes émergeront.

### En suivant MacArthur et Wilson

Parmi les visions les plus importantes de la biogéographie ce trouve celle contenue dans le livre publié en 1967 *The Theory of Island Biogeography*, produit de la fructueuse rencontre du mathématicien et biologiste Robert MacArthur et du myrmécologue Edward Wilson<sup>3</sup>. A partir d'un grand nombre de données sur les faunes insulaires de diverses régions du monde, ces auteurs ont construit un cadre théorique puissant pour expliquer les variations du nombre d'espèce trouvé sur ces îles (MacArthur and Wilson, 1967). Leur démarche théorique permet de lier des observations à un modèle mathématique donnant une explication simple et convaincante des variations étudiées. Ils font ainsi basculer la discipline dans une ère nouvelle, ce dont les auteurs étaient conscients, en atteste le premier paragraphe du dernier chapitre de leur livre :

« Biogeography has long remained in a natural history phase, accumulating information about the distribution of species and higher taxa and the taxonomic composition of biotas. Interpretative reasoning has been largely directed to the solution of special problems connected with the histories of individuals taxa and biotas. Without doubt this descriptive activity will continue to be of fundamental

---

audacieux pour l'époque mais bien en-dessous de l'âge des plus anciens fossiles découverts à ce jour qui est estimé à 3.4 milliards d'années (Wacey et al., 2011).

3. Cet actuel professeur émérite à l'université d'Harvard est reconnu pour ses apports en biologie et en sociologie, il est notamment l'auteur de 32 livres. C'est pour son immense connaissance des fourmis que j'ai choisi l'adjectif de myrmécologue.

importance to the science, one of the most physically adventurous of all scientific enterprises and, in the richness of the detail it unfolds, esthetically pleasing. But biogeography is also in a position to enter an equally interesting experimental and theoretical phase. »

Dans cet extrait, MacArthur et Wilson affirment que l'étude de la distribution des espèces doit sortir du royaume des contingences historiques pour devenir un objet de science au sens d'être manipulé aussi bien expérimentalement que par l'abstraction mathématique. La validation expérimentale de la théorie a d'ailleurs été menée par Wilson et son étudiant au doctorat de l'époque Daniel Simberloff devenu depuis un célèbre écologue ([Simberloff and Wilson, 1969](#)). Le travail d'abstraction mathématique a été conduit par MacArthur dans le livre de 1967 et prolongé dans les annexes de son livre de 1972 ([MacArthur, 1972](#)). Ces auteurs proposent une explication de la variation spécifique des îles fondée sur deux processus opposés : la colonisation d'espèce depuis le continent qui augmente le nombre d'espèce sur l'île et un processus d'extinction locale qui diminue ce nombre. C'est en reliant ces processus aux propriétés physiques de l'île (aire et isolation) et en interprétant la richesse spécifique des îles en terme d'équilibre entre ces deux processus que les auteurs parviennent à expliquer de manière convaincante les relations observées entre richesse spécifique, taille de l'île et isolement. Dans le troisième temps de cette introduction, je reviens amplement sur cette théorie nommée théorie de la biogéographie des îles que je noterai TIB dans la suite.

Le paradigme de la TIB est un legs qui a eu un impact considérable sur les développements théoriques en écologie ([Warren et al., 2015](#)). Au centre du projet de la TIB se loge la volonté de mettre l'espèce au cœur de la biogéographie afin de permettre à la discipline de s'enrichir des mécanismes biologiques qui sont un moteur essentiel de la variation dans la distribution des espèces. L'intérêt de leur *biogéographie de l'espèce* (terme donné à l'avant-dernière phrase de leur livre) est dans l'affirmation qu'il faut regarder les contraintes conjointes de l'évolution (qui met un certain nombre d'espèces avec des caractéristiques propres en présence) et du contexte écologique qui détermine les conditions d'extinction. Cette intrication de

l'écologie et de l'évolution est bien inscrite dans la pensée de MacArthur et Wilson même si la puissance de leur vision réside dans le fait de les occulter en partie.

Près de 50 ans après la parution de leur livre, une des clef en biologie semble être la compréhension des rétroactions qu'il existe entre écologie et évolution dans les variations spatiales et temporelles de la biodiversité. Je reprend ci-dessous trois aphorismes cités par [Schoener \(2011a\)](#) concernant les liens entre biologie, écologie et évolution :

« Nothing in biology makes sense except in the light of evolution. » ([Dobzhansky, 1973](#))

« Nothing in evolutionary biology makes sense except in the light of ecology. » ([Grant and Grant, 2008](#))

« Nothing in evolution or ecology makes sense except in the light of the other. » ([Pelletier et al., 2009a](#))

La chronologie de ces citations est un indice de la reconnaissance actuel du besoin (de la nécessité ?) de croiser écologie et évolution. Un parallèle avec les sciences humaines me semble possible dans lequel l'écologie serait à la biologie ce que la géographie est aux sciences humaines et l'évolution serait à la biologie ce que l'histoire est aux sciences humaines. Nous pouvons certes étudier l'une sans l'autre, mais le dialogue entre les deux disciplines est indispensable. En son absence, les disciplines avancent en trainant avec elles des hypothèses fortes sur l'autre qui finiront éventuellement par nuire à une compréhension plus profonde de la biologie. Par exemple, supposer que les variations démographiques ont des origines purement écologiques devient problématique si les variations génétiques sont suffisantes pour expliquer qu'une partie importante de cette variation comme cela l'a été montré sur une population de mutons de Soay ([Pelletier et al., 2007](#)). Je ne cherche pas à nier l'utilité des savoirs acquis de manière autonome par un champ disciplinaire, j'insiste simplement sur l'importance de mettre ces connaissances en commun dans une synthèse indispensable pour

décrypter l'information contenue dans les distributions d'espèce.

### **Quelles informations renferment les distributions d'espèces ?**

Cette question est non seulement une invitation à découvrir les raisons de la présence de tel ou tel organisme en un lieu donné du globe, mais elle suggère aussi que certaines informations ne peuvent pas être obtenues par l'analyse de la répartition géographique des espèces. Les auteurs mentionnés dans les paragraphes précédents y ont apporté des éléments de réponse cruciaux : Wallace a montré que les distributions géographiques reflétaient en partie les liens de parenté entre les espèces, quant à MacArthur et Wilson, ils ont suggérés que ces distributions étaient le résultat de processus écologiques dynamiques. Examiner les aires de répartition, en détailler la géométrie exacte et les variations spatio-temporelle, faire des recoupements entre les répartitions géographiques de différentes espèces ou encore avec la distribution de variables abiotiques sont des démarches fondamentales pour en apprécier les mécanismes sous-jacents.

Dans son ouvrage de 1972, MacArthur discute de l'ensemble de ces mécanismes, il considère aussi bien le rôle que peuvent jouer les variables climatiques que celui des interactions écologiques. En plus des exemples concrets amenés pour illustrer ses propos, MacArthur développe des modèles mathématiques pour prolonger la discussion. Au chapitre 2, il formalise l'impact de la compétition sur la coexistence des espèces aboutissant ainsi sur un principe de ségrégation spatiale des espèces liées par ce type de relation : deux compétiteurs ne peuvent pas co-occurrencer (résider durablement au même endroit) sauf éventuellement sur zone très restreinte de leur distribution (MacArthur, 1972). Toujours dans cet ouvrage, MacArthur évoque la distribution en damier (*checkerboard*) que peuvent générer des espèces en compétition. La discussion sur ce type de distribution sera approfondie par Jared Diamond (Diamond, 1975) dont les travaux déclencheront un débat important sur la détermination de modèle nul de co-occurrence (Connor and Simberloff, 1979) et sur laquelle ma thèse apporte quelques éléments nouveaux.

L'étude sur un grand nombre d'espèce de leurs limites spatiales permet d'y déceler des généralités quant aux mécanismes qui les déterminent (MacArthur, 1972). L'examen des variations spatio-temporelles apporte une information très utile sur l'importance relative des divers mécanismes. Le contexte des changements climatiques est une bonne illustration de ce principe car les bouleversements actuels des répartitions géographiques permettent en effet de pointer le rôle majeur de certains mécanismes évolutifs auparavant sous-estimés (Lavergne et al., 2010). Enfin, l'examen des distributions doit aussi être un examen des co-distributions, il faut s'intéresser à l'information de sous ensembles d'espèces et notamment les espèces en interaction afin de tester si la biologie laisse des empreintes dans la géométrie des aires de répartition. Par exemple, dans ma thèse je propose de regarder l'intersection des l'aire associée à un ensemble de proies pour savoir ce qu'elle nous apprend sur la distribution de leur prédateur.

### **Enjeux de la connaissance de la répartition géographique des espèces**

Les observations et la compréhension des causes profondes de la géométrie et la dynamique des aires de répartitions des espèces ont déjà amené à des découvertes majeures en écologie et en évolution. La phase d'expérimentation et de théorisation de la biogéographie décrite par MacArthur et Wilson se poursuit et se tourne vers un objectif très ambitieux : faire de la biogéographie une discipline prédictive, pourvoyeuse de prédictions fiables sur les aires de répartitions futures de n'importe quelle espèce. Ce problème est d'autant plus présent dans la littérature récente que nous sommes dans un contexte où ces aires sont profondément bouleversées. En biogéographie, les changements climatiques ont en effet canalisés l'attention des chercheurs qui constatent avec stupeur l'ampleur à laquelle la biodiversité mondiale est affectée (Koh et al., 2004; Bellard et al., 2012). La volonté d'anticiper la localisation future des espèces a également engendré des efforts conséquents pour développer des outils statistiques essentiellement centrés sur la corrélation entre les variables abiotiques et les données de présence (d'occurrence) des espèces (Elith et al., 2006).

En ciblant l'étude de la distribution de certaines espèces, la biogéographie rencontrent des enjeux socio-économiques majeurs. Ainsi, pour un pays comme la France, la restriction des zones favorables à la croissance de la vigne envisagée à l'aide des scénarios de changements climatique (Hannah et al., 2013) pourrait conduire à des pertes économiques importantes et un bouleversement identitaire des grandes régions viticoles. De plus, détecter aujourd'hui un potentiel viticole futur dans des régions où cette production n'existe pas peut conduire à des augmentations drastiques du prix des terres agricoles. En guise de second exemple, je pose la question suivant : où seront les érablières de demain ? La réponse est donnée par la détermination de la répartition future des aires favorables à la croissance de l'érable à sucre (*Acer saccharum*) et de sa capacité à les atteindre afin de s'y établir. Je termine avec un troisième exemple : la perte des pollinisateurs et notamment des abeilles. Pas moins de quatre grandes classes de facteurs d'origine anthropique les mettent en danger : les changements climatiques, le changement dans l'utilisation des terres<sup>4</sup>, l'apparition de nouveaux pathogènes (dont l'acarien parasite *Varroa destructa* vecteur de nombreux virus) et l'arrivée d'espèces invasives (comme le frelon asiatique) (Vanbergen, 2013). Le défi actuel est de prédire la distribution future des pollinisateurs en intégrant ces multiples aspects et leurs interactions. De plus, dans le cas des espèces invasives, il faut comprendre comment une espèce peut sortir de son aire de répartition naturelle et en établir une nouvelle.

Actuellement, les outils de prédictions des aires de répartition future reposent essentiellement sur les scénarios de changements climatiques. La démarche est cohérente : la connaissance basée sur les corrélations de variables climatiques permet d'établir une relation climat-présence. En utilisant les résultats des climatologues qui dérivent les variations climatiques associées à des scénarios d'émission de gaz à effet de serre par les activités humaine, les biogéographes établissent les probabilités de présence des espèces dans les conditions climatiques futures. Cependant, la relation climat-présence n'est qu'une facette du lien qui

---

4. Les changements dans l'utilisations des terres sont accompagnés, entre autres, de l'utilisation parfois massive de pesticides de la famille des néonicotinoïdes affaiblissant les colonies.

unissent les espèces à l'espace et chaque nouvelle invasion nous montre à quel point il est difficile de prédire les aires de répartitions. Ces problèmes de qualité de prédictions sont le reflet de lacunes théoriques qui amènent plusieurs chercheurs à se positionner en faveur d'un renouvellement des fondations théoriques pour édifier une biogéographie plus intégrative (Lomolino, 2000b; Beck et al., 2012; Thuiller et al., 2013). Bien sur, ces appels soulèvent des défis importants dont on ne peut qu'espérer qu'ils soient relevés au plus vite pour faire face à l'urgence.

### **Travail théorique et modélisation**

Avant d'énumérer, avec des exemples concrets, l'ensemble des forces qui régissent la répartition géographique d'une espèce, je souligne dans cette partie l'importance du travail de théorie et de modélisation qui joue un rôle prépondérant dans ma thèse.

### **Rassembler et intégrer des faits**

Le travail de théorie est avant tout la mise en cohésion d'un certain nombre de faits, d'observations. Dans la TIB, par exemple, MacArthur et Wilson proposent une explication cohérente de l'augmentation de la richesse spécifique dans les îles de plus grande taille. Trois principes encadrent la construction d'une théorie scientifique :

1. la théorie doit pouvoir être testées (par une expérience ou par la récolte de données),
2. la théorie doit être falsifiable : la théorie demeure valide tant qu'elle n'est pas prouvée fausse, tant qu'une théorie alternative ne la supprime pas,
3. la théorie doit être parcimonieuse, ne pas invoquer de multiples processus sans raison (c'est-à-dire sans une augmentation du nombre de faits expliqués), c'est un principe qui



est aussi connu sous le nom de Rasoir d'Ockham.

Une boutade, dont je ne suis pas capable de me souvenir son auteur, énonce que les physiciens expliquent 95% de l'univers avec 5 règles alors que les économistes expliquent 5% des phénomènes qu'ils étudient avec 95 règles<sup>5</sup>. Le problème n'est pas tant de dénigrer une discipline mais de constater d'un côté la puissance prédictive d'une théorie mature et de l'autre, les problèmes posés par une théorie lacunaire. En biogéographie, j'ai le sentiment que les théories manquent de maturité, la TIB donne certes une vision cohérente de la richesse spécifique insulaire mais c'est une théorie peu précise : prédire un nombre d'espèce n'aide que partiellement à comprendre le monde qui nous entoure. Pour faire un peu de prospective, une théorie qui donnerait des prédictions sur la topologie des réseaux et la composition en masse des espèces présentes supplanterait la TIB car elle expliquerait davantage de faits au prix probable d'une complexité supérieure.

### **Des modèles pour explorer et tester la théorie**

Le terme de modèle signifie simplement que l'objet en question a des propriétés bien connues. Un organisme modèle, par exemple, est un organisme souvent facile à élever et manipuler pour lequel beaucoup de connaissances ont été acquises, il sert souvent d'unité empirique pour un ou plusieurs groupes de recherche. Les modèles statistiques sont des outils pour tester des relations basées sur des hypothèses issues de théories. De même, pour un travail de modélisation mathématique, la description du modèle est contenue dans une série d'équations dérivée d'une théorie. A travers les modèles, quelle qu'en soit leur nature on explore et on teste une théorie que l'on a éventuellement participé à établir.

Les modèles sont souvent décrits comme une simplification de la réalité : comment,

---

5. Une variante indique que les économistes ont prédit 12 des trois dernières crises économiques. Je pense que pour ce qui est de nos capacités de prédictions, la biogéographie est plus proche de l'économie que la de la physique..

en effet, prétendre que les mécanismes biologiques décelés chez *Arabidopsis Thaliana*<sup>6</sup> sont les mêmes à l'oeuvre pour l'ensemble des plantes à fleurs ? pour combien de systèmes proie-prédateur le modèle de Lotka-Volterra est-il pertinent ? Les limites des modèles doivent être reconnues mais il ne faut pas nier l'apport de ces derniers. Les modèles sont autant de chance pour explorer une ou plusieurs prédiction d'une théorie. Le choix du modèle employé est lié à l'histoire du chercheur qui l'utilise, à ses propensions mentales à utiliser avec succès telle ou telle démarche scientifique, c'est ce que rappelle Kevin McCann dans la préface de son livre *Food Webs* (McCann, 2011) :

« It just so happens that some people find it easier to think about things in terms of x's and y's, and other in terms rabbits of and lynx. »

En d'autres termes, certaines personnes ont plus de facilités pour penser en termes d'abstraction mathématique alors que d'autres font meilleur usage de manipulations plus concrètes. Je suis plutôt dans la première catégorie de personne, je pense que les mathématiques sont un cadre de penser très puissant comme l'indique le grand écologue Robert May (May, 2004) :

« The virtue of mathematics in such a contexte is that it forces clarity and precision upon the conjecture, thus enabling meaningful comparison between the consequences of basics assumptions and the empirical facts. Here mathematics is seen in its quintessence : no more, but no less, than a way to think clealy. »

Dans ma thèse, j'ai essayé d'utiliser les mathématiques pour développer des modèles dont le point de départ a été une réflexion collective autour du rôle que pouvaient jouer les interactions dans la répartition géographique des espèces. J'ai alors établi un cadre théorique

---

6. Il s'agit de la plante modèle par excellence dont le génome fut le premier à être séquencé chez les plantes (Arabidopsis Genome Initiative, 2000).

avec lequel j'ai dérivé des prédictions dont certaines semblent être vérifiées dans les données empiriques.

## Nouvelles prédictions

Après l'établissement d'une théorie expliquant un certain nombre de faits et pour laquelle un certain nombre de tests ont été réalisés, le raisonnement fondé sur celle-ci peut conduire à la production de nouvelles prédictions dont la vérification la renforceront. En revanche, l'apparition des faits expérimentaux en désaccord avec cette théorie demanderont des réponses qui se traduiront soit par une meilleur compréhension de la limite d'application de la théorie soit par l'émergence d'une théorie nouvelle qui expliquera ces faits nouveaux tout en couvrant le rayon de compréhension de la théorie précédente. Ces dernières années, la physique nous a donné deux exemples très probants du pouvoir de l'imagination avec la vérification expérimentale de théories énoncées bien avant que les outils permettant de la vérifier existent. En 2012, c'est la détection du Boson de Higgs dont l'existence fut prédite en 1964<sup>7</sup>. Cette année, c'est la détection des ondes gravitationnelles soit 100 ans après qu'Einstein en ait prédit l'existence (Waldrop, 2016). En biogéographie, une théorie devrait être capable, par exemple, de dresser des cartes d'invasibilité à l'échelle mondiale pour l'ensemble des espèces. Je pense que nous en sommes encore loin, néanmoins, le chemin pour y parvenir passe par une connaissance approfondie de l'ensemble des mécanismes qui interviennent dans le tracé des aires de répartition, c'est-à-dire connaître leur nature, la portée de leur action mais aussi leurs interactions et leurs importances relatives.

---

7. Pour plus de détail, je réfère le lecteur au bulletin du CERN disponible en ligne <http://cds.cern.ch/journal/CERNBulletin/2012/28/News%20Articles/1459456?ln=fr>

## Les processus qui façonnent les aires de répartition

### Biogéographie historique

Il s'agit de la compréhension des impacts sur les êtres vivants des événements de grande amplitude temporelle (allant de quelques milliers d'années à plusieurs millions d'années). C'est dans l'étude de la proximité des taxons mais aussi des fossiles éventuels que l'on peut déchiffrer les mouvements de colonisation des différentes branches de l'arbre du vivant. Pour prendre un exemple de phénomène de très grande amplitude, on peut faire appel à la théorie de la dérive des continents établie par Alfred Lothar Wegener<sup>8</sup> qui implique que des groupes éventuellement proches il y a des millions d'années ont été séparés et ont donné naissance à des lignées différentes. Aujourd'hui, nous sommes capables de retracer ces liens de parenté à l'aide de phylogénies moléculaires qui sont des outils très efficaces pour estimer le temps que sépare différents. Ainsi, par la comparaison des génomes mitochondriaux, il a été montré récemment que les lémuriens (primates malgaches) ont été séparés de toute autre lignée de primates il y a 60 millions d'années environs ([Finstermeier et al., 2013](#)). Cette séparation questionne bien sûr la série d'événements qui ont conduit à l'isolation de ce groupe de singes à Madagascar et à la construction des communautés que nous y observons actuellement ([Razafindratsima et al., 2013](#)).

Les processus de grandes amplitudes temporelles sont cependant dominés par leur composante historique (et donc non reproductible) et prédire des phénomènes tel que l'extinction des dinosaures est, dans le meilleur des cas, très compliqué. Néanmoins, dans les mouvements de grandes amplitudes se manifestent des processus qui sont en permanence à l'oeuvre. Ainsi, l'étude de la diversification des bousiers entreprise par Joachim Hortal et collègues ([Hortal et al., 2011](#)) montre que la dernière glaciation a laissé des empreintes encore visibles dans

---

8. La similarité des fossiles trouvés sur des continents très éloignés a été un des arguments en faveur de cette théorie.

la carte de répartition de la diversité de ce groupe : la limite de la thermocline 0°C durant le dernier maximum glaciaire (il y a 21000 ans environs) sépare les zones de fortes diversifiées en bousiers des autres. De plus, ils montrent que la diversité phylogénique des espèces nordiques, c'est-à-dire plus tolérantes au froid, est un sous-ensemble phylogénétique bien identifié, par conséquent peu de branches des bousiers sont à l'origine des colonisations nordiques. Ainsi, après une contraction de la zone favorable au développement des bousiers, les mouvements de colonisation ont marqué à la fois la carte de répartition de la richesse spécifique de ce groupe mais aussi la carte de la répartition des différentes branches de l'arbre phylogénétique des bousiers européens ([Hortal et al., 2011](#)).

### **Capacités de dispersion**

La remontée nordique des bousiers depuis le dernier maximum glaciaire est le résultat d'événements de dispersion individuel. Au cours de leur vie, les bousiers parcourent de grandes distances à la recherche de nourriture, on peut imaginer qu'au fil des générations, si les conditions environnementales le permettent, certains individus établissent des populations de plus en plus nordiques. Ce qui est vrai pour ce groupe d'espèce mobile l'est aussi pour des espèces sessiles comme les plantes qui possèdent également des capacités de dispersion du fait de la dissémination de leurs semences par des mécanismes très diversifiés. Ce rapport à l'espace des différents organismes est une forme de diffusion : des mouvements stochastiques conduisent à une augmentation de la répartition (c'est une question de probabilité), mais cette diffusion n'est pas totalement libre.

Plusieurs types de contrainte limitent l'élargissement de l'aire de répartition d'une espèce. Pour les espèces terrestres, les mers et les océans sont des obstacles majeurs à la colonisation de nouveaux territoires. A l'échelle régionale, les rivières, les hauts reliefs peuvent fortement limiter la dispersion d'une espèce. De même, pour les plantes dont la stratégie de dissémination est l'anémochorie, la vitesse et la direction des vents sont des facteurs primordiaux pour comprendre la propagation de l'espèces. Enfin, à l'échelle du paysage, il existe

très souvent une mosaïque d'habitats plus ou moins favorables à la dispersion d'une espèce. Toutes ces possibilités sont complexes à intégrer et c'est en partie pour cela que la théorie en biogéographie a été fondée sur les îles, les flux de colonisation y étant relativement faciles à identifier : de la côte la plus proche vers l'île.

Dans l'expérience historique de Simberloff et Wilson qui valida la TIB, les chercheurs ont éradiqué la faune de six îlots de mangrove rouge dans la Baie de Floride et ils ont alors observé qu'en une année, la richesse spécifique en insecte était similaire à celle constatée avant de commencer l'expérience (Simberloff and Wilson, 1969). Ainsi, les événements de colonisation, bien qu'individuel, peuvent être assez fréquents pour conduire rapidement à l'établissement de populations et même d'une communauté locale d'insecte. A l'échelle d'un continent, malgré les divers obstacles physiques, il est très probable qu'une espèce donnée puisse, en un temps plus ou moins long, atteindre n'importe quelle zone du continent. Cependant, le plus souvent, les aires de répartition des espèces sont limitées à une portion du continent. Pour comprendre ces restrictions, il faut invoquer les performances des espèces dans des conditions environnementales données.

### **Contraintes abiotiques et niche écologique**

Dans le chapitre 6 de son livre de 1972 *Geographical Ecology* (MacArthur, 1972) illustre l'importance des contraintes climatiques avec l'exemple de l'aire de répartition du cactus Saguaro (*Cereus giganteus* en 1972 mais aujourd'hui *Carnegiea gigantea*). Ce résident des hauteurs du désert de Sonora (bordé à l'ouest par l'océan pacifique) est sensible au gel et ne peut guère résister à une exposition de quelques dizaines d'heures au gel. Cette contrainte physiologique explique bien les limites nord et est de sa répartition. Pour la limite sud, il semblerait que l'abondance des pluies hivernales qu'il y trouve lui soit défavorables. En s'appuyant sur les conditions climatiques actuelles dans lesquelles le cactus se développe, des résultats récents prédisent que dans le cadre des changements climatiques, *Carnegiea gigantea* trouvera refuge à des altitudes supérieures mais que ce mouvement pourrait être

entravé par l'augmentation de la fréquence des feux ([Springer et al., 2015](#)).

Cette démarche de croisement de la limite des aires de répartition avec des variables climatiques est une forme répandue de la détermination de la niche écologique d'une espèce. Ce concept de niche est très débattu en écologie et son caractère éluif s'accompagne d'un certain nombre de problèmes de définition<sup>9</sup>. Afin d'éviter ces problèmes, je parlerai de la niche au sens de Joseph Grinnel qui en tentant d'expliquer la restriction de la répartition du Moqueur de Californie écrit :

« An explanation of this restricted distribution is probably to be found in the close adjustment of the bird in various physiological and psychological respects to a narrow range of environmental conditions. »

Dans ses travaux, Grinnel montre que la présence du Moqueur de Californie est fortement corrélée à des conditions de températures et d'humidité assez élevées ([Grinnell, 1917](#)). Ainsi la niche écologique au sens de Grinnel est un ensemble de conditions environnementales dans laquelle une espèce donnée est trouvée. Si on ne se restreint pas aux observations *in situ* et que l'on détermine l'ensemble des conditions d'existence possibles, alors on caractérise une niche écologique théorique appelée niche fondamentale. Cette caractérisation expérimentale a été poussée à son paroxysme dans l'article de Michael Kearney et Warren Porter sur le gecko nocturne australien *Heteronotia binoei* ([Kearney and Porter, 2004](#)). Ils ont montrés qu'en combinant des mesures physiologiques (dont le taux métaboliques au repos, la température cumulée nécessaire au bon développement des oeufs et des mesures de températures caractéristiques) avec des données climatiques, ils obtenaient une bonne concordance des probabilités d'occurrence et des observations, justifiant ainsi la démarche

---

9. En 1957, Hutchinson propose de voir la niche écologique comme un hyperespace (un espace d'un grand nombre de dimension) dans lequel une espèce peut se développer. Le problème est de savoir quelles sont les dimensions et notamment si les autres espèces sont parmi ces dimensions. Pour essayer d'avoir une définition plus claire de la niche écologique, certains auteurs proposent de parler de la niche comme un espace où le taux de croissance net de l'espèce est supérieur à 0 ([Chase and Leibold, 2003](#)). En dépit de l'aspect plus quantitatif de cette définition, un problème subsiste, celui de trouver une méthode générale pour trouver cet espace.

prédictive en s'appuyant sur des scénarios de changements climatiques pour aller essayer de comprendre les répartitions futures. De manière générale, cette méthode est la recherche de facteurs abiotiques limitant le développement d'une espèce et donc sa répartition géographique. Au niveau du Panama, par exemple [Engelbrecht et al. \(2007\)](#) ont montrés que les distributions locales et régionales de 48 espèces d'arbres s'expliquent par la sensibilité à la sécheresse, donc à une variation dans la disponibilité d'une ressource. Ces corrélations convaincantes fondent les modèles de distributions d'espèces (SDM en référence au terme anglais utilisé souvent dans le reste de la thèse) qui sont des solutions techniques (statistiques) pour l'application de la méthode générale que je viens de décrire ([Elith et al., 2006](#); [Elith and Leathwick, 2009](#)).

L'engouement actuel autour de ces modèles est lié à l'espoir de pouvoir faire des prédictions fiables sur la géographie de la biodiversité mondiale de demain dans un contexte de changement climatique. Cette démarche s'est appliquée avec succès à différents cas, par exemple en 2009, Tingley et collègues ont ainsi montré que sur 53 espèces d'oiseaux étudiés dans la Sierra Nevada, 48 ont colonisé de nouveaux sites où les conditions de température et de précipitations leur étaient plus favorables ([Tingley et al., 2009](#)). Une autre justification de l'utilisation abondant des SDMs est la relative facilité de leur mise en application grâce à l'abondance des données climatiques et des données d'occurrence et au partage des implémentations numériques de ces méthodes statistiques. Pour le premier type de données, WorldClim illustre cette facilité d'accès en proposant des données à l'échelle mondiale gratuitement téléchargeables (voir <http://worldclim.org> [Hijmans et al., 2005](#)). Pour les données d'occurrence, plusieurs initiatives offrent des données gratuites dont les plus exhaustives sont celles disponibles sur le portail de données sur la biodiversité à l'échelle mondiale GBIF (*Global Biodiversity Information Facility*, voir <http://www.gbif.org>) qui présentent cependant des biais liés à l'inégalité d'échantillonnage des régions du globe ([Beck et al., 2014](#)). Enfin pour ce qui est le partage des implémentations des SDM, on peut évoquer le logiciel libre R ([R Core Team, 2015](#)) qui a des paquets dédiés à l'utilisation des SDMs et qui sont largement utilisés dans la communauté scientifique.



Un des principaux problèmes posés par l'utilisation massive de ces approches est le manque est la faible remise en question des hypothèses sur lesquelles elles reposent. Le message délivré par les SDMs doit être pris comme une potentialité : étant donné les conditions actuelles dans lesquelles une espèce est trouvée et sachant les variations climatiques donnés par les modèles climatologiques, s'il n'existe pas d'obstacle majeur au mouvement de l'espèce en question, alors il est probable que celle-ci se déplace en suivant les conditions climatiques qui sont similaires à celles dans laquelle elle est actuellement trouvée, ce qui nous permet de savoir où sera l'espèce demain. Ce message est délivré en supposant que 1- une forme d'équilibre de la distribution des espèces et des conditions climatiques actuelles et 2- que les espèces sont indépendantes (Jeschke and Strayer, 2008). Ces deux hypothèses sont très fortes et demandent un examen approfondi. Etant donné que ma thèse porte sur la seconde, je propose de la discuter dans le paragraphe suivant.

### **Réseaux d'interactions : interdépendance des espèces**

Au chapitre 6 de son livre *Geographical Ecology*, MacArthur parle précisément du rôle que peut avoir la compétition dans la distribution des espèces (MacArthur, 1972). Il reprend l'exemple donné par Brown en 1971 de l'exclusion compétitive de deux espèces de tamias, *Eutamias dorsalis* et *E. umbrinus*, dans les forêts d'altitude (au-dessus des déserts) de pins et de genévriers (*pinyon-juniper woodland*) du sud ouest des Etats-Unis. L'article de Brown montre bien comment une différence comportementale peut engendrer une séparation des distributions locales. Ainsi, l'agressivité de *Eutamias dorsalis* lui est favorable dans les forêts clairsemées de basse-altitude où son compétiteur doit dépenser beaucoup d'énergie pour lui échapper en se réfugiant dans un arbre, elle devient pénalisante lorsque l'abondance des arbres augmente car cela facilite la fuite de *E. umbrinus* (Brown, 1971). La ségrégation locale des deux espèces reflète donc bien une interaction biotique, il y a donc une information comportementale dans ces aires de répartition.

Au-delà de la compétition, l'écologie des réseaux nous montre aujourd'hui la difficulté

de concevoir les espèces comme étant des entités indépendantes, elles sont reliées par des relations de natures très diverses. Les relations trophiques sont les plus évidentes, il existe cependant une myriade d'interactions non trophiques qui affectent aussi la démographie des espèces (voir [Kéfi et al. \(2012\)](#) pour une réflexion sur le sujet et une classification de ces interactions). De plus, aucun argument théorique ne justifie actuellement la primauté d'un type d'interaction sur les autres. Récemment, les interactions trophiques et non-trophiques ont été exhaustivement analysées pour 104 espèces des écosystèmes intertidaux rocheux de la partie centrale de la côte chilienne révélant ainsi que les interactions non-trophiques y étaient globalement plus abondantes et concentrées sur les bas niveau trophiques ([Kéfi et al., 2015](#)).

L'écologie des réseaux est traversée de débats dont le plus important est sans doute celui de la relation qu'il existe entre la diversité spécifique d'un écosystème et sa stabilité ([May, 1973](#); [McCann, 2000](#)). Autour de cette question, l'écologie s'est considérablement enrichi en terme d'outils mathématiques. Une preuve récente réside dans la mise en évidence par Stefano Allesina et Si Tang du caractère déstabilisant des interactions de compétition et de mutualisme et du rôle stabilisant des relations trophiques ([Allesina and Tang, 2012](#)). Ce résultat est en effet la mise en application directe d'un résultat mathématique récent établi par Terence Tao et Vam Vu ([Tao et al., 2010](#)). Les réseaux contiennent de nombreuses informations sur l'écologie des population et à mon avis, ils doivent être placés au centre d'une théorie intégrative de la biogéographie. Cette idée n'est pas seulement la mienne, MacArthur et Wilson l'ont suggérée au dernier paragraphe de leur théorie de la biogéographie avec ces mots ([MacArthur and Wilson, 1967](#)) :

« In short, biogeography appears to us to have developed to the extent that it can be reformulated in terms of the first principles of population ecology and genetics.

»

Et pour appuyer cette phrase dans son entièreté, je développe un certain nombre d'idées relatives à l'importance des échanges génétiques.

## Echanges d'informations génétiques et processus micro-evolutifs

La vie, telle que nous la connaissons, pérennise l'information accumulée au cours du temps via à un support moléculaire, l'ADN. J'ai déjà évoqué que les informations véhiculées par cette molécule pouvaient permettre d'établir des relations de parenté entre les espèces. Cette possibilité est rendue possible par les mécanismes qui la modifient. L'information génétique d'un individu est un ensemble de bases dont la séquence renferme l'ensemble de l'information pour assurer le développement de l'individu. Néanmoins, le code génétique de certaines cellules de l'individu peut être modifiées (des mutations) et si ces cellules sont celles qui seront transmises à la descendance, alors ces modifications peuvent être transmises à la génération suivante. Sous certaines conditions, la mutation peut rester dans la population, c'est le moteur de la variation à l'échelle populationnelle du code génétique. Bien loin d'être une combinaison précise de paires de bases, l'ADN d'une espèce est en effet un ensemble de possibilités, un ensemble de versions du code possible mais contraint par certaines règles. Pour schématiser, les échanges de gènes doivent rester possibles entre individus d'une même espèce. A l'échelle des populations, tant que les échanges d'informations sont importants, la compatibilité est assurée mais lorsque ces échanges diminuent ou même cessent, les supports d'information peuvent alors diverger au point d'empêcher les échanges, ce qui conduit à la distinction de deux espèces. Bien que cette vision soit très simplifiée, elle permet de comprendre que l'ADN de deux espèces puissent refléter leur lien de parenté qu'il permet l'établissement d'une phylogénie moléculaire.

Cela étant dit, les causes de la divergence de l'ADN sont multiples et ce qui m'intéresse ici, ce sont que ces variations puissent engendrer un différentiel démographique positive dans un milieu nouvellement exploré par une population alors que cette même variation dans un autre milieu ne l'était pas. La vitesse des mécanismes mis en jeu semble bien plus rapide au point que ceux-ci puissent jouer des rôles prépondérants dans la réponse des espèces aux changements climatiques ([Lavergne et al., 2010](#)). En 2009, Joan Balanyá et collègues publient un article dans lequel ils comparent la composition génétique de la mouche *Drosophila*

*subobscura* entre des échantillons contemporains et des échantillons prélevé 24 années auparavant en Europe et en Amérique (où elle a été introduite accidentellement). Leurs résultats montrent que dans les zones de réchauffement climatique avéré, il y a aussi un changement de la composition génotypique avec une plus grande importance des génomes adaptés aux températures chaudes (Balanyá et al., 2006).

Les preuves récentes de l'impact des variations génétiques rapides sur la démographie des espèces populations poussent les chercheurs à se demander si négliger ces processus dans les travaux de dynamique des populations est une hypothèse raisonnable (Pelletier et al., 2009b; Post and Palkovacs, 2009; Schoener, 2011b). Takehito Yoshida et collègues ont montré en 2003 que la réponse des algues vertes unicellulaires *Chlorella vulgaris* aux rotifères *Brachionus calyciflorus* conduit à un changement dans la fréquence et la phase des cycles de la dynamiques proie-prédateur (Yoshida et al., 2003). En 2009, une étude basée sur un suivi de plus de 20 ans d'une population de moutons Soay sur l'île d'Hirta dans l'archipel de Saint-Kilda (au nord-est de l'Écosse), Fanie Pelletier et collègues établissent les variations dans la taille corporelle des ovins, d'origine génétique, et les variations dans leur survie et leur reproduction, ils démontrent alors que les facteurs génétiques peuvent contribuer jusqu'à 20% dans la croissance de la population certaine année. Les conséquences des dynamiques eco-evolutives et l'intégration des flux d'information génétique sont certainement capitaux pour comprendre la biodiversité de demain (Sexton et al., 2009; Lavergne et al., 2010). Nous sommes face à un enjeu appliqué capital et pourtant nos connaissances fondamentales restent insuffisantes. Pour illustrer ces lacunes et l'urgence dans laquelle nous nous trouvons, je discute d'un exemple concret : l'invasion européenne du frelon asiatique.

### **L'invasion européenne du frelon asiatique**

*Vespa velutina* est une espèce présente depuis le nord-est de l'Inde jusqu'à l'est de la Chine et de la péninsule et de l'indochinoise à l'archipel indonésien (Villemant et al., 2006). Dix sous-espèces sous identifiées dont *Vespa velutina nigrithorax* qui a été observé pour la

première fois en France en 2004 dans le Lot-et-Garonne chez un producteur de bonsaï qui importe régulièrement des poteries du Yunnan (Villemant et al., 2006). Ce frelon généraliste se nourrit notamment des abeilles domestiques et les conséquences sur les récoltes de miel sont désastreuses et ce même dans les zones d'origine où l'abeille asiatique (*Apis cerana*) est pourtant capable de tuer efficacement le frelon. Pour ce faire, les abeilles forment une boule autour du frelon et battent des ailes pour augmenter la température corporelle de leur prédateur, ce qui conduit à la mort de ce dernier. L'abeille européenne (*Apis mellifera*) est capable d'utiliser la même stratégie de défense mais avec une efficacité moindre (Villemant et al., 2006). Ce frelon représente un danger pour l'entomofaune européenne et aussi menace l'apiculture qui s'ajoute aux nombreuses autres que connaît actuellement le secteur (Vanbergen, 2013). Plusieurs éléments sont remarquables dans ce cas d'invasion : c'est un cas unique (première colonisation avec succès d'une nouvelle espèce frelon en France), la rapidité de propagation de ce frelon, le besoin urgent d'anticiper sa répartition dans les prochaines années pour mettre le plus rapidement en place les mesures d'éradication.

Après son arrivée en 2004, le frelon s'étendait déjà en 2006 largement sur l'Aquitaine avec une aire de répartition française constituée d'une bande de 300km du nord au sud et de 150 km d'est en ouest (Villemant et al., 2006) et cela malgré l'éradication systématique des nids détectés. Alors que 2 nids étaient observés en 2004, 1636 nids ont été observés en 2009 et en 2013 près des trois quarts des départements étaient affectés (Robinet et al., 2016). Des travaux récents tentent de caractériser les conditions climatiques favorables au développement de l'espèce (Villemant et al., 2011) et révèlent alors qu'une large partie de l'Europe occidentale est une zone de développement probable. Un autre phénomène intéressant lié à cette invasion est que dans la même période de la colonisation européenne, le frelon est arrivé la Corée du Sud où sa propagation est cependant bien moins rapide (Villemant et al., 2011). L'explication plausible de la différence de succès de la même espèce est une différence dans la composition en espèce proche des deux régions : en Europe, il n'y a qu'une espèce de frelon *Vespa crabro*, alors qu'il y en a de six en Corée du Sud dont *Vespa mandarinia* qui est une meilleur compétitrice (Villemant et al., 2011). Cette nécessité de faire appel à la compo-

sition biologique pour comprendre les raisons d'un changement d'aire de répartition est ce qui donne tout l'intérêt du travail théorique mené durant ma thèse.

## Cadre théorique de la thèse

Les développements entrepris durant ma thèse sont des tentatives d'encrage des interactions écologiques dans la TIB. Je vais maintenant revenir sur cette théorie plus en détail pour expliquer pourquoi elle a marqué durablement l'écologie. Je signale d'ailleurs que ces idées étaient partagées par d'autres écologues et qu'il y a, à ma connaissance, deux autres découvertes indépendantes des idées qui ont conduit à la théorie. La première découverte est attribuée au spécialiste des lépidoptères Eugene Gordon Munroe qui a formulé dès 1948, des idées similaires dans 5 des 555 pages de sa dissertation de graduation ([Brown and Lomolino, 1989](#); [Lomolino and Brown, 2009](#)). La seconde est celle de Richard Levins et Harold Heatwole qui publie en 1963, soit la même année que l'article fondateur de la TIB, l'idée d'un équilibre de la richesse spécifique régit par les mêmes processus que ceux décrits par MacArthur et Wilson ([Levins and Heatwole, 1963](#)). Néanmoins, ce sont sans aucun doute MacArthur et Wilson qui ont marqués les écologues par l'ensemble des développements présentés dans leur livre de 1967, *The Theory of Island Biogeography* ([MacArthur et al., 1967](#)).

## Une vision puissante de la dynamique des distributions d'espèces

Dans la préface de l'ouvrage de 1967, MacArthur et Wilson doutent que les idées proposées résisteraient longtemps à l'essor de la biogéographie expérimentale dont ils furent des acteurs de premier plan :

« We do not seriously believe that that the particular formulations advanced in in the chapters to follow will fit for very long the exacting results of future empirical invesitgation. »

Et pourtant près de 50 ans après la parution de ce livre, leurs travaux sont le fondement de nombreux développements récents, en témoigne le livre paru en 2010 *The Theory of Island Biogeography Revisited* (Losos and Ricklefs, 2010) et l'article de perspectives publié récemment par Ben Warren et collègues dans *Ecology Letters* (Warren et al., 2015). L'idée majeure de la TIB est simple et puissante : étant donné une île colonisable par un ensemble d'espèces depuis un continent voisin, la diversité locale résulte de la balance entre 1- des événements de colonisation depuis le continent et 2- des extinctions locales. La TIB est une métaphore, le cas simple d'un territoire isolé (l'île) où les flux d'individus depuis le pool d'espèces régionales (le continent) sont facilement représentables. Le modèle peut donc être étendu à de nombreux cas où un territoire isolé est colonisé par les organismes à proximité, par exemple après un incendie ou une fragmentation de l'habitat (Cook et al., 2002). Au chapitre 5 de son livre de 1972, MacArthur prend notamment l'exemple des îlots de paramo (végétation andine située au-dessus des forêts mais en-dessous des neiges éternelles). De manière générale, le modèle est acceptable est très adaptable au prix d'un certains nombre d'hypothèse notamment une certaine rigidité du réservoir d'espèces régional (au moins en nombre d'espèce) et une absence de rétroaction dans la communauté locale sur celui-ci.

Il y a une forme de hasard et de nécessité qui fait écho à l'oeuvre de Jaques Monod (Monod, 1970). Ce prix Nobel de médecine présente les mutations au niveau de l'ADN comme une source de hasard dont la persistance n'est rendu possible que dans un cadre physico-chimico-évolutifs précis, la nécessité. Dans les travaux de MacArthur et Wilson, l'événement de colonisation peut être interprété comme un pourvoyeur de stochasticité alors que les contraintes écologiques régissent l'organisation des communautés. Outre le fait que la prédiction de la colonisation ne peut se faire qu'en terme de fréquence, le caractère stochastique de cette dernière donne une dimension historique aux assemblages insulaires. L'arrivée d'une espèce est en fait un tirage aléatoire (éventuellement pondéré par les capacités respectives de dispersion) dans un réservoir régional d'une singularité historique car l'espèce en question à une histoire évolutive propre et des singularités physiologiques qui en découlent. A son arrivée sur l'île, son éventuelle insertion est déterminée par la rencontre des singula-

rités de l'espèce et du contexte biotique et abiotique de l'île. Les espèces installées sur une île ont ainsi été passées au crible des contraintes écologiques, de cette forme de nécessité qui est renouvelée à chaque nouvelle insertion. C'est ainsi que l'on peut décrire le moteur de la reconfiguration perpétuelle des réseaux écologiques locaux. Une telle dynamique peut être également analysée comme une imbrication de deux échelles de processus : régionalement, le réservoir d'espèce est façonné par une histoire évolutive de grande amplitude lié à des processus climatiques eux aussi de grande échelle, alors que les événements insulaires relèvent de processus de plus courte portée ([Ricklefs, 1987](#)).

Enfin, la TIB, bien que cela soit rarement souligné, fait l'hypothèse de l'équivalence écologique des espèces considérées : il n'y a ni plantes ni animaux, ni proies ou prédateurs, simplement des espèces qui compte pour un. Étant donné les exemples choisis par les auteurs on peut néanmoins penser que la théorie est développée pour des groupes d'espèce au rôle écologique similaire et phylogénétiquement proches. Ainsi, le premier exemple donné est pour l'herpétofaune (amphibiens et reptiles) et non sur un inventaire exhaustif de toutes les espèces de l'île ([MacArthur and Wilson, 1967](#)). Cette hypothèse est à relier aux objectifs des auteurs notamment celui d'expliquer les relations constatées entre la taille des îles et leur richesse spécifique, pour y arriver réduire les espèces à deux caractéristiques est suffisant et commode. La démarche peut néanmoins être perçue comme antithétique pour des auteurs qui cherchent à formuler une « biogéographie de l'espèce » et de surcroît quand on connaît la qualité de ces deux naturalistes ([Lomolino and Brown, 2009](#)). Cependant, la forme d'équivalence amenée par MacArthur et Wilson ne nie la diversité et la complexité, elle est plutôt une abstraction nécessaire pour capturer les processus essentiels, pour aller au-delà des singularités des êtres vivants, vers des généralisations ([Lomolino and Brown, 2009](#)).



## Le modèle mathématique et les prédictions de la TIB

Je ne rentre pas ici dans les détails mathématiques du modèle, ils sont néanmoins abordés dans le premier chapitre et aussi dans les deux annexes de la thèse<sup>10</sup>. J'écris ci-dessous l'équation qui résume à elle seule le paradigme livré par la TIB : les  $P$  espèces d'un continent colonisent l'île avec un taux individuel  $c$ , ce qui en augmente la richesse spécifique  $S$  mais augmente les risques d'extinctions dont le taux par espèce est noté  $e$  :

$$\frac{dS}{dt} = c(P - S) - eS \quad (1)$$

La dynamique ainsi engendrée conduit  $S$  jusqu'à un équilibre  $S_{eq}$  pour lequel les variations temporelles s'annulent, qui est donné par :

$$S_{eq} = P \frac{c}{c + e} \quad (2)$$

Cet équilibre est une prédiction très importante de la théorie, c'est même le point de départ des développements mathématiques dans le livre de 1967 (MacArthur and Wilson, 1967). L'existence d'un tel équilibre a été validée par l'expérience de défaunation de Simerloff et Wilson mentionnée plus haut (Simberloff and Wilson, 1969). Une seconde prédiction de la TIB est la variation de cet équilibre avec les caractéristiques de l'île. Dès leur article de 1963, MacArthur et Wilson présentent la taille de l'île comme un facteur affectant le taux d'extinction : plus l'île est grande, moins le risque d'extinction est grand (MacArthur

---

10. La première annexe est un article de vulgarisation qui aborde de manière didactique la formulation la plus simple du modèle. La seconde annexe est abordée des aspects plus techniques qui ont été l'objet d'un article dont je suis co-auteur.

and Wilson, 1963). De même, ils supposent que l'isolement de l'île en affecte le flux de migrants : plus l'île est isolée moins les événements de colonisation sont fréquents. J'ai résumé la vision classique de la TIB sur la figure 1 en y ajoutant les graphiques de l'article de 1963. Cette prédiction de la théorie en est aussi l'origine : MacArthur et Wilson expliquent avec ces mécanismes que les îles de plus grandes tailles est plus d'espèces mais aussi que des exceptions liées à l'isolement puisse exister. Cette relation est d'ailleurs présentée dès le début du chapitre 2 de la TIB avec l'augmentation linéaire du nombre d'espèce de l'herpétofaune avec le logarithme de la surface des îles de l'ouest des Caraïbes.

De manière plus générale, la TIB fournit une explication à la relation aire-espèce qui est un des objets les plus discutés de l'écologie (Lomolino, 2000a). Il s'agit de la courbe d'augmentation de la richesse spécifique ( $S$ ) avec la surface d'échantillonnage ( $A$ ). La question soulevée par l'étude de ces courbes porte sur la nature des mécanismes qui régissent les variations régionales. La TIB propose une explication à cette relation et supporte une courbe de la forme  $S = CA^z$  avec les observations présentées (MacArthur and Wilson, 1967). La relation aire-espèce est surtout connue pour ses applications dans le domaine de la conservation<sup>11</sup>. Elle permet d'estimer la taille qu'une zone de protection doit avoir pour atteindre un objectif de sauvegarde chiffré en nombre d'espèce (Neigel, 2003; Desmet and Cowling, 2004). La relation peut être aussi utilisée dans le sens inverse pour apprécier les taux d'extinction liés à une dégradation d'habitat (He and Hubbell, 2011).

---

11. Récemment Wilson a répondu à une entrevue dans laquelle il se base sur cette relation pour indiquer la proportion de la Terre qu'il faudrait épargner afin de maximiser la sauvegarde des espèces sans pour autant empêcher le développement humain <http://www.nytimes.com/2016/03/13/opinion/sunday/the-global-solution-to-extinction.html>.




Figure 1: **La Théorie de la biogéographie des Iles.** (A) illustre l'évolution des taux de colonisation et d'extinction est présentée pour deux îles aux caractéristiques différentes. Les tailles relatives des îles et les distances qui les séparent du continent sont schématisées sur la droite, les couleurs associent les îles à leurs courbes respectives. Le réservoir d'espèce régional ( $P$ ) est constitué de 100 espèces, les taux de colonisation et d'extinction sont exprimés en terme de probabilité d'évènement (de colonisation ou d'extinction). Les points marquent les intersections entre les courbes d'extinction et de colonisation c'est-à-dire lorsque ces processus s'équilibrent. L'abscisse de ces point indique les richesses spécifiques de l'île à l'équilibre  $S_{eq}$ . (B) et (C) sont respectivement les figures 4 et 5 extraites de l'article de 1963 de MacArthur et Wilson qui livre essentiellement le même message que celui illustré en (A) (MacArthur and Wilson, 1963). La forme convexe des courbes de 1963 sont justifiées par des facteurs biologiques qui ne sont pas intégrés dans l'équation qui confère une forme concave aux courbes comme vu en (A).

## L'importance de la TIB dans des développements théoriques plus récents

### La théorie des métapopulations

Bien que ne représentant que cinq pour-cents des terres émergées, ce sont bien les observations de la faune des îles qui ont mené à une vision paradigmatique de la biogéographie. L'importance des îles s'expliquent par leur relative abondance, leur disparité, leur diversité, la relative simplicité des assemblages biologiques qu'on y trouve et aussi, comme je l'ai évoqué précédemment, par la clarté des flux de migrations ([Simberloff, 1974](#)). Cette dernière propriété est souvent absente pour des populations continentales<sup>12</sup>. La théorie des métapopulations s'intéresse justement aux populations reliées entre elles par des flux de migrations ([Hanski, 2010](#)). Le premier modèle de métapopulations a été proposé par Levins<sup>13</sup> lors d'une réflexion sur le contrôle démographique des ravageurs dans les cultures ([Levins, 1969](#)). Pour un ravageur donné, les îlots de culture sont autant de patches où une population peut se maintenir et disperser dans les autres patches alentour. Levins montre alors que les mesures de la lutte biologique doivent être conduites à large échelle pour en augmenter les probabilités de succès, c'est-à-dire d'extinction régional du ravageur ([Levins, 1969](#)). Le modèle est simple et très proche de celui de la TIB : l'évolution de la proportion  $p$  est aussi gouvernée par des événements de colonisation  $c$  et d'extinction  $e$  :

$$\frac{dp}{dt} = cp(1 - p) - ep \quad (3)$$

La différence fondamentale avec la TIB est que la migration dépend de la proportion de patches occupés : plus elle est importante plus la migration est importante. Parmi

---

12. Les îles sont cependant souvent dans des archipels où la lecture de ces flux n'est pas si simple.

13. Richard Levins qui avec Heatwole est un des co-découvreurs des idées de la TIB.

les démonstrations il y a les travaux menés notamment par Ikkha Hanski sur les population du Mélitée du plantain (*Melitaea cinxia*) au sud-ouest de la Finland ([Hanski, 1998](#)). En plus de données un cadre de penser plus réaliste en terme de configuration spatiale, les dynamiques populationnelles associées sont bien comprises et mènent à des risques d’extinction mieux évalués ([Hanski, 1998](#)). C’est aussi un cadre approprié pour insérer l’étude des flux génétiques liés à l’arrangement spatial des populations. Ainsi, toujours sur ces mêmes populations de papillon, Ilik Saccheri et collègues montrent qu’en ajoutant le degrés d’hétérozygotie, ils obtiennent des prédictions précises quant l’extinction locale des populations ([Saccheri et al., 1998](#)). Les travaux théoriques autour du concept de métapopulations proposent un certain nombre de paradigmes qui permettent d’évaluer le rôle que jouent les processus de colonisation et d’extinction dans les variations spatio-temporelles de la démographie d’une espèce ([Leibold et al., 2004](#)). La prépondérance de ces mécanismes qui font la force de la TIB et de la théorie des métapopulations a été poussée à son paroxysme dans la théorie neutre de la biogéographie.

### **La théorie neutre de la biogéographie et le débat qu’elle soulève**

La théorie neutre postule l’équivalence écologique entre les différents individus d’espèces éventuellement différentes et décrit les dynamiques populationnelles reposant sur les différences d’abondance relative à l’échelle régionale et locale. Ainsi, en 1997, dans l’article fondateur de la théorie neutre, Stephen Hubbell décrit un modèle dans lequel le remplacement d’un individu mort dans une communauté locale est le résultat d’un tirage aléatoire : le nouvel individu peut soit être recruté localement et la probabilité que l’individu soit d’une espèce donnée dépend de l’abondance relative de cette dernière dans la communauté locale soit le nouvel individu peut-être un immigrant dont l’identité de l’espèce à laquelle il appartient est liée à l’abondance à l’échelle régionale de celle-ci ([Hubbell, 1997](#)). En plus des exemples donnés dans l’article de 1997, Hubbell montre de manière convaincante que dans la forêt tropicale du Panama, à la suite d’un chablis, le recrutement de l’arbre n’est pas prévisible par ces ca-

ractéristiques et que le recrutement est similaire à la composition alentour (Hubbell, 1999). La dynamique engendrée est appelée la dérive écologique, elle dominée par la stochasticité qui conduit presque certainement à l’extinction de toutes les espèces sauf une, ce qui est contrebalancée par l’apparition d’espèces nouvelles (Hubbell, 2010; Ricklefs, 2003).

La théorie neutre partage beaucoup de caractéristiques avec la TIB : les mécanismes fondamentaux sont l’extinction et la colonisation, l’hypothèse d’équivalence écologique et l’imbrication des échelles régionales et locales. Comme le fait remarquer Hubbell en 2010 dans le chapitre qu’il écrit dans *The Theory of Island Biogeography Revisited*, la théorie neutre place l’équivalence écologique au niveau des individus et non plus au niveau des espèces (Hubbell, 2010). Une conséquence directe revendiquée par Hubbell est que cette hypothèse explique la forme convexe des courbes de colonisation et d’extinction décrites par MacArthur et Wislon mais que n’explique pas leur modèle (voir 1 et Hubbell (2010)). Le principe d’équivalence et la place importante que prend le hasard dans cette théorie a soulevé de très vif débats et des démonstrations à charge contre la véracité de la théorie (voir par exemple McGill and Collins (2003) et Ricklefs (2003)). L’équivalence écologique doit, à mon sens, être comprise comme une abstraction de la singularité des espèces, une simplification de la diversité des systèmes biologiques, nécessaire à l’isolation d’une portion restreinte des phénomènes en cause dans la répartition géographique des espèces pour en évaluer le pouvoir explicatif. Bien qu’un certain nombre de cas d’études permettent de rejeter cette théorie (McGill and Collins, 2003; John et al., 2007), les défenseurs de la théorie neutre affirment qu’elle est tout aussi utile quand une étude en démontre la fausseté (Rosindell et al., 2012). La théorie neutre peut en effet être présentée comme une jauge qui mesure sur l’importance des processus de différenciation de niches (Wennekes et al., 2012). Ainsi, pour certaines communauté la dérive écologique est plus importante que dans d’autre et du point du vue de formalisme des solutions ont déjà été proposée pour dresser un continuum de la théorie neutre vers la théorie de la niche écologique (Gravel et al., 2006). Malgré les possibilités offertes par ces deux théories, elles occultent largement les interactions écologiques qui sont factuelles ; si les observations donnent crédit à ces théories, une théorie intégrative de la biogéographie doit

expliquer pourquoi.

## **Le rôle des interactions dans la distribution des espèces**

Ma thèse a pour objectif de trouver des leviers pour intégrer les interactions écologiques dans la TIB et comprendre comment elles affectent la répartition géographique des espèces et de comprendre où chercher les traces qu'elles pourraient éventuellement laisser dans les données d'occurrence des espèces. Comme je l'ai mentionné plus haut cette idée est très ancienne, Wallace le remarque dans son livre publié en 1881 :

« Both competition and predation appear now to be much more important in biogeography than people had formerly guesses »

Le problème de ces relations écologiques est leur spécificité, l'unicité de chacune d'entre elles, dont découle nos difficultés pour les prévoir bien que des travaux récents explorent des pistes prometteuses pour les prédire notamment sur la base de relations allométriques entre proie et prédateur ([Gravel et al., 2013](#)). Au point de vue théorique et à l'examen des chapitres du dernier livre de MacArthur ([MacArthur, 1972](#)), il apparaît que l'intégration des interactions est une étape clef pour aller vers une biogéographie intégrative et c'est dans cette direction que j'ai mené ma thèse, en essayant d'apporter des pistes pour arriver à une telle intégration.

## **Importance des interactions dans la distribution**

Dans la théorie de la biogéographie des îles, les interactions sont en fait omniprésentes car elles sont une des composantes principales du processus d'extinction. Cependant, dans la formulation du modèle, elles ne sont jamais mentionnées explicitement, cachées dans le taux d'extinction  $e$ . Comme je le montre à la figure 1, la différence dans l'allure des courbes

dessinées par MacArthur et Wilson et celles obtenues en supposant un taux d'immigration et de colonisation sont différentes. D'après les auteurs, l'immigration devient plus difficile lorsque les espèces s'accumulent sur l'île et les extinctions sont de plus en plus fréquentes dues à l'intensification des interactions. Pour parler en terme de réseau d'interaction, l'accumulation d'espèces sur l'île sature le réseau local et rend difficile l'intégration d'une nouvelle espèce qui le rend par ailleurs de plus en plus instable. Une interprétation en terme de communauté de la TIB est tout à fait possible mais les liens entre les espèces ne sont pas formulés mathématiquement en 1967.

Depuis les années 60, la littérature théorique n'a cessé de discuter le rôle joué par les interactions intra- et inter-spécifiques dans la distribution spatiale des espèces. Il est reconnu que l'interdépendance des espèces détermine le caractère favorable de l'environnement au sens large (biotique et abiotique). En 2009, Robert Holt et Michael Barfield discutent de l'impact de la prédation sur la répartition d'espèces en compétition insistant alors sur le rôle majeur des interactions dans le dessin des aires de répartition ([Holt and Barfield, 2009](#)). En 2012, William Godsoe et Luke Harmon Godsoe introduisent les interactions dans un modèle simple de distribution d'espèce et montre comment la probabilité de présence d'une espèce peut être affectée par la distribution d'une seconde et concluent alors que cela doit affecter vraisemblablement la qualité de prédictions des SDMs ([Godsoe and Harmon, 2012](#)). La remise en cause actuelle de ces modèles est une étape fondamentale car leur succès depuis la fin du siècle dernier a relégué les interactions écologiques au second plan en démontrant que la corrélation avec les variables climatiques étaient peut-être suffisante, au moins en première approximation pour expliquer les aires de répartitions ([Pearson and Dawson, 2003](#)). Pourtant, dès 1998, le travail précurseur d'Andrew Davis et collègues ([Davis et al., 1998](#)) avait fortement remis en question l'hypothèse d'indépendance des espèces ([Jeschke and Strayer, 2008](#)). L'expérience dont les résultats sont publiés en 1998 est une analyse d'abondance de trois espèces de drosophile le long d'un gradient de température. Les comparaisons d'abondance sont menées pour toutes les combinaisons possibles de ces trois mouches (assemblages à 1, 2 ou 3 espèces) mais aussi en présence ou en absence d'un parasitoïdes. La démonstration est



sans appel, la compétition et parasitisme affectent drastiquement la survie le long du gradient de température, les interactions affectent donc très probablement les réponses au changement climatique.

Plus récemment, on constate une grande motivation pour intégrer les relations écologiques dans les modèles de distribution d'espèces ([Kissling et al., 2012](#); [Guisan and Rahbek, 2011](#)). Une méthodologie récente appelée JSDM intègre les corrélations dans la présence des espèces pour améliorer les prédictions ([Pollock et al., 2014](#)). Néanmoins, ces efforts se heurtent à un manque de maturité des modèles et théories qui cherchent à rassembler distribution et interactions. Parmi les travaux récents, Franck Jabot et Jordi Bascompte ont rassemblé métacommunauté et écologie des réseaux souligner importance des relations écologiques dans la répartition géographique des espèces ([Jabot and Bascompte, 2012](#)). De même, Dominique Gravel et collègues introduisent en 2011 l'interdépendance proie-prédateur dans le modèle de la TIB menant aux prémices d'une théorie trophique de la biogéographie des îles ([Gravel et al., 2011](#)) préfigurée par Holt ([Holt, 2009](#)). Ces travaux essayent de dépasser l'hypothèse d'équivalence écologique en vue de faire des prédictions plus précises sur les compositions spécifiques attendues localement.

C'est dans la lignée de ces développements théoriques récents que s'inscrit mon premier chapitre de thèse. J'y ai montré comment l'intégration du concept de réseau écologique dans la TIB était possible tout en ajoutant la reconnaissance de performances plus ou moins importantes des espèces dans un contexte abiotique donné (niche écologique). Pour y arriver, j'ai montré l'intérêt de considérer les espèces sous la forme d'assemblage plutôt qu'une à une. Du point de vue technique, mon travail montre aussi qu'un retour aux processus stochastiques tels que ceux présentés en 1967 est une démarche puissante pour ajouter de nouveaux mécanismes dans la TIB.




Figure 2: **Intégration des interactions et des contraintes abiotiques dans la TIB.** Pour intégrer les interactions j'ai considéré non pas un ensemble d'espèces indépendant mais des espèces au sein d'un réseau décrit à l'échelle régional (a). Comme dans la TIB ces espèces peuvent être colonisée l'île (b), mais dans le modèle que j'ai développé, les taux de colonisation varient avec le long d'un gradient environnemental (c). Enfin, les interactions influencent les taux d'extinction locaux (d).

## Un problème d'échelle ?

En repartant de l'exemple classique de la ségrégation spatiale des tamias *Eutamias dorsalis* et *E. umbrinus* (Brown, 1971), j'ai précédemment mis en évidence qu'une information sur les interactions est contenue dans les aires de répartitions de ces espèces. Il y a cependant deux caractéristiques qui peuvent conduire à la rareté de ce type de lecture : la singularité de l'interaction et son caractère locale. Je reviens un peu plus bas sur la première propriété et m'arrête ici sur la seconde. Une idée dominante en biogéographie est que les interactions ont des rôles majeurs à l'échelle locale mais que leurs conséquences sont de moins en moins perceptible à mesure que l'on considère des échelles spatiales de plus en plus grandes (voir l'unique figure de McGill (2010)). Du point de vue théorique, c'est tout à fait ce qui est décrit dans la TIB car c'est à l'échelle locale que les interactions influencent l'extinction. Néanmoins, ces conséquences locales sont présentes sur l'ensemble de la distribution de l'espèce, il est alors pertinent de se demander pourquoi nous ne sommes pas capables de détecter les interactions en examinant les distributions d'espèces. En réalité, bien que cela soit rare, nous avons des preuves que cela est possible dans certains cas. En 2010, Nicholas Gotelli et collègues divisent l'avifaune danoise en différentes catégories fondées sur la similarité écologique et démontrent que les espèces d'une même catégorie sont très souvent significativement spatialement ségréguées (Gotelli et al., 2010). De même, en 2007, Risto Heikkinen et collègues avaient obtenu des performances accrues de leurs modèles statistiques par l'utilisation de la répartition de six espèces de pics pour expliquer la présence de quatre espèces de hiboux (Heikkinen et al., 2007). Dans cette même étude, le signal est plus fort quand les données sont sur de 10x10km que 40x40km en faveur d'une dépendance à l'échelle, récemment supportée par d'autres travaux (Belmaker et al., 2015). Ce qui est remarquable dans les travaux de Gotelli et de Heikkinen est que l'utilisation d'une connaissance biologique et écologique a permis de révéler une trace des interactions dans la distribution d'espèces.

La dépendance spatiale de la détection des interactions est facile à comprendre : en

examinant des données de présence à des échelles spatiales de plus en plus larges, le nombre d'espèces s'accumule (c'est le principe de la relation aire-espèce) menant à la dégradation de l'information potentielle. Cela signifie que l'information nécessaire pour déceler des empreintes laissées par les interactions sera fournie par des données à l'échelle relativement fine. Cependant, cela ne permet pas de conclure sur le rayon d'action de ces interactions. Pour dépasser la question spatiale, il faut aussi envisager l'impact de la nature des interactions sur la répartition géographique. Ainsi, en 2014, Miguel Araújo et Alejandro Rozenfeld ont prouvé théoriquement que les interactions positives (mutualisme) se propageaient davantage que les interactions négatives ([Araújo and Rozenfeld, 2014](#)). Par conséquent, la nature de la relation qui unit des espèces peut influencer la perte d'information contenue dans les aires de répartition. Suite à mes travaux sur l'intégration des interactions, je me suis penché sur un autre aspect qui peut influencer la perte d'information dans les données de présence : l'abondance des interactions. Au chapitre 2, je montre que les interactions directes et indirectes affectent les données de distributions mais aussi que l'abondance des interactions rend difficile de distinguer la co-occurrence d'espèces en interactions d'une co-occurrence aléatoire. Ce qui est encore plus intéressant, c'est que j'ai accumulé un certain nombre d'indices dans des données de présence et d'absence réelle qui semblent confirmer nos prédictions. Je discute de ces résultats dans le troisième chapitre de ma thèse.

En constatant que l'abondance des interactions peut justifier l'hypothèse d'indépendance des espèces, je soulève le même paradoxe que celui relevé par MacArthur dans son œuvre de 1972 ([MacArthur, 1972](#)) :

« A few decades ago it was fashionable for ecologists to study communities in the arctic on the grounds that these would be very simple communities and hence easy to understand. Many excellent ecologists still follow this belief, but there are others who feel that it may be easier to understand the extremely complex communities. This sounds paradoxical : How can more complex communities be easier to understand ? A possible answer might be that a complex community has

strong interactions among species so that the lives of the separate species are less independent than in a simple community. Where there is greater interdependence, patterns may be more conspicuous. »

Encore une fois, je déplace le problème car si l'interdépendance est importante pour des système simple, le problème est de prédire quand ces systèmes le sont. Autrement dit, il serait peut-être pertinent de situer les prédictions en biogéographie au niveau du réseau écologique. Le problème d'échelle n'est plus seulement spatial et temporel il est aussi un problème d'échelle biologique : individus, population, communauté ou réseaux ?

### **Vers une biogéographie énergétique**

Le problème d'échelle biologique est aussi un problème de catégorisation des espèces. J'ai suggéré que les prédictions étaient plus faciles pour des espèces généralistes que pour des espèces spécialistes. Malheureusement, le spectre est très large et plutôt balancé avec un continuum entre des espèces hyperspécialistes de d'autres très généralistes (Poisot et al., 2015a). On peut néanmoins espérer que la réduction des espèces à un certains nombres de traits (McGill et al., 2006; Poisot et al., 2015b) doublée d'une réduction des des réseaux à un certains nombre de propriétés puissent permettre des généralisations utiles dans notre compréhension de la distribution des communautés. Il m'apparaît aujourd'hui urgent que le niveau bon niveau de détail dans nos descriptions des systèmes écologiques soit trouvé afin de renforcer les fondements théoriques de la dynamique des aires de répartition.

Une piste prometteuse pour prolonger la recherche des propriétés est me semble-t-il de s'appuyer sur la nature profonde des espèces : des systèmes énergétiques qui se perpétuent. La lecture de la théorie de la dynamique du budget énergétique de Sebastian Kooijman (Kooijman, 2000) m'a été très profitable pour cerner les possibilités offertes par une telle approche. S'il est possible, comme le suggèrent les travaux de Kooijman, de dériver de manière précise un grand nombre de propriétés énergétiques des espèces sur leur masse et leur forme, alors

les espoirs sont grands de pouvoir trouver des règles d'assemblages fiables des communautés et donc de comprendre d'un point de vue mécaniste les extinctions locales. Ce sont les mêmes espoirs que ceux nourrit par la théorie métabolique de l'écologie qui rassemble des relations entre la taille des espèces et différentes de leurs propriétés ([Brown et al., 2004](#)) qui montrent en somme qu'il est possible d'aller au-delà de l'espèce ([Poisot et al., 2015b](#)). Mes réflexions sur l'intersection entre la TIB et une vision énergétique de l'écologie sont présentées au chapitre 4 de ma thèse, dans un chapitre qui se veut aussi comme une ouverture vers les projets de recherche que j'aimerais mener dans un futur proche.

## ARTICLE 1

### A PROPOS DES INTERACTIONS BIOTIQUES ET DES CONTRAINTES ENVIRONNEMENTALES À L'ÉCHELLE BIOGÉOGRAPHIQUE

#### 1.1 Résumé en français

##### 1.1.1 Contexte scientifique

En 1967, R. MacArthur et E. O. Wilson publient leur théorie de la biogéographie des îles. Comme mentionner dans l'introduction, la TIB est toujours le support de nombreux travaux de recherche récents, ce dont témoigne le livre édité par J. Losos et R. Ricklefs : *The Theory of Island Biogeography Revisited* publié en 2010. Ce livre souligne l'importance des travaux de R. MacArthur et E. O. Wilson et fait l'inventaire des questions qui restent à explorer. Parmi ces interrogations, celles qui concernent le rôle des relations trophiques dans la théorie, développée au sixième chapitre par Robert D. Holt.

C'est précisément sur ce sujet que portent les travaux de D. Gravel et collègue présentés dans l'article *Trophic Theory of Island Biogeography* publiée dans *Ecology Letters* en 2011. Dans cet article, les auteurs montrent comment les résultats de la théorie classique sont modifiés par la prise en compte des liens écologiques unissant proies et prédateurs. Cet article est également le point de départ de mon premier article de thèse. L'objectif fixé était de 1- généraliser à tous types d'interaction le travail de Gravel et collègues et 2- introduire les contraintes environnementales afin de comprendre dans quelle mesure les prédictions de la théorie classique en étaient affectées.

Pour y parvenir, la clé de mon travail a été de considérer les espèces de manière indépendantes, mais comme des assemblages, des communautés possibles. J'ai alors été capable de bâtir des probabilités de survie qui étaient dépendantes du réseau écologique présent

sur l'île. De même, les probabilités de colonisation des espèces du continent ont été reliées aux conditions environnementales de l'îles. Après avoir montré comment le modèle a été construit et donné des prédictions simples, nous nous sommes intéressés à des scénarios portant sur 10 espèces et pour des types d'interactions différents : mutualistes, prédation et compétitions le long de gradients environnementaux. Ce qui apparaît ressort de nos simulations est un portrait des impacts potentielles des interactions sur la distributions des espèces. Dépendamment de leur nature et de leur nombre, les interactions peuvent changer drastiquement la biodiversité attendue dans le cadre de la théorie classique. Cela pourrait avoir des conséquences majeures sur nos prévisions de richesse spécifique dans le contexte actuel des changements globaux.

### 1.1.2 Publication associée

Le travail réalisé a donné lieu à un article intitulé *On the integration of biotic interaction and environmental constraints at the biogeographical scale*. Il fut accepté pour publication au printemps 2015 dans le journal *Ecography*. La conception de l'article est le résultat de nombreux échanges entre tous les auteurs de l'article. J'ai développé le modèle et l'ensemble des scripts pour aboutir aux résultats finaux. Dominique Gravel a supervisé l'ensemble des étapes et est devenu le dernier auteur. David Mouillot et Nicolas Mouquet ont grandement contribué à la rédaction du manuscrit. Afin de promouvoir notre publication, j'ai rédigé un article court sur le blog de la revue *Ecography* accessible en ligne à l'adresse suivante: <http://www.ecography.org/blog/towards-integrated-theory-biogeography>.

### 1.1.3 Traduction du résumé de l'article publié

La biogéographie est concernée en premier lieu par la répartition spatiale de la biodiversité dont elle produit les scénarios de variation dans un contexte de changement environnemental. Les efforts déployés pour développer les modèles de distribution d'espèce



ont donné des outils qui, bien que prédictifs, sont restés corrélatifs et ont largement ignorés les interactions biotiques. Dans cet article, nous utilisons la théorie de la biogéographie des îles en tant que première approximation d'une dynamique d'assemblage des communautés locales dans un contexte de métacommunautés. Nous y superposons l'ensemble des types d'interactions et nous introduisons les contraintes environnementales sur la dynamique de colonisation et d'extinction. Nous développons une approche probabiliste reposant sur les chaînes de Markov et nous calculons les probabilités de réalisation des assemblages spécifiques plutôt que les probabilités de présence des espèces une à une. Nous considérons la distribution de richesse spécifiques pour les différents types d'interactions écologiques. De plus, nous illustrons également le potentiel de notre approche en étudiant l'interaction entre les différences de besoins écologiques, les interactions et la distribution de la biodiversité le long d'un gradient environnemental. Notre approche supporte l'idée que les recherches futures en biogéographie requièrent une intégration cohérente de plusieurs concepts écologiques dans une théorie unique afin de mener vers des innovations méthodologiques et conceptuelles comme le changement de distributions espèce-centrée vers des distributions communauté-centrée.

*Les sections qui suivent sont celles de l'article publié.*

## **1.2 Title**

On the integration of biotic interaction and environmental constraints at the biogeographical scale.

## **1.3 Authors**

Kévin Cazelles, Nicolas Mouquet, David Mouillot, Dominique Gravel.

## **1.4 Abstract**

Biogeography is primarily concerned with the spatial distribution of biodiversity, including performing scenarios in a changing environment. The efforts deployed to develop species distribution models have resulted in predictive tools, but have mostly remained correlative and have largely ignored biotic interactions. Here we build upon the theory of island biogeography as a first approximation to the assembly dynamics of local communities embedded within a metacommunity context. We include all types of interactions and introduce environmental constraints on colonization and extinction dynamics. We develop a probabilistic framework based on Markov chains and derive probabilities for the realization of species assemblages, rather than single species occurrences. We consider the expected distribution of species richness under different types of ecological interactions. We also illustrate the potential of our framework by studying the interplay between different ecological requirements, interactions and the distribution of biodiversity along an environmental gradient. Our framework supports the idea that the future research in biogeography requires a coherent integration of several ecological concepts into a single theory in order to perform conceptual and methodological innovations, such as the switch from single-species distribution to community distribution.

## 1.5 Introduction

Biogeography is concerned with the description of the distribution of biodiversity and understanding its underlying processes. The discipline is central to the simulation of future scenarios of biodiversity under climate change ([Thuiller et al., 2013](#)). The extensive development of statistical models of species distributions based on actual ranges and environmental data have provided valuable knowledge and predictions ([Kearney and Porter, 2004](#)), but often remain purely correlative. There is now consensus that future developments in biogeography will require solving critical limitations of species distribution models ([Kissling et al., 2012](#)) and incorporating explicitly biotic interactions and dispersal ([Lavergne et al., 2010](#)). This effort must be supported by theory in order to guide model development, maintain tractability and manage complexity. Developing a mechanistic theory of species distribution will require an integration of three fundamental principles and their interplay ([Thuiller et al., 2013](#)): 1) how local and regional dynamics are linked, 2) how species interact with the abiotic environment and 3) how they are embedded in a network of biotic interactions. Each of these principles are discussed in detail below.

A cornerstone of biogeography is the recognition of the contribution of regional-scale processes such as disturbances, historical contingencies (e.g. macro evolutionary history or glaciations) and dispersal limitations to local community dynamics ([Ricklefs, 1987](#)). The metacommunity concept has been proposed as a simple framework to link different spatial scales in ecology ([Leibold et al., 2004](#)). It emphasizes reciprocal feedbacks between local scale processes, such as competitive interactions and local adaptation, and regional scale processes such as dispersal, gene flow, and speciation. A central concept of metacommunity ecology is the idea that local communities are highly dynamic owing to colonization events and local interaction, resulting in a spatial mosaic of assemblages sampled non-randomly from the regional species pool. As the concept matures there are new themes emerging, such as the investigation of evolution in metacommunities ([Urban et al., 2008](#)), and spatial food webs ([Massol et al., 2011](#); [Gravel et al., 2011](#)). The field provides remarkable concepts and

tools to build an integrated theory for biogeography.

Species distribution is also constrained by physiological requirements, which is at the core of the niche concept (Peterson et al., 2011). The niche is usually defined as a N-dimensional environmental and resource hyper-volume within which a species is able to maintain a viable population over the long term (Chase and Leibold, 2003). Recent developments refined this definition based on demography and metapopulation dynamics (Holt and Barfield, 2009). The abiotic niche, often referred as the Grinnelian niche, has been central to the development of species distribution models (SDMs, Jeschke and Strayer, 2008). Despite all of its criticisms, SDMs remain remarkably popular and operational for conservation ecology (Guisan et al., 2013). Recent attempts to improve the quantification of the niche include the addition of experimental assessments of the fundamental physiological constraints, as well as dispersal and proxies of biotic interactions (Boulangeat et al., 2012). The search for the most adequate set of environmental variables explaining diversity should be continued despite criticisms of the actual SDMs, and most of all must constitute a central principle of a general theory for biogeography.

Finally, species are not isolated, they are embedded within complex networks of ecological interactions. While interactions define community ecology, they are less informative for biogeography (Peterson, 2003). Theory predicts that interactions in small community modules (2-4 species) should influence range limits (Gilman et al., 2010), but there is no extension to highly diverse communities. It has been hypothesized that factors determining distribution are hierarchical, such that climate would govern the distribution at the regional scale while biotic interactions would be more important at the local scale (Araújo and Rozenfeld, 2014). However an increasing number of studies emphasizes the role of local interactions as a major factor influencing geographical ranges (Jabot and Bascompte, 2012; Gotelli et al., 2010). The representation of interactions in a network is a convenient method to summarize the type and strength of interactions among species, their organization (Proulx et al., 2005) and their consequences on dynamics (Allesina and Tang, 2012). Food webs were first considered in the

development of a trophic theory of biogeography (Gravel et al., 2011), where it was shown that a diversity of interactions enhance persistence. Networks are however more than food webs and are rarely made of a single type of interaction (Kéfi et al., 2012). Mutualism, competition and indirect effects (Wootton, 1994), for instance, also impact local environmental suitability (Godsoe and Harmon, 2012). Tools and knowledge acquired through the study of local ecological networks, such as the community matrix and metrics of structure (Allesina and Tang, 2012), must now be incorporated into a theory for biogeography.

These three principles should be all mixed together to provide an integrated assessment of their relative contribution to species distribution. To do so, the theory of Island Biogeography (hereafter referred as TIB) (MacArthur and Wilson, 1967; Warren et al., 2015) is a convenient starting point. The TIB describes variations of species richness among islands as a dynamic equilibrium between two opposite processes, colonization and extinction, directly linked with island characteristics. The TIB is a metaphor that goes beyond the intrinsic interest of islands; it serves as a first approximation to understanding the assembly of local communities embedded in a metacommunity context with straightforward species flux. The simplicity of the model and the relevance of its predictions demonstrate after more than 50 years since its publication it is still a useful tool in ecology and conservation (Cook et al., 2002; Warren et al., 2015). The TIB emphasizes the role of regional processes to local community assembly. Indeed it can be regarded as the simplest representation of metacommunity dynamics (Leibold et al., 2004). Furthermore, the model is easily expandable. Following (Holt and Barfield, 2009), Gravel et al. (2011) introduced trophic interactions in the TIB (hereafter the trophic TIB, TTIB;). Species interactions were found to be a key factor to understand species distributions, as the probability of finding any species in a locality increases with the generality of its diet and decreases with trophic rank.

We propose to generalize of the TIB by integrating the three principles described above. The TIB already explicitly includes the effect of regional processes (colonization and extinction dynamics) on local community assembly, and the TTIB includes predator-

prey interactions. We extend this framework to all potential interactions, thus resulting in a general model of metacommunity dynamics, akin to the Lotka-Volterra equations for local community dynamics. We also incorporate abiotic constraints on colonization and extinction dynamics. Hence we integrate the ingredients we believe are essential to model biodiversity distribution at the biogeographical scale. With this model in hand we then describe species distribution along environmental gradients. We use the mathematical formalism of Markov Chains ([Kemeny and Snell, 1983](#); [Black and McKane, 2012](#)) to derive expected assemblages and co-distribution at both the local and the regional scale. We illustrate how the interplay between biotic interactions and environmental requirements can affect the distribution of biodiversity over environmental gradients. Our results support the idea that the future research in biogeography require a consistent integration of several ecological concepts into a single framework to build promising approaches such as the switch from single-species distributions to community distributions.

## 1.6 The model

### 1.6.1 A simple probabilistic biogeographical model

The challenge of adding species interactions within the classical model of the TIB is gaining generality without losing simplicity. Following MacArthur and Wilson's theory, we model the dynamics of occurrence probability of a species  $i$  in a local community. Species occurrence is the result of a balance between colonization and extinction dynamics, which occur at rates  $c_i$  and  $e_i$  respectively,. Local species richness is given by the sum of occurrence probabilities over all species of the regional species pool  $P$ , here simply defined as the set of all species whose propagules (as defined in [Simberloff and Wilson, 1969](#)) can land on the island considered. The model thereby takes into account local (extinction) and regional (colonization) processes. More precisely, the dynamics of occurrence probability of species

i,  $p_i$ , follows:

$$\frac{dp_i}{dt} = c_i(1 - p_i) - e_i p_i \quad (1.1)$$

Here,  $c_i$  and  $e_i$  are constant and a property of species  $i$ . In this widespread version of the TIB, also called the linear version of the TIB ([Schoener, 2010](#)), the equilibrium occurrence probability of a species  $i$  is given by  $p_{i,eq} = \frac{c_i}{e_i + c_i}$ . Also, species are assumed to be independent, therefore, the richness  $S_{eq}$  is given by the sum of the  $P$  different  $p_{i,eq}$ . The linear TIB can be modified to include trophic interactions (after [Gravel et al., 2011](#)) and we propose to extend it to all types of interactions. To reach that goal, the first step is to find a way to derive the expected species composition at any time. This composition can actually be depicted at any time by a vector of  $P$  zeroes and ones indicating, respectively, presences and absences of each species considered, these combinations will be referred as assemblages. Following Mac-Arthur and Wilson, we use a stochastic modelling approach to describe the dynamics of assemblages. The simplest scenario is the one species case. Here there are only two assemblages for the locality: one with species  $i$  present and the other without. Let  $X_i$  be a random variable describing the occurrence of species  $i$ . When species  $i$  is present in the locality,  $X_i$  is 1, when it is absent  $X_i$  is 0;  $X_i$  is then a Bernoulli variable. We define this random variable at any time  $t$  which describes a stochastic process we denote  $\mathbf{X}_{i,t>0}$ . The occurrence probability of species  $i$  at time  $t + dt$  ( $dt$  being a very small time step) is then given as follows:

$$\begin{aligned} \mathbb{P}(X_{i,t+dt} = 1) &= \mathbb{P}(X_{i,t+dt} = 1 | X_{i,t} = 1) \mathbb{P}(X_{i,t} = 1) \\ &\quad + \mathbb{P}(X_{i,t+dt} = 1 | X_{i,t} = 0) \mathbb{P}(X_{i,t} = 0) \end{aligned} \quad (1.2)$$

$\mathbb{P}(X_{i,t+dt} | X_{i,t})$  is the conditional probability describing  $X_{i,t+dt}$  stating  $X_{i,t}$ . As  $X_{i,t+dt}$  solely depends on  $X_{i,t}$  (not on other earlier time steps) we have a discrete-time Markov chain. In this process, species  $i$  will be present in a locality at time  $t + dt$  if it was already present at time  $t$  and persisted (meaning it did not go extinct, with probability  $(1 - e_i dt)$ ), or if it was absent

and colonized the community from the mainland (with probability  $c_i dt$ ). Note that  $dt$  is small enough to get  $0 < c_i dt < 1$  and  $0 < e_i dt < 1$ . Hence, equation (1.2) becomes:

$$\mathbb{P}(X_{i,t+dt} = 1) = c_i dt \mathbb{P}(X_{i,t} = 0) + (1 - e_i dt) \mathbb{P}(X_{i,t} = 1) \quad (1.3)$$

This equation leads to (1.1) when  $dt$  tends to zero. This formulation keeps the simplicity of the original MacArthur and Wilson model, but can also more generally be used to consider the probability of any given assemblage.  $\mathbb{P}(X_{i,t+dt}|X_{i,t})$  defines the rules to switch from one assemblage to one another during the interval  $dt$ . There are  $P$  occurrence probabilities we gather within  $\mathbf{Y}_{t>0} = (\mathbf{X}_{1,t>0}, \mathbf{X}_{2,t>0}, \dots, \mathbf{X}_{P,t>0})$  which leads to the description of  $2^P$  assemblages depicted by a given collection of zeros and ones. The conditional probabilities provide the transition from one local assemblage  $k$  to any other  $l$  during  $dt$ . For any species  $i$  there are only four possible cases: at time  $t$  either species  $i$  is locally absent and colonizes the locality ( $I_1$ ) or not ( $I_2$ ) during  $dt$ , either species  $i$  is present and goes extinct ( $I_3$ ) or survives ( $I_4$ ) during  $dt$ . The conditional probabilities between two communities states ( $l$  and  $k$ ) can then be simply derived from these four probabilities:

$$\begin{aligned} \mathbb{P}(\mathbf{Y}_{t+dt} = \text{"state k"} | \mathbf{Y}_t = \text{"state l"}) &= \prod_{i_1 \in I_1} c_{i_1} dt \prod_{i_2 \in I_2} (1 - c_{i_2} dt) \\ &\quad \prod_{i_3 \in I_3} e_{i_3} dt \prod_{i_4 \in I_4} (1 - e_{i_4} dt) \end{aligned} \quad (1.4)$$

We now apply the complete probability formula as defined in (1.2) to get the probability of observing one assemblage at  $t + dt$  given its state at  $t$ . This is where the main benefit of Markov chain models is: it allows us to derivate exact solutions for the probabilities for assemblages, instead of a set of independent occurrence probabilities for each species. This approach is promising for building joint species distribution models (see Discussion). This property will be fully explored in the next section to include interactions.

Consider as an example a pool of two species ( $P = 2$ ) for which we find four assemblages: at any time  $t$ , a locality can contain either two species ( $X_{1,t} = 1, X_{2,t} = 1$ ), only one



species ( $X_{1,t} = 1, X_{2,t} = 0$ ) and ( $X_{1,t} = 0, X_{2,t} = 1$ ), or none of them ( $X_{1,t} = 0, X_{2,t} = 0$ ). The transition from one local assemblage to another is then easily obtained. Table 1 presents these conditional probabilities (application of (1.4)). This is actually the transition matrix of a Markov chain we solve (by calculating one eigen value, see below). To illustrate the dynamics expected in TIB from our assemblage point of view, we simulate the model as follows:  $c_1 = c_2 = 0.15$ ,  $e_1 = e_2 = 0.05$ ,  $\mathbb{P}(X_{1,0} = 0, X_{2,0} = 0) = 0.6$  and  $\mathbb{P}(X_{1,0} = 1, X_{2,0} = 0) = 0.4$ , so species 2 is absent at time  $t = 0$ . Just as for the single species situation, the probabilities of observing each community tend to an equilibrium (Fig.3, panel A). By summing the previous probabilities where a given species (1 or 2) is present (the conditional probabilities) we get its overall occurrence probability (marginal probability, Fig.3-B). Finally, we can calculate the expected number of species in a locality (Fig.3-C), in agreement with the TIB. Interestingly, this calculation is often achieved in the other way. Firstly, the presence probability of all species are computed:  $\mathbb{P}(X_i) = \frac{c_i}{c_i + e_i}$ . Then the richness is obtained under the assumption that species are independent and so  $P(X_i, X_j) = \mathbb{P}(X_i)\mathbb{P}(X_j)$ . We show below that occurrence probabilities of each assemblage is a key to introduce interactions among species.

### 1.6.2 Integrating biotic interactions

We start by representing the interaction network by a community matrix  $\mathbf{A}$  of  $P$  species that we incorporate into the Markovian TIB chain model. The elements  $\alpha_{i,j}$  of  $\mathbf{A}$  quantify the effect of species  $j$  on the dynamics of species  $i$ . We first consider that interactions could alter both the colonization and the extinction probabilities (Gravel et al., 2011). When  $\alpha_{i,j}$  is negative, the colonization probability of species  $i$  decreases and/or its extinction probability increases when  $j$  is found locally. Inversely, when  $\alpha_{i,j}$  is positive, the colonization probability increases and/or the extinction probability decreases. Note that diagonal elements provide the extinction probability per time unit when no other species is present.

The elements of the community matrix  $\mathbf{A}$  represent the pairwise effects of ecological interactions on transition probabilities. To account for the cumulative effects of local interac-

$(\mathbf{X}_{1,t}, \mathbf{X}_{2,t})$	$(\mathbf{X}_{1,t+dt}, \mathbf{X}_{2,t+dt})$			
	(0,0)	(0,1)	(1,0)	(1,1)
(0,0)	$(1 - c_1dt)(1 - c_2dt)$	$(1 - c_1dt)c_2dt$	$c_1dt(1 - c_2dt)$	$c_1dte_2dt$
(0,1)	$(1 - c_1dt)e_2dt$	$(1 - c_1dt)(1 - e_2dt)$	$c_1dte_2dt$	$c_1dt(1 - e_2dt)$
(1,0)	$e_1dt(1 - c_2dt)$	$e_1dte_2dt$	$(1 - e_1dt)(1 - c_2dt)$	$(1 - e_1dt)c_2dt$
(1,1)	$e_1dte_2dt$	$e_1dt(1 - e_2dt)$	$(1 - e_1dt)e_2dt$	$(1 - e_1dt)(1 - e_2dt)$

Table 1: **Conditional probabilities between potential assemblages.** At any time  $t$  we calculate all the possible conditional probabilities between the four potential assemblages for a two species regional pool. These probabilities are derived by multiplying probabilities of single species events defined in equation (1.4). By doing so, we build the transition matrix of our markov chain where species are assumed to be independent. We release this hypothesis by linking extinction coefficients and species assemblages.

tions on transition probabilities, we make colonization and extinction probabilities community dependent. As explained above, at a time  $t$ , the  $\mathbf{Y}_t$  vector gives the local assemblages. We calculate the sum of interactions at any time and for each species as  $\mathbf{v} = \mathbf{A}\mathbf{Y}_t^T$  (where  $T$  denotes the transpose operator). Our approach can be interpreted as a spatial analogue to the generalized Lotka-Volterra model because it takes into account the impact of the whole network of interactions on each species dynamics and can deal with any type of interaction. We denote the coefficients of  $\mathbf{v}$  by  $v_i$ , they are species-specific parameters (weighted by parameter  $d_i$ ) of two species-specific functions:  $f_i$  and  $g_i$ , respectively, standing for extinction and colonization probabilities for species  $i$ . Note that at this stage we do not define any specific function relating interactions to colonization ( $f_i$ ) and extinction probabilities ( $g_i$ ), to keep the description of the model general (see below for some proposed functions). At each time step, the local community composition impacts: i) the colonization probability of species present in the regional pool but absent from the local community, and ii) the extinction probability of species present on the local community.

If we expand the two species example (labelled 1 and 2, Table 1), according to the general model, we define two  $f$  functions ( $f_1$  and  $f_2$ ) linking interaction and extinction and two  $g$  functions linking extinction and colonization ( $g_1, g_2$ ). At this stage, to reduce the model's

complexity, we consider that interactions solely impact extinction probabilities. This assumption is reasonable if we consider that local interactions impact mostly demography (possibly leading to extinction) and that colonization success solely depends on the first propagule (interactions occur after arrivals). Therefore  $g_1$  and  $g_2$  are constant functions, respectively, returning  $c_1$  and  $c_2$ . The functions  $f$  are assumed to have a sigmoid shape (1.5). There are many reasons such a function is of interest: 1) we get a clear link with the basic extinction probability, i.e.  $e_i$  for an interaction strength of 0; 2) we define both a minimum and a maximum extinction probability; 3) the first interactions to occur are the most influential (Gravel et al. (2011) considered that at least one interaction was required to persist, which is very similar).


$$f_i(\mathbf{v}) = f(\mathbf{v}, (e_i, e_{i,min}, e_{i,max}, d_i))$$

$$= e_{i,min} + \frac{1}{\frac{1}{e_{i,max}-e_{i,min}} + \left( \frac{1}{e_i-e_{i,min}} - \frac{1}{e_{i,max}-e_{i,min}} \right) \exp(d_i * v_i)} \quad (1.5)$$

$$g_i(\mathbf{v}) = c_i \quad (1.6)$$

To illustrate how interactions modify occurrence probabilities, we simulate the model for two networks:  $A_1$  where all interactions are negative and  $A_2$  where they are all positive. We consider null diagonal elements for both networks. Consequently, there is no difference with the model without interaction when one species is alone in the locality. Simulation results are presented at Figure 4. Panel A presents the functions  $f_1$  and  $f_2$  we chose for our two species example. For networks  $A_1$  and  $A_2$ , we show how interactions alter the probabilities of observing different assemblages (respectively, Fig.4-B and Fig.4-C). The assemblage with both species present (solid red lines) is the most affected by interactions, switching from an occurrence probability of 0.2 (for negative interactions) to 0.8 (for positive interactions). Positive interactions enhance, as expected, co-occurrence while negative interactions prevent species from being found on the same island. Consequently, occurrence probabilities of single species states are lower in  $A_2$  than in  $A_1$ . According to a defined network, occurrence probabilities of the different assemblages are then modified, which affect the expected species

richness (Fig.4-D).



**Figure 3: Dynamics of the community assembly.** As a direct development of MacArthur and Wilson model, we simulate for two species, the dynamics of the four communities states with different initial conditions associated (A). By summing every states where one given species is present we get the occurrence probability of two considered species (B). Finally by summing the four states probabilities weighted by their species richness, we get the classical model of MacArthur and Wilson (C). The calculation of (B) and (C) does not require species being independent while classical approaches focus on (B) to derive (A) and (C) under this assumption of independence.

### 1.6.3 Integrating environmental gradients

We now introduce the effect of abiotic conditions, such as climatic variables, on transition probabilities. We denote the vector of  $n$  environmental conditions by  $\mathbf{w}$ :  $\mathbf{w} = (w_1, w_2, \dots, w_n)$ . We first assume that physiological constraints can affect both colonization and extinction probabilities through the functions  $f_i$  and  $g_i$  (affecting, respectively, extinction and colonization rates). Again the model in its general formulation does not presume any shape for these functions. We now have all the ingredients of an integrated model of biogeography as the transition probabilities at a location depend on 1) species-specific colonization and existence probabilities, 2) the network of interactions, 3) local community composition, and 4) local environmental conditions. In the general formulation of the model, functions  $f_i$  and  $g_i$  are functions of multiple variables ( $\mathbf{v}$  and  $\mathbf{w}$ ).

At any time  $t$ , for a regional pool of  $P$  species among which interactions are summarized by the community matrix  $\mathbf{A}$ , in an environment characterized by  $\mathbf{w}$ , we can derive all transition probabilities. These constitute a transition matrix of a Markov chain that we denote  $\mathbf{M}(\mathbf{v}, \mathbf{w})$ . Its elements,  $\mu_{k,l}(\mathbf{v}, \mathbf{w})$ , give the probability the locality in assemblage  $k$  turns into assemblage  $l$  (left side of equation (1.4)):

$$\begin{aligned} \mu_{k,l}(\mathbf{w}, \mathbf{v}) = & \prod_{i_1 \in I_1} g_{i_1}(\mathbf{v}, \mathbf{w}) dt \prod_{i_2 \in I_2} (1 - g_{i_2}(\mathbf{v}, \mathbf{w}) dt) \\ & \prod_{i_3 \in I_3} f_{i_3}(\mathbf{v}, \mathbf{w}) dt \prod_{i_4 \in I_4} (1 - f_{i_4}(\mathbf{v}, \mathbf{w}) dt) \end{aligned} \quad (1.7)$$

Note that the dimension of  $\mathbf{M}(\mathbf{w})$  will increase as a power of the number of species  $P$  and thus can rapidly becomes large. Let  $\mathbf{C}_t$  be the line vector of the probability of observing each assemblage, defined by:  $\mathbf{C}_t = (\mathbb{P}(\mathbf{Y}_t = \text{"state 1"}), \mathbb{P}(\mathbf{Y}_t = \text{"state 2"}), \dots, \mathbb{P}(\mathbf{Y}_t = \text{"state } 2^P\text{"}))$ . The Markov Chain formalism defines the probability of the future community composition at time  $t + dt$  as  $\mathbf{C}_{t+dt} = \mathbf{C}_t \mathbf{M}$ .  $\mathbf{C}_t$  asymptotically reaches the  $\mathbf{C}_{eq}$  after a certain number of time steps.  $\mathbf{C}_{eq}$  is given by the normalized left eigenvector associated to the first left eigenvalue.

$$\lim_{\substack{l \rightarrow +\infty \\ l \in \mathbb{N}}} \mathbf{C}_0 \mathbf{M}^l = \mathbf{C}_{eq} \quad (1.8)$$

$\mathbf{C}_{eq}$  contains the probability of all assemblages at the equilibrium. The occurrence probability of a given species, is provided by the sum of all probabilities of assemblage where that species is present. The richness at the equilibrium  $S_{eq}$  is the sum of  $\mathbf{C}_{eq}$  elements weighted by the number of species found in the associated assemblages.

For the sake of illustration, we further reduce the complexity of our model. We have previously removed the interactions ( $\mathbf{v}$ ) from colonization ( $g$ ) functions; we now state that extinction does not depend on environmental variables and so we remove the abiotic environment ( $\mathbf{w}$ ) from extinction functions ( $f$ ). This can be interpreted as the effects of the abiotic environment on extinction rate being included within  $e_i$  (i.e. extinction rate without interaction). Furthermore, we assume solely one environmental variable and a Gaussian shape for


$g_i$  functions (1.9). A simple function with a clear optimum and very low colonization for extreme environment values is.

$$g_i(w_1) = g(w_1, (c_i, h_i, r_i)) = c_i * \exp\left(-\left(\frac{w_1 - h_i}{r_i}\right)^2\right) \quad (1.9)$$

This enables us to define an environmental optimum ( $h_i$ ), a colonization probability per time unit ( $c_i$ ) and also suitable range ( $r_i$ ) for each species. Figure 5 presents the interplay between the three components of the integrated biogeographical model. The chosen functions for the environment-colonization relationship are illustrated in Panel A. For the two previous networks ( $A_1$  and  $A_2$ ; illustrated in Fig 4) we now compute the probabilities of observing the different assemblages at equilibrium, along the environmental gradient (Panel B and C). When interactions are negative (network  $A_1$ ), species repulse each other and rarely co-occur, whatever the environment is. Most of their occurrence follow their abiotic niche (blue and green lines) as they are barely found together. Inversely, when interactions are positive (for  $A_2$  network) they often co-occur where their abiotic niches overlap, thereby decreasing the probability of an empty community (Panel D, solid grey line). Finally, we present how interactions modify the resulting community composition along the environmental gradient (Panel D). Species richness is constrained by the distribution of abiotic niches and the sign of the interactions. As expected, the role of interactions is strongest when abiotic niches largely overlap.

## 1.7 Exploring the model

In our exploration, we choose a regional pool  $P$  of 10 species to keep the number of assemblages reasonable ( $2^{10} = 1024$ ) and to numerically compute the exact solution of the equilibrium distribution  $\mathbf{C}_{eq}$ . We consider four types of interaction matrices  $\mathbf{A}$ . The first situation corresponds to the classical MacArthur and Wilson model, where the  $\mathbf{A}$  matrix is null (no interactions). For the three other scenarios we generate random matrices with fixed



**Figure 4: Effects of biotic interactions on colonization-extinction dynamics.** For any species  $i$ , the extinction probability  $e_i$  is related to the strength of the interaction as shown in (A). The intersections of extinction curves with the grey dotted lines indicate the potential values of  $e_i$  according to the different biotic context ( $A_1$ ,  $A_2$  and without interaction). We set the other parameters as follows:  $c_1 = c_2 = 0.15$ ,  $\mathbb{P}(X_{1,0} = 1, X_{2,0} = 0) = 0.4$ ,  $\mathbb{P}(X_{1,0} = 0, X_{2,0} = 0) = 0.6$ ,  $\mathbb{P}(X_{1,0} = 0, X_{2,0} = 1) = \mathbb{P}(X_{1,0} = 1, X_{1,0} = 1) = 0$ . We then simulate the model for two simple networks  $A_1$  and  $A_2$  and present community assembly dynamics associated ((B) and (C)). Finally we compare the expected species richness on the locality (D) for our two networks and for the case without interaction which corresponds to the widespread linear version of the TIB.

connectance (number of existing links divided by the number of potential links). The coefficients within  $\mathbf{A}$  are drawn uniformly within  $[0, 1]$  and the sign of the interaction is determined by the action of one species on another, for instance, a predator has a negative impact on its prey leading to a negative  $\alpha$  coefficient; in return, a prey has a positive effect on its predators. The intensity of the interaction is then determined by the  $d$  coefficient of extinction functions (see equation (1.8)). We assume that the distribution of the links are given by the niche model (Williams and Martinez, 2000). This model is simple and provides relevant random food webs with the same number of positive and negative interactions. For the two last scenarios, we keep the rules to distribute the links, but turn all the coefficients in  $\mathbf{A}$  positive to generate a mutualism network, or negative for competition networks. Although these basic structures with exclusive interaction types are not realistic, they facilitate comparison among results. Hence, the scenarios simply differ by the sign distribution within the matrix  $\mathbf{A}$ : (i) no interaction  $\mathbf{A}$  is null, (ii) predation mixes both signs “+/-”, (iii) mutualism only “+”, (iv)-competition, only “-”. With these scenarios in hands, we 1) present the assemblages probabilities associated with a given level of species richness and 2) we look at the species richness expected along an environmental gradient. For all figures presented hereafter we used 1000 randomly-generated  $\mathbf{A}$  matrices.

### Assemblage probabilities

First, we illustrate how interactions affect richness of species assemblages. To do so, we build the Markov chains for all the 1000  $\mathbf{A}$  matrices generated (connectance set to 0.2) and we calculate the vector  $\mathbf{C}_{eq}$ . This is a vector of 1024 occurrence probabilities (as we consider 10 species). Then we sum all the probabilities that correspond to assemblages of the same richness. We do so for three values of  $d$  coefficient (0.1, 1 and 10); that is, we look at how the strength of interaction affect community richness predictions. Figure 6 presents the results of such investigation, with Panels A to C corresponding to the results for the three different values of the  $d$  parameter.




As expected, positive interactions increase local species richness by diminishing extinction probabilities, while negative interactions weaken large communities (see the contrast between blue and red symbols on Fig. 6). This is stressed as interaction strengths increase, that is for increasing values of  $d$ . Indeed, when  $d$  is low, there is almost no difference among scenarios because interactions do not impact strongly colonization and extinction dynamics; occurring species can be regarded as mostly independent. All scenarios converge to the classical TIB scenario (no-interaction, grey symbols), the resulting species richness distribution is binomial (here for all species  $p_{i,eq} = 0.5$  as  $c_i = e_i = 10^{-5}$ ). Differences between interaction types increase with  $d$ . Species rich mutualistic communities are more likely to occur since positive interactions tends to promote co-occurrence. Therefore species occurrence can be dramatically affected by the strength of interactions: for  $d = 10$  (Panel C in Fig. 6), the species richness is 9.46 for positive interactions (red symbols), 2.24 for the negative ones (blue symbols) and 5 without interactions. When positive and negative interactions are mixed (our predation scenario, green symbols on Fig. 6), it seems that the negative effect of predators on their prey prevails and so predation reduces species richness, but less than for competitive networks.

As we introduce variability through the use of randomly-generated matrices, we also compute the standard deviation associated with occurrence probabilities. The variability is provided as the coloured vertical bars found in Fig. 6 which stand for 50% of the total standard deviation. Clearly, variability increases with (i) the strength of interaction and (ii) the occurrence probability. Although this can simply reflect the variability of values found in  $\mathbf{A}$  matrices, this could potentially be caused by the variability of the location of non-zero values in  $\mathbf{A}$  matrices; that is, the structure of the networks we use.

### 1.7.1 Biodiversity distribution over environmental gradients

In this section, we introduce an environmental gradient to emphasize the interplay between interactions species-specific requirements along an environmental gradient. Our en-



**Figure 5: Equilibrium for interacting species along an environmental gradient.** The colonization probability of species  $i$ ,  $c_i$ , is related to the environment variable  $\mathbf{w}$  according to species-specific requirements (A). The intersection of the colonization curve of species  $i$  with the grey dotted lines represents the value of  $c_i$  associated with its environmental optimum  $h_i$ . We compute equilibrium occurrence probabilities for the different assemblages along the environmental gradient, for the networks  $A_1$  with negative interactions (B) and  $A_2$  with positive one (C). We calculate the expected species richness on the locality for the two networks and without interaction (D).

environmental gradient takes values from 0 to 30, for each of them we calculate the expected species richness associated to all scenario. To do so, we start by computing the colonization functions ( $g_i$  functions): species optima  $h_i$  are drawn from a uniform distribution from the range [10, 20] and the widths of the abiotic niches are kept constant for all the simulations  $r_i = 5$ . Then we build the Markov chains for the different values of the environmental gradient and for the different  $\mathbf{A}$  matrices. Again, we derive the vector  $\mathbf{C}_{eq}$  and we sum its elements,


i.e. occurrence probabilities of assemblage community, weighted by the species richness to which they refer. We repeat the procedure for an increasing value of connectance of **A** matrices: from 0 to 0.4. For this section, the parameter  $d$  is set to 10, also extinction parameters are set as follows:  $e_i = 10^{-5}$ ,  $e_{i,min} = 10^{-3}e_i$ ,  $e_{i,max} = 10^3e_i$  and  $c_i = 10^{-5}$ . Like so we obtain the profile of species richness we report on Figure 7.

For all scenarios, the richness is maximal at the center of the environmental gradient (Fig. 7). This is due to the distribution of species optima in the range [10, 20]. Also this is the range of environmental values for which the effect of interaction are the most important. Indeed, the higher the colonization probabilities, the higher interactions occur, therefore, interactions strongly impact species richness for favourable abiotic conditions. We also find that changes in species richness increase with connectance, as depicted by the colour of the solid lines for the three panels of Fig. 7: from black (without interaction) to the lightest blue (connectance set to 0.4).

Species richness is inversely related to connectance when interactions are negative (Panel A in Fig. 7). Moreover, when abiotic conditions are favourable, the number of species expected tends to 1. At the centre of the gradient, even though colonization probabilities are maximal, many species colonize but likely go extinct because of competition. We expect the locality to be most often occupied by species that are not affected by competition. Alternatively, in the case of positive interactions (Panel B in Fig. 7), the expected species richness is strongly enhanced by interactions even for low connectance. The expected species richness tends to reach the total number of species from the most favourable to semi-harsh abiotic conditions. As the connectance increases the Gaussian shape of the richness profile turns into a hat shape, which has one major consequence: from favourable to semi-harsh conditions, the species richness is maintained thanks to positive interaction, but it also quickly collapses as the environment becomes slightly harsher.

Finally, when positive and negative interactions are mixed, the higher the connectance, the flatter the richness profile (Panel B in Fig. 7). The expected species richness declines as

connectance increases but far less than it does for negative interactions only. We think this is caused by the colonization of numerous prey that promote the survival of predators which in turn prevent assemblages to be as large as they can be without interaction (as predators reduce the persistence of prey). Conversely, from harsh to intermediate environmental conditions, mixed sign interactions positively affect the species richness. We explain this as the consequence of the benefit predators take from the preys presence. Assemblages with few predators, promoted by positive effect of the prey on their predators, may be relatively stable. Since colonization is low, this assemblage may enhance species richness over time but they may also collapse as soon as an extra predator colonizes the island.



**Figure 6: Probabilities of species richness for different types of interaction.** We compute expected species richness at the equilibrium with the following set of parameters:  $e_i = 10^{-5}$ ,  $e_{i,min} = 10^{-3}e_i$ ,  $e_{i,max} = 10^3e_i$  and  $c_i = 10^{-5}$ . We do so for three different interaction weights:  $d = 0.1$  (A),  $d = 1$  (B),  $d = 10$  (C). In each panel, the four colours stand for the following types of networks: no interaction (grey), predation (green), competition (blue) and mutualism (red). Probabilities associated to coloured points are the means calculated for 1000 interaction matrices randomly-generated according to the niche model (Williams and Martinez, 2000) with a connectance set to 0.2. Additionally, vertical bars represent 50% of the standard deviations associated to these means. To facilitate comparisons among panels, we do not represent the occurrence probability of the 10 assemblages community in panel C for mutualism, which is 0.66 (the standard deviation associated is 0.33).

## 1.8 Discussion

Understanding how colonization-extinction dynamics influence species distribution and community structure remains a major challenge in biogeography (Wiens, 2011; Jabot and Bascompte, 2012; Godsoe and Harmon, 2012). Here, we build upon the simplicity of the Theory of Island Biogeography (TIB) to integrate crucial ecological processes, namely biotic and abiotic dimensions of the niche. Using the formalism of Markov chains, we derive an exact general solution for the occurrence probabilities of all possible assemblages that we calculate numerically (up to 10 species). Our approach is in stark contrast to the classic TIB (MacArthur and Wilson, 1967) where environmental gradients were not introduced and the co-occurrence among species was not modelled, despite empirical evidence of their impact (Diamond and Gilpin, 1982). By taking these constraints together we reveal how they interplay and affect species richness. We believe our approach offers new perspectives on the theory of biogeography and will support the development of species distribution models with the addition of species interactions.

In our model, we introduce the effect of biotic interactions as an ecological process affecting colonization/extinction probabilities. This has already been considered in many ways in the literature. For instance, more than forty years ago, Levins and Culver introduced extinction and migration rates affected by competition and showed analytically how it reduces co-occurrence (Levins and Culver, 1971). More recently, Jabot and Bascompte introduced production of eggs and seeds affected by interaction in an individual-based, meta-community framework and, hence, highlighted the potential effects of interactions on local diversity (Jabot and Bascompte, 2012). Also, Calcagno and colleagues demonstrated that tuning extinction and colonization rates based on the trophic relationships among species could explain the limited length of food chain (Calcagno et al., 2011). In contrast with previous studies, our approach is fully rooted on the TIB which yields well-defined null predictions (adding neither interaction nor environmental gradients), focuses on assemblages, and allows the investigation of the impact of any kind of network, including mixed interactions.




Figure 7: **Biodiversity distribution along environmental and connectance gradients.** We compute the expected species richness along an environmental gradient for competition (A), mutualism (B) and predation (C). We do so for different values of connectance depicted by the shades of blue. Species richness profile associated with the scenario without interaction is provided in each panel by the darkest solid line (connectance set to 0). Abiotic niches do have the same range for all species ( $r_i = 5$ ) and the optima are randomly drawn in the interval  $[10, 20]$ . The interaction weight ( $d$ ) is set to 10. The extinction parameters are set as follows:  $e_i = 10^{-5}$ ,  $e_{i,min} = 10^{-3}e_i$ ,  $e_{i,max} = 10^3e_i$  and  $c_i = 10^{-5}$ .

Networks are convenient representations of the structure of ecological communities to study persistence and resilience (Thébault and Fontaine, 2010). A strength of our model is that it not only takes all direct interactions into account, but also indirect ones (Wootton, 1994). For instance, in a linear trophic chain of three species, the occurrence of the top predator depends not only on the presence of its prey but also on the species at the bottom of the chain (Gravel et al., 2011). This means that the distribution of the top predator will be influenced not only by its own abiotic requirements, but also by those of its prey and the species at the bottom of the chain. The signature of such indirect interactions should be common in co-occurrence networks. This property comes from the assumption that interactions change extinction rates and the Markov chain formalism employed. Our formalism therefore provides a tool, similar to the general Lotka-Volterra equations for the local scale, that could be used to study the emergence of indirect interactions in networks at the large spatial scale.

The challenge of developing joint species distribution models (Pollock et al., 2014; Pel-

[lissier et al., 2013](#)) have recently motivated researchers to investigate co-occurrence ([Araújo and Rozenfeld, 2014](#); [Veech, 2013](#)). Our framework helps to disentangle the two main processes by which non-random species associations (co-occurrence) can arise. First, two species not interacting with each other could be non-randomly co-distributed because of similar or antagonistic ecological requirements. As we introduced an abiotic constraint on the colonization probability, some assemblages will be more likely than others on a given environment simply because some species are favoured and others filtered out. We thus expect to find a signature of the covariance in species response to the environment on these assemblage probabilities. Secondly, non-random co-distribution will arise from ecological interactions. We considered an additive impact of all ecological interactions a species is experiencing from the community. Species interact in various ways, but at the end all interactions do impact demography by definition. This reality enters the model by either enhancing or decreasing extinction probabilities. In other words, the occurrence of a single species is derived from the expectation of observing all other species in the community.

Our framework therefore provides a formalism to investigate the relationship between co-occurrence networks ([Araújo et al., 2011](#)) and interaction networks. There is a significant amount of information contained in the data of co-occurrence, which is overlooked by most current methods of community analysis. Standard species distribution models are fitted to univariate presence/absence data, neglecting the information contained in the distribution of associated taxa. Multivariate statistics summarize the spatial structure of ecological communities, but they are essentially limited to the description of co-occurrence, they are not meant to predict species distributions conditional on other species. Most analyses of co-occurrence aggregate pairwise observations into a single index for the whole community, thereby missing substantial information pertaining to the consequences of biotic interactions ([Boulangeat et al., 2012](#)). This situation is not surprising given there is no general theory for co-occurrence. Current hypotheses are mostly limited to negative interactions, leading to negative co-occurrence (repulsion), or positive interactions, leading to positive co-occurrence (attraction). Many theoretical achievements are required to study co-occurrence for more

complex assemblages, mixing positive, negative and antagonistic interactions. In addition, the impact of indirect interactions emerging in interaction networks on species distribution is ignored. Our approach provides a formal framework to overcome these limitations as we calculate assemblage probability at biogeographical scale and then derive co-occurrence. It also allows the decomposition of the strength of pairwise associations between abiotic and biotic drivers, opening the way for novel statistical developments of species distribution models taking into account this multi-occurrence information. We propose that studying the role of biotic factors at large scale requires us to introduce them as assemblages instead of adding species as factors which likely leads to non-equivocal conclusion ([Araújo and Luoto, 2007](#)). In addition, our approach is not limited to species pairs, the assemblage probabilities provide a valuable tool to the co-occurrence of groups of species such as motifs ([Stouffer et al., 2007](#)).

The importance of interactions across different scales is still debated ([McGill, 2010](#); [Araújo and Rozenfeld, 2014](#)). A common assumption is that interactions are negligible at large spatial scales, based on the rationale that abiotic filters primarily determine the composition of assemblages ([Pearson and Dawson, 2003](#)). This argument persists even though theoretical ([Gravel et al., 2011](#)) and empirical ([Gotelli et al., 2010](#)) evidence suggest the opposite. The key issue to solve this debate is thus to know how interactions can influence species assemblages with increasing spatial scale. Although the TIB still provide insights into the assembly of natural communities, the success of recent approaches integrating interactions strongly support their relevance at large scales. Indeed the addition of network structure ([Pellissier et al., 2013](#)) or correlation between species ([Pollock et al., 2014](#)) as proxies for interactions have adequately improved forecast accuracy. Here we do not solve this fundamental issue, however our model illustrates how species distribution at large scale will be impacted by the kind of interaction, their numbers and their distribution.

Although our framework is not readily applicable to real datasets, it nonetheless provides a theoretical foundation for the derivation of new statistical modelling approaches. We propose a different perspective which is rooted on theory, in contrast with what is usually done



with phenomenological model representing the structure of the data (Thuiller et al., 2013). There are nonetheless significant challenges to apply our framework to empirical data. First, we must find a way to deal with large numbers of species. At present, given  $n$  species in the regional pool, we compute an eigen vector of  $2^n$  probabilities from a  $2^n * 2^n$  transition matrix of a Markov chain. Moreover, in its current formulation, it requires us to evaluate a very large amount of data including a description of network of the same species across time and space to get accurate estimations. Solving this issue will requires a rational to reduce the number of species considered. This could be achieved either by inference of the relevant interactions, or alternatively by pooling species into groups. A systematic and rigorous method to build meaningful groups of interacting species from proxies such as traits and phylogenies remains to be developed, but there are nonetheless promising avenues (Baskerville et al., 2011). The relatively small number (from 3 to 7) of dimensions to ecological networks, i.e. the number of trait-axes required to properly infer interactions (Eklöf et al., 2013), supports its feasibility. A second challenge is to account for spatial structure that constrains population flux. Despite the theoretical developments, applied approaches to model species distribution struggle to introduce it efficiently (Boulangeat et al., 2012). The island-mainland approximation remains elegant but might be too simple for applied situations. One solution may be to identify source and sink localities, (Boulangeat et al., 2012). This requires us to consider i) species abundances and ii) spatial structures which would strongly increase the complexity of the model. One first step forward could be to apply the Levins model rather than the island-mainland model as Levins and Culver did to study the impact of competition (Levins and Culver, 1971)).

Despite our call for a new integrated theory of biogeography, we acknowledge the limitations of the framework. Recent studies aimed at integrating population dynamics, for instance, using approximations from the metabolic theory of ecology. This is a hopeful direction to assess local extinction risk, accounting for network structure, body size and abundance (Schneider et al., 2012). Beyond body size, other functional traits (McGill et al., 2006) could help us to escape from species singularity toward more general rules. Hence, approximating

food web structure could be done using traits ([Gravel et al., 2013](#)) and energetic requirements can be easily quantified through body size and local temperature constraints ([Brown et al., 2004](#)). Moreover considering changes in traits over time may be a key to introduce evolutionary processes. This would help us to release one strong assumption of our work: no speciation processes are taken into account. Although it might not matter for short time periods, having a pool of species unchanged becomes a major issue for time scales that exceed by far the lifespans of species we consider. Further, our framework could be applied to investigate diversification dynamics on remote areas, with a particular emphasis on the effect of ecological interactions on adaptive radiations. Despite the complexity of such model, it would very likely provide valuable insights on the future of biodiversity under current global change.

Since the seminal work of Davis et al. ([Davis et al., 1998](#)), there is growing evidence that the response of species to climate change must be studied at the community scale ([Suttle et al., 2007](#)). Even though species respond individually to climate change, they are constrained by complex direct and indirect biotic interactions emerging from large scale organization ([Lavergne et al., 2010](#)). The study of Cahill and colleagues ([Cahill et al., 2013](#)) has revealed the difficulties to link climate changes and species extinction. Even when the climate is expected to drive local extinctions, it actually implies a chain of perturbations amidst which biotic factors prevail (*e.g.* loss of prey ([Durance and Ormerod, 2010](#))). For instance, species contributing to the persistence of plant-pollinator networks are paradoxically the most vulnerable to extinction ([Saavedra et al., 2011](#)), highlighting the risk of extinction cascades. As Säterberg et al. expressed, “the species to be the first to go extinct is not the one whose mortality rate is increased but instead some other species in the food web”, thereby suggesting that perturbations which affect species differently also spread over the network making extinction difficult to predict ([Säterberg et al., 2013](#)). Although this is fully understandable as species interact, this makes forecasting of future species distributions more complicated. Therefore the challenge of proposing biodiversity scenarios to global change requires new approaches integrating ecological processes over time and spatial scales, and to disentangle their rela-

tive contribution ([Lavergne et al., 2010](#)). We think that the assemblage-based approach we propose here is a promising perspective to introduce interactions in biogeographical models.

## **1.9 Acknowledgment**

We thank Chantal L. Hutchison for insightful comments on the manuscript. KC was supported by a grant from the Ministry of Higher Education and Research of France. DG was supported by a NSERC Discovery grant and the Canadian Research Chair program. NM was supported by the CNRS.

## ARTICLE 2

### THEORIE DE CO-OCCURRENCE DES ESPECES DANS LES RESEAUX D'INTERACTION

#### 2.1 Résumé en français du deuxième article

Au chapitre précédent, j'ai mis en évidence l'impact potentielle des interactions sur la présence locale des espèces dans le cadre de la théorie de la biogéographie des îles. Afin de faire un pas en avant faire le traitement de données typique du biogéographe : les données de présence et d'absence à large échelles des espèces, je me suis confronté aux problèmes de co-occurrence des espèces en interactions.

La co-occurrence observée dans les données est simplement le nombre total de sites où les espèces sont présentes ensemble rapporté au nombre totale de sites étudiés. Pour pouvoir aller plus loin, nous avons défini une mesure de co-occurrence sous l'hypothèse d'indépendance de deux espèces. Cela signifie que nous prenons l'occurrence respective des deux espèces et que nous les multiplions pour obtenir notre seconde mesure de co-occurrence sous cette hypothèse. Grâce à la comparaison entre ces deux valeurs nous avons pu illustrer, dans l'article présenté ci-dessous, cinq grands principes relatifs à la co-occurrence des espèces en interaction :

**1. Les interactions directes entre deux espèces affectent la co-occurrence.** C'est une transposition directe du chapitre précédent sur la mesure de co-occurrence : s'il existe un lien entre deux espèces, leur probabilité d'être présentes simultanément dans une localité diffère de la probabilité attendue si elle se rencontraient aléatoirement.

**2. Les interactions indirectes modifient la probabilité de co-occurrence.** Malgré l'absence d'interaction directe entre deux espèces, ces dernières peuvent néanmoins être

liées par une ou plusieurs autres espèces, l'interaction est alors qualifiée d'indirecte. Si les conséquences des interactions directes se propagent à travers le réseau via ces relations indirectes, il est alors possible que répartition d'une espèce soit affectée par une autre espèce avec laquelle aucune interaction directe n'est constatée.

**3. L'effet des interactions sur la co-occurrence n'est pas symétrique.** Il n'existe a priori aucune raison pour que ces effets soient symétriques. Néanmoins, en utilisant la mesure de co-occurrence telle que décrite ci-dessus, nous la considérons comme telle. Nous montrons alors comment les probabilités conditionnelles peuvent prendre en compte l'asymétrie des effets des interactions.

**4. La force d'association entre deux espèces diminue avec la longueur du plus court chemin entre deux espèces.** Plus les espèces sont éloignées dans le réseau, moins les conséquences des interactions indirectes seront perceptibles, nous illustrons donc que les effets des interactions directes diminuent lors de leur propagation dans le réseau.

**5. La force d'une association avec une autre espèce diminue avec le nombre d'interaction qu'elle entretient.** Si une espèce a de nombreux liens dans le réseau (par exemple, un prédateur généraliste), alors celle-ci sera moins dépendante d'une espèce en particulier et de fait la relation qu'elle entretient avec les espèces se rapprochera de la co-occurrence sous hypothèse d'indépendance.

### 2.1.1 Publication associée

Pour ce second article, le contexte est particulier : Dominique Gravel a été invité à un numéro spécial de la revue scientifique *Theoretical Ecology* sur les réseaux écologiques. Dominique Gravel m'a alors proposé de travailler sur le prolongement de la réflexion menée au premier chapitre et de l'appliquer sur les données de co-occurrence. J'ai alors conceptualisé un modèle probabiliste pour tenter de comprendre comment les interactions peuvent affecter la co-occurrence. Je me suis occupé de toute la partie modèle et des figures. Avec Dominique

Gravel, nous avons rédigé l'essentiel du manuscrit avec des apports significatifs de Nicolas Mouquet et de Miguel B. Araújo qui sont alors devenus co-auteurs.

### **2.1.2 Traduction du résumé de l'article publié**

L'étude de la co-occurrence des espèces a été essentielle en écologie des communautés depuis la fondation de la discipline. Cependant, les données de co-occurrence sont une source d'information négligée par les modèles de distribution d'espèces et les biogéographes se demandent encore quels sont les effets des interactions aux différentes échelles biogéographiques. Nous soutenons l'idée qu'une théorie de la co-occurrence des espèces au sein d'un réseau écologique est nécessaire pour mieux interpréter les données de co-occurrence, pour formuler des hypothèses sur les mécanismes d'assemblage des communautés et pour étendre l'analyse des distributions d'espèces qui se concentre actuellement sur la relation entre les données d'occurrences et les facteurs abiotiques. L'objectif principal de cet article est de fournir les premières briques d'une théorie générale de la co-occurrence des espèces. Nous formalisons le problème avec différentes probabilités étudiées dans un contexte d'analyse de co-occurrence. Nous analysons les modules de trois espèces en interactions et nous conduisons des simulations pour documenter cinq principes influençant les associations au sein d'un réseau écologique : i) les interactions directes affectent la co-occurrence ii); les interactions indirectes affectent la co-occurrence; iii) la co-occurrence est rarement symétrique; iv) la force d'une association diminue avec la longueur du chemin le plus court entre deux espèces; v) la force d'une association diminue avec le nombre d'interaction qu'une espèce entretient. Nos analyses révèlent la difficulté d'interprétation des interactions entre espèces avec des données de co-occurrence. Nous discutons la possibilité d'inférence de la structure des réseaux d'interaction depuis ces données. Nous argumentons également que les modèles de distributions peuvent bénéficier de l'incorporation des probabilités conditionnelles des interactions en vue d'intégrer la contribution des interactions biotiques dans la distribution d'une seule espèce.

*Les sections qui suivent sont celles de l'articles publié.*

## 2.2 Titre

A Theory for species co-occurrence in interaction networks

## 2.3 Auteurs

Kévin Cazelles, Miguel Araújo, Nicolas Mouquet et Dominique Gravel

## 2.4 Résumé en anglais

The study of species co-occurrences has been central in community ecology since the foundation of the discipline. Co-occurrence data are, nevertheless, a neglected source of information to model species distributions and biogeographers are still debating about the impact of biotic interactions on species distributions across geographical scales. We argue that a theory of species co-occurrence in ecological networks is needed to better inform interpretation of co-occurrence data, to formulate hypotheses for different community assembly mechanisms, and to extend the analysis of species distributions currently focused on the relationship between occurrences and abiotic factors. The main objective of this paper is to provide the first building blocks of a general theory for species co-occurrences. We formalize the problem with definitions of the different probabilities that are studied in the context of co-occurrence analyses. We analyse three species interactions modules and conduct multi-species simulations in order to document five principles influencing the associations between species within an ecological network: i) direct interactions impact pairwise co-occurrence; ii) indirect interactions impact pairwise co-occurrence; iii) pairwise co-occurrence rarely are symmetric; iv) the strength of an association decreases with the length of the shortest path between two species; v) the strength of an association decreases with the number of interactions a species is experiencing. Our analyses reveal the difficulty of the interpretation of species



interactions from co-occurrence data. We discuss whether the inference of the structure of interaction networks is feasible from co-occurrence data. We also argue that species distributions models could benefit from incorporating conditional probabilities of interactions within the models as an attempt to take into account the contribution of biotic interactions to shaping individual distributions of species.

## 2.5 Mot-clef

Co-occurrence, Ecological networks, Biogeography, Indirect interactions, Null models

## 2.6 Introduction

Understanding of the processes driving the assembly of communities has been a central theme of ecology since the foundation of the discipline. How do we start from a regional species pool to assemble a structured community? Why are some species associated with each other? The work of [Diamond \(1975\)](#) pioneered the analysis of species co-occurrence in geographical space and, together with the controversy triggered by [Connor and Simberloff \(1979\)](#), it stimulated the development of a new field of research in numerical ecology ([Stone and Roberts, 1990](#); [Gotelli and Graves, 1996](#); [Legendre and Legendre, 2012](#)). The foundational work on species co-occurrences also led to the development of a rich array of methodological tools designed to test null hypotheses in ecology. Even if null models could be achieved numerically (*e.g.*, [Araújo et al., 2011](#)), typically they are based on permutations of distribution data. Null models have been used to infer the role of biotic interactions between pairs of species on their individual distributions. Studying the different drivers of species co-occurrence is not only of theoretical interest for improving understanding of the mechanisms of community assembly. It is also instrumental in predictive ecology, because a considerable amount of information is contained in species distributions data.

Despite its historical importance for community ecology, co-occurrence data remain a neglected source of information in models of species distributions. Biogeographers are still debating the impact of biotic interactions on species distributions ([Guisan and Thuiller 2005](#); [Gotelli et al. 2010](#); [Kissling et al. 2012](#); [Pellissier et al. 2013](#)). The distribution of a species is thought to be first influenced by its physiological tolerance to environmental conditions, but also by interactions with other species ([Hutchinson, 1957](#); [MacArthur, 1972](#); [Peterson et al., 2011](#); [Boulangeat et al., 2012](#)). The question of whether such interactions leave imprints in the distributions of individual species at biogeographical scales is still open to debate (*e.g.* [Davis et al. 1998](#)), but recent empirical [Gotelli et al. 2010](#), modeling (*e.g.*, [Araújo and Luoto, 2007](#)), and theoretical ([Araújo et al., 2011](#)) evidence invites the interpretation that this might indeed be the case.

The overwhelming majority of species distributions modelling applications, nonetheless, neglect information contained in joint distributions. Even multivariate analysis of community data (*e.g.* redundancy analysis - ([Legendre and Legendre, 2012](#))) do not use co-occurrence in geographical space to condition individual species response to environmental variation. There has been a recent rise of interest however in joint species distribution modelling ([Clark et al., 2014](#); [Harris, 2015](#); [Pollock et al., 2014](#)). These methods estimate the distribution of all species from a pool simultaneously and allow to condition the presence of a species on all other ones. However, estimated relationships are inferred from co-occurrence in environmental space rather than geographical space. That is, joint responses to the environment are inferred rather than biotic interactions themselves ([Baselga and Araújo, 2009](#)). JSDMs are, nonetheless, a first step towards developing a next generation of models accounting for the impact of biotic interactions on the distributions of species. They are, however, purely empirically driven and carry no specific hypotheses about how interactions can affect distributions. An exception is the recent attempt to model the effects of predator-prey dynamics on distributions and abundances using a meta- community framework coupled with phenomenological species distributions models ([Fordham et al., 2013](#)). The problem with such approaches is that data to parametrize interactions mechanistically are generally lacking

([Morales-Castilla et al., 2015](#)); therefore, they are hardly applied in most circumstances. It follows that we are faced with at least two major problems: i) understanding of the ecological interactions underlying the distributions of species is limited; and ii) knowledge of interactions is typically limited to net interactions, mixing both direct and indirect interactions. A theory of species co-occurrences in ecological networks is, therefore, needed to help interpret co-occurrence data, to formulate hypotheses for different community assembly mechanisms, and to extend the analysis of species distributions currently focused on the relationship between occurrences and abiotic factors.

The analysis of species co-occurrences starts with a matrix representing the presence and absence of each species over a set of sites. There are two aspects to the quantitative study of co-occurrence. The first is the choice of the metric used to quantify the strength of associations (relationships between species occurrences) between pairs of species. The simplest measure of species co-occurrence is the number of species combinations, as defined by [Pielou and Pielou \(1968\)](#). A second index is the count of checkerboards [Diamond \(1975\)](#): “In such a pattern, two or more ecologically similar species have mutually exclusive but interdigitating distributions in an archipelago, each island supporting only one species” (p. 32). Another popular index of co-occurrence is the C-score ([Stone and Roberts, 1990](#)). This index is similar to the count of checkerboards; it measures the average association or repulsion between pairs of species.

The second aspect of the analysis of species co-occurrence is the formulation of a null model. The controversy generated in ([Connor and Simberloff, 1979](#)) was partly (and rightly) based on the absence of a valid null hypothesis in Diamond’s analysis. Subsequent debates were mostly concerned with the formulation of the null hypothesis (*e.g.*, [Diamond and Gilpin, 1982](#)). Thanks to the theoretical work of [Gotelli and Graves \(1996\)](#), there is now a clear understanding of the different null models that can be constructed from the community matrix. New indices are constantly proposed, such as in ([Boulangeat et al., 2012](#); [Veech, 2013](#)); see also Table 2 in ([Ulrich and Gotelli, 2013](#)) for a description of 15 indices for co-occurrence

analysis. A promising avenue is the one proposed by [Araújo et al. \(2011\)](#) for the study of the matrix of species co-occurrence with tools borrowed from network theory.

Surprisingly, there is currently no theory for co-occurrence in multi-species communities. The basic hypotheses are that pairwise negative interactions result in repulsion, while pairwise positive interactions result in attraction. Attraction and repulsion are assessed by a comparison of the number of co-occurrence events to the number expected under a totally independent distribution. Similar environmental requirements between species could also result in attraction, even in the absence of interactions, if the sampling is conducted across heterogeneous environmental conditions. This theory is limited to pairwise and symmetric interactions; there is nothing for antagonistic and indirect interactions. Food web ecologists were among the first to recognize the important effect of indirect interactions on abundance ([Wootton, 1994](#)). For instance, plant and carnivore abundances are expected to correlate across a productivity gradient ([Hairston et al., 1960](#); [Oksanen et al., 1981](#)) because of top-down control on the herbivore population. Similarly, the propagation of indirect interactions has been studied in more complex interaction networks ([Yodzis, 1988](#)). Indirect interactions could reverse the net interaction in a surprising way, such that predator-prey abundances could be positively related ([Montoya et al., 2009](#)). Empirical analysis of co-occurrence for several taxa has shown that they are usually asymmetric ([Araújo et al. 2011](#)), such that a species distribution tended to be nested within the distribution of other (*e.g.* predator-prey distributions [Holt and Barfield, 2009](#); [Gravel et al., 2011](#)). In such a case, even if the co-distribution signature is quite understood, available methods will likely fail at using this piece of information to improve forecasts.

The main objective of this paper is to provide the first building blocks of a general theory of species co-occurrences. We formalize the proposed theory with definitions of different quantities that are studied in the context of co-occurrence analyses. Herewith, we analyse three species interactions modules in order to document five principles influencing the association between pairs of species from an ecological network: i) direct interactions impact

pairwise co-occurrence; ii) indirect interactions impact pairwise co-occurrence; iii) pairwise co-occurrence does not have to be symmetric; iv) the strength of an association decreases with the length of the shortest path between two species; v) strength of an association decreases with the number of interactions a species is experiencing. We base our mathematical argument on a general model of species distributions that is free of any assumption about how the ecological interactions operate. Finally we extend our analysis with simulations of multi-species networks in order to analyse how these mechanisms scale up in species rich communities.

## 2.7 Definitions

We start with definitions to formalize the quantities that can be computed from species distribution data and be used in the context of co-occurrence analyses. Let  $X_i$  be the random variable representing the presence of species  $i$ .  $X_i = 1$  when species  $i$  is present,  $X_i = 0$  otherwise. Then  $X_{i,t>0}$  is the random process associated, giving the value that  $X_{i,t}$  takes at any time  $t$ . Let  $p_{i,t}$  standing for the probability  $\mathbb{P}(X_{i,t} = 1)$ . Also, to illustrate the definitions, we derive the quantities for a simple presence/absence dataset (see Table 2).

The **marginal occurrence probability**  $\mathbb{P}(X_{i,\infty} = 1) = p_i^*$  represents the occurrence probability of species  $i$  when the system is at equilibrium, in the sense of the classical theory of island Biogeography [MacArthur and Wilson \(1967\)](#). As we assume so for all species, we drop the  $*$  and the  $\infty$  for the sake of clarity. The marginal occurrence probability is the sum of the occurrence of the species across all possible set of species in the data. In other words, it corresponds to the sum of the column of the site  $\times$  species table, divided by the total number of sites  $N$ . Marginal occurrence probabilities for species in Table 2 are:  $p_1 = 0.6$ ,  $p_2 = 0.6$  and  $p_3 = 0.4$ .

The **observed co-occurrence** between species  $i$  and  $j$  is the joint probability  $p_{i,j} = \mathbb{P}(X_i = 1 \cap X_j = 1)$ . It represents the number of sites where the two species are found together,

across all possible set of species in the data (in other words, it is a marginal probability with respect to other species), divided by  $N$ . In our dataset, for instance, we have  $p_{1,2} = 0.3$  and  $p_{1,3} = 0.2$ .

The **conditional co-occurrence** between species  $i$  and  $j$  is  $p_{i|j} = \mathbb{P}(X_i = 1|X_j = 1)$ . It represents the probability of observing species  $i$ , knowing that species  $j$  is already present. This quantity is close to the measure of association between two species because it is independent of the marginal occurrence probability of both species. The problem is that, as soon as there are other species present, the conditional co-occurrence as expressed here is marginalized over the set of all other species from the community  $K$ . For instance, for three species, we have:  $p_{1|2} = \mathbb{P}(X_1 = 1|X_2 = 1, X_3 = 1) + \mathbb{P}(X_1 = 1|X_2 = 1, X_3 = 0)$ . It, therefore, includes both the effect of *direct* and *indirect* associations between species, e.g. the direct association of species 1 with species 2 or the indirect association of species 3 with 1 via its effect on 2. Consequently, the measure of pairwise association should be:  $p_{i|j,\bar{K}} = \mathbb{P}(X_i = 1|X_j = 1, X_K = 0)$ , where the horizontal bar over  $K$  denotes absence of all other species. We name this the **fundamental conditional co-occurrence**. For instance, in Table 2, we get  $p_{1|2} = \frac{p_{1,2}}{p_2} = 0.5$  and  $p_{1|2,\bar{3}} = \frac{p_{1,2,\bar{3}}}{p_{2,\bar{3}}} = \frac{0.2}{0.3} = 0.67$ .

Following the same logic, we define the **fundamental occurrence** as  $p_{i|\bar{K}} = \mathbb{P}(X_i = 1|X_K = 0)$ . The fundamental occurrence is conceptually equivalent to the fundamental niche of Hutchinson (1957) and represents the probability of observing a species in the absence of biotic interactions, i.e., when all other species are absent. By analogy, the marginal occurrence should be interpreted as the realized distribution. For species 1 in Table 2 we calculate  $p_{1|\bar{23}} = \frac{p_{1,\bar{2},\bar{3}}}{p_{\bar{2},\bar{3}}} = \frac{0.2}{0.3} = 0.67$ .

Finally, we define the **independent co-occurrence** as  $p_{i,j,IND} = \mathbb{P}(X_i = 1)\mathbb{P}(X_j = 1)$ . It represents the co-occurrence between any pairs of species expected in absence of any association between them. In ecological terms, it would represent the co-occurrence when ecological interactions and habitat filtering do not impact species distribution. It also represents the null model against which observed co-occurrence is usually compared to. Note the independent

co-occurrence is different from the one expected under a neutral model ([Stephen P. Hubbell, 2001](#)). Firstly because strong competitive interactions in the neutral model forces repulsion and, secondly, because dispersal limitation also causes spatial aggregation and thus a non-random distribution of co-occurrence ([Bell, 2005](#)). In our example, we obtain, for instance,  $p_{1,2;IND} = 0.36$  and  $p_{2,3;IND} = 0.24$ .

### Direct association between two species

We start with the analysis of a two species situation, labeled species 1 and species 2, in order to understand direct associations between species pairs. A third species, 3, will be introduced in the next section to study indirect associations. The model we develop is general, as we do not specify the type of ecological interactions involved. It therefore accounts for all possible mechanisms from which an association between a pair of species could arise, such as trophic interactions involving energy fluxes, non-consumptive interactions, parasitism, direct interference, territoriality, space pre-emption, niche construction, etc. The impact of predator-prey interactions in a metapopulation setting with colonization and extinction dynamics will be considered for the multi-species simulations.

As we are willing to understand the role played by interactions in co-occurrence, we start by defining marginal co-occurrence probabilities of our two species by a decomposition into conditionnal co-occurrences. By the formula of total probability we have:

$$\begin{aligned}
 p_1 &= \mathbb{P}(X_1 = 1 \cap X_2 = 1) + \mathbb{P}(X_1 = 1 \cap X_2 = 0) \\
 &= \mathbb{P}(X_1 = 1 | X_2 = 1) \mathbb{P}(X_2 = 1) \\
 &\quad + \mathbb{P}(X_1 = 1 | X_2 = 0) \mathbb{P}(X_2 = 0)
 \end{aligned} \tag{2.1}$$

We do the same for species 2. Using the notation described above, [\(2.1\)](#) could be

rewritten as:

$$\begin{cases} p_1 = p_{1|2}p_2 + p_{1|\bar{2}}(1 - p_2) \\ p_2 = p_{2|1}p_1 + p_{2|\bar{1}}(1 - p_1) \end{cases} \quad (2.2)$$

where the vertical bar denotes the absence of a species. By solving the latter system, we get:

$$\begin{cases} p_1 = \frac{p_{1|\bar{2}} + p_{2|\bar{1}}(p_{1|2} - p_{1|\bar{2}})}{1 - (p_{2|1} - p_{2|\bar{1}})(p_{1|2} - p_{1|\bar{2}})} \\ p_2 = \frac{p_{2|\bar{1}} + p_{1|\bar{2}}(p_{2|2} - p_{2|\bar{1}})}{1 - (p_{2|1} - p_{2|\bar{1}})(p_{1|2} - p_{1|\bar{2}})} \end{cases} \quad (2.3)$$

When species are independent, we have  $p_{1|\bar{2}} = p_{1|2} = p_1$  and  $p_{2|\bar{1}} = p_{2|1} = p_2$ , then we logically find (2.1) again. Then, we can deduce the following interpretation of the impact of **direct interactions** on co-occurrence:

- i if species 1 cannot persist in absence of 2 (e.g., a parasite requiring its host), then  $p_{1|\bar{2}} \rightarrow 0$ , therefore  $p_1 \rightarrow p_{1|2}p_2$
- ii if species 1 depends strongly on 2 thereby perfectly tracking its distribution 2, the  $p_{1|\bar{2}} \rightarrow 0$  and  $p_{1|2} \rightarrow 1$ , and therefore  $p_1 \rightarrow p_2$
- iii if species 2 excludes 1, then  $p_{1|2} \rightarrow 0$  and  $p_{2|1} \rightarrow 0$ , so  $p_1 = \frac{p_{1|\bar{2}} - p_{2|\bar{1}}p_{1|\bar{2}}}{1 - p_{2|\bar{1}}p_{1|\bar{2}}}$  and  $p_2 = \frac{p_{2|\bar{1}} - p_{2|\bar{1}}p_{1|\bar{2}}}{1 - p_{2|\bar{1}}p_{1|\bar{2}}}$ .  
Therefore, if  $p_{1|\bar{2}} \rightarrow 1$ , then  $p_1 \rightarrow 1$  and  $p_2 \rightarrow 0$ .

### Co-occurrence in three-species modules

Now, we consider the co-occurrence between three species. We start with a general derivation of co-occurrence and then interpret the results for particular modules in order to reveal fundamental principles underling co-occurrence in ecological networks. Our solution provides insights to decipher the solution of species-rich networks since the three-node



connected subgraphs are fundamental building blocks of larger networks (Milo et al. 2002; Stouffer et al. 2007; Stouffer and Bascompte 2010). We use the same approach as in (2.1) and get the subsequent equation:

$$\begin{aligned} p_1 = & \mathbb{P}(X_1 = 1 \cap X_2 = 1 \cap X_3 = 1) + \mathbb{P}(X_1 = 1 \cap X_2 = 0 \cap X_3 = 1) \\ & + \mathbb{P}(X_1 = 1 \cap X_2 = 1 \cap X_3 = 0) + \mathbb{P}(X_1 = 1 \cap X_2 = 0 \cap X_3 = 0) \end{aligned} \quad (2.4)$$

As  $\{X_3 = 1, X_3 = 0\}$  forms a partition we get:

$$p_1 = \mathbb{P}(X_1 = 1|X_3 = 1)p_3 + \mathbb{P}(X_1 = 1|X_3 = 0)(1 - p_3) \quad (2.5)$$

This equation is analogous to the two-species interactions equation but enables the study of networks involving three species interactions, with species 2 being hidden by marginalization. We split the three species problem in two distinct two-interactions species problems. Firstly, we solve the equation for sites without species 3 and get:

$$p_{1|\bar{3}} = \mathbb{P}(X_1 = 1|X_3 = 0) = \frac{p_{1|\bar{2}\bar{3}} + p_{2|\bar{1}\bar{3}}(p_{1|2\bar{3}} - p_{1|\bar{2}\bar{3}})}{1 - (p_{2|1\bar{3}} - p_{2|\bar{1}\bar{3}})(p_{1|2\bar{3}} - p_{1|\bar{2}\bar{3}})} \quad (2.6)$$

which is similar to equation (2.3) but with an explicit absence of species 3. We do similarly for the conditional occurrence of 1 on species 3:

$$p_{1|3} = \mathbb{P}(X_1 = 1|X_3 = 1) = \frac{p_{1|\bar{2}3} + p_{2|\bar{1}3}(p_{1|23} - p_{1|\bar{2}3})}{1 - (p_{2|13} - p_{2|\bar{1}3})(p_{1|23} - p_{1|\bar{2}3})} \quad (2.7)$$

Doing so, we get the following set of equations describing the marginal occurrence probabilities for the three species:

$$\begin{cases} p_1 = p_{1|3}p_3 + p_{1|\bar{3}}(1 - p_3) \\ p_2 = p_{2|3}p_3 + p_{2|\bar{3}}(1 - p_3) \\ p_3 = p_{3|2}p_2 + p_{3|\bar{2}}(1 - p_2) \end{cases} \quad (2.8)$$

Note that we could have chosen a different set of equations depending on the way we split the problem, for instance, we could have started by considering the occurrence of species 1 given the occurrence of species 2 instead of species 3. Now, we solve the above linear system of three equations with three unknowns and find that:

$$\begin{cases} p_1 = \frac{p_{1|\bar{3}} + p_{3|\bar{2}}(p_{1|3} - p_{1|\bar{3}}) + (p_{3|2} - p_{3|\bar{2}})(p_{1|3}p_{2|\bar{3}} - p_{1|\bar{3}}p_{2|3})}{1 - (p_{2|3} - p_{2|\bar{3}})(p_{3|2} - p_{3|\bar{2}})} \\ p_2 = \frac{p_{2|\bar{3}} + p_{3|\bar{2}}(p_{2|3} - p_{2|\bar{3}})}{1 - (p_{2|3} - p_{2|\bar{3}})(p_{3|2} - p_{3|\bar{2}})} \\ p_3 = \frac{p_{3|\bar{2}} + p_{2|\bar{3}}(p_{3|2} - p_{3|\bar{2}})}{1 - (p_{2|3} - p_{2|\bar{3}})(p_{3|2} - p_{3|\bar{2}})} \end{cases} \quad (2.9)$$

Conditional probabilities of the right-hand sides can all be derived as we did for  $p_{1|3}$  in equation (2.7).

## Community modules

We now interpret these equations with examples of well-studied food web modules in community ecology: 1) linear food chain, 2) exploitative competition and 3) apparent competition. To do so, we consider matrices of direct associations representing the conditional co-occurrence probabilities among all pairs of species (see Table 3).

We are interested by the *observed co-occurrence* because this is the quantity that is

easily measurable from species distributions data, thus being the one that is typically studied. We consider that the marginal occurrence is also a known quantity and, therefore, we examine the effect of particular conditional co-occurrence arrangements on observed co-occurrences. We will not provide derivations for each module, but focus on particular pairs to illustrate two of the five principles.

Sites	Species 1	Species 2	Species3
1	0	1	1
2	0	1	1
3	1	1	0
4	1	0	1
5	0	0	0
6	1	1	1
7	0	1	0
8	1	0	0
9	1	0	0
10	1	1	0

Table 2: Presence/absence dataset for three species and 10 sites.

**Indirect interactions** . The comparison between the observed co-occurrence and the conditional co-occurrence reveals the role of indirect interactions on species associations. Based on (2.9) and (2.6) we get the association between species  $i$  and  $k$ :

$$\begin{aligned}
 p_{i,k} &= p_i - p_{i,\bar{k}}(1 - p_k) \\
 p_{i,k} &= p_i - \frac{p_{i|\bar{j}\bar{k}} + p_{j|\bar{i}\bar{k}}(p_{i|\bar{j}\bar{k}} - p_{i|\bar{j}\bar{k}})}{1 - (p_{j|\bar{i}\bar{k}} - p_{j|\bar{i}\bar{k}})(p_{i|\bar{j}\bar{k}} - p_{i|\bar{j}\bar{k}})}(1 - p_k)
 \end{aligned} \tag{2.10}$$

Therefore the observed co-occurrence between species  $i$  and  $k$  depends on their respective interaction with species  $j$  ( $p_{j|\bar{i}\bar{k}}, p_{j|\bar{i}k}$  and  $p_{j|ik}$ ). The conditional co-occurrence between two species could be null, but their observed co-occurrence be non-independent because of a shared interaction. This principle is best illustrated by the co-occurrence between a carnivore and a plant (species 3 and 1, respectively) in a linear food chain. In this situation, according to Table 3, we find that the observed co-occurrence between the plant and the carnivore is:

$$p_{1,3} = p_1 - \frac{p_{1|\bar{2}3}}{1 - p_{2|1\bar{3}}(p_{1|\bar{2}3} - p_{1|2\bar{3}})}(1 - p_3) \quad (2.11)$$

It is clear from this equation that there is a significant association between the carnivore and the plant, despite the conditional co-occurrence of the two species being totally independent. The indirect association gets stronger with the strenght of both conditional co-occurrence.

Similar observations could be made by studying the observed co-occurrence between consumers (species 2 and 3) in the exploitative competition module:

$$p_{2,3} = p_2 - \frac{p_{1|\bar{2}3}p_{2|1\bar{3}}}{1 - (p_{1|\bar{2}3} - p_{1|2\bar{3}})p_{2|1\bar{3}}}(1 - p_3) \quad (2.12)$$

And between resources in the apparent competition module (species 1 and 2):

$$p_{1,2} = p_1 - \frac{p_{1|\bar{2}3}}{1 - p_{3|1\bar{2}}(p_{1|\bar{2}3} - p_{1|2\bar{3}})}(1 - p_2) \quad (2.13)$$

**Associations do not have to be symmetrical.** Many studies of co-occurrence assume pairwise associations to be symmetrical (but see [Araújo et al., 2011](#); [Boulangeat et al., 2012](#)). The reason is simple, usually the observed co-occurrence is compared to the independent co-occurrence. These two metrics of association are perfectly symmetrical. This information is

providing us an inappropriate interpretation of the effect of interactions on species distribution. If we consider for instance the association between the two consumers (species 2 and 3) competing for a single resource (species 1), we have the observed co-occurrence at (2.12), which is symmetrical by definition. The proportion of the area occupied by species 2 where species 3 is also present is not however equivalent to the proportion of the areas occupied by species 3. Rephrasing the problem, we find that using (2.7) and (2.12),  $p_{2,3}/p_2$  is not equal to  $p_{2,3}/p_3$ . One species could have a stronger impact on the distribution of the other one. Predator distribution for instance tends to be nested within the distribution of the prey (Gravel et al., 2011), and consequently the predator has a high conditional co-occurrence with the prey, and alternatively the prey has a low conditional co-occurrence with the predator.

General case	Linear chain
$\begin{pmatrix} p_{1 \bar{2}\bar{3}} & p_{1 2\bar{3}} & p_{1 \bar{2}3} \\ p_{2 \bar{1}\bar{3}} & p_{2 \bar{1}3} & p_{2 \bar{1}\bar{2}} \\ p_{3 \bar{1}\bar{2}} & p_{3 \bar{1}2} & p_{3 \bar{2}3} \end{pmatrix}$	$\begin{pmatrix} p_{1 \bar{2}\bar{3}} & p_{1 2\bar{3}} & p_{1 \bar{2}3} \\ p_{2 \bar{1}\bar{3}} & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}$
Exploitative competition	Apparent competition
$\begin{pmatrix} p_{1 \bar{2}\bar{3}} & p_{1 2\bar{3}} & p_{1 \bar{2}3} \\ p_{2 \bar{1}\bar{3}} & 0 & 0 \\ p_{3 \bar{1}\bar{2}} & 0 & 0 \end{pmatrix}$	$\begin{pmatrix} p_{1 \bar{2}\bar{3}} & p_{1 2\bar{3}} & p_{1 \bar{2}3} \\ p_{2 \bar{1}\bar{3}} & p_{2 \bar{1}3} & p_{2 \bar{1}\bar{2}} \\ p_{3 \bar{1}\bar{2}} & p_{3 \bar{1}2} & 0 \end{pmatrix}$

Table 3: Direct associations between pairs of species for different modules. Entries indicate the fundamental conditional probabilities of occurrence of species  $i$  given the presence of species  $j$  and the absence of species  $k$ . *Linear chain*: 1 is the resource, 3 the top predator ; *Exploitative competition*: 2 and 3 are the consumers; *Apparent competition*: 1 and 2 are the resources. When  $p_{i|\bar{j}\bar{k}} = 0$ , it means that species  $i$  cannot be found without  $k$ . When two species  $i$  and  $j$  do not interact directly, if the absence of species  $k$  do not impact species  $i$  survival then :  $p_{i|\bar{j}\bar{k}} = p_{i|\bar{j}k}$ . For apparent competition, if species 1 and 2 are interchangeable for species 3 then :  $p_{3|\bar{1}\bar{2}} = p_{3|\bar{1}2}$ .

## Multi-species simulations

Now we move to multi-species simulations of more complex networks to reveal the last two principles of our theory. To do so, we run simulations of the model of trophic island biogeography developed by [Gravel et al. \(2011\)](#). The model describes the occurrence of a  $S$  species regional network. Species stochastically colonize islands with probability  $c$  and go extinct with probability  $e$ , as in the original model of [MacArthur and Wilson \(1967\)](#). Interactions are introduced with three additional assumptions: i) a consumer species could colonize an island only if it has at least one prey present (for simplicity, we consider producers to be resident permanently on the island); ii) a consumer species goes extinct if it loses its last prey species and iii) the presence of at least one predator species increases the extinction probability by  $e_d$ . The consequence of these assumptions is a sequential build-up of the food web on the island, starting with low trophic level species with a general diet. Small and isolated islands promote selection in favor of the most generalist species. The predictions converge to the classic island biogeography theory for highly connected regional food webs and large and connected islands (details in [Gravel et al., 2011](#)).

As mentioned above, there is a strong dependence of the predator occurrence on the presence of its preys. Alternatively, when  $e_d$  is sufficiently large, the preys will tend to avoid locations with the predator present. We consequently expect a strong signature of the network of interactions on the co-occurrence matrix. We are however concerned that indirect associations could emerge, as exemplified with the analysis of three species modules above, and thereby mask the signal of conditional co-occurrences.

We simulated complex networks from 5 to 100 species using the niche model of food web structure ([Williams and Martinez, 2000](#)). The diversity of primary producers was fixed at 2, and their niche position was drawn randomly between 0 and 1 according to a uniform distribution. We fixed connectance at  $C = 0.1$ . Colonization probability was set at  $c = 0.1$ , baseline extinction probability at  $e = 0.2$  and predator-dependent additional extinction

probability at  $e_d = 0.2$ . Simulations were run for  $10^7$  time steps to evaluate the conditional occurrence probabilities, and 100 replicated networks were simulated for each level of species richness.




Figure 8: Co-occurrence in multi-species networks. (A) The disparity between observed co-occurrence ( $P_{i,j}$ ) and independent co-occurrence ( $P_{i,j;IND}$ ) decreases with the path length between nodes (species). The envelopes are drawn around the 5% and 95% quantiles of all of the data, from 100 replicated simulations for every species richness value (5 to 100 species). (B) The strength of co-occurrence ( $\log(P_{i,j}/P_{i,j;IND})$ ) decreases with the number of interactions of a species  $i$  (*i.e.* the degree of a node). Points represent the mean for a particular degree of node value (1 to 60). The solid line represents the overall trends and the grey envelopped reflects the variance associated. At least 3000 values were used for each point. (C) The standard deviation of the strength of association ( $sd(P_{i,j}/P_{i,j;IND})$ ) and thus the variance decreases with species richness. Taken together, (B) and (C) imply that species distributions converge to independence with increasing species richness.

**Distance decay of observed co-occurrence.** The distribution of observed co-occurrence is illustrated for pairs of species separated by different path lengths at Figure 1A. The observed co-occurrence is presented as a function of the expected co-occurrence under the hypothesis of independent distributions. The strongest associations (given by the distance between the observed and the independent co-occurrence) are observed among pairs of species directly interacting with each other. The variance of the distribution reduces from direct to first order indirect interactions, and from first-order to higher interactions. We conclude that indirect non-independent co-occurrences are possible in complex networks, but their magnitude decreases as the number of links between two nodes decreases. This result is similar to the

observation of a distance decay of indirect interactions in food webs (Berlow et al., 2009).

**Strength of co-occurrence decreases with degree and species richness.** We performed simulations with a gradient of species richness and observed that the variance of observed co-occurrence also decreases with the degree of a species, i.e. the number of direct interactions a species is experiencing (Fig. 1B). We illustrated the relationship between the degree of a species and the observed co-occurrence for pairs of species with a direct association (Fig. 1C). This phenomenon has the consequence that the strength of observed co-occurrence reduces with species richness. The niche model has a constant connectance (Williams and Martinez, 2000), which has for consequence an increase of the degree with species richness. We find that the strength of co-occurrence decreases with the degree. This result is straightforward to interpret: the more diverse are the interactions, the weaker the impact of each pairwise direct interaction on the species distribution. Again, this result is similar to the observation of a scaling relationship between pairwise interactions and food web diversity (Berlow et al., 2009).

## Discussion

We first develop a probabilistic species distribution model constrained by biotic interactions using conditional probabilities of co-occurrence. We then illustrate five general principles underlying the impact of ecological interactions on co-occurrence and that should be considered for the formulation of a general theory of species co-occurrence. Two of them have been widely noted before: **i)** direct interactions affect species distributions and generate deviations in co-occurrences from that expected if distributions of species were independent from each other; **ii)** the effect of direct associations is often asymmetric, as envisioned in trophic metacommunity ecology (Holt and Barfield, 2009). We also illustrate principles that have been overlooked in most studies of co-occurrence (Araújo et al., 2011); **iii)** indirect interactions generate deviate co-occurrence from expectation under independence assumption;



iv) the strength of indirect associations decreases with the length of the shortest path distance between species pairs in a network; while v) also decreasing with the number of interactions a species is experiencing. We started with the analysis of three species modules to document these principles and then showed their applicability in multi-species networks. We find that the above principles also apply in larger networks, but that the strength of pairwise associations weakens as the number of species increases.

Our results have considerable implications for interpretation of co-occurrence data. Firstly, they demonstrate the considerable variety of mechanisms causing pairwise associations. Such variety of mechanisms makes interpretation aggregated indices of co-occurrence, such as the C-score, very difficult (see also [Araújo and Rozenfeld, 2014](#)). Previous studies already made the argument that positive and negative interactions could balance each other ([Boulangeat et al., 2012](#)) and consequently associations should be studied on a pairwise basis ([Veech, 2013](#)). At least, some measure of the variability of the associations is required, and at best metrics such as network analyses ([Araújo et al., 2011](#)) should be used to characterize their complex structure. But most importantly, our analyses reveal the difficulty to infer species interactions from co-occurrence matrices. Associations are not symmetric and, therefore, indices that are capable of dealing with them are required. Null model testing is not sufficient; significance is assessed from the difference between observed co-occurrence and co-occurrence expected under independent distributions and is, consequently, symmetric. In addition, statistically significant associations cannot be interpreted as evidence of direct interactions. Our results also show that indirect interactions, and not only second order interactions, contribute to generate apparent non-independent co-occurrence. These indirect associations could be of any kind and are impossible to detect solely based on knowledge of direct interactions.

Null models of species associations should, thus, be used only to reveal the structure of co-occurrence data. The lack of an association between a pair of species is no unequivocal evidence of absence of direct interactions. It must be interpreted as the absence of a net effect

in the spatial co-occurrence arising from pairwise interaction alone. For instance, in the case species A is competing with species B and species C, and B with C, it is possible that A and C could be independently co-occurring if there is a strong indirect positive interaction A-C arising from the A-B and B-C direct interactions. Null model testing is consequently subject to important type I (false interpretation of a significant association) and type II errors (false interpretation of an absence of association). The problem itself does not come from the statistical method per se, the description of co-occurrence in the data will be right provided that the technique is adequate, but from the interpretation of the null model analysis.

Should we, therefore, abandon joint species distributions modelling (JSDM) and all of the information contained in co-distribution data? While our results might lead to such an interpretation, there is still some value in species co-occurrence data that could be used in distribution models. The appropriate use of JSDMs is to remove biases in the evaluation of species-specific relationship with the environment. Accounting for joint distribution will contribute to the evaluation of the conditional distribution of a species when all other species are absent. In other words, they should be used to improve the evaluation of the fundamental niche. The JSDMs will, however, fail to predict the right occurrence probability of a species for communities that have no analogue to the training dataset. JSDMs are using only the net associations between pairs of species and are not meant to recover the direct pairwise conditional co-occurrences. For instance, a JSDM evaluated for a plant, an herbivore and a carnivore will provide the correct description of the joint distribution of all three species, but will be of limited use to predict the distribution of the plant and the herbivore if the carnivore disappears from the system. Further developments are, consequently, required to solve the issue and account for both direct and indirect interactions. One possible solution would be to constrain JSDMs with a prior expectation of the underlying structure of direct interactions. It is also valuable to ask whether the inference of the structure of interaction networks is feasible from the observation of co-occurrences (as they result from many ecological processes). There is growing interest in inferring ecological network structure from alternative sources of information ([Gravel et al., 2013](#); [Morales-Castilla et al., 2015](#)). This problem is challenging

because of the multiple influences on co-occurrence. Our analysis of three species modules with conditional probabilities revealed it is feasible numerically, to obtain an estimate of all pairwise conditional probabilities when accounting for higher order interactions. Known quantities are the marginal probabilities and observed co-occurrence. The parameters to be evaluated are all fundamental conditional probabilities, representing the direct associations between pairs of species (the  $p_{i,j,\bar{K}}$ ). This is a  $S \times S$  problem to solve and thus requires a significant amount of data. It might, however, be solved with large datasets where the number of sites  $N$  is much larger than  $S$ . There might also be methods to reduce the dimension of the problem because usually only a small fraction of potential interactions are met in a network (corresponding to the connectance  $C$ ). While a net interaction network is likely to be fully connected ( $S \times S$  links), the direct interaction network has still only a fraction  $C$  of these links realized. Bayesian approach with latent variables could even further help reducing the dimension of the problem (*e.g.* [Rohr et al., 2010](#); [Ovaskainen et al., 2010](#)). In such methods, latent variables are evaluated for each species to represent the underlying structure of the ecological network. It was found that between two and four parameters per species would be required to successfully represent more than 80% of interactions in a predator-prey network ([Rohr et al., 2010](#)). This approach could, therefore, be used to represent the underlying structure of direct interactions and to evaluate numerically the non-null conditional probabilities. Note that these pairwise direct interactions should be interpreted specifically with reference to spatial dynamics because they would still represent phenomenologically the consequences of interactions, not the mechanisms of interactions. The next step in the development of a theory of species co-occurrence (and of species distribution) is the addition of environmental constraints. Our approach assumed a homogeneous environment, mainly for tractability of equations. We acknowledge that non-independent co-occurrence could also arise because of shared environmental requirements. The addition of environmental constraints would be easy to implement in our framework by simply making the conditional probability in absence in absence of interactions a function of the environment. Every quantity we derive after would be conditional on the environment. What would be more challenging but, nonetheless, feasi-

ble numerically, would be to make the direct interaction itself a function of the environment. There is now growing evidence that ecological interactions are context dependent ([Chamberlain et al., 2014](#); [Poisot et al., 2012](#)). We view this integration as the next step to the derivation of a theory-driven species distribution model taking into account biotic interactions ([Thuiller et al., 2013](#)).

## 2.8 ACKNOWLEDGMENTS

This work was inspired by discussions with T. Poisot, D. Stouffer, A. Cyrtwill and A. Rozenfeld. Thanks to Matt Talluto and Isabelle Boulangeat for helpful comments on a previous version of the manuscript. Financial support was provided by the Canada Research Chair program and a NSERC-Discovery grant to D. Gravel. M. Araújo acknowledges support from Imperial College's Grand Challenges in Ecosystems and Environment Initiative.

## 2.9 Context

Biogeographers have long been fascinated by the picture of species distributions and questioned how it could have been made, i.e. searching for the processes shaping biodiversity on Earth ([Engelbrecht et al., 2007](#); ?). Starting from the clear relationship between abiotic variables and the physiological constraints of organisms, large-scale studies have been conducted in a pattern-driven perspective making Biogeography a realm of correlations (?). Such an approach has provided many valuable knowledge along with the development of efficient statistical tools. However, in the context of global changes, many researchers claim for strengthening the theoretical foundations of the field towards a biogeography mechanism-driven providing reliable biodiversity forecasts ([Lomolino, 2000b](#); [Thuiller et al., 2013](#); ?).

The importance of biotic constraints on species distribution are one of the many concerns regarding this request ([Godsoe and Harmon, 2012](#); [Araújo and Rozenfeld, 2014](#)). In

order to test whether interactions influence species distributions, the simplest avenue is to investigate the species co-distribution in light of their ecological relationships. Such investigation started with Diamond's original study stating that species interacting by competition should avoid each other in space, leading to a 'checkerboard' distribution (Diamond, 1975). This idea was rapidly criticized for the lack of an adequate null hypothesis (Connor and Simberloff, 1979; ?). Nevertheless, the resulting debate captured the attention of biogeographers as it must unravel whether co-occurrence data are more than the sum of occurrence information of a set of species (Clark et al., 2014). The answer to this question has direct and major consequences: a negative one would support the use of classical species distribution models (hereafter SDMs, ?) that postulate the independence among species whereas a positive one would give credit to methods taking co-occurrence information as a proxy for ecological interactions (Morales-Castilla et al., 2015) and would support the development of methods including network into species distribution models (Ovaskainen et al., 2010; Pollock et al., 2014).

Recent theoretical developments have proposed mechanisms explaining how ecological interactions must affect the fundamental niche (see Box 1) and how they could impact occurrence data (Holt and Barfield, 2009; Araújo and Rozenfeld, 2014). However, ranges of species are very often inferred from the realized niche which includes the impact of abiotic and biotic factors. Therefore, finding evidences of interaction signals may prove difficult which would explain the scarcity of studies reporting such effect (but see Gotelli et al. (2010)). Fortunately, the co-occurrence theory in interaction networks has been formalized and suggests that the repercussion of interactions in co-occurrence data depends on the structure of the network (?). Notably, the higher the degree of a species, *i.e.* the number of species which with it interacts, the trickier the detection of a signal of co-occurrence. Finding good correspondence between network topology and co-occurrence signal in empirical data would support the idea that interactions affect geographic ranges, even if for many pairs of interacting species no significant co-occurrences are found.

Here, we examined occurrence data in the light of these recent theoretical developments for five different datasets for which interactions are observed or assessed to determine whether ecological interactions impact the distribution of species. We report that the analysis of co-occurrence failed to clearly reveal a difference between pairs of interacting species and pairs of not-interacting species. However our results suggest that the degree of species influences our ability to detect significant association making co-occurrence information more than a collection of co-occurrence only for species with a limited number of link. Moreover, we discover a clear relationship between the co-occurrence strength of a species and the cumulated occupancy of the entire set of species with which a species interact. Interestingly, we point out that the relation vanishes when we used classical SDMs which questions the capacity of these approaches in capturing adequately species interactions. Our results reveal the existence of conditions under which interaction may or may not be detected in static co-occurrence data. This lead us to think that interaction in biogeography is not only a matter of scale but also a matter of relevant biological unit for which occurrence must be predicted.

## **2.10 Material and Methods**

### **2.10.1 Datasets**

We analyzed five datasets spawning a large range of environmental conditions (see Fig S1 and SI Text), a large diversity of organisms and covering all fundamental type of interactions (see ?). Four of them came with observed interactions: the Willow Leaf Network (WLN), the Pitcher Plants Network (PPN), the Caribbean Hummingbirds Network (CHN), the French Breeding Birds Survey (FBBS). For these network, we derived metawebs, *i.e.*, the matrix recording all interactions, and computed three metrics: (1) the connectance of metawebs (proportion of links), the degree of species (number of links for one species) and the shortest-path between all pairs of species (the minimum number of link from one species to the other, see SI Text). For the North American Trees (NAT) datasets, we derive a distance

based on functional traits (see table S1 and Fig S2). For the FBBS dataset, we also derived different trait-based distance (see table S2). For all datasets, we kept only species that were present at least on 1% of the total number of sites (see SI Text).

### 2.10.2 Measures of co-occurrence

For each pair of species, we determined the number of observed co-occurrence  $O_{i,j}$  and we calculated the expected co-occurrence values  $E_{i,j}$  and its standard deviation  $SD_{i,j}$  to compute the following Z-score  $(O_{i,j} - E_{i,j}) / SD_{i,j}$  (?). For this standardized metric of co-occurrence, positive (negative) values indicates more (less) co-occurrence than randomly expected. Expectations were derived using three different methods. First, we assumed that all sites were equivalent, meaning we occulted the potential influence of abiotic conditions. The distribution of co-occurrence for a limited number of sites have been already studied elsewhere (?Veech, 2013), therefore, we used an hypergeometric distribution (see SI Text for further details). For the two other expected values, we used two different classical SDMs, namely, Generalized Linear Model (hereafter GLM) and Radom Forest (hereafter RF) in order to assign a probability of being presence in a given site for all species (see SI Text for more details and Fig S3 for the assessment of models' performances). With these two last scenario, we actually hypothesize that species may often co-occur simply because they have similar abiotic requirements which is basically what SDMs do.

## 2.11 Results

For two out of four datasets for which interactions were known, we obtained a clear difference between interacting and not-interacting species (10 panels A to D). Therefore, when integrating all pairs of species we did not obtain a clear evidence that direct interaction species co-occur differently from not-interacting one. For the WLN, distinguishing herbivore-willow interactions from herbivore-parasitoids revealed that the strength of co-occurrence

was stronger for the former than the latter ones (11 A-B). Interestingly, we noticed that the higher the mean degree of species in the dataset, the more difficult the detection of a signal of interactions in co-occurrence was (11 A-D).

For the NAT and FBBS datasets for which we inferred a distance based on functional traits, we found that co-occurrence was higher for pairs of similar species (10 panels E and F). As similarity could be taken as a proxy for competition strength (Morales-Castilla et al., 2015), this result suggests that competition is poorly detectable at large scale which is theoretically supported (Araújo and Rozenfeld, 2014). Therefore, co-occurrences of similar species are likely driven by the similarity of their abiotic requirements. The results for the FBBS dataset were identical irrespective of the type of traits examined (Fig S4). This hypothesis was further supported by the decrease of the Z-score with the distance for both datasets (fig S6 A and D).

For all datasets, we report that the distribution is concentrated but the Wilcoxon rank test we used (see SI Text) lead us to the distribution is symmetric about 0 taking environmental context into account shifts the distribution toward 0 and diminishes the variance associated (10). This trend is more pronounced for RF than for SDM. Hence, assuming that sites are not identical for species due to the abiotic context reduces the signal of co-occurrence which could even disappear. In the PPN dataset, we found that the signal is even reversed (10) but the quality of the SDM approaches used were low (fig S3 B).

When examining the Z-scores against the shortest path between preys (hosts) versus predators (pollinators), we report that the higher the number of links the less distinguishable observed co-occurrence are from random ones. For shortest paths higher than one, a large part of co-occurrence are non significant (11 A-D) as predicted by the theory, but the observed decay is steeper (?). We conclude that species separated by more than one link may be treated as independent species. The decay is similar when all pairs of species were taken into account (see Fig S5). Moreover when we calculated the mean degree in of predators (pollinators), we demonstrate the signal of co-occurrence for low shortest path is stronger for specialists (11



A) than for generalists (11 C) suggesting that the abundance of interactions may prevent us from detecting them in static occurrence data.

Finally, we analyze the mean Z-score of a predator (pollinator), *i.e* Z-scores averaged over all the set of its preys (host plants), against the total number of site covered by its preys (hots) (12). We report that when abiotic context is not taken into account, there is a clear negative relationship explaining up to .69 of the variance (12 panels A, D, G and J). Therefore, when a predator feeds on a set of preys that jointly cover a large part of the geographic range studied, the impact of species interactions is undetectable, but when the joint repartition of the prey is restricted, the imprint of interactions remains appreciable. Additionally, the associated linear regression outperformed the one using the degree of the species that has been envisioned by the theory (Fig S7 panels A, D, G and J). Also, we show this relation asymmetric: the decay is less convincing when the mean Z-scores of the preys are plotted against the cumulated range of their predators (fig S8 panels A, D, G and J). Hence the imprints of interactions in static occurrence are appreciated once relevant pair of species are studied. When species are highly linked with other species and when these species have ranges that do not completely overlap, we cannot make clearly co-occurrence to interactions. This suggests that the range of the set of species should be examined rather than individual range of prey. Interestingly, we found that using the presence of the whole set of prey as predictor to assign the presence of species outperformed GLMs (see Fig S9) but not RFs.

All these last results are no longer true once abiotic constraints are taken into account, the clear relationship obtained is either weakened or even reversed (12 B,C,E,F,H,I,K,L) meaning the signals of co-occurrence for specialists are no longer different from the one of generalists. If we consider that this relationship is biologically meaningful as biotic constraints may be softer for generalists than for specialists we must then question to what extent inferring species distribution fail in capturing occurrence properties exhibit by species assemblage.

### 2.11.1 Discussion

Our study demonstrate how the information of ecological networks can shed light upon static occurrence data over broad spatial scales. Although we noted that we cannot assert in general that interacting species co-occur differently from non-interacting one, we highlight the possibilities offer by asking alternative questions such as the impact of the abundance of interactions and the relative positions of two species within the network on their co-occurrence. Our results demonstrate that there is an imprint of ecological interactions on occurrence data but also that there are specific conditions under which they can be detected. Hence, we show that the signal of co-occurrence is blurred by the abundance of interactions. This has been envisioned by a recent theoretical work (?) and stress the usefulness of such studies to propose ways to revisit co-occurrence data with additional information. Moreover, we also prove relevant the examination of a the range of a given predator against the range of the set of its preys rather than individual prey. This indicates that the role of ecological interactions may not only be a matter of spatial scale (McGill, 2010) but also a matter of consistent biological unit to be predicted at over large spatial scales.

Classical SDMs approaches are based on the assumption that species are independent and focus one abiotic variables (Jeschke and Strayer, 2008) to propose scenarios of tomorrow's biodiversity. Emerging approaches propose to release this strong hypothesis by predicting a set of species as assemblage among which correlations are accounted for (Pollock et al., 2014; ?). The development of more powerful statistical tools is fundamental, however it can only provides a partial solution to improve range forecasting. A substantial part of the solution lies in the understanding of the role of ecological and evolutionary mechanisms shaping species distributions (Thuiller et al., 2013). Here, we underline the need for not hypothesizing *a priori* the independence among species. Rather, the assumption must be proved to apply, otherwise a relevant assemblage must be modeled. Here, we suggest that for generalist species, the assumption of independence is reasonable while for specialists the relative position with the other species within the network should help deciding which set of

species are to be modeled. We claim for the integration of ecological information at the core of SDMs and we suggest that the amount of information to integrate should be a reasonable. Therefore, once identified, such information should greatly strengthen our predictions for a fair additional complexity into our SDMs.

Our results lead us to conclude that the abundance of interaction renders the assumption of independence valid which therefore simplifies the predictions. Interestingly, it has been recently highlighted that some allometric features of networks are better predicted when the species richness increases ([Berlow et al., 2009](#)). More than forty years ago, MacArthur similarly noted the following paradox “How can a more complex community be easier to understand? A possible answer might be that the complex community has strong interactions among species so that the lives of the separate species are less independent than in a simple community. Where there is greater interdependence, patterns may be more conspicuous.” ([MacArthur, 1972](#), p.199). In our study, the strong interactions among species ease the forecasting of biodiversity. However, under the ongoing mass extinction, a myriad of links vanishes while new ones emerge with the changes in the composition of local communities. Therefore, even if under current conditions the assumption of independence may be valid, under dramatic modifications of ecosystem as currently observed, it may often prove false. As a step forward, new biodiversity scenarios must not solely map the future ranges of individual species but the entire community including the consequences of potential extinction in term of community structure.

### 2.11.2 Box 1

The fundamental niche is here described as the occurrence probability under the assumptions that (1) biotic factors are not limiting occupancy and (2) that dispersion is unconstrained. In this case, only abiotic factors (such as water availability, temperature variability and edaphic variables) limits survival and/or reproduction success, and then the occurrence probability. Consequently, predators occupancy is computed assuming that preys are abundant enough all along the environmental gradient. Similarly, the fundamental niche of any prey is not influenced neither by predators nor by competitors.

For a three species network made of one predator and its two preys, we derive the three fundamental niches  $f_i$  (9 A). Regarding the predator (species 3), we assume its prey are equivalent and that the presence of at least one prey is sufficient to release all the biological constraints:

$$f_3(w) = P(X_3 = 1 | X_2 + X_1 > 0, G = w)$$

where  $G$  denotes the environmental gradient and  $X_i$  is the random variable associated to the presence of species  $i$ . Similarly,  $f_1$  and  $f_2$  are obtained assuming that 3 is absent :

$$f_2(w) = P(X_2 = 1 | X_3 = 0, G = w)$$

and:

$$f_1(w) = P(X_1 = 1 | X_3 = 0, G = w)$$

Once projected on a map, the fundamental niche unravels the potential distribution of a species (Kearney and Porter, 2004). The expected distribution can be compared to real ob-

servations and could reveal whether dispersal limits and ecological interactions are prevalent in the occupancy dynamic of studied species. The realized niche (9 B) includes these factors. In our simplified example, fundamental and realized niches of preys are identical. The realized niche of the predator,  $r_3$ , is controlled by the joint realized niches of its preys:

$$r_3(w) = f_3(w) (1 - (1 - r_1(w))(1 - r_2(w)))$$

The above expression may often be more complicated due to the size and the structure of the network. For instance, we do not consider the apparent competition between 1 and 2 although it must affect the distribution of all species. Integrating the impact of many interactions may be possible using occurrence probabilities of species assemblages rather than single species (?). Integrating network information to shed light upon species distribution is also crucial to understand what kind of co-occurrence is biologically relevant. Consider as an example the co-occurrence between species 1 and 3: the co-occurrence may be strong if we restrict the analysis to the suitable conditions for species 1 but it must be weak if the entire environmental gradient is sampled. However, if we examine the co-occurrence between 3 and the assemblage made of species 1 plus 2, the co-occurrence may always be strong. Although this is meaningful in a biological point of view, co-occurrence studies often remain focus on pairs of species.

### 2.11.3 Tables

Table 4: Data sets analyzed in this article.

Type	No. of sites	No. of species	Interaction type	Observed	Traits	Connectance	References
Willow Leaf Network	374	156	Trophic / Parasitism	yes	no	0.042	unpublished
Pitcher Plants Network	39x20	53	Trophic	yes	no	0.44	?
Caribbean Humming- birds Network	32	62	Mutualist	yes	no	0.011	?, ?, ?
North American Trees	128891	31	Competition	yes	no	-	unpublished
French Breeding Birds Survey	2354	179	Trophic	yes	no	0.018	?

## 2.12 Figures




Figure 9: **Probabilistic description of fundamental and realized niches** For a three species network all the occurrence probabilities are derived along an environmental gradient assuming that A interactions are not limiting the distribution and B that species 3 needs at least of one of its preys, *i.e.* species 1 or 2. Horizontal dotted lines stand for the occurrence probabilities reached at an environmental optimum.

## 2.13 Material and methods

In this section, we present in more details, the datasets and the methodology we used. All analyses have been performed using R environment software (table S1 includes functions and packages we used).

### 2.13.1 Datasets

Sites for the five datasets are reported on five maps gathered in Fig S1. Below, we describe in more details the five datasets. The total number of species, the number of species present in at least 1% of the total number of site and the number of species for which traits information were available are reported in table S2.

#### 2.13.1.1 Willows leafs network


#### 2.13.1.2 Pitcher plants network

#### 2.13.1.3 Caribbean Hummingbirds-Plant network

#### 2.13.1.4 North American Trees datasets

**2.13.1.4.1 Traits-based distance** We used a distance built upon nine functional traits whose values were retrieved from (?), see **Supplementary Table 3** available at <http://onlinelibrary.wiley.com/doi/10.1111/j.1466-8238.2010.00592.x/supinfo>. Each of the nine selected variables were centered and scaled (R functions used reported in table S1) then used as is to derive Euclidean distances for all pairs of species. Then, we use agglomeration clustering with Ward's method (implemented in the *hclust()* function we used, see Table S1) to obtain the dendrogram presented in 14.





**Figure 10: Co-occurrence of interacting versus not-interacting pairs of species** Figures under each groups of boxplots indicate the number of pairs to which the Z-score distributions refer. The light grey rectangle corresponds to the 95% confidence interval for the standard normal distribution which gives insight into the proportion of pairs of species significantly different from 0. The comparison made in panels A to D is based on direct interactions observed. For panels E and F, similar species are defined as the species for which the trait-based distance is less than or equal to the lower decile of this distance distribution. Note that outliers are not displayed. P values were computed using the Wilcoxon rank sum test, to compare interacting versus not-interacting Z-score distribution calculated for the three different methods (black symbols) and to show whether the distribution is symmetric about 0 (light grey symbols).





Figure 11: **Co-occurrence signal decays when the shortest path between a pair of species decay** The Z-score distribution are plotted against the shortest path for A willows-herbivores interactions, B herbivores-parasitoids interactions, C birds-plants interactions and D the pitcher plants network. First figures under each grouped boxplots indicate the shortest path associated while the figures below provide the number of pair to which the distribution refers. Note that we used the same y-axis for panels A and B as they regard two different kind of interaction of the same dataset.

### 2.13.1.5 French Breeding Birds Survey datasets

**2.13.1.5.1 Traits-based distance** We used 73 traits that are boolean variable (see Table S4) we kept as is to derive Euclidean distances for all pairs of species.

### 2.13.2 Building metawebs

For four datasets, we built network based on all observed interactions and derived associated quantities, *i.e.* the connectance of the metawebs, the degrees of species and the shortest-path, using the R package “igraph” (table S1).



**Figure 12: Co-occurrence significance decreases as the cumulated occupancy increases**  
 For a given species, Z-scores are averaged over the all set species it interacts with and plotted against the joint distribution of the same set of species. We do so for the herbivores in the willows leaves network (panels A to C), the parasitoids in the willow leaves network (panels D to F), the hummingbirds in the Caribbean hummingbirds datasets (panels G to I) and all species in the pitcher plants network that consume other species (panels J to L). The x-axis is expressed as a log proportion of the total number of sites. Black symbols are mean Z-scores significantly different from 0 (see SI Text). In each panel, the dotted line represents the linear regression  $y = ax + b$  for which the  $R^2$  is provided. The size of circles reflects the degree of species for which the Z-score was calculated, the relation size-degree for each row is given in the middle panel. For the hummingbirds dataset (panels G to I), the triangle represent the values obtained for the former distribution of a species already analyzed (see SI text).

### 2.13.3 Co-occurrence measurement

For a given pair of species  $i$  and  $j$ , we examined the relationship between the observed co-occurrence  $O_{i,j}$  and the expected co-occurrence  $E_{i,j}$ . Here, we provide more information about the three methods we used to analyse co-occurrence.

### 2.13.4 Hypergeometric distribution

This distribution has been mentioned in a different context (see ?) and have been fully exploited in ? despite the author never mentioned it is a classical distribution. To clarify this, we start from the distribution written in equation (1) in [Veech \(2013\)](#). We consider the co-occurrence of two species on  $n$  sites. Species 1 is present in  $n_1$  while species 2 is present in  $n_2$  sites. The probability of having  $j$  co-occurrence,  $p_j$  is:

$$p_j = \frac{\binom{n}{j} \binom{n-j}{n_2-j} \binom{n-n_2}{n_1-j}}{\binom{n}{n_2} \binom{n}{n_1}}$$

if  $\max 0, n_1 + n_2 - n \leq j \leq \min n_1, n_2$  and 0 otherwise. The expression above yields:

$$p_j = \frac{n!}{(n-j)!j!} \frac{(n-j)!}{(n-j-n_2+j)!(n_2-j)!} \frac{(n-n_2)!}{(n-n_2-n_1+j)!(n_1-j)!} \frac{(n-n_1)!n_1!}{n!} \frac{1}{\binom{n}{n_2}}$$

by rearrangement:

$$p_j = \frac{1}{j!} \frac{1}{(n_2-j)!} \frac{1}{(n-n_2-n_1+j)!(n_1-j)!} \frac{(n-n_1)!n_1!}{1} \frac{1}{\binom{n}{n_2}}$$

once sorted out, this results in:

$$p_j = \frac{\binom{n_1}{j} \binom{n-n_1}{n_2-j}}{\binom{n}{n_2}}$$

Thus, the number of co-occurrence follows a hypergeometric distribution of parameters  $(n, n_1, n_2)$  we used to calculate the expected co-occurrence  $E_{i,j}$  under the hypothesis that all sites were identical for all species.

### 2.13.5 GLM and RF

For GLM and RF,  $E_{i,j}$  correspond to probabilities of occurrence computed based on climatic data. R functions are reported in Table S1.

#### 2.13.5.1 Climatic data

We used the global climate layers provided data WorldClim, version 1.4, available at <http://www.worldclim.org> (?). For each dataset, we performed a principal component analysis and kept as many axes as needed to explain 90% of the total inertia. We used these axes in GLM and RF.

#### 2.13.5.2 Generalized Linear Model

For all datasets, we performed a Generalized Linear Model (?) using all the axes provided by the PCA as polynomials of degree 2. To constrain the number of parameters, we did not evaluate the interactions among axes. We also performed a selection model based on the Akaike's information criterion (AIC) C in a Stepwise Algorithm. R functions used to carry out the analyses are indexed in table SX.

### 2.13.5.3 Random Forests

Random Forests (?) were performed using the same formula as for GLMs. For all species, 10000 trees were computed and the probability for a species being in a given site were calculated based on the number of votes the sites were granted.

### 2.13.5.4 Evaluating the models

For all species, we assess the performance of the Species Distribution Models we used, *i.e.* Generalized Linear Model and Random Forest, using the Area Under the Receiver Operating Characteristic (AUROC) (?). We present the results as a cumulative sum of frequencies corresponding to the score for all species for each of the four ecological systems we studied (see Fig ?).

## 2.14 Supporting Tables

### 2.14.0.1 R packages used

Analysis	Function	Package name	Version	Citation
Scaling and Centering	scale	base	3.3.1	<a href="#">R Core Team (2015)</a>
Euclidean distance	dist	stats	3.3.1	<a href="#">R Core Team (2015)</a>
Clustering	hclust	stats	3.3.1	<a href="#">R Core Team (2015)</a>
PCA	dudi.pca	ade4	1.7.4	?
GLM	glm	stats	3.3.1	<a href="#">R Core Team (2015)</a>
GLM Selection	step	stats	3.3.1	<a href="#">R Core Team (2015)</a>
Degree of species	degree	igraph	1.0.1	?
Shortest Paths	shortest.paths	igraph	1.0.1	?
AUROC	somers2	Hmisc	3.17.2	?

Analysis	Function	Package name	Version	Citation
TSN retrieving	get_tsn	taxize	0.7.4	?, ?
Wilcoxon tests	wilcox.test	stats	3.3.1	<a href="#">R Core Team (2015)</a>

**Supplementary Table 1:** R and packages used for the analyses. GLM: Generalized Linear Model, PCA: Principal Component Analysis. AUROC: Area Under the Receiver Operating Characteristic. {#tbl:code}

Type	Total	Selected	Traits
Willow Leaf Network	274	156	-
Pitcher Plants Network	91	53	-
Caribbean Hummingbirds Network	62	62	-
North American Trees	31	31	31
French Breeding Birds Survey	340	179	321

**Supplementary Table 2:** For each datasets the total number of species (column *Total*), the number of species present in more that 1% of the total number of sites (column *Selected*), and the number of species for which traits information are available (column *Traits*). The symbol ‘-’ means ‘not relevant’. {#tbl:numsp}

Species	TSN	maxH	GR	WD	TolS	TolD	AM	EM	LMA	Nmass
<i>Abies balsamea</i>	18032	25	1	0.34	5.0	1.0	0	1	151.00	1.66
<i>Acer negundo</i>	28749	20	3	0.44	3.5	3.0	1	0	37.04	2.50
<i>Acer rubrum</i>	28728	25	3	0.49	3.4	1.8	1	0	71.09	1.91
<i>Acer saccharum</i>	28731	35	1	0.56	4.8	2.3	1	0	70.63	1.83
<i>Betula alleghaniensis</i>	19481	25	3	0.55	3.2	3.0	0	1	46.08	2.20
<i>Betula papyrifera</i>	19489	25	3	0.48	1.5	2.0	0	1	77.88	2.31

Species	TSN	maxH	GR	WD	TolS	TolD	AM	EM	LMA	Nmass
<i>Carpinus caroliniana</i>	19504	8	1	0.58	4.6	2.0	0	1	49.05	2.15
<i>Carya cordiformis</i>	19227	25	1	0.60	2.1	4.0	0	1	44.05	2.60
<i>Fagus grandifolia</i>	19462	25	1	0.56	4.8	1.5	0	1	61.22	2.04
<i>Fraxinus americana</i>	32931	30	2	0.55	2.5	2.4	1	0	76.75	2.12
<i>Fraxinus nigra</i>	32945	20	2	0.45	3.0	2.0	1	0	71.94	2.10
<i>Fraxinus pennsylvanica</i>	32929	25	3	0.53	3.1	3.9	1	0	87.72	1.80
<i>Larix laricina</i>	183412	25	3	0.48	1.0	2.0	0	1	120.00	1.36
<i>Ostrya virginiana</i>	19511	12	1	0.63	4.6	3.3	1	0	37.04	2.20
<i>Picea glauca</i>	183295	25	1	0.35	4.2	2.9	0	1	302.86	1.28
<i>Picea mariana</i>	183302	20	1	0.41	4.1	2.0	0	1	294.12	1.12
<i>Picea rubens</i>	18034	25	2	0.38	4.4	2.5	0	1	304.67	1.15
<i>Pinus banksiana</i>	183319	20	3	0.42	1.4	4.0	0	1	243.90	1.24
<i>Pinus resinosa</i>	183375	25	3	0.39	1.9	3.0	0	1	294.12	1.17
<i>Pinus strobus</i>	183385	30	3	0.36	3.2	2.3	0	1	121.92	1.42
<i>Populus balsamifera</i>	22453	25	3	0.37	1.3	1.8	1	1	83.46	1.95
<i>Populus grandidentata</i>	22463	20	3	0.39	1.2	2.5	1	1	70.45	2.50
<i>Populus tremuloides</i>	195773	25	3	0.37	1.2	1.8	1	1	82.02	2.16
<i>Prunus pensylvanica</i>	24799	12	3	0.36	1.0	2.0	1	1	50.00	2.40
<i>Quercus alba</i>	19290	35	1	0.60	2.9	3.6	0	1	81.21	2.39
<i>Quercus macrocarpa</i>	19287	15	1	0.58	2.7	3.9	0	1	92.74	2.27
<i>Quercus rubra</i>	19408	25	2	0.56	2.8	2.9	0	1	84.20	2.06
<i>Thuja occidentalis</i>	505490	15	1	0.30	3.5	2.7	1	0	223.00	1.02
<i>Tsuga canadensis</i>	183397	30	1	0.40	4.8	1.0	0	1	122.55	0.99
<i>Ulmus americana</i>	19049	35	3	0.46	3.1	2.9	1	0	79.47	2.07
<i>Ulmus rubra</i>	19050	25	3	0.48	3.3	3.0	1	0	59.88	2.50



**Supplementary Table 3:** Tree species and traits used. Abbreviations are as follows: TSN - Taxonomic Serial Number defined by Integrated Taxonomic Information System (ITIS), maxH - Average maximum height, GR - Growth rate, WD - Wood Density, TolS - Shade tolerance, TolD - Drought tolerance, AM - Arbuscular mycorrhiza (Endomycorrhiza), EM - Ectomycorrhiza, LMA - Leaf mass per area, Nmass - Nitrogen content per leaf mass unit(?) available at <http://onlinelibrary.wiley.com/doi/10.1111/j.1466-8238.2010.00592.x/supinfo.{#tbl:trees}>

Category	Trait name
Activity	Nocturnal
Activity	Crepuscular
Activity	Diurnal
Diet	Seeds, nuts or grain
Diet	Fruits / frugivory
Diet	Vegetative
Diet	invert
Diet	fish
Diet	Very small mammals
Diet	Large mammals
Diet	Herptile
Diet	Small birds
Diet	Long birds
Diet	Vertebrate
Diet	Bones
Diet	Carrion
Feeding behavior	Pursuit (air and/or aquatic)
Feeding behavior	Sally
Feeding behavior	Foliage gleaning
Feeding behavior	Pouncing


Category	Trait name
Feeding behavior	Grazing
Feeding behavior	Picking, pecking or stabbing
Feeding behavior	Digging
Feeding behavior	Overturning
Feeding behavior	Probing
Feeding behavior	Filtering
Feeding habitat	Water-surface
Feeding habitat	Underwater
Feeding habitat	Water
Feeding habitat	Mud
Feeding habitat	Ground
Feeding habitat	Canopy
Feeding habitat	Shrub (low and high)
Feeding habitat	Vegetation
Feeding habitat	Air
Foraging habitat	Wet grassland, meadows, fens, sedges or tundra
Foraging habitat	Dry grassland
Foraging habitat	Rocky slope
Foraging habitat	Fast river/stream
Foraging habitat	Slow river/stream
Foraging habitat	Shore (marine)
Foraging habitat	Salt marsh
Foraging habitat	Mud or silt
Foraging habitat	Sandy gravel/beach
Foraging habitat	Reed marshes
Foraging habitat	Conifer
Foraging habitat	Mixed forest

Category	Trait name
Foraging habitat	Deciduous
Foraging habitat	Mediterranean oak or other
Foraging habitat	Open/low forest
Foraging habitat	Forest or habitat edge
Foraging habitat	Shrub/bush
Foraging habitat	Urban
Foraging habitat	Garden
Foraging habitat	High air
Nesting habitat	Wet grassland, meadows, fens, sedges or tundra
Nesting habitat	Dry grassland
Nesting habitat	Banks/sand/mud
Nesting habitat	Rock surface/outcrops
Nesting habitat	Near water/shore/island
Nesting habitat	Sand gravel/beach
Nesting habitat	Reed marshes
Nesting habitat	Conifer
Nesting habitat	Mixed forest
Nesting habitat	Deciduous
Nesting habitat	Mediterranean oak and other
Nesting habitat	Open/low forest
Nesting habitat	Shrub/bush
Nesting habitat	Urban
Nesting habitat	Garden
Nesting location	Elevated
Nesting location	Tree hole
Nesting location	Ground

**Supplementary Table 4:** List of the Boolean traits used to derive Euclidean distances between all pairs of species in the French Breeding Birds Survey.

## **2.15 Supporting Figures**

# A. Willow Leaf Comm



[Contact GitHub](#) [API Training](#) [Shop](#) [Blog](#) [About](#) © 2016 GitHub, Inc. [Terms](#) [Privacy](#)  
[Security](#) [Status](#) [Help](#)




Figure 14: **Dendrogram representing the trait-based distances between the 31 species studied in the North American tree datasets.** Names of angiosperm species are written in dark grey while names of Gymnosperm species are in a lighter grey.






Figure 15: **Evaluation of the SDM approaches** For each dataset, the distributions of performance of generalized linear models (light grey symbols) and random Forest (dark grey symbols) for all species are presented.




Figure 16: \*\* Co-occurrence and the nature of the trait-based distance in the FBBS dataset\*\*  
 The different panels correspond to four different set of trait upon which for different distance are built. Similar species are defined as the species for which the trait-based distance is less than or equal to the lower decile of this distance distribution. Note that outliers are not displayed. The light grey rectangle corresponds to the 95% confidence interval for the standard normal distribution which gives insight into the proportion of pairs of species significantly different from 0. P values were computed using the Wilcoxon rank sum test, to compare interacting versus not-interacting Z-score distribution calculated for the three different methods (black symbols) and to show whether whether Z-score were greater for hypergeometric versus GLM and whether GLM versus RF (grey symbols).




Figure 17: **Co-occurrence signal decays when the shortest path between a pair of species decay** Distribution of Z-scores for all interactions are grouped by shortest-path indicated by the first numbers below boxplots. The other figures below stand for the number of pairs of species included within the distributions.





Figure 18: *Changes co-occurrence signal when increasing the distance between two species*  
 Points represent the result for all pairs of interaction for two datasets: the North American Tree dataset (A=C) and the FBBS (D-I). For the latter, we used the trait-based distance computed with all available traits (D-F) and the body-size ratios (the lighter species over the heavier, panels G-I). In each panel, the equation on the bottom-left corner indicated the results of the linear regression depicted by the dotted line.



**Figure 19: The degree of species partially explains the decrease of the co-occurrence strength** For the herbivores (A-C) and the parasitoids in the willow leaves network datasets (D-F), the hummingbirds in the Caribbean hummingbirds datasets (G-I) and all species in the pitcher plants network that consume other species (J-L) the mean Z-score is plotted against the degree of the species. Black symbols are mean Z-scores significantly different from 0 (see SI Text). In each panel, the dotted line represents the linear regression  $y = ax + b$  for which the  $R^2$  is provided.




Figure 20: \*Reversed figure 4\*\* This figures correspond to the figure 4 in the main text but the Z-score are calculated for preys (host plants) rather than for predators 9pollinators). Mean Z-score are computed for willows (A-C) and herbivores (based on the herbivores-parasitoids only, D-F) of the willows leaves network, the hosts plants in the Caribbean hummingbirds datasets (G-I) and species that feed on the detritus in the pitcher plants network (panels J-L). The x-axis is expressed as a log proportion of the total number of sites included in the considered dataset. Black symbols are mean Z-scores significantly different from 0 (see SI Text). In each panel, the dotted line represents the linear regression  $y = ax + b$  for which the  $R^2$  is provided. The size of circles reflects the degree of species for which the Z-score was calculated, the relation size-degree for each row is given in the middle panel.




Figure 21: **Predicting herbivore distribution based on the distribution of willows** For the herbivores in the willow leaves network dataset, we compared the AUC obtained when using willow it does not interact with (circles) a willow in interacts with (squares) and the set of willow it interacts with (triangles) to AUC obtained for GLM (A) and RF (B). Positive values indicated that species based model outperformed the SDM model.

## ARTICLE 3

### TOWARDS A METABOLIC THEORY OF BIOGEOGRAPHY

#### 3.0.1 Résumé en français du troisième article

Cet article est à la fois une présentation d'une extension directe du modèle de TTTb et d'un cas particulier de mon premier chapitre mais aussi le prendre pas vers un espoir motivé par plusieurs choses. En 2015 j'ai commencé à m'intéresser au problème d'énergie mais pas assez et puis j'ai commencé des réflexions pour lever le nombre d'espèces dans mon modèle mais aussi à essayer d'aller vers quelque chose qui est plus réaliste. Quand on pense à une pyramide du vivant il y a la base des producteurs qui de l'énergie inorganique jusqu'au top prédateurs sont capables de faire de l'énergie. Comme souligné par le papier de Gravel et al. il n'y a pas le simple fait qu'un prédateur mais aussi il y a une différence de triplement dans un top prédateur et un herbivore dans la répartition de l'énergie. J'ai essayé de me confronter à ces problèmes mais j'ai dû commencer par comprendre l'état du savoir et les enjeux. Il m'apparaît que 2 théories intéressantes plus ou moins mécaniques n'ont pas tant de problème. Mais comment comprendre que certaines relations soient visibles à l'échelle est difficile à comprendre ou sont la preuve que certaines hypothèses n'ont pas la saturation.. Ce modèle offre des perspectives nouvelles et un point d'ancrage concret sur plein de belles choses.

#### 3.0.2 Publication envisagée

Cet article traite de deux aspects les perspectives qu'il offre une approche et propose un prendre pas pour s'élancer vers les populations depuis le haut. Le modèle qui y est présenté est simple mais innovant et offre la possibilité pour explorer des hypothèses. Le chapitre est en cours de développement. L'avancement est indiqué dans le rapport du troisième compte-



rendu de comité de thèse. Il est la base d'un projet que je viens de proposer pour le post doc.  
Miguel Araujo et Loic Pelissier. cool

### 3.1 Introduction

Disentangling the respective contribution of processes shaping species distribution remains the central tenet of biogeography. While biogeographers clearly envision the list of ingredients that are needed to understand species distribution ([Thuiller et al., 2013](#)), they are currently lacking a recipe to mix them in the right proportions toward forecasting community assembly. Therefore, in the current context of global changes, we likely fail to predict accurately biodiversity responses to global changes as we keep focusing on abiotic factors, *i.e.* temperature and precipitations, and overlooking biotic interactions and short-term evolutionary responses [[Lavergne et al. \(2010\)](#)]. At the core of this issue is the need for a renewal of theoretical foundations of the field that could be started by a synthesis of recent enrichments of the model of the island-based theory of biogeography ([Lomolino, 2000a](#); [Warren et al., 2015](#)).

Among main theoretical challenges toward more realistic models, biotic interactions should be integrated as a constraint for species co-existence in meta-communities. It is a truism of community ecology that species interact and that their persistence in communities rely on these relationships. Consequently, ecological interactions may explain, at least partially, the dynamics of local extinctions which in turn must explain some properties of the geometrical shape of the ranges of species ([Holt and Barfield, 2009](#); ?) even if we have poor evidence of such effect at large spatial scales (but see [Gotelli et al., 2010](#)). However, two of the most influential models in biogeography assume ecological equivalence of species. Indeed, the Theory of Island Biogeography of MacArthur and Wilson (hereafter TIB, [MacArthur and Wilson, 1967](#)) focuses on the species richness on islands according to properties of islands and overlook the variation among species characteristics, especially their interactions

within food-webs. Second, in his neutral theory Hubbell assumes that individuals of different species are ecologically equivalent and predicts the distribution of abundance without considering interactions (Hubbell, 1997). These two theoretical models have been proved relevant for certain groups of species and inadequate for others, but none of them was intended to describe exhaustively the different components of communities on islands. To take leap toward predictions at the community scale, the hypothesis of ecological equivalence must be released and ecological interactions explicitly integrated (?).

The TIB is well-suited to explore the consequences of the integration of ecological interaction at broad spatial scale as it includes two fundamental processes of biogeography, namely immigration and extinction, while being very simple and readily expandable (Losos and Ricklefs, 2010; Warren et al., 2015). Building upon the classical models, recent studies have included interactions in the TIB (Gravel et al., 2011; ?). In such approaches, the pool of species that can colonize the island becomes a metaweb which makes species interdependent entities with specificities (*e.g.* a given trophic level) rather than indistinguishable unities of a species quantity. As a important consequence of such consideration, colonization and extinction rates vary with respect to the species identity and the local community. Five years ago, Gravel et al. (2011) propose that predators can locally survive as long as they find at least one prey and prevent their colonization if the island is prey-free. More generally, ? presented a Lotka-Volterra like model in which the composition of the local community determines the extinction rates. Although extinctions are impacted by the local community, these approaches overlook where energy is limited, the consumption of any populations matter and for a predator, preys are not equivalent. Here, we propose to extent the TIB with the explicit integration of energy as a fundamental driver of extinction in local ecological network.

Species do not escape from thermodynamics laws that have shaped the pyramid of life we currently observe (?). As a dramatic evidence, despite the complexity of evolutionary trajectories, many of key ecological properties of species scale with body mass (?) founding the metabolic theory of ecology (Brown et al., 2004). Among the major results of this theory

is the scaling of the metabolic rates (?) than often scale with the power function of the body mass often between  $2/3$  and  $3/4$  (?). Even if all the relationships are not well-understood (see the case of abundances reviewed in ?, and the recent relationship between prey and predator biomasses ?), the commonness of allometric relationships promises to lower the complexity of ecosystems by using body mass distribution to describe many of its properties. Allometric relationships and energy flows are also a way to revisit models of populations dynamics as envisioned by ? who considered species as energy processors to derive a bioenergetic model of population dynamics where energy uptake is based on allometric relationships. Recent development have convincingly shown that allometric is a key to analysis the properties of networks (such as stability ?), the role of species within (Schneider et al., 2012) and some authors have even proposed to infer ecological interactions with promising results (Gravel et al., 2013; ?).

At large spatial scales, variations of energy availability explain the latitudinal gradient of species richness. As a first approximation, energy availability could be estimated by the productivity of primary producers which relies on temperature, water availability and stoichiometry (?). Several mechanisms have been proposed to explain the positive relationship between energy and species diversity (see ?, for a review) among which two involve a change in the breadth of ecological niche or the width of the ecological space. Recently, based on the analysis of 196 empirical food webs, ? have shown that the link density remains constant over a latitude gradient for most ecosystems supporting that the more the energy the wider the niche space rather than an increase of niche width poleward. The Ecological Limits hypothesis posits Interestingly ?. Turnover of species ... gives credit fluctuation and ecological debts . Damuth showed based that the well correlation of can be is a there is a share we propose here to test such idea and propose to see consequence in term of biomass allocation. . .

o This hypothesis the functional evolution saturated the space and imprint such hypothesis is important and island where turnover forced while is not here we propose to look at that !

? predicted the signal of as expected under the constraint when energy is limiting and as an indirect effects notably their example. ..

Ability of local population to cover their energy demand is of fundamental importance and when they cannot they must certainly go locally extinct. On the other hand, the need of a population at a locality cannot sustain. Evidence of the relative constance of the niche breadth indirectly support the idea of a This recent results hypothesis partially support the idea that as suggested by in the TIB TIB equilibrium dynamic where energy constrain total diversity but also we assume MVP to derive energy uptake of species and constrain species richness. We also... energy is a good rationale for extinction. As remembered by ? average per capita growth rates of all coexisting species must be positive at low density and strong enough to overcome negative random effects. Our model : -The proposed conceptual framework that links: - extinction dynamics - ecological networks - energetic constraints allometry - simple model constrains - simple calculation - more than probability of extinction - highlight turnover with degree

We show that : - dynamics - species - turnover increase with max

Our model sheds light upon and is so hopeful... - species turnover - energy effect - population fluctuations

### 3.2 Model

The model developed by MacArthur and Wilson in their TIB is a colonization and extinction dynamics linked to island characteristics, predicting the species richness on the island according to island size and its distance from the mainland (MacArthur and Wilson, 1967). One promising direction to extend the theory of biogeography is to include ecological interactions into the classical model (?Gravel et al., 2011; ?). Following this avenue, we considered explicit interactions among the pool of species based on allometric relationships. Body masses of species are furthermore exploited to determine the quantity of energy a species requires to maintain a local population on the island. Once the island cannot sustain a minimal population for all local species, then extinctions occur. Therefore, in the model

described below, we extent the TIB with purely stochastic colonization event together with deterministic extinction based on a energy rational.

### 3.2.1 Primary producers and habitat heterogeneity

An island is assumed to be a patch of land covered by a maximal quantity of primary producers constituting an amount of energy available upon which a food web can be built. The maximal amount of energy available for herbivores is noted  $E_0$  and varies with island area, which is basically the assumption behind the drop of extinction rates with an increase in the size of the island in the classical theory (MacArthur and Wilson, 1967; ?). In our model, when  $E_0$  increases, irrespective of the nature of primary producers, the energy available to sustain herbivore populations raises. We further postulate that the habitat heterogeneity increases with the partitioning of  $E_0$  in a vector of different categories of primary producers. Hence, for  $p$  different habitats, the vector  $(E_1, E_2, \dots, E_p)$  represent the energy partitioning among species such that  $\sum_i E_i = E_0$ . The simplification made here is twofold: (1) the diversity of primary producers is not taken into account, (2) the production is constant over the time.

### 3.2.2 Metawebs

The regional pool of species in the TIB is basically a number of species  $P$  reflecting the regional diversity. Here, we not only consider a fixed number of species  $P$  but we also include trophic relationships among species. Following ?, we built regional metawebs of  $P$  species using the niche model Williams and Martinez (2000). We furthermore posit the niche axis as the body size species of species and also species without any links in the metaweb are assumed to be herbivores. Primary producers are not included in the niche model, and species without link are assumed to be herbivores. Apart from primary producers regarded as a quantity of available energy, the model we use exhibits strong correlations between

trophic level and body mass: herbivores are often the smallest and top predators. Although this assumption is an oversimplification for ecosystems in general, it remains reasonable for marine ecosystems (?) in which such allometric relationships have been used to infer the structure of food webs (Gravel et al., 2013).

### 3.2.3 Migration of species from the metaweb

An island is made of one or more habitats made of primary producers. Here, we focus on the migration of herbivores and species of higher trophic levels. Following Gravel et al. (2011), we assume that the colonization of herbivores/predators is successful only if they find at least one of their habitat/prey on the island. Moreover, in the model we propose, arrivals of new species are always possible until the energetic requirements to maintain local populations exceed the primary production in which case energetic constraints and networks topology determine the identity of species to go extinct. Therefore, colonization events are assumed to be purely stochastic, extinctions are more deterministic (this difference in stochastic nature between these fundamental processes of biogeography has been recently supported in ?).

### 3.2.4 Energetic constraints on local food webs

The energetic rationale of the model is simple: local populations need a certain amount of energy to maintain a minimum of population under which the species goes extinct. Under this constraint, the species richness locally increases until the energy production is no longer sufficient for all populations. For a given species  $i$ , energy requirements of any individual is derived from allometric relationships proposed by the metabolic theory of ecology (Brown et al., 2004), that is a consumption of the form  $c_i m_i^b$  where  $b$  is often set to .75. We do not integrate variance among individual of a species,  $m_i$  is therefore a constant for a species and the energy uptake associated to a minimal viable population (hereafter MVP) of  $n_i$  individuals becomes  $n_i m_i^b$ . ? defined the MVP as “[...] the smallest isolated population having a 99%

chance of remaining extant for 1000 years despite the foreseeable effects of demographic, environmental and genetic stochasticity, and natural catastrophe”, highlighting that the smaller the population, the higher the extinction risk which is assessed by the time to extinction. Building upon this idea, ? showed that the time to extinction is also affected by the mean population growth rate underlining that species characteristics may lead to a heterogeneity in MVP. Moreover, ? have developed a metabolic framework within which they proved the growth rate to be proportional to  $m_i^{-b}$ . Based on these results, we explore two simple cases: 1) MVP is equal for all species:  $n_i = n_0$  ; 2) MVP scales with the growth rate  $n_i = n_0 m_i^{-b}$ . For both scenarios, species  $i$  can survive only if the energy expenditures can be covered, *i.e* if the energy available is greater than:  $n_i c_i m_i^b$ .

### 3.2.5 Energy fluxes and transfer efficiency

Although the expression of energy consumption is similar among species, they obtain it from different sources: herbivores feed on primary producers, whereas predators feed on a set of preys. The primary production must therefore be split properly through the entire community. To deal with this, we propose to convert the energy costs for maintaining predator populations into additional populations of herbivores to be maintained. To exemplify this idea, we start with the simplest trophic network where a predator  $j$  feeds upon a herbivore  $i$ . The cost to maintain the MVP of  $i$  is  $n_i c_i m_i^b$  and  $n_j c_j m_j^b$  for  $j$ . For the latter, we convert  $n_j$  into an extra population of  $i$ ,  $n_{i,j}$  herbivore individuals dedicated to  $j$  consumption. Furthermore, we account for the energy loss the conversion begets by including a transfer efficiency  $\tau$ . Hence, the conversion from  $n_j$  to  $n_{i,j}$  is given by the following equation:

$$\tau n_{i,j} c_i m_i^b = n_j c_j m_j^b$$

{#eq:id1}

which yields:

$$n_{i,j} = \frac{n_j c_j}{\tau c_i} \left( \frac{m_j}{m_i} \right)^b$$

{#eq:id2}

In our study, we postulate that the transfer efficiency is constant across trophic levels, which is likely an oversimplification of the reality as suggested by the sparse empirical data available (??). We now add a new predator  $k$  feeding on  $j$  to the insular food web. According to our reasoning, we must covert  $n_k$  into  $n_{i,k}$ . To do so, we start by converting  $n_k$  into a population of  $j$ :

$$n_{j,k} = \frac{n_k c_k}{\tau c_j} \left( \frac{m_k}{m_j} \right)^b$$

{#eq:id2}

We now turn  $n_{j,k}$  into a herbivore population:

$$n_{i,k} = \frac{\frac{n_k c_k}{\tau c_j} \left( \frac{m_k}{m_j} \right)^b c_j}{\tau c_i} \left( \frac{m_j}{m_i} \right)^b$$

{#eq:id3}

which gives:

$$n_{i,k} = \frac{n_k c_k}{\tau^2 c_i} \left( \frac{m_k}{m_i} \right)^b$$

{#eq:id3b}

In a similar fashion, for a linear trophic chain, we can demonstrate that the additional population of herbivore  $i$  to be produced to maintain predator  $j$  of level  $l$  is:



$$n_{i,k} = \frac{n_j c_j}{\tau^l c_i} \left( \frac{m_j}{m_i} \right)^b$$

{#eq:id4}

In many cases, a predator feeds on a set of prey rather than a single one. In such case, the energy uptake should be shared by the different sources. Basically, the split of energy is the realm of populations dynamics as the population consumption depends on individuals number. To overcome the complexity such consideration would bring, we assume that energy costs associated with the local maintaining of a predator are minimal. Therefore,  $n_j$  is converted into  $i$ , the herbivore linked to  $j$  for which (?) is minimal. Basically,  $i$  should be a large and separated from  $j$  by a low number of species. Hence, on the island, the species richness increases as long as the inequality below holds true:

$$\sum_i c_i n_i m_i^b + \sum_j \frac{n_j c_j}{\tau^{l_j} c_i} \left( \frac{m_j}{m_{i_j}} \right)^b < E_0$$

{#eq:id5}

The first term is the energy cost for maintaining populations of herbivores, the second term is associated to higher trophic levels: predator  $j$  is converted into herbivore  $i_j$  from which it is separated by  $l_j$  links. For the sake of simplicity, we make an extra assumptions:  $c_i$  values are constant among species and set to 1. Therefore is we assume that MVP is constant (scenario 1), inequality ? becomes:

$$\sum_i m_i^b + \sum_j \frac{1}{\tau^{l_j}} \left( \frac{m_j}{m_{i_j}} \right)^b < \frac{E_0}{n_0}$$

{#eq:id5a}

Similarly, if we assume  $n_i = n_0 m_i^{-b}$  and  $n_j = n_0 m_j^{-b}$  (scenario 2), then:

$$\sum_i 1 + \sum_j \frac{1}{\tau_j} \left( \frac{1}{m_{ij}} \right)^b < \frac{E_0}{n_0}$$

{#eq:id5b}

The left side of this equation provides the minimal energy needed to sustain all populations while the right side is the total amount of energy available. Therefore, the maximal population of species  $i$  without any extinction is given by converting the extra amount of energy available into an additional population of species  $i$ , the population of  $i$  we get is denote  $n_{i,max}$ . The range  $[n_i, n_{i,max}]$  is then the range of possible fluctuations of species  $i$  without any new extinction event.

### 3.2.6 Extinctions

When the local primary production cannot sustain the establishment of a new immigrant, *i.e.* when inequality ? is no longer verified, its arrival is either impossible or lead to extinction of other species. In the latter situation, we must determine the identity of species to go extinct. This is a challenge that remains we will not take up here especially given its complexity highlighted by recent theoretical studies (Säterberg et al., 2013; ?). To overcome this difficulty, we use two scenarios: 1) *random extinction*: any species can go extinct and 2) *costs-based extinctions*: the probability of extinction is proportional to the energetic costs of the species. Once a extinction happened we ensure that all species remained link to at least on herbivore, otherwise unlinked species go extinct. Furthermore extinctions occur until ? is satisfied.

Although the assumptions we have made here are likely unrealistic, they all intend to examine the model under the special case of an optimal energy allocation on island. As long as we focus on the qualitative consequences in term of community dynamics, these assumption remain acceptable. As an important remark, in our model the energy cost associated to one predator is not an inherent properties as it is determined by the identity of preys available

on the island.

### **3.2.7 Simulations**

## **3.3 Results**

## CONCLUSION GÉNÉRALE

### Introduire davantage d'écologie des communautés dans les modèles de distribution d'espèce

#### L'information des réseaux

L'effort essentiel de ma thèse porté sur la caractérisation du questionnement sur les interactions d'espèces m'a conduit à constater qu'il y a un problème d'échelle biologique.

Dans le projet ici présenté, nous proposons de construire des modèles théoriques plus intégratifs en repartant d'un modèle théorique classique, celui de la théorie de la biogéographie des îles proposée par MacArthur et Wilson [MacArthur and Wilson \(1967\)](#). Dans un premier temps, nous y ajoutons les interactions entre espèces et une relation explicite avec l'environnement abiotique au travers d'une approche communauté centrée qui étend le modèle classique. Dans un second temps, nous combinons une approche population centrée et les processus évolutifs pour une biogéographie insulaire plus mécaniste. Enfin, au regard des enjeux que soulève le rôle des interactions entre espèces dans la construction de la biodiversité, nous réfléchissons sur l'inférence d'espèces interdépendantes.

différentes théories pour différentes échelles ??

De par son pouvoir explicatif et son élégance, le modèle de MacArthur et Wilson est un point de départ approprié pour construire des modèles plus intégratifs en intégrant explicitement des processus écologiques et évolutifs. Cette idée n'est pas nouvelle et les auteurs de la TIB ont étudié un certain nombre de processus écologiques. Notamment, ils ont intégré les phénomènes de spéciation [MacArthur and Wilson \(1967\)](#) et réfléchis sur l'importance des interactions quant à la répartition des espèces ?. Néanmoins, dans le modèle classique, l'ensemble de ces aspects sont absents, l'idée que les processus écologiques importent peu aux larges échelles domine. Nous allons, dans ce projet, à l'encontre de cette idée et proposons de construire des modèles intégratifs qui étendent la TIB.

### 3.3.1 Interaction des différents mécanismes

L'interaction de différents L'ensemble des trois éléments jusqu'ici évoqués (environnement abiotique, interaction, évolution) peuvent également être étroitement associés. Grant et Grant 2006 rapportent le cas de la compétition entre trois espèces de pinsons (dits de Darwin) sur l'île de Daphne (Galapagos) qui engendre une modification de la taille de leurs becs. Cette évolution liée à la compétition est elle-même reliée à l'environnement abiotique car, par l'abondance ou l'absence de précipitations, il détermine la disponibilité des ressources et donc l'intensité de la compétition ?. A travers cet exemple, nous comprenons l'importance d'inclure l'ensemble des différents processus pour construire un modèle intégratif en biogéographie. Un tel modèle serait capable, par exemple, de renseigner les risques d'exclusion compétitive dans l'exemple décrit par Grant et Grant.

### Vers une théorie en intégrative de la biogéographie

Dans la réédition de 2001 [] Wilson rappelle que le problème :

“The flaws of the book lie in its oversimplification and incompleteness, which are endemic to most efforts at theory and synthesis.”

Suite à mes travaux présentés dans cette thèse, je pense avoir réellement compris les lacunes théoriques de la Biogéographie relevées il y a plus de 15 ans par Mark Lomolino [Lomolino2000]. En ouverture d'un numéro spécial dédié à la biogéographie des îles, il insistait sur le besoin d'introduire davantage d'écologie dans la biogéographie. Je me range de son côté et de celui de différents auteurs pour voir un modèle en biogéographie articuler autour de trois composantes fondamentales de la biogéographie : migration, extinction et évolution. Comment le reste de l'écologie affecte et influence l'interplay de ces processus...

L'effort théorique nécessaire en biogéographie porte sur l'intégration ordonnée de concepts clés issus de différents champs de l'écologie [Thuiller et al. \(2013\)](#). Ainsi, alors que les condi-

tions climatiques et plus généralement la géographie physique sont classiquement évoquées pour expliquer la répartition des espèces [Kearney and Porter \(2004\)](#), les interactions entre espèces sont quant à elles souvent occultées. De même, bien que les processus évolutifs soient souvent évoqués comme déterminants majeurs de la diversité des espèces, leurs effets à court terme sont souvent ignorés dans les scénarios décrivant la biodiversité de demain [Lavergne et al. \(2010\)](#). La difficulté principale est alors de produire des modèles (théoriques en première instance) qui intègrent l'ensemble des processus et les relations qu'ils entretiennent [Thuiller et al. \(2013\)](#) tout en gardant une relative simplicité. Une théorie intégrative en biogéographie pourrait être le meilleur point d'ancrage pour construire de nouvelles approches appliquées. Avec une telle théorie en main, nous pourrions aller vers l'enjeu majeur de ces dernières années en biogéographie : relâcher les hypothèses que les modèles classiques de répartition des espèces d'aujourd'hui utilisent (notamment en occultant les interactions) pour prédire la biodiversité de demain [Guisan and Rahbek \(2011\)](#).

## **Des contraintes physiologiques**

### **3.3.2 L'abstraction des espèces**

L'abstraction de l'espèce @Poisot2015 pour des questions centrales : - quelles espèce va interagir avec qui ?%

## **Traits fonctionnels**

Les traits fonctionnels sont des propriétés mesurables sur les organismes en relation avec leurs performances et leur rôle dans l'écosystème [McGill et al. \(2006\)](#). Les traits étudiés peuvent être de différentes natures, 1-morphologiques : taille de différentes parties du corps, position des yeux, taille des oeufs chez les organismes ovipares, taille des graines pour les végétaux, 2- physiologiques : taux métaboliques de bases, stœchiométrie (rapport de la

concentration entre divers éléments qui compose l'organismes) [McGill et al. \(2006\)](#); ?); ?. Un ensemble approprié de ces propriétés peut être un outil puissant pour décrire un ensemble d'espèce dans un même espace. Leur proximité dans l'espace des traits est alors un indice précieux d'une proximité fonctionnelle. Ainsi, à l'aide de 13 traits ecomorphologiques, Albouy *et al.* 2011 parviennent à prédire les guildes trophiques de 35 espèces de poissons de la Méditerranée ?. Edwards *et al.* 2013 montrent que l'effet saisonnier sur une communauté de phytoplancton dans la Manche peut être capturé à l'aide de traits décrivant : le taux maximal de croissance, la compétitivité pour la lumière et l'azote ?. La distribution des traits fonctionnels au sein de la biodiversité est aussi une entrée de choix pour réfléchir quand à la fragilité potentielle des fonctions remplies par les écosystèmes ?. %DG : je comprends cette citation de Mouillot, mais juste une mise en garde contre ce type de référence. Mouillot se base sur l'hypothèse que les traits nous informent du fonctionnement, sans jamais documenter cette relation. Ce qui est souvent le cas, et par conséquent contribue à bâtir des mythes dans la littérature qui à l'occasion ne sont pas toujours bien appuyés. L'approche par traits est un bel exemple, on a édifié rapidement une structure conceptuelle sur les traits, mais on n'a pas solidement appuyé le concept sur de bonnes bases empiriques.

L'approche de la biodiversité par les traits fonctionnels est plus quantitative que l'approche taxonomique et permet de déduire un grand nombre de propriétés en se passant de la connaissance de leur identité. Ainsi McGill, dans son article d'opinion de 2006, propose une approche nouvelle de l'écologie des communautés qui transforme les questions centrées autour des espèces par des questions qui interrogent la répartition et la variabilité des traits [McGill et al. \(2006\)](#). L'emploi des traits fonctionnels est en fait un appel à une écologie plus mécaniste, qui se penche sur la physiologie des organismes, en prend les faits les plus importants (relativement au problème traité) pour les placer dans un espace de traits commun. Cette approche est aussi en lien avec la controversée théorie métabolique en écologie [Brown et al. \(2004\)](#); ?. Dans cette théorie un certain nombre de grandeurs (comme le taux métabolique) sont reliées à la biomasse corporelles de l'adulte, fournissant ainsi en un seul trait de nombreuses relations pour des groupes d'organismes très différents. Par ces nouvelles approches,

l'espérance de s'extraire de la seule identité des espèces est accrue, l'idée d'avoir des règles générales se concrétise.

Dans une théorie intégrative de la biogéographie, les traits fonctionnels peuvent être un pivot très intéressant pour rassembler les différents concepts que nous avons développés dans les paragraphes précédents. Les traits peuvent tout d'abord être mis en relation avec le milieu abiotique. Le taux métabolique ou encore la sensibilité à la sécheresse sont des indices performant pour décrire la survie dans un milieu donné [Kearney and Porter \(2004\)](#); [Engelbrecht et al. \(2007\)](#) que l'on peut capturer sous forme de traits. Kearney *et al.* 2010 propose une approche prometteuse dans laquelle, l'environnement physique, la disponibilité des ressources et la dynamique énergétique sont reliées par les traits fonctionnelles le tout aboutissant à un modèle de distribution très mécanistes. La structure d'un réseaux peut également être dérivée à partir de l'espace des traits. Dans leur méthode proposée cette année, Gravel *et al.* infèrent les paramètres du modèle de niche de Williams et Martinez [Williams and Martinez \(2000\)](#) à partir des relations de masse du corps entre proie et prédateurs [Gravel et al. \(2013\)](#). Ils sont alors en mesure de dériver un réseau global pour un ensemble d'espèce donné. Enfin, en tant qu'expression phénotypique, les traits fonctionnels sont soumis aux processus évolutifs. Sur les temps longs, l'expression de l'évolution résulte en la modification progressive des traits qui se répercute sur l'ensemble des propriétés qui en découle. Ainsi la considération d'une modification des traits est une approche simple et réaliste pour introduire les processus évolutifs et leurs conséquences ??.

### **Des données nouvelles**

Comme souvent en écologie / science nous avons besoin de données, mais ce n'est pas une question vaine, L'Accumulation des données doit se faire avec une certaine normalisation pour utiliser les. Il est souvent difficile et la conséquence c'est de trouver des difficultés pour réintégrer des anciennes données @Tingley2009b celle des muséum @Shaffer1998 Malgré les espoirs des remplacer les ordinateurs pour formuler les hypothèse, toujours besoin d'un



développement théorique plus de que de corrélations essayer d'estimer aujourd'hui en utilisant le plus près possible la méthode d'hier pour savoir quel biais probable il y avait. Ici si on détecte beaucoup plus bas qu'avant avec la même méthode, alors on peut se dire que le fait que ce soit des fausses absences est faible. Par contre si on essaie d'avoir des comparaisons et que les résultats sont dus à la période de l'année... C'est plus compliqué ! Aller vers des occupancy models

## **Les défis à relever dans un monde en changement**

### **Une érosion de la biodiversité affolante**

L'érosion de la biodiversité exerce une certaine nostalgie qui parfois conduit à une forme de fatalisme chez certains experts. Relevons la tête il va falloir trouver les solutions dans le mimétisme ?

Allant jusqu'à des problèmes de santé La tique la souris le réservoir et des hommes des problèmes de productions

Sommes nous en train de biaiser le signal phylogénétique ? (cf article Thuillier)

### **3.3.3 Avons-nous des espoirs vains ?**

Le royaume de la contingence du à l'impact historique de l'histoire évolutive. Alors comment trouver des espoirs de généralité quand le moteur repose sur de la stochasticité Mais cette loi mène à des prédictions exoécologie Les bactéries mais comment généraliser alors que l'évolution a fait émerger bon nombre d'organismes qui en soi loin quoique complètement imbriqués on a plus de micro-organismes que de cellules...

inertie historique comment imaginer des plantes sans mycorrhizées mais d'autres systèmes auraient pu marcher En fait quand on pense à la plante on pense à l'immunité de la plante +

mycorrhizales et quand on pense à un vertébré on inclut toutes ces bactéries on ne peut certes pas comprendre comment l'un marche sans l'autre mais pour on a pas besoin de tout connaître c'est un problème de rupture de symétrie.

Les conséquences sont compliquées des changements climatiques sont nombreuses et certaines espèces voient le range grandir d'autres diminuer pour certaines espèces de co-existent et donc à un changement prononcé de la morphologie des communautés alors que le nombre d'espèces peut être peu affecté @Moritz2008

On nous fait miroiter que finalement que l'érosion de la biodiversité est dramatique et le ressort actuel pour faire un levier face à cela c'est les services écosystémiques qui sont actuellement l'argument choc pour renforcer la production de la nature. Il y a un côté pervers qui est la financiarisation et la substituabilité l'argent peut alors être utilisée pour intervertir ou alors remplacer un type d'écosystème par un autre ailleurs... En fait on a l'impression que c'est plus un principe de précaution qui est invoqué et ultimement il est vraisemblable que la destruction de la nature telle que nous la connaissons soit dans le futur un générateur de conflit... et ultimement on a à craindre de faire une planète invivable pour nous même. Mais les changements sont des remplacements et pour la conservation on peut se demander la stratégie. Dans son article 'Don't judge a species on their origin' Mark Davis prend à revers un certain nombre d'idées reçues et souligne que les effets des envahisseurs peuvent être positifs @Davis2011.

### **Des écosystèmes bouleversés**

Reconfiguration des écosystèmes naturels il y a eu d'autres crises avant. Finalement avec du catastrophisme, la question s'est si nous on ira mal. On est gros pour la taille de la planète peut être plus susceptible à l'extinction que l'on pense. Mais ce n'est pas le pessimisme qui m'importe. - Et si on faisait rien pour le frelon asiatique ?

## Transients

### Vers une écologie prédictive ?

#### Un espoir

La défense des modèle climqtaur bioclimate enveloppe de Pearson comme une dpremière approximation utilise se faitt sur 3 exemple de plantes @Pearson2003

## Catégories

L'écologie ne traite pas de telle ou telle manière les différentes espèces. Il y a des champs entier dédié à des classes d'espèces. Par exemple, on traite mcro-organisme ou me-taphyte vs. metazoaire. Les échelles et la proximité nous biaise fortement la vision de la biodiversité. Il est difficile d'appliquer les théories à l'ensemble des espèces, peut être seule-ment la théorie de l'évolution mais se qui est intéressant c'est que ce processus est finalemnt l'essence de la vie. L'aphrisne d e Dobzansky à mon avis ne devrait simpemtn dire que la bio-logie est à regarder à la lumière mais que le vivant est une ensemble dde moécule organique dont l'organisation est soumise à l'évolution. Il existe des sepécificités des êtres.

Malgré les apparences, La TIB n'est pas formulée pour l'ensemble des espèces. Le premier exemple du livre herpetofaune puis les fourmis de la famille des Ponerinae [@Ma-cArthur1967]. De même la théorie neutre a été classique tropical les arbres dans els forêts tropicaux et les coraux. Biensur ces théories sont liées à la connaissance fine mais biaisée du vivant (comment avoir une connaissance exhaustive du vivant...). Aisin, certaine théorie s'applique à certaine partie du monde vivant, il y aurait donc une classification à faire ou une compréhension du côté de pourquoi telle ou telle propriété est Ok pour telle ou telle partie diu vivant. Cela conduit à une intérogation sur les règles possibles de composition des écosystèmes.

Wallace n'aurait-il pas eu plus de mal à comprendre les zones aujourd'hui. Si naïvement on réduit aux villes, l'homogénéité ++ mais avec les espèces invasive le signal est fortement brouillé aussi ! Je pense qu'on est à un tournant de la biologie vers un changement de paradigme communautaire centré qui ne nait pas les travaux précédant mais les suit.

Information dans les distributions gecko australien généraliste *Heteronotia binoei* => alors peut être que ça marche bien mais sur une espèce spécialiste ??

## Les produits de l'évolution

Quelles hypothèses pouvons nous faire sur les produits de l'évolution ? Si on peut supposer qu'il y a des compétitions ou la règle est le changement cette même propriété a-t-elle des propriétés sur le long terme. Peut-on affirmer que les produits de l'évolution dans un environnement stable amène à des entités qui optimisent l'utilisation de l'énergie. Si oui, que dire des produits de l'évolution dans avec variation. Si on peut faire des hypothèses comment les tester. Dans l'article de

Si l'évolution est imprévisible si au delà d'un certain temps on ne peut presque rien dire... Si la chance de des abeilles européennes changeait comment prédire cela changement de comportement mais que nous sommes dans l'incapacité de le prédire que pensent du status de l'écologie et de l'évolution en tant que science. Si la composant historique domine le royaume de la biologie devons-nous nous contenter de le décrire. L'espoir mais la publication de Ian Hattin eût faire douter de l'absence de l'absolue de règle. Comment croire qu'il n'y a pas des principes d'ordre énergétique là-dessous. Convergence...

2014, Hurlbert et Stegen propose une série d'hypothèses pour mettre en évidence l'impact de l'énergie sur l'évolution la troisième hypothèse est temps suffisant pour l'équilibre. Une telle hypothèse une forme de maximisation de la production de la biomasse et l'utilisation qui est peut être. Peut-être que les différents mécanismes en jeu dans les processus évolutifs amène probablement à une forme de stationarité...

### Quels types de prédictions pouvons-nous faire ?

Quels objet qu'est ce qui peut faire l'objet d'une prédiction / qu'est ce qui ne peut pas ?

#### 3.3.3.1 Une question d'échelle ?

L'écologie porte sur l'ensemble du monde vivant quel que soit leur taille mais les différents champs ne sont pas toutes relatives à la même échelle alors il y a bien des échelles de temps, les échelles spatiales mais il y a le niveau d'organisation. Il est bien important de comprendre cela !

Un schéma avec des variables qui émergent avec différents paramètres et quelques exemples de théorie ! (DEB Evolution foodweb... ) et l'action de

Repartition des espèces des passages historiques dans l'origine des espèces et dans Wallace. Le principe même de l'écologie (la définition de l'écologie). On arrive à l'idée de la niche. Exemple historique. Dans son ouvrage, le grand biogéographe Wallace reconnaît en introduction le caractère fascinant de la répartition de la biodiversité des îles avec des faits intrigants quant à la faune et la flore. Ainsi il constate qu'il peut y avoir plus de différence entre île très éloignée et deux îles très proches. Il écrit que la faune et la flore sont plus dissimilaires entre les deux îles des Galapagos Bali et Lombok qu'entre Hokkaido (Japon) et La Grande Bretagne ou encore la Nouvelle Zélande et l'Australie,

Exemple classique de Grinnell et des Trasher + évolution avec les caractéristiques de déplacement.

Nous accumulons des évidences quand à l'impact du changement anthropique. A différentes échelles la diminution de la biodiversité, changement en composition @Taranu2015 @DeRoos2008

La biogéographie avec au moins 3 problèmes d'échelles => spatiale => temporelle plus on augmente plus l'empreinte historique est forte => grands événements géologiques (l'extinction)

mouvement des plaques) biogéographies historiques mais aussi forme un pool d'espèces => Mais aussi l'échelle taxonomique : la relation aire espèce est décrite à l'intérieur des taxons les relations allométriques à l'intérieur des taxons E O Wilson a commencé à rapporter des relations sur les formes les exemples du livre sont herpète faune (reptile plus amphibien) mécanisme => diversité de milieu

### **3.3.3.2 Prédire des communautés**

#### **Les dangers d'aller trop vite**

There is also a danger that predictions grow faster than our understanding of ecological systems, resulting in a gap between the scientists generating the predictions and stakeholders using them

“Predictive ecology in a changing world” [@Mouquet2015]

### **3.3.4 Contraintes énergétiques**

Moi je pars vers ça..

=> des interactions changer de paradigme => abstraction des espèces

## ANNEXE I

### COMMENT LA BIODIVERSITÉ S'INSTALLE EN TERRITOIRES ISOLÉS

#### Préambule

La biogéographie est la science qui s'interroge sur les causes de la répartition de la biodiversité dans les différentes parties du globe (voir encadré 1). Le modèle déterministe de MacArthur et Wilson décrit l'évolution de la richesse spécifique sur les îles vers un équilibre, mais la migration des espèces vers des îles et leur extinction potentielle sont des phénomènes aléatoires. Comment un modèle déterministe peut-il renfermer ce hasard ? C'est ce que nous allons voir.

#### 4.4 Le modèle de MacArthur et Wilson

Un des modèles les plus puissants en biogéographie est celui proposé par MacArthur et Wilson dans leur passionnante théorie de la biogéographie des îles. Ces deux illustres prédécesseurs nous ont livré un paradigme simple et puissant pour envisager la construction de la biodiversité sur une île. Notons dès maintenant que l'île n'est pas nécessairement un monticule de sable fin au milieu de l'océan mais plutôt -et plus généralement- un territoire isolé. Le modèle permet de comprendre l'impact des capacités de dispersion et de survie des espèces sur la biodiversité  $S$  de l'île considérée. Considérons que cette île est accessible depuis un continent qui présente un ensemble constant d'espèces  $P$ . Les  $P$  espèces sont potentiellement celles que nous retrouverons sur l'île, nous avons donc  $S \leq P$ . À un temps

$t$  donné, l'ensemble des  $S$  espèces de l'île peuvent s'éteindre avec un taux  $e$ . De même, les  $P - S$  espèces du continent (absentes de l'île) peuvent coloniser l'île avec un taux  $c$ . En langage mathématique, nous obtenons l'équation différentielle déterministe suivante pour décrire l'évolution temporelle de  $S$  :

$$\begin{aligned}\frac{dS}{dt} &= c(P - S) - eS \\ &= cP - (c + e)S\end{aligned}\tag{4.1}$$

Cette équation est linéaire, nous pouvons la résoudre facilement par la méthode de la variation de la constante (essayez puis regardez l'encadré 2 !). Pour les valeurs arbitraires  $c = 0.2$ ,  $e = 0.1$  et la condition initiale suivante : à  $t = 0$ , l'île est inhabitée ( $S(0) = 0$ ), nous représentons graphiquement la solution de l'équation à la figure 22. Pour un temps infini, la biodiversité tend vers la valeur :  $S_{eq} = P\left(\frac{c}{c+e}\right)$ , interprétée comme le nombre d'espèces provenant du continent et présentes sur l'île à l'équilibre.

## 4.5 Comment un modèle déterministe émerge-t-il de phénomènes aléatoires

### 4.5.1 Modèle stochastique ?

Il est très intéressant de réaliser que l'équation (4.1), d'apparence déterministe, renferme un modèle stochastique. "Stochastique" est un terme signifiant aléatoire, avec une part de hasard. Un modèle stochastique est donc un modèle dont l'issue contient une part de hasard ; deux réalisations du modèle ne donneront donc pas nécessairement le même résultat. Pour comprendre où se cache le hasard dans le modèle que nous étudions, nous avons besoin d'objets mathématiques particuliers appartenant au domaine des probabilités : variables aléatoires et processus stochastiques (reportez-vous à l'encadré 3).



#### 4.5.2 Les objets mathématiques requis

Comment savoir combien d'espèces sont présentes sur l'île ? Il suffit de compter 1 pour chaque espèce présente sur l'île. Nous introduisons donc  $X_i$ , la variable aléatoire de présence sur l'île de l'espèce  $i$  choisie parmi les  $P$  espèces du continent.  $X_i$  est égale à 0 si l'espèce n'est pas sur l'île et 1 si elle est sur l'île. C'est une variable aléatoire de Bernoulli (encadré 4). Ainsi, le nombre d'espèces présentes sur l'île,  $S$ , est égale à la somme des  $X_i$ . Nous allons encore plus loin en enregistrant ces valeurs au cours du temps. C'est ainsi que nous définissons le processus stochastique  $X_{i,t}$ , qui n'est autre qu'une succession de 1 et de 0 indiquant à chaque instant si oui ou non l'espèce  $i$  est sur l'île. A priori, les suites  $X_{i,t}$  ne sont pas prévisibles, ce qui n'exclut pas que la fonction  $S$  présente, en moyenne, un comportement tout à fait régulier.

#### 4.5.3 Les briques élémentaires du modèle stochastique

Au temps  $t = 0$ , l'espèce  $i$  n'est pas sur l'île. Pour construire la suite de son histoire, il nous faut une règle simple pour décrire l'évolution de sa présence entre deux pas de temps très proches. Le modèle de MacArthur et Wilson postule que si l'espèce considérée est sur l'île au temps  $t$ , elle s'éteint avec un taux  $e$  ; si elle est sur le continent, elle le colonise avec un taux  $c$ . Pour simplifier, nous supposons que les taux d'extinction et de colonisation sont les mêmes pour toutes les espèces. Nous donnons maintenant une signification en terme de probabilité à ces taux :  $c$  est la probabilité de colonisation par unité de temps (on parle de densité du processus),  $e$  est la probabilité d'extinction par unité de temps. Ainsi,  $edt$  désigne la probabilité d'extinction pendant l'intervalle de temps  $dt$ . De même,  $cdt$  est la probabilité de colonisation pendant  $dt$ . En faisant appel aux probabilités conditionnelles (encadré 5), en

supposant  $dt$  assez petit, nous différencions quatre cas :

$$\forall t \in \mathbb{R}^+, P(X_{i,t+dt} = 1 | X_{i,t} = 0) = cdt \quad (4.2)$$

$$P(X_{i,t+dt} = 0 | X_{i,t} = 1) = edt \quad (4.3)$$

$$P(X_{i,t+dt} = 0 | X_{i,t} = 0) = (1 - cdt) \quad (4.4)$$

$$P(X_{i,t+dt} = 1 | X_{i,t} = 1) = (1 - edt) \quad (4.5)$$

Il faut comprendre la première équation ainsi : sachant que l'espèce  $i$  était absente au temps  $t$ , la probabilité qu'elle soit sur l'île au temps  $t+dt$  est égale à la probabilité qu'elle colonise l'île durant  $dt$ , c'est-à-dire  $cdt$ . Vous pourriez avancer que durant  $dt$ , une espèce peut coloniser, s'éteindre et re-coloniser et que nous n'en parlons pas ! Tous ces évènements sont en fait beaucoup moins probables et nous pouvons les ignorer complètement à la limite quand  $dt$  tend vers 0. Les trois autres équations s'interprètent avec un raisonnement similaire. Ces quatre équations sont les briques du modèle stochastique que nous construisons. Nous devons maintenant les assembler correctement.

#### 4.5.4 Assemblons les briques !

L'assemblage consiste en l'utilisation de la formule des probabilités totales expliquée dans l'encadré 5. Nous pouvons en déduire la probabilité de présence de l'espèce  $i$  à l'instant  $t + dt$  :

$$P(X_{i,t+dt} = 1) = P(X_{i,t+dt} = 1 | X_{i,t} = 0)P(X_{i,t} = 0) + P(X_{i,t+dt} = 1 | X_{i,t} = 1)P(X_{i,t} = 1) \quad (4.6)$$

Il s'agit d'une somme de deux termes couvrant toutes les possibilités puisque, à l'instant  $t$ , l'espèce  $i$  était soit absente de l'île soit présente, c'est un système complet d'évènements (encadré 5). Si l'espèce  $i$  était absente (premier terme) à  $t$ , elle sera présente à  $t + dt$  si elle colonise pendant  $dt$  (4.2). Si elle était déjà sur l'île à  $t$ , elle s'y maintient à condition de ne

pas s'éteindre (4.5). En remarquant que  $P(X_{i,t} = 0) = 1 - P(X_{i,t} = 1)$ , nous pouvons écrire :

$$P(X_{i,t+dt} = 1) = cdt(1 - P(X_{i,t} = 1)) + (1 - edt)P(X_{i,t} = 1) \quad (4.7)$$

Pour simplifier l'écriture,  $s_{i,t}$  représente  $P(X_{i,t} = 1)$  :

$$s_{i,t+dt} = cdt(1 - s_{i,t}) + (1 - edt)s_{i,t} \quad (4.8)$$

Divisons par  $dt$  :

$$\frac{s_{i,t+dt} - s_{i,t}}{dt} = c(1 - s_{i,t}) - es_{i,t} \quad (4.9)$$

Nous faisons alors tendre  $dt$  vers 0. Nous obtenons alors :

$$\frac{ds_i}{dt} = c(1 - s_i) - es_i \quad (4.10)$$

C'est l'équation que nous avons résolu dans l'encadré 1, avec  $P = 1$  :

$$s_i(t) = \frac{c}{c + e} (1 - \exp(-(c + e)t)) \quad (4.11)$$

#### 4.5.5 Retrouvons le modèle classique

Nous avons presque retrouvé la solution de l'équation déterministe ; vous pourriez penser qu'il faut simplement multiplier par le nombre d'espèces  $P$ . L'idée est bonne, mais demande justification ! Nous allons effectivement considérer non pas une mais bien les  $P$  espèces du continent pour lesquelles l'équation (4.11) est valable. Nous allons alors définir un nouveau processus stochastique qui est simplement la somme des  $P$  processus de présence que nous supposons indépendants  $X_{i,t>0}, Y_t > 0 = X_{1,t>0} + X_{2,t>0} + \dots + X_{P,t}$ . Le problème est alors de connaître à tout instant  $t$  la probabilité d'avoir un nombre donné  $k$  d'espèces présentes sur l'île. À un instant donné  $t$  nous faisons une somme de variable aléatoires de Bernoulli. La

somme de  $P$  variables de Bernoulli indépendantes est une variable aléatoire suivant une loi binomiale (encadré 5). Nous avons donc :

$$\forall t \in \mathbb{R}^+, P(Y_t = k) = \binom{P}{k} s(t)^k (1 - s(t))^{P-k} \quad (4.12)$$

De plus, on peut démontrer les résultats suivants :

$$\forall t \in \mathbb{R}^+, E(Y_t) = Ps(t) \quad (4.13)$$

$$\forall t \in \mathbb{R}^+, V(Y_t) = Ps(t)(1 - s(t)) \quad (4.14)$$

Grâce à ces résultats de probabilité, nous avons donc :

$$\forall t \in \mathbb{R}^+, E(Y_t) = P \frac{c}{c + e} (1 - \exp(-(c + e)t)) \quad (4.15)$$

Nous retrouvons la solution déterministe ! Ainsi, le modèle déterministe peut être considéré comme reflétant l'évolution de l'espérance des variables  $Y_t$  pour  $t > 0$ , c'est donc l'espérance du processus stochastique. Nous pouvons alors simuler le modèle stochastique grâce au modèle déterministe : à chaque instant  $t$ , il suffit de simuler une loi binomiale de paramètres  $P$  et  $E(Y_t)$ . Regardez la figure 23, il y a des différences fondamentales entre les deux approches. Le modèle déterministe donnera des valeurs continues et, pour des paramètres donnés, toujours le même résultat. De son côté, le modèle stochastique livrera des valeurs discrètes. De plus, deux simulations du modèle aléatoire ne donneront pas nécessairement les mêmes courbes.

## 4.6 Applications et perspectives

Bien que datant du milieu des années 1960, le modèle de MacArthur et Wilson demeure très précieux, notamment pour étudier les habitats fragmentés (des îles !) et prendre des

décisions relatives à la conservation des espèces. Prenons un exemple : vous êtes le gestionnaire d'une nouvelle zone protégée que nous assimilerons à une île. Cette zone a longtemps été exploitée par l'homme de sorte qu'au temps  $t = 0$  le nombre d'espèces est  $S = 0$ . Dans la région, la richesse en espèces est de  $P = 100$ , c'est-à-dire que vous pouvez rencontrer jusqu'à 100 espèces dans votre aire de gestion. À la suite de MacArthur et Wilson, nous allons supposer que la valeur du taux de colonisation  $c$  dépend de la difficulté d'accès. De plus, la survie d'une espèce sur cette zone dépend de sa superficie : ainsi plus l'aire de conservation est grande, moins les espèces s'éteignent ;  $e$  est donc plus faible pour les grandes aires de protection. Prenons trois situations :

- Grande zone et accès aisé :  $e=0.01$  et  $c=0.05$
- Petite zone et accès aisé :  $e=0.05$  et  $c=0.05$
- Petite zone et accès délicat :  $e=0.05$  et  $c=0.01$

Pour mettre en place un suivi et savoir comment se comporte votre réserve, il faut avoir une idée de la dynamique attendue ! Ce que nous permet le modèle étudié, que nous simulons pour les trois situations, avec les deux approches du modèle. Les résultats sont donnés par la figure 1. Cette simple démarche vous permet d'estimer l'évolution de la biodiversité au sein de votre réserve. La paramétrisation est plus compliquée dans les faits, mais l'idée est très utilisée pour apprécier la santé des réserves naturelles.

Dans cet exemple, le modèle stochastique n'a finalement que peu d'intérêt. Il est cependant très utile si vous êtes intéressé par la variance du modèle (que nous obtenons avec l'équation (4.14)). C'est aussi un bon point de départ pour les démarches statistiques, indispensables pour le suivi de votre réserve. De plus, l'approche stochastique révèle toute sa force pour enrichir le modèle. En partant de l'équation (4.1), si vous souhaitiez introduire un effet des espèces les unes sur les autres (les interactions entre espèces), vous auriez quelques difficultés. Avec le travail réalisé ici vous pouvez trouver et surtout justifier des approches fertiles en réfléchissant en terme de probabilité. Ces approches pourraient vous amener à résoudre des systèmes d'équations différentielles déterministes un peu plus compliqués !

**Encadré 1 : la Biogéographie**

La biogéographie est la science qui s'interroge sur les causes de l'agencement spatial des espèces à la surface de notre planète. Comment, par exemple, pouvons-nous expliquer la répartition et les différences entre les grands biomes terrestres ? Le premier élément de réponse est l'implication des facteurs climatiques. La température, l'humidité, les précipitations sont autant de variables qui contraignent les conditions d'existence des espèces qui composent et modèlent les paysages terrestres. C'est avec ces contraintes que les écologues élaborent des modèles dits "bioclimatiques" afin de décrire l'évolution de la biodiversité avec les contraintes climatiques de demain. Cependant, les facteurs qui expliquent la distribution des espèces ne sont pas toujours climatiques, les causes sont multiples. Les mouvements des espèces et leurs histoires évolutives sont ainsi des moteurs fondamentaux en biogéographie. Les capacités de dispersion des espèces leur donnent accès à de nouveaux territoires qui, avec le temps peut-être, les verront disparaître. C'est ainsi qu'un territoire voit sa composition en espèces évoluer, avec l'arrivée de nouvelles espèces et l'extinction d'autres. C'est autour de cette idée majeure que s'articule le modèle présenté.

**Encadré 2**

Il est facile de vérifier que la solution générale de l'équation homogène :

$$\frac{dS}{dt} = -(c + e)S$$

est donnée par :

$$\forall t \in \mathbb{R}^+, S(t) = C \exp(-(c + e)t)$$

où  $C$  est une constante, la solution de l'équation non homogène est alors de la forme :

$$S(t) = f(t) \exp(-(c + e)t)$$

En remplaçant dans l'équation

différentielle on obtient :

$$f'(t) \exp(-(c + e)t) = cP$$

ce qui nous donne :

$$f'(t) = cP \exp((c + e)t)$$

En intégrant on obtient :

$$f(t) = P \frac{c}{c + e} \exp((c + e)t) + K$$

où  $K$  est une constante arbitraire.

$$S(t) = P \frac{c}{c + e} + K \exp(-(c + e)t)$$

Avec  $S(0) = 0$ ,  $K = -P \frac{c}{c + e}$ , nous obtenons donc :

$$S(t) = P \frac{c}{c + e} (1 - \exp(-(c + e)t))$$

### Encadré 3

Les **variables aléatoires** sont des fonctions définies sur l'ensemble des résultats possibles d'un événement aléatoire. Comme le résultat de l'événement est aléatoire, on associe à chaque valeur de la variable sa probabilité, de sorte que la somme des probabilités sur l'ensemble des valeurs possibles soit égale à 1. Dans notre exemple,  $X_i$  est une variable aléatoire, 0 et 1 sont ses valeurs, et nous nous intéressons à leur probabilité  $P(X_i = 1)$  et  $P(X_i = 0)$  avec  $P(X_i = 1) + P(X_i = 0) = 1$ .

Les **processus aléatoires ou stochastiques** sont des collections de variables aléatoires indexées par le temps  $t$ , c'est-à-dire ordonnées. Dans notre exemple, l'indexation est faite selon le temps  $t$ . Nous suivons ainsi l'évolution de  $X_i$  au cours du temps d'où la notation  $X_{i,t>0}$ . Attention !  $X_{i,t>0}$  est un processus aléatoire mais  $X_{i,t}$  est l'une des variables aléatoires du processus.

### Encadré 4

Une variable aléatoire  $X$  suit un **schéma de Bernoulli** lorsqu'elle prend uniquement deux valeurs 0 ou 1. Si nous notons  $p = P(X = 1)$ , nous avons :

$$— P(X = 0) = 1 - p$$

$$— E(X) = p$$

$$— V(X) = p(1 - p)$$

$X$  est alors une variable aléatoire de Bernoulli de paramètre  $p$ .

Une variable aléatoire  $X$  suit une **loi Binomiale** de paramètres  $n \in \mathbb{N}$  et  $p \in [0, 1]$  lorsqu'elle est la somme de  $n$  variables de Bernoulli indépendantes de paramètre  $p$ .

On peut alors montrer les résultats suivants :

$$— \forall k \leq n, P(X = k) = \binom{n}{k} p^k (1 - p)^{n-k}$$

$$— E(X) = np$$

$$— V(X) = np(1 - p)$$



### Encadré 5

Une **probabilité conditionnelle**, est une probabilité qu'un évènement se réalise à condition qu'un autre soit déjà réalisé. Soit deux évènements  $A$  et  $B$ , la probabilité de  $A$  sachant  $B$ , notée ici  $P(A|B)$  est la probabilité que  $A$  se réalise lorsque  $B$  est réalisé.

Un **système complet d'évènements** est un ensemble d'évènements exclusifs dont la somme des probabilités est 1. Par exemple, l'évènement  $B$  et son complémentaire  $\bar{B}$  forme un tel système (voir figure ??).

La **formule des probabilités totales** nous donne la probabilité de réalisation d'un évènement à partir de la connaissance de probabilités conditionnelles sous réserve que l'ensemble des évènements du conditionnement forment un système complet d'évènements. Ainsi pour connaître la probabilité de  $A$ , nous pouvons écrire :

$$\begin{aligned} P(A) &= P(A \cap B) + P(A \cap \bar{B}) \\ &= P(A|B)P(B) + P(A|\bar{B})P(\bar{B}). \end{aligned}$$




Figure 22: Représentation graphique de la solution de l'équation différentielle (4.1) pour les conditions suivantes :  $c = 0.04$ ,  $e = 0.01$  et  $S(0) = 0$ . La ligne en pointillés représente la valeur de la biodiversité  $S_{eq}$  que nous obtenons à un temps infini.




Figure 23: Dynamique de la biodiversité de trois zones protégées. Les caractéristiques de la zone protégée sont données par les valeurs de  $c$  et  $e$ . Les symboles rouges sont relatifs à une zone protégée de grande taille et facilement accessible, en vert, ils font référence à une petite zone facilement accessible ; enfin, en bleu, sont présentées les résultats pour une petite zone difficilement accessible. Les courbes striées en couleurs pastelées sont les réalisations du modèle stochastique. Les courbes en traits pleins, font référence au modèle déterministe qui est également la moyenne du modèle stochastique. Enfin, les courbes en pointillés représentent la moyenne du modèle stochastique plus ou moins l'écart type.

## RÉFÉRENCES

- Allesina, S., Tang, S., mar 2012. Stability criteria for complex ecosystems. *Nature* 483 (7388), 205–208.  
URL <http://www.ncbi.nlm.nih.gov/pubmed/22343894> 19, 45, 46
- Arabidopsis Genome Initiative, dec 2000. Analysis of the genome sequence of the flowering plant *Arabidopsis thaliana*. *Nature* 408 (6814), 796–815.  
URL <http://www.ncbi.nlm.nih.gov/pubmed/11130711><http://www.nature.com/doifinder/10.1038/35048692> 11
- Araújo, M. B., Luoto, M., nov 2007. The importance of biotic interactions for modelling species distributions under climate change. *Global Ecology and Biogeography* 16 (6), 743–753.  
URL <http://doi.wiley.com/10.1111/j.1466-8238.2007.00359.x> 65, 75
- Araújo, M. B., Rozenfeld, A., dec 2014. The geographic scaling of biotic interactions. *Ecography* 37 (5), 406–415.  
URL <http://doi.wiley.com/10.1111/j.1600-0587.2013.00643.x> 37, 45, 64, 65, 90, 93, 94, 97
- Araújo, M. B., Rozenfeld, A., Rahbek, C., Marquet, P. A., dec 2011. Using species co-occurrence networks to assess the impacts of climate change. *Ecography* 34 (6), 897–908.  
URL <http://doi.wiley.com/10.1111/j.1600-0587.2011.06919.x> 64, 74, 75, 77, 85, 89, 90
- Balanyá, J., Oller, J. M., Huey, R. B., Gilchrist, G. W., Serra, L., sep 2006. Global genetic change tracks global climate warming in *Drosophila subobscura*. *Science (New York, N.Y.)* 313 (5794), 1773–5.  
URL <http://www.ncbi.nlm.nih.gov/pubmed/16946033> 21
- Baselga, A., Araújo, M. B., apr 2009. Individualistic vs community modelling of species distributions under climate change. *Ecography* 32 (1), 55–65.  
URL <http://doi.wiley.com/10.1111/j.1600-0587.2009.05856.x> 75
- Baskerville, E. B., Dobson, A. P., Bedford, T., Allesina, S., Anderson, T. M., Pascual, M., dec 2011. Spatial guilds in the Serengeti food web revealed by a Bayesian group model. *PLoS computational biology* 7 (12), e1002321.  
URL <http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=3248389&tool=pmcentrez&rendertype=abstract> 66
- Beck, J., Ballesteros-Mejia, L., Buchmann, C. M., Dengler, J., Fritz, S. A., Gruber, B., Hof, C., Jansen, F., Knapp, S., Kreft, H., Schneider, A.-K., Winter, M., Dormann, C. F., may 2012. What's on the horizon for macroecology? *Ecography* 35 (March), 001–011.  
URL <http://doi.wiley.com/10.1111/j.1600-0587.2012.07364.x> 9

- Beck, J., Böller, M., Erhardt, A., Schwanghart, W., jan 2014. Spatial bias in the GBIF database and its effect on modeling species' geographic distributions. *Ecological Informatics* 19, 10–15.  
URL <http://dx.doi.org/10.1016/j.ecoinf.2013.11.002><http://linkinghub.elsevier.com/retrieve/pii/S1574954113001155> 17
- Bell, G., jul 2005. The co-distribution of species in relation to the neutral theory of community ecology. *Ecology* 86 (7), 1757–1770.  
URL <http://www.esajournals.org/doi/abs/10.1890/04-1028> 80
- Bellard, C., Bertelsmeier, C., Leadley, P., Thuiller, W., Courchamp, F., jan 2012. Impacts of climate change on the future of biodiversity. *Ecology letters* 15, 365–377.  
URL <http://www.ncbi.nlm.nih.gov/pubmed/22257223> 7
- Belmaker, J., Zarnetske, P., Tuanmu, M.-N., Zonneveld, S., Record, S., Strecker, A., Beaudrot, L., jul 2015. Empirical evidence for the scale dependence of biotic interactions. *Global Ecology and Biogeography* 24 (7), 750–761.  
URL <http://doi.wiley.com/10.1111/geb.12311> 36
- Berlow, E. L., Dunne, J. A., Martinez, N. D., Stark, P. B., Williams, R. J., Brose, U., 2009. Simple prediction of interaction strengths in complex food webs. *Proceedings of the National Academy of Sciences of the United States of America* 106 (1), 187–191. 89, 100
- Black, A. J., McKane, A. J., jun 2012. Stochastic formulation of ecological models and their applications. *Trends in ecology & evolution* 27 (6), 337–345.  
URL <http://www.ncbi.nlm.nih.gov/pubmed/22406194> 47
- Boulangeat, I., Gravel, D., Thuiller, W., jun 2012. Accounting for dispersal and biotic interactions to disentangle the drivers of species distributions and their abundances. *Ecology letters* 15 (6), 584–593.  
URL <http://www.ncbi.nlm.nih.gov/pubmed/22462813> 45, 64, 66, 75, 76, 85, 90
- Brown, J. H., mar 1971. Mechanisms of Competitive Exclusion Between Two Species of Chipmunks. *Ecology* 52 (2), 305–311.  
URL <http://doi.wiley.com/10.2307/1934589> 18, 36
- Brown, J. H., Gillooly, J. F., Allen, A. P., Savage, V. M., West, G. B., jul 2004. Toward a metabolic theory of ecology. *Ecology* 85 (7), 1771–1789.  
URL <http://www.esajournals.org/doi/abs/10.1890/03-9000> 39, 67, 131, 135, 144
- Brown, J. H., Lomolino, M. V., dec 1989. Independent Discovery of the Equilibrium Theory of Island Biogeography. *Ecology* 70 (6), 1954–1957.  
URL <http://doi.wiley.com/10.2307/1938125> 23

Cahill, A. E., Aiello-Lammens, M. E., Fisher-Reid, M. C., Hua, X., Karanewsky, C. J., Ryu, H. Y., Sbeglia, G. C., Spagnolo, F., Waldron, J. B., Warsi, O., Wiens, J. J., jan 2013. How does climate change cause extinction? *Proceedings. Biological sciences / The Royal Society* 280 (1750), 20121890.

URL <http://www.ncbi.nlm.nih.gov/pubmed/23075836http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=3574421{&}tool=pmcentrez{&}rendertype=abstract> 67

Calcagno, V., Massol, F., Mouquet, N., Jarne, P., David, P., oct 2011. Constraints on food chain length arising from regional metacommunity dynamics. *Proceedings. Biological sciences / The Royal Society* 278 (1721), 3042–3049.

URL <http://www.ncbi.nlm.nih.gov/pubmed/21367786> 62

Chamberlain, S. A., Bronstein, J. L., Rudgers, J. A., jul 2014. How context dependent are species interactions? *Ecology Letters* 17 (7), 881–890.

URL <http://www.ncbi.nlm.nih.gov/pubmed/24735225http://doi.wiley.com/10.1111/ele.12279> 93

Chase, J. M., Leibold, M. A., 2003. *Ecological niches : linking classical and contemporary approaches*. The University of Chicago Press, Chicago.

URL [http://www.americanbanker.com/issues/179{ }124/which-city-is-the-next-big-fintech-hub-new-york-stakes-its-claim-1068345-1.html\\$\\delimiter"026E30F\\$nhhttp://www.ncbi.nlm.nih.gov/pubmed/15003161\\$\\delimiter"026E30F\\$nhhttp://cid.oxfordjournals.org/lookup/doi/10.1093/cid/cir991\\$\\delimiter"026E30F\\$nhhttp://www.scielo.cl/pdf16,45](http://www.americanbanker.com/issues/179{ }124/which-city-is-the-next-big-fintech-hub-new-york-stakes-its-claim-1068345-1.html$\\delimiter)

Clark, J. S., Gelfand, A. E., Woodall, C. W., Zhu, K., jul 2014. More than the sum of the parts: forest climate response from joint species distribution models. *Ecological Applications* 24 (5), 990–999.

URL <http://www.ncbi.nlm.nih.gov/pubmed/25154092http://doi.wiley.com/10.1890/13-1015.1> 75, 94

Connor, E. F., Simberloff, D., dec 1979. The Assembly of Species Communities: Chance or Competition? *Ecology* 60 (6), 1132.

URL <http://www.jstor.org/stable/1936961?origin=crossref> 6, 74, 76, 94

Cook, W. M., Lane, K. T., Foster, B. L., Holt, R. D., sep 2002. Island theory, matrix effects and species richness patterns in habitat fragments. *Ecology Letters* 5 (5), 619–623.

URL <http://doi.wiley.com/10.1046/j.1461-0248.2002.00366.x> 24, 46

Davis, A. J., Jenkinson, L. S., Lawton, J. H., Shorrocks, B., Wood, S., feb 1998. Making-mistakes when predicting shifts in species range in response to global warming. *Nature* 391 (6669), 783–786.

URL [www.blight.comhttp://www.ncbi.nlm.nih.gov/pubmed/280836http://](http://www.blight.comhttp://www.ncbi.nlm.nih.gov/pubmed/280836http://)

[www.ncbi.nlm.nih.gov/pubmed/9486646](http://www.ncbi.nlm.nih.gov/pubmed/9486646)<http://www.nature.com/doi/10.1038/35842> 33, 67, 75

Desmet, P., Cowling, R., 2004. Using the species-area relationship to set baseline targets for conservation. *Ecology And Society* 9 (2), 1–39.

URL <http://www.ecologyandsociety.org/vol9/iss2/art11> 27

Diamond, J. M., 1975. Assembly of species communities. In: Cody, M. L., Diamond, J. M. (Eds.), *Ecology and Evolution of Communities*. Harvard University Press, Cambridge, Massachusetts, USA., pp. 342–444. 6, 74, 76, 94

Diamond, J. M., Gilpin, M. E., jan 1982. Examination of the “null” model of connor and simberloff for species co-occurrences on Islands. *Oecologia* 52 (1), 64–74.

URL <http://link.springer.com/10.1007/BF00349013> 62, 76

Dobzhansky, T., mar 1973. Nothing in Biology Makes Sense except in the Light of Evolution. *The American Biology Teacher* 35 (3), 125–129.

URL <http://abt.ucpress.edu/cgi/doi/10.2307/4444260> 5

Durance, I., Ormerod, S. J., dec 2010. Evidence for the role of climate in the local extinction of a cool-water triclad. *Journal of the North American Benthological Society* 29 (4), 1367–1378.

URL <http://www.bioone.org/doi/abs/10.1899/09-159.1> 67

Eklöf, A., Jacob, U., Kopp, J., Bosch, J., Castro-Urgal, R., Chacoff, N. P., Dalsgaard, B., de Sassi, C., Galetti, M., Guimarães, P. R., Lomáscolo, S. B., Martín González, A. M., Pizo, M. A., Rader, R., Rodrigo, A., Tylianakis, J. M., Vázquez, D. P., Allesina, S., may 2013. The dimensionality of ecological networks. *Ecology letters* 16 (5), 577–583.

URL <http://www.ncbi.nlm.nih.gov/pubmed/23438174> 66

Elith, J., H. Graham, C., P. Anderson, R., Dudík, M., Ferrier, S., Guisan, A., J. Hijmans, R., Huettmann, F., R. Leathwick, J., Lehmann, A., Li, J., G. Lohmann, L., A. Loiselle, B., Manion, G., Moritz, C., Nakamura, M., Nakazawa, Y., McC. M. Overton, J., Townsend Peterson, A., J. Phillips, S., Richardson, K., Scachetti-Pereira, R., E. Schapire, R., Soberón, J., Williams, S., S. Wisz, M., E. Zimmermann, N., apr 2006. Novel methods improve prediction of species’ distributions from occurrence data. *Ecography* 29 (2), 129–151.

URL <http://doi.wiley.com/10.1111/j.2006.0906-7590.04596.x> 7, 17

Elith, J., Leathwick, J. R., dec 2009. Species Distribution Models: Ecological Explanation and Prediction Across Space and Time. *Annual Review of Ecology, Evolution, and Systematics* 40 (1), 677–697.

URL <http://www.annualreviews.org/doi/abs/10.1146/annurev.ecolsys.110308.120159> 17

- Engelbrecht, B. M. J., Comita, L. S., Condit, R., Kursar, T. a., Tyree, M. T., Turner, B. L., Hubbell, S. P., may 2007. Drought sensitivity shapes species distribution patterns in tropical forests. *Nature* 447 (7140), 80–82.  
URL <http://www.ncbi.nlm.nih.gov/pubmed/17476266> 17, 93, 145
- Finstermeier, K., Zinner, D., Brameier, M., Meyer, M., Kreuz, E., Hofreiter, M., Roos, C., 2013. A Mitogenomic Phylogeny of Living Primates. *PLoS ONE* 8 (7), 1–10. 13
- Fordham, D. A., Akçakaya, H. R., Brook, B. W., Rodríguez, A., Alves, P. C., Civantos, E., Triviño, M., Watts, M. J., Araújo, M. B., jul 2013. Adapted conservation measures are required to save the Iberian lynx in a changing climate. *Nature Climate Change* 3 (10), 899–903.  
URL <http://www.nature.com/doifinder/10.1038/nclimate1954> 75
- Gilman, S. E., Urban, M. C., Tewksbury, J., Gilchrist, G. W., Holt, R. D., jun 2010. A framework for community interactions under climate change. *Trends in ecology & evolution* 25 (6), 325–331.  
URL <http://www.ncbi.nlm.nih.gov/pubmed/20392517> 45
- Godsoe, W., Harmon, L. J., sep 2012. How do species interactions affect species distribution models? *Ecography* 35 (9), 811–820.  
URL <http://doi.wiley.com/10.1111/j.1600-0587.2011.07103.x> 33, 46, 62, 93
- Gotelli, N. J., Graves, G. R., 1996. Null models in ecology. Vol. 14. Smithsonian Institution Press, Washington, D.C.  
URL <http://books.google.com/books?id=vjsBAAAAAAAJ&printsec=frontcover> 74, 76
- Gotelli, N. J., Graves, G. R., Rahbek, C., mar 2010. Macroecological signals of species interactions in the Danish avifauna. *Proceedings of the National Academy of Sciences* 107 (11), 5030–5035.  
URL [http://www.pnas.org/content/107/11/5030.abstract\\$\\delimiter\"026E30F\\$\\nfile://c:/DocumentsandSettings/Cristina/Meusdocumentos/MyDropbox/MeuDocumentos/Papers/2010/Gotellieta12010PNAS.pdfhttp://www.pnas.org/cgi/doi/10.1073/pnas.0914089107](http://www.pnas.org/content/107/11/5030.abstract$\\delimiter\) 36, 45, 65, 75, 94, 130
- Grant, P. R., Grant, B. R., 2008. How and Why Species Multiply: The Radiation of Darwin's Finches. Princeton Series in Evolutionary Biology. Princeton University Press.  
URL <https://books.google.ca/books?id=h5urQiWUB7AC> 5
- Gravel, D., Canham, C. D., Beaudet, M., Messier, C., apr 2006. Reconciling niche and neutrality: the continuum hypothesis. *Ecology letters* 9 (4), 399–409.  
URL <http://www.ncbi.nlm.nih.gov/pubmed/16623725> 31



- Gravel, D., Massol, F., Canard, E., Mouillot, D., Mouquet, N., 2011. Trophic theory of island biogeography. *Ecology Letters* 14 (10), 1010–1016. [34](#), [44](#), [46](#), [48](#), [50](#), [52](#), [63](#), [65](#), [77](#), [86](#), [87](#), [131](#), [133](#), [135](#)
- Gravel, D., Poisot, T., Albouy, C., Velez, L., Mouillot, D., 2013. Inferring food web structure from predator-prey body size relationships. *Methods in Ecology and Evolution* 4 (11), 1083–1090.  
URL <http://timotheepoisot.fr/reprints/gravel{ }mee{ }niche.pdf> [32](#), [67](#), [91](#), [132](#), [135](#), [145](#)
- Grinnell, J., oct 1917. The Niche-Relationships of the California Thrasher. *The Auk* 34 (4), 427–433.  
URL <http://www.jstor.org/stable/4072271><http://www.jstor.org/stable/info/10.2307/4072271> [16](#)
- Guisan, A., Rahbek, C., aug 2011. SESAM - a new framework integrating macroecological and species distribution models for predicting spatio-temporal patterns of species assemblages. *Journal of Biogeography* 38 (8), 1433–1444.  
URL <http://doi.wiley.com/10.1111/j.1365-2699.2011.02550.x> [34](#), [143](#)
- Guisan, A., Thuiller, W., sep 2005. Predicting species distribution: offering more than simple habitat models. *Ecology Letters* 8 (9), 993–1009.  
URL <http://doi.wiley.com/10.1111/j.1461-0248.2005.00792.x> [75](#)
- Guisan, A., Tingley, R., Baumgartner, J. B., Naujokaitis-Lewis, I., Sutcliffe, P. R., Tulloch, A. I. T., Regan, T. J., Brotons, L., McDonald-Madden, E., Mantyka-Pringle, C., Martin, T. G., Rhodes, J. R., Maggini, R., Setterfield, S. a., Elith, J., Schwartz, M. W., Wintle, B. a., Broennimann, O., Austin, M., Ferrier, S., Kearney, M. R., Possingham, H. P., Buckley, Y. M., oct 2013. Predicting species distributions for conservation decisions. *Ecology letters* 16, 1424–1435.  
URL <http://www.ncbi.nlm.nih.gov/pubmed/24134332> [45](#)
- Hairston, N. G., Smith, F. E., Slobodkin, L. B., jan 1960. Community Structure, Population Control, and Competition. *The American Naturalist* 94 (879), 421.  
URL <http://www.journals.uchicago.edu/doi/abs/10.1086/282146> [77](#)
- Hannah, L., Roehrdanz, P. R., Ikegami, M., Shepard, A. V., Shaw, M. R., Tabor, G., Zhi, L., Marquet, P. a., Hijmans, R. J., apr 2013. Climate change, wine, and conservation. *Proceedings of the National Academy of Sciences* 110 (17), 6907–6912.  
URL <http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=3637704&tool=pmcentrez&rendertype=abstract><http://www.pnas.org/cgi/doi/10.1073/pnas.1210127110> [8](#)
- Hanski, I., 1998. Metapopulation dynamics. *Nature reviews* 396 (November), 41–49. [30](#)

- Hanski, I., 2010. The Theories of Island Biogeography and Metapopulation Dynamics. In: The Theory of Island Biogeography Revisited. Princeton University Press, Princeton, NJ, p. 476. [29](#)
- Harris, D. J., 2015. Estimating species interactions from observational data with Markov networks. bioRxiv preprint. [75](#)
- He, F., Hubbell, S. P., may 2011. Species-area relationships always overestimate extinction rates from habitat loss. *Nature* 473 (7347), 368–71.  
URL <http://www.ncbi.nlm.nih.gov/pubmed/21593870> [27](#)
- Heikkinen, R. K., Luoto, M., Virkkala, R., Pearson, R. G., Körber, J.-H., nov 2007. Biotic interactions improve prediction of boreal bird distributions at macro-scales. *Global Ecology and Biogeography* 16 (6), 754–763.  
URL <http://doi.wiley.com/10.1111/j.1466-8238.2007.00345.x> [36](#)
- Hijmans, R. J., Cameron, S. E., Parra, J. L., Jones, P. G., Jarvis, A., 2005. Very high resolution interpolated climate surfaces for global land areas. *International Journal of Climatology* 25 (15), 1965–1978. [17](#)
- Holt, R. D., nov 2009. Bringing the Hutchinsonian niche into the 21st century: ecological and evolutionary perspectives. *Proceedings of the National Academy of Sciences of the United States of America* 106 Suppl, 19659–65.  
URL <http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=2780934&tool=pmcentrez&rendertype=abstract> [34](#)
- Holt, R. D., Barfield, M., apr 2009. Trophic interactions and range limits: the diverse roles of predation. *Proceedings. Biological sciences / The Royal Society* 276 (1661), 1435–1442.  
URL <http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=2677217&tool=pmcentrez&rendertype=abstract> [33](#), [45](#), [46](#), [77](#), [89](#), [94](#), [130](#)
- Hortal, J., Diniz-Filho, J. A. F., Bini, L. M., Rodríguez, M. Á., Baselga, A., Nogués-Bravo, D., Rangel, T. F., Hawkins, B. A., Lobo, J. M., aug 2011. Ice age climate, evolutionary constraints and diversity patterns of European dung beetles. *Ecology Letters* 14 (8), 741–748.  
URL <http://doi.wiley.com/10.1111/j.1461-0248.2011.01634.x> [13](#), [14](#)
- Hubbell, S. P., jun 1997. A unified theory of biogeography and relative species abundance and its application to tropical rain forests and coral reefs. *Coral Reefs* 16 (5), S9–S21.  
URL <http://link.springer.com/article/10.1007/s003380050237http://link.springer.com/10.1007/s003380050237> [30](#), [131](#)
- Hubbell, S. P., jan 1999. Light-Gap Disturbances, Recruitment Limitation, and Tree Diversity in a Neotropical Forest. *Science* 283 (5401), 554–557.  
URL <http://www.sciencemag.org/cgi/doi/10.1126/science.283.5401.554> [31](#)

- Hubbell, S. P., 2010. Neutral Theory and the Theory of Island Biogeography. In: Losos, J. B., Ricklefs, R. E. (Eds.), *The Theory of Island Biogeography Revisited*. Princeton University Press, Princeton, NJ, p. 479. 31
- Hutchinson, G. E., 1957. Concluding remarks. *Cold Spring Harbor Symposium. Quantitative Biology* 22, 415–427. 75
- Jabot, F., Bascompte, J., mar 2012. Biotrophic interactions shape biodiversity in space. *Proceedings of the National Academy of Sciences of the United States of America* 109 (12), 4521–4526.  
URL <http://www.ncbi.nlm.nih.gov/pubmed/22393006><http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=3311373>{&}tool=pmcentrez{&}rendertype=abstract 34, 45, 62
- Jeschke, J. M., Strayer, D. L., jan 2008. Usefulness of bioclimatic models for studying climate change and invasive species. *Annals of the New York Academy of Sciences* 1134, 1–24.  
URL <http://www.ncbi.nlm.nih.gov/pubmed/18566088> 18, 33, 45, 99
- John, R., Dalling, J. W., Harms, K. E., Yavitt, J. B., Stallard, R. F., Mirabello, M., Hubbell, S. P., Valencia, R., Navarrete, H., Vallejo, M., Foster, R. B., jan 2007. Soil nutrients influence spatial distributions of tropical tree species. *Proceedings of the National Academy of Sciences* 104 (3), 864–869.  
URL [C:\\\$\\delimiter"026E30F\\$PDFs\\$\\delimiter"026E30F\\$2006{ }287.pdf\\$\\delimiter"026E30F\\$nhhttp://www.pnas.org/lookup/doi/10.1073/pnas.0604666104](C:\$\\delimiter)<http://www.pnas.org/lookup/doi/10.1073/pnas.0604666104> 31
- Kearney, M., Porter, W. P., nov 2004. MAPPING THE FUNDAMENTAL NICHE: PHYSIOLOGY, CLIMATE, AND THE DISTRIBUTION OF A NOCTURNAL LIZARD. *Ecology* 85 (11), 3119–3131.  
URL <http://www.esajournals.org/doi/abs/10.1890/03-0820><http://doi.wiley.com/10.1890/03-0820> 16, 44, 101, 143, 145
- Kéfi, S., Berlow, E. L., Wieters, E. A., Joppa, L. N., Wood, S. A., Brose, U., Navarrete, S. A., jan 2015. Network structure beyond food webs: mapping non-trophic and trophic interactions on Chilean rocky shores. *Ecology* 96 (1), 291–303.  
URL <http://doi.wiley.com/10.1890/13-1424.1> 19
- Kéfi, S., Berlow, E. L., Wieters, E. A., Navarrete, S. A., Petchey, O. L., Wood, S. A., Boit, A., Joppa, L. N., Lafferty, K. D., Williams, R. J., Martinez, N. D., Menge, B. A., Blanchette, C. A., Iles, A. C., Brose, U., apr 2012. More than a meal... integrating non-feeding interactions into food webs. *Ecology Letters* 15 (4), 291–300.  
URL <http://www.ncbi.nlm.nih.gov/pubmed/22313549><http://doi.wiley.com/10.1111/j.1461-0248.2011.01732.x> 19, 46

- Kemeny, J. G., Snell, J. L., 1983. Finite Markov Chains: With a New Appendix "Generalization of a Fundamental Matrix". Undergraduate Texts in Mathematics. Springer New York.  
URL <https://books.google.ca/books?id=0bTK5uWzbYwC> 47
- Kissling, W. D., Dormann, C. F., Groeneveld, J., Hickler, T., Kühn, I., McInerny, G. J., Montoya, J. M., Römermann, C., Schiffers, K., Schurr, F. M., Singer, A., Svenning, J.-C., Zimmermann, N. E., O'Hara, R. B., dec 2012. Towards novel approaches to modelling biotic interactions in multispecies assemblages at large spatial extents. *Journal of Biogeography* 39 (12), 2163–2178.  
URL <http://doi.wiley.com/10.1111/j.1365-2699.2011.02663.x> 34, 44, 75
- Koh, L. P., Dunn, R. R., Sodhi, N. S., Colwell, R. K., Proctor, H. C., Smith, V. S., sep 2004. Species coextinctions and the biodiversity crisis. *Science (New York, N.Y.)* 305 (5690), 1632–4.  
URL <http://www.ncbi.nlm.nih.gov/pubmed/15361627><http://www.sciencemag.org/cgi/doi/10.1126/science.1101101> 7
- Kooijman, S. A. L. M., 2000. Dynamic Energy and Mass Budgets in Biological Systems. Cambridge University Press, Cambridge.  
URL <http://ebooks.cambridge.org/ref/id/CB09780511565403> 38
- Lavergne, S., Mouquet, N., Thuiller, W., Ronce, O., dec 2010. Biodiversity and Climate Change: Integrating Evolutionary and Ecological Responses of Species and Communities. *Annual Review of Ecology, Evolution, and Systematics* 41 (1), 321–350.  
URL <http://www.annualreviews.org/doi/abs/10.1146/annurev-ecolsys-102209-144628> 7, 20, 21, 44, 67, 68, 130, 143
- Legendre, P., Legendre, L. F. J., 2012. Numerical Ecology. Developments in Environmental Modelling. Elsevier Science.  
URL <https://books.google.ca/books?id=6ZBOA-iDviQC> 74, 75
- Leibold, M. a., Holyoak, M., Mouquet, N., Amarasekare, P., Chase, J. M., Hoopes, M. F., Holt, R. D., Shurin, J. B., Law, R., Tilman, D., Loreau, M., Gonzalez, A., jun 2004. The metacommunity concept: a framework for multi-scale community ecology. *Ecology Letters* 7 (7), 601–613.  
URL <http://doi.wiley.com/10.1111/j.1461-0248.2004.00608.x> 30, 44, 46
- Levins, R., sep 1969. Some Demographic and Genetic Consequences of Environmental Heterogeneity for Biological Control. *Bulletin of the Entomological Society of America* 15 (3), 237–240.  
URL [http://izt.ciens.ucv.ve/ecologia/Archivos/ECOLOGIA{ }DE{ }POBLACIONES{ }Hasta2004/ECOL{ }POBLAC{ }Hasta2004{ }\(H-N\)/Levins\(yotrodeRabinovich\).pdf](http://izt.ciens.ucv.ve/ecologia/Archivos/ECOLOGIA{ }DE{ }POBLACIONES{ }Hasta2004/ECOL{ }POBLAC{ }Hasta2004{ }(H-N)/Levins(yotrodeRabinovich).pdf)<http://besa.oxfordjournals.org/cgi/doi/10.1093/besa/15.3.237> 29

- Levins, R., Culver, D., jun 1971. Regional Coexistence of Species and Competition between Rare Species. *Proceedings of the National Academy of Sciences* 68 (6), 1246–1248.  
URL [papers://aa15ed4a-8b41-4036-84a6-41087bba0cd6/Paper/p811http://www.pnas.org/cgi/doi/10.1073/pnas.68.6.1246](http://papers://aa15ed4a-8b41-4036-84a6-41087bba0cd6/Paper/p811http://www.pnas.org/cgi/doi/10.1073/pnas.68.6.1246) 62, 66
- Levins, R., Heatwole, H., 1963. On the distribution of organisms on islands. *Caribbean Journal of Science* 3, 173–177. 23
- Lomolino, M., 2000a. Ecology's most general, yet protean pattern : the species area relationship. *Journal of Biogeography* 27 (1), 17–26.  
URL <http://onlinelibrary.wiley.com/doi/10.1046/j.1365-2699.2000.00377.x/full> 27, 130
- Lomolino, M. V., jan 2000b. A call for a new paradigm of island biogeography. *Global Ecology and Biogeography* 9 (1), 1–6.  
URL <http://onlinelibrary.wiley.com/doi/10.1046/j.1365-2699.2000.00185.x/full><http://doi.wiley.com/10.1046/j.1365-2699.2000.00185.x> 9, 93
- Lomolino, M. V., Brown, J. H., 2009. The reticulating phylogeny of island biogeography theory. *Q. Rev. Biol.* 84 (4), 357–390. 23, 25
- Losos, J. B., Ricklefs, R. E. (Eds.), 2010. *The Theory of Island Biogeography Revisited*. Princeton University Press, Princeton, NJ. 24, 131
- MacArthur, R. H., 1972. *Geographical Ecology: Patterns in the Distribution of Species*. Biology / [Princeton University Press]. Princeton University Press.  
URL <https://books.google.co.in/books?id=3NAYEKc--WAC> 4, 6, 7, 15, 18, 32, 37, 75, 100
- MacArthur, R. H., Wilson, E. O., 1963. An equilibrium theory of insular zoogeography. *Evolution* 17 (4), 373–387.  
URL <http://www.jstor.org/stable/2407089> 26, 28
- MacArthur, R. H., Wilson, E. O., 1967. *Theory of Island Biogeography*. Vol. 1 of *Princeton Landmarks in Biology*. Princeton University Press, Princeton, NJ. 3, 19, 25, 26, 27, 46, 62, 78, 87, 130, 133, 134, 141
- MacArthur, R. H., Wilson, E. O., MacArthur, W., 1967. The theory of island biogeography.  
URL <http://books.google.com/books?hl=en&lr=&id=a10cdkywhVgC&pgis=1> 23
- Massol, F., Gravel, D., Mouquet, N., Cadotte, M. W., Fukami, T., Leibold, M. A., mar 2011. Linking community and ecosystem dynamics through spatial ecology. *Ecology letters* 14 (3), 313–323.  
URL <http://www.ncbi.nlm.nih.gov/pubmed/21272182> 44

- May, R. M., 1973. Stability and complexity in model ecosystems. *Monographs in population biology* 6, 1–235. [19](#)
- May, R. M., feb 2004. Uses and abuses of mathematics in biology. *Science* (New York, N.Y.) 303 (5659), 790–3.  
URL <http://www.ncbi.nlm.nih.gov/pubmed/14764866> [11](#)
- McCann, K. S., may 2000. The diversity-stability debate. *Nature* 405 (6783), 228–33.  
URL <http://www.ncbi.nlm.nih.gov/pubmed/10821283> [19](#)
- McCann, K. S., 2011. *Food Webs. Monographs in Population Biology*. Princeton University Press.  
URL <https://books.google.ca/books?id=kgPeM8E8GVMC> [11](#)
- McGill, B., Collins, C., 2003. A unified theory for macroecology based on spatial patterns of abundance. *Evolutionary Ecology Research* 5 (4), 469–492. [31](#)
- McGill, B. J., apr 2010. Matters of Scale. *Science* 328 (5978), 575–576.  
URL <http://www.ncbi.nlm.nih.gov/pubmed/20431001><http://www.sciencemag.org/cgi/doi/10.1126/science.1188528> [36](#), [65](#), [99](#)
- McGill, B. J., Enquist, B. J., Weiher, E., Westoby, M., apr 2006. Rebuilding community ecology from functional traits. *Trends in ecology & evolution* 21 (4), 178–185.  
URL <http://www.ncbi.nlm.nih.gov/pubmed/16701083> [38](#), [66](#), [143](#), [144](#)
- Milo, R., Shen-Orr, S., Itzkovitz, S., Kashtan, N., Chklovskii, D., Alon, U., oct 2002. Network motifs: simple building blocks of complex networks. *Science* (New York, N.Y.) 298 (5594), 824–7.  
URL <http://www.ncbi.nlm.nih.gov/pubmed/12399590> [82](#)
- Monod, J., 1970. *Le hasard et la nécessité*. Editions Du Seuil.  
URL <https://books.google.fr/books?id=-sVLAAAAMAAJ> [24](#)
- Montoya, J., Woodward, G., Emmerson, M. C., Solé, R. V., sep 2009. Press perturbations and indirect effects in real food webs. *Ecology* 90 (9), 2426–2433.  
URL <http://www.ncbi.nlm.nih.gov/pubmed/19769121><http://doi.wiley.com/10.1890/08-0657.1> [77](#)
- Morales-Castilla, I., Matias, M. G., Gravel, D., Araújo, M. B., 2015. Inferring biotic interactions from proxies. *Trends in Ecology & Evolution* 30 (6), 347–356.  
URL <http://linkinghub.elsevier.com/retrieve/pii/S0169534715000774> [76](#), [91](#), [94](#), [97](#)
- Neigel, J., feb 2003. Species-area relationships and marine conservation. *Ecological Applications* 13 (sp1), 138–145.  
URL [http://www.esajournals.org/doi/abs/10.1890/1051-0761\(2003\)013\[0138:SARAMC\]2.0.CO;2](http://www.esajournals.org/doi/abs/10.1890/1051-0761(2003)013[0138:SARAMC]2.0.CO;2) [27](#)

- Oksanen, L., Fretwell, S. D., Arruda, J., Niemela, P., aug 1981. Exploitation Ecosystems in Gradients of Primary Productivity. *The American Naturalist* 118 (2), 240–261.  
URL <http://www.journals.uchicago.edu/doi/10.1086/283817> 77
- Ovaskainen, O., Hottola, J., Shtonen, J., 2010. Modeling species co-occurrence by multivariate logistic regression generates new hypotheses on fungal interactions. *Ecology* 91 (9), 2514–2521. 92, 94
- Pearson, R. G., Dawson, T. P., sep 2003. Predicting the impacts of climate change on the distribution of species: are bioclimate envelope models useful? *Global Ecology and Biogeography* 12 (5), 361–371.  
URL <http://doi.wiley.com/10.1046/j.1466-822X.2003.00042.x> 33, 65
- Pelletier, F., Clutton-Brock, T., Pemberton, J., Tuljapurkar, S., Coulson, T., mar 2007. The Evolutionary Demography of Ecological Change: Linking Trait Variation and Population Growth. *Science* 315 (5818), 1571–1574.  
URL <http://www.sciencemag.org/cgi/doi/10.1126/science.1139024> 5
- Pelletier, F., Garant, D., Hendry, A., 2009a. Eco-evolutionary dynamics. *Philosophical Transactions of the Royal Society B: Biological Sciences* 364 (1523), 1483–1489.  
URL <http://rstb.royalsocietypublishing.org/cgi/doi/10.1098/rstb.2009.0027> 5
- Pelletier, F., Garant, D., Hendry, a. P., jun 2009b. Eco-evolutionary dynamics. *Philosophical transactions of the Royal Society of London. Series B, Biological sciences* 364 (1523), 1483–9.  
URL <http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=2690510&tool=pmcentrez&rendertype=abstract> 21
- Pellissier, L., Rohr, R. P., Ndiribe, C., Pradervand, J.-N., Salamin, N., Guisan, A., Wisz, M., nov 2013. Combining food web and species distribution models for improved community projections. *Ecology and evolution* 3 (13), 4572–4583.  
URL <http://doi.wiley.com/10.1002/ece3.843><http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=3856755&tool=pmcentrez&rendertype=abstract> 63, 65, 75
- Peterson, A. T., dec 2003. Predicting the Geography of Species' Invasions via Ecological Niche Modeling. *The Quarterly Review of Biology* 78 (4), 419–433.  
URL <http://www.ncbi.nlm.nih.gov/pubmed/14737826><http://www.jstor.org/stable/10.1086/378926> 45
- Peterson, A. T., Soberon, J., Pearson, R. G., Martinez-Meyer, E., 2011. *Ecological Niches and Geographic Distributions*. Vol. 2011. Princeton University Press, Princeton, NJ.  
URL <http://books.google.com/books?id=Q{ }h9FlvgM6wC{ }pgis=1> 45, 75



- Pielou, D. P., Pielou, E. C., nov 1968. Association among species of infrequent occurrence: the insect and spider fauna of *Polyporus betulinus* (Bulliard) Fries. *Journal of theoretical biology* 21 (2), 202–16.  
URL <http://www.ncbi.nlm.nih.gov/pubmed/5700435> 76
- Poisot, T., Canard, E., Mouillot, D., Mouquet, N., Gravel, D., Jordan, F., dec 2012. The dissimilarity of species interaction networks. *Ecology letters* 15 (12), 1353–61.  
URL <http://www.ncbi.nlm.nih.gov/pubmed/22994257> 93
- Poisot, T., Kéfi, S., Morand, S., Stanko, M., Marquet, P. A., Hochberg, M. E., 2015a. A continuum of specialists and generalists in empirical communities. *PLoS ONE* 10 (5), 1–12. 38
- Poisot, T., Stouffer, D. B., Gravel, D., 2015b. Beyond species: why ecological interactions vary through space and time. *Oikos* 124, 243–251.  
URL <http://biorxiv.org/lookup/doi/10.1101/001677> 38, 39
- Pollock, L. J., Tingley, R., Morris, W. K., Golding, N., O'Hara, R. B., Parris, K. M., Vesk, P. A., McCarthy, M. A., may 2014. Understanding co-occurrence by modelling species simultaneously with a Joint Species Distribution Model (JSDM). *Methods in Ecology and Evolution* 5 (5), 397–406.  
URL <http://doi.wiley.com/10.1111/2041-210X.12180> 34, 63, 65, 75, 94, 99
- Post, D. M., Palkovacs, E. P., jun 2009. Eco-evolutionary feedbacks in community and ecosystem ecology: interactions between the ecological theatre and the evolutionary play. *Philosophical transactions of the Royal Society of London. Series B, Biological sciences* 364 (1523), 1629–40.  
URL <http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=2690506&tool=pmcentrez&rendertype=abstract> 21
- Proulx, S. R., Promislow, D. E. L., Phillips, P. C., jun 2005. Network thinking in ecology and evolution. *Trends in ecology & evolution* 20 (6), 345–353.  
URL <http://www.ncbi.nlm.nih.gov/pubmed/16701391> 45
- R Core Team, 2015. R: A Language and Environment for Statistical Computing.  
URL <https://www.r-project.org/> 17, 111, 112
- Razafindratsima, O. H., Mehtani, S., Dunham, A. E., 2013. Extinctions, traits and phylogenetic community structure: Insights from primate assemblages in Madagascar. *Ecography* 36 (1), 047–056. 13
- Ricklefs, R. E., jan 1987. Community Diversity: Relative Roles of Local and Regional Processes. *Science* 235 (4785), 167–171.  
URL <http://www.sciencemag.org/cgi/doi/10.1126/science.235.4785.167> 25, 44



- Ricklefs, R. E., 2003. A comment on Hubbell's zero-sum ecological drift model. *Oikos* 100(1), 185–192. [31](#)
- Robinet, C., Suppo, C., Darrouzet, E., jul 2016. Rapid spread of the invasive yellow-legged hornet in France: the role of human-mediated dispersal and the effects of control measures. *Journal of Applied Ecology*.  
URL <http://doi.wiley.com/10.1111/1365-2664.12724> [22](#)
- Rohr, R. P., Scherer, H., Kehrli, P., Mazza, C., Bersier, L.-F., aug 2010. Modeling food webs: exploring unexplained structure using latent traits. *The American naturalist* 176 (2), 170–7.  
URL <http://www.ncbi.nlm.nih.gov/pubmed/20540609> [92](#)
- Rosindell, J., Hubbell, S. P., He, F., Harmon, L. J., Etienne, R. S., 2012. The case for ecological neutral theory. *Trends in Ecology and Evolution* 27 (4), 203–208.  
URL <http://dx.doi.org/10.1016/j.tree.2012.01.004> [31](#)
- Saavedra, S., Stouffer, D. B., Uzzi, B., Bascompte, J., oct 2011. Strong contributors to network persistence are the most vulnerable to extinction. *Nature* 478 (7368), 233–235.  
URL <http://www.ncbi.nlm.nih.gov/pubmed/21918515> [67](#)
- Saccheri, I., Kuussaari, M., Kankare, M., Vikman, P., Fortelius, W., Hanski, I., 1998. Inbreeding and extinction in a butterfly metapopulation. *Nature* 392 (6675), 491–494. [30](#)
- Säterberg, T., Sellman, S., Ebenman, B., jul 2013. High frequency of functional extinctions in ecological networks. *Nature* 499 (7459), 468–70.  
URL <http://www.ncbi.nlm.nih.gov/pubmed/23831648> [67](#), [139](#)
- Schneider, F. D., Scheu, S., Brose, U., may 2012. Body mass constraints on feeding rates determine the consequences of predator loss. *Ecology letters* 15 (5), 436–443.  
URL <http://www.ncbi.nlm.nih.gov/pubmed/22380767><http://doi.wiley.com/10.1111/ele.12333> [66](#), [132](#)
- Schoener, T. W., 2010. The MacArthur-Wilson Equilibrium Model. In: Losos, J. B., Ricklefs, R. E. (Eds.), *The Theory of Island Biogeography Revisited*. Princeton University Press, Princeton, NJ, Ch. 2, pp. 52–87. [48](#)
- Schoener, T. W., 2011a. The Newest Synthesis : Understanding Ecological Dynamics. *Science* 331 (January), 426–429. [5](#)
- Schoener, T. W., jan 2011b. The newest synthesis: understanding the interplay of evolutionary and ecological dynamics. *Science (New York, N.Y.)* 331 (6016), 426–9.  
URL <http://www.ncbi.nlm.nih.gov/pubmed/21273479> [21](#)
- Sexton, J. P., McIntyre, P. J., Angert, A. L., Rice, K. J., dec 2009. Evolution and Ecology of Species Range Limits. *Annual Review of Ecology, Evolution, and Systematics* 40 (1), 415–436.

- URL <http://www.annualreviews.org/doi/abs/10.1146/annurev.ecolsys.110308.120317> 21
- Simberloff, D. S., nov 1974. Equilibrium Theory of Island Biogeography and Ecology. *Annual Review of Ecology and Systematics* 5 (1), 161–182.  
URL <http://www.annualreviews.org/doi/abs/10.1146/annurev.es.05.110174.001113> 29
- Simberloff, D. S., Wilson, E. O., mar 1969. Experimental Zoogeography of Islands: The Colonization of Empty Islands. *Ecology* 50 (2), 278–296.  
URL <http://doi.wiley.com/10.2307/1934856> 4, 15, 26, 47
- Springer, A., Swann, D., Crimmins, M., 2015. Climate change impacts on high elevation saguaro range expansion. *Journal of Arid Environments* 116, 57–62. 16
- Stephen P. Hubbell, 2001. The unified neutral theory of Biodiversity and Biogeography. Princeton University Press, Princeton, NJ. 80
- Stone, L., Roberts, A., nov 1990. The checkerboard score and species distributions. *Oecologia* 85 (1), 74–79.  
URL <http://link.springer.com/10.1007/BF00317345> 74, 76
- Stouffer, D. B., Bascompte, J., feb 2010. Understanding food-web persistence from local to global scales. *Ecology Letters* 13 (2), 154–161.  
URL <http://www.ncbi.nlm.nih.gov/pubmed/19968697> 82
- Stouffer, D. B., Camacho, J., Jiang, W., Amaral, L. A. N., aug 2007. Evidence for the existence of a robust pattern of prey selection in food webs. *Proceedings. Biological sciences / The Royal Society* 274 (1621), 1931–1940.  
URL <http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=2275185&tool=pmcentrez&rendertype=abstract> 65, 82
- Suttle, K. B., Thomsen, M. A., Power, M. E., feb 2007. Species interactions reverse grassland responses to changing climate. *Science* 315 (5812), 640–2.  
URL <http://www.ncbi.nlm.nih.gov/pubmed/17272720> 67
- Tao, T., Vu, V., Krishnapur, M., sep 2010. Random matrices: Universality of ESDs and the circular law. *The Annals of Probability* 38 (5), 2023–2065.  
URL <http://projecteuclid.org/euclid.aop/1282053780> 19
- Thébault, E., Fontaine, C., aug 2010. Stability of ecological communities and the architecture of mutualistic and trophic networks. *Science* 329 (5993), 853–856.  
URL <http://www.ncbi.nlm.nih.gov/pubmed/20705861> 63

- Thuiller, W., Münkemüller, T., Lavergne, S., Mouillot, D., Mouquet, N., Schiffrers, K., Gravel, D., may 2013. A road map for integrating eco-evolutionary processes into biodiversity models. *Ecology Letters* 16, 94–105.  
URL <http://doi.wiley.com/10.1111/ele.12104> 9, 44, 66, 93, 99, 130, 142, 143
- Tingley, M. W., Monahan, W. B., Beissinger, S. R., Moritz, C., nov 2009. Birds track their Grinnellian niche through a century of climate change. *Proceedings of the National Academy of Sciences of the United States of America* 106 Suppl, 19637–43.  
URL <http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=2780944&tool=pmcentrez&rendertype=abstract> 17
- Ulrich, W., Gotelli, N. J., 2013. Pattern detection in null model analysis. *Oikos* 122 (1), 2–18.  
76
- Urban, M. C., Leibold, M. A., Amarasekare, P., De Meester, L., Gomulkiewicz, R., Hochberg, M. E., Klausmeier, C. A., Loeuille, N., de Mazancourt, C., Norberg, J., Pantel, J. H., Strauss, S. Y., Vellend, M., Wade, M. J., jun 2008. The evolutionary ecology of metacommunities. *Trends in ecology & evolution* 23 (6), 311–317.  
URL <http://www.ncbi.nlm.nih.gov/pubmed/18439707> 44
- Vanbergen, A. J., 2013. Threats to an ecosystem service: Pressures on pollinators. *Frontiers in Ecology and the Environment* 11 (5), 251–259. 8, 22
- Veech, J. A., feb 2013. A probabilistic model for analysing species co-occurrence. *Global Ecology and Biogeography* 22 (2), 252–260.  
URL <http://doi.wiley.com/10.1111/j.1466-8238.2012.00789.x> 64, 76, 90, 96, 109
- Villemant, C., Barbet-Massin, M., Perrard, A., Muller, F., Gargominy, O., Jiguet, F., Rome, Q., 2011. Predicting the invasion risk by the alien bee-hawking Yellow-legged hornet *Vespa velutina nigrithorax* across Europe and other continents with niche models. *Biological Conservation* 144 (9), 2142–2150.  
URL <http://dx.doi.org/10.1016/j.biocon.2011.04.009> 22
- Villemant, C., Haxaire, J., Streito, J., 2006. Premier bilan de l'invasion de *Vespa velutina* Lepeletier en France (Hymenoptera, Vespidae). *Bulletin de la Société entomologique de France* 111 (4), 535–538. 21, 22
- Wacey, D., Kilburn, M. R., Saunders, M., Cliff, J., Brasier, M. D., 2011. Microfossils of sulphur-metabolizing cells in 3.4-billion-year-old rocks of Western Australia. *Nature Geoscience* 4 (10), 698–702.  
URL <http://dx.doi.org/10.1038/ngeo1238> 3
- Waldrop, M. M., feb 2016. The hundred-year quest for gravitational waves — in pictures. *Nature*.  
URL <http://www.nature.com/doifinder/10.1038/nature.2016.19340> 12

- Wallace, A. R., 1858. On the Tendency of Varieties to depart indefinitely from the Original Type. *Proceedings of the Linnean Society Of London* 3 (July), 53–62. [2](#)
- Wallace, A. R., feb 1860. On the Zoological Geography of the Malay Archipelago. *Journal of the Proceedings of the Linnean Society of London. Zoology* 4 (16), 172–184.  
URL <http://www.biodiversitylibrary.org/item/35034><http://doi.wiley.com/10.1111/j.1096-3642.1860.tb00090.x> [2](#)
- Wallace, A. R., 1881. *Island Life: Or, The Phenomena and Causes of Insular Faunas and Floras, Including a Revision and Attempted Solution of the Problem of Geological Climates.* Harper & brothers.  
URL <https://books.google.ca/books?id=ghcPAAAYAAJ> [1](#), [2](#)
- Warren, B. H., Simberloff, D., Ricklefs, R. E., Aguilée, R., Condamine, F. L., Gravel, D., Morlon, H., Mouquet, N., Rosindell, J., Casquet, J., Conti, E., Cornuault, J., Fernández-Palacios, J. M., Hengl, T., Norder, S. J., Rijdsdijk, K. F., Sanmartín, I., Strasberg, D., Triantis, K. a., Valente, L. M., Whittaker, R. J., Gillespie, R. G., Emerson, B. C., Thébaud, C., feb 2015. Islands as model systems in ecology and evolution: prospects fifty years after MacArthur-Wilson. *Ecology Letters* 18 (2), 200–217.  
URL <http://doi.wiley.com/10.1111/ele.12398> [4](#), [24](#), [46](#), [130](#), [131](#)
- Wennekes, P. L., Rosindell, J., Etienne, R. S., sep 2012. The Neutral—Niche Debate: A Philosophical Perspective. *Acta Biotheoretica* 60 (3), 257–271.  
URL <http://link.springer.com/10.1007/s10441-012-9144-6> [31](#)
- Wiens, J. J., aug 2011. The niche, biogeography and species interactions. *Philosophical transactions of the Royal Society of London. Series B, Biological sciences* 366 (1576), 2336–2350.  
URL <http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=3130432&tool=pmcentrez&rendertype=abstract> [62](#)
- Williams, R. J., Martinez, N. D., mar 2000. Simple rules yield complex food webs. *Nature* 404 (6774), 180–183.  
URL <http://www.ncbi.nlm.nih.gov/pubmed/10724169> [57](#), [61](#), [87](#), [89](#), [134](#), [145](#)
- Wootton, J. T., jan 1994. Predicting Direct and Indirect Effects: An Integrated Approach Using Experiments and Path Analysis. *Ecology* 75 (1), 151.  
URL <http://www.jstor.org/stable/1939391?origin=crossref> [46](#), [63](#), [77](#)
- Yodzis, P., apr 1988. The Indeterminacy of Ecological Interactions as Perceived Through Perturbation Experiments. *Ecology* 69 (2), 508–515.  
URL <http://www.jstor.org/stable/1940449?origin=crossref> [77](#)

Yoshida, T., Jones, L. E., Ellner, S. P., Fussmann, G. F., Hairston, N. G., jul 2003. Rapid evolution drives ecological dynamics in a predator-prey system. *Nature* 424 (6946), 303–6.

URL <http://www.ncbi.nlm.nih.gov/pubmed/12867979> 21